

**Original citation:**

Kleinman, Monica E., Perkins, Gavin D., Bhanji, Farhan, Billi, John E., Bray, Janet E., Callaway, Clifton W., de Caen, Allan, Finn, Judith C., Hazinski, Mary Fran, Lim, Swee Han, Maconochie, Ian, Nadkarni, Vinay, Neumar, Robert W., Nikolaou, Nikolaos, Nolan, Jerry P., Reis, Amelia, Sierra, Alfredo F., Singletary, Eunice M., Soar, Jasmeet, Stanton, David, Travers, Andrew, Welsford, Michelle and Zideman, David (2018) ILCOR Scientific knowledge gaps and clinical research priorities for cardiopulmonary resuscitation and emergency cardiovascular care : a consensus statement. *Circulation* . CIR.0000000000000561.  
doi:10.1161/CIR.0000000000000561

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/101654>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**A note on versions:**

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)

**Knowledge Gaps in Resuscitation Science Identified During and After the 2015  
International Liaison Committee on Resuscitation Evidence Evaluation Process:  
A Consensus Statement**

Monica E. Kleinman, MD; Gavin D. Perkins, MBChB, MD, FRCP, FERC, FFICM; Farhan Bhanji, MD, M.Sc.(Ed), FRCPC; John E. Billi, MD; Janet Bray, RN, PhD; Clifton W. Callaway, MD, PhD; Allan de Caen, MD; Judith Finn, PhD, RN; Mary Fran Hazinski, RN, MSN; Swee Han Lim, MBBS, FRCSEd (A&E), FRCP Edin, FAMS; Ian Maconochie, FRCPCH, FRCEM, FRCPI, PhD; Peter Morley, MBBS, FRACP, FANZCA, FCICM, AFRACMA, FERC; Vinay Nadkarni, MD, FAHA; Robert Neumar, MD, PhD, FACEP, FAHA; Nikolaos Nikolaou; Jerry P. Nolan, FRCA, FCEM; Amelia Reis, MD, PhD; Alfredo F. Sierra, U MD MAdm; Eunice M. Singletary, MD; Jasmeet Soar, MA, MB, BChir, FRCA; David Stanton, CCA; Andrew Travers, MD, MSc, FRCPC; Michelle Welsford, BSc, MD, FACEP, CCPE, FRCPC; David Zideman, LVO, QHP(C), BSc, MBBS, FRCA, FRCP, FIMC, FERC.

## **Abstract**

Despite significant advances in the field of resuscitation science, important knowledge gaps persist. Current guidelines for resuscitation are based on the International Liaison Committee on Resuscitation (ILCOR) 2015 Consensus on Science, which includes treatment recommendations supported by the available evidence. The writing group developed this consensus statement with the goal of focusing future research by addressing the knowledge gaps identified during and after the 2015 ILCOR evidence evaluation process. Key publications since the 2015 Consensus on Science are referenced, along with known ongoing clinical trials that are likely to impact future guidelines.

# 1 [h1]Introduction

2

3 Since 1992, when the International Liaison Committee on Resuscitation (ILCOR) was created,  
4 resuscitation experts from around the world have regularly convened to evaluate the existing  
5 evidence, achieve consensus on science, and, when appropriate, provide treatment  
6 recommendations for cardiopulmonary resuscitation (CPR) and emergency cardiovascular care.  
7 Between 2012 and 2015, the 7 ILCOR Task Forces identified 165 resuscitation-related questions  
8 by using the PICO (population, intervention, comparator, outcome) format.<sup>1</sup> Topic experts  
9 performed systematic reviews based on a formal search of the published literature,<sup>2</sup> and the  
10 strength of recommendation and quality of evidence were reported by using the Grading of  
11 Recommendations, Assessment, Development, and Evaluations (GRADE) methodology and the  
12 GRADE Guideline Development Tool.<sup>3</sup> A standardized evidence-to-recommendation framework  
13 was used to guide the Task Forces when writing treatment recommendations.<sup>4</sup>

14

15 *The 2015 International Consensus on CPR and Emergency Cardiovascular Care Science With*  
16 *Treatment Recommendations (CoSTR)* publication was a focused update of key topics and, as  
17 with prior evidence review cycles, was notable for the limited number of treatment  
18 recommendations supported by high-quality science.<sup>5,6</sup> To promote transparency, each  
19 systematic review was accompanied by a list of knowledge gaps identified by the Task Forces as  
20 they wrote treatment recommendations that were frequently qualified as “weak” on the basis of  
21 “low-quality evidence.” When there was insufficient or conflicting evidence that prohibited  
22 formulation of a treatment recommendation, the following language was used: “The confidence  
23 in effect estimates is so low that the panel feels a recommendation to change current practice is

1 too speculative.” The 7 ILCOR member resuscitation councils used the resultant CoSTR to  
2 develop council-specific, evidence-based resuscitation guidelines and educational materials.

3  
4 The objective of this Consensus Statement is to provide the resuscitation science community  
5 with a focused account of knowledge gaps identified during and after the 2015 ILCOR evidence  
6 evaluation process. Each major content area (Basic Life Support [BLS]; Advanced Life Support  
7 [ALS]; Pediatric Life Support; Education, Implementation, and Teams; and First Aid) includes a  
8 synopsis of recent data and a summary of current evidence to characterize the existing state of  
9 the field. The writing group members recognize that this statement does not exhaustively review  
10 the scientific literature since publication of the October 2015 CoSTR, nor does it include every  
11 knowledge gap listed, because prioritization by the Task Force members led to the exclusion of  
12 some gaps in favor of others. Efforts were made to reference sentinel papers and high-quality  
13 investigations in progress (eg, randomized controlled trials [RCTs]).

14  
15 The statement’s writing group members were selected from the ILCOR Task Forces involved in  
16 the 2015 evidence evaluation and consensus on science processes. We sought to achieve  
17 representation from each of the 7 member councils and included experts from the ILCOR  
18 steering committee as well as individuals involved in managing conflict of interest. The writing  
19 group was asked to query the remaining Task Force members, and, with the use of a survey  
20 process, specific gaps were rated in terms of *impact* and *priority*, reflecting the potential to save  
21 lives or years of life, and *feasibility*, indicating the potential to design a study to answer the  
22 specific question. The figure represents the selected knowledge gaps as they correspond to the  
23 links in the Chain of Survival, reflecting a wide range of opportunities to answer critical

1 questions. It is our hope that directing research efforts toward areas of high priority and impact  
2 will drive future progress in resuscitation outcomes.

3

#### 4 **[h1]Scope of the Problem**

5

6 Data from the World Health Organization show that ischemic heart disease remains the leading  
7 single cause of mortality worldwide, accounting for approximately 13% of all deaths. In 2008,  
8 this equated to an estimated 17.3 million lives lost.<sup>7</sup> In Europe alone, there are an estimated 4.1  
9 million deaths from cardiovascular disease annually.<sup>8</sup>

10

11 Global estimates of the incidence of out-of-hospital cardiac arrest (OHCA) show marked  
12 variation between countries, making the total number of cardiac arrests worldwide difficult to  
13 quantify.<sup>9</sup> According to the Resuscitation Outcomes Consortium (ROC) Epistry, approximately  
14 347 000 adults in the United States experienced emergency medical services (EMS)–assessed  
15 OHCA in 2015; of the approximately 52% who were treated by EMS, the survival rate to  
16 hospital discharge was 11.4%.<sup>10,11</sup> In the Cardiac Arrest Reporting to Enhance Survival  
17 (CARES) Registry, among EMS-treated adults with nontraumatic OHCA, the survival rate to  
18 hospital discharge was 10.6% and the survival with favorable neurologic outcome was 8.6%.<sup>10</sup>  
19 This equates to an estimated US annual loss of 311 000 lives and 3.3 million life-years.<sup>12</sup> In  
20 England, 2014 data indicated there were 28 729 EMS-treated cardiac arrests (53 per 100 000  
21 population), and the survival to hospital discharge rate was 7.9%.<sup>13</sup> In Japan, a nationwide  
22 OHCA registry is used to track the incidence of cardiac arrest. During calendar years 2005 to

1 2009, there were 547 143 episodes of OHCA recorded, with the annual incidence rate increasing  
2 from 80.7 to 90.4 per 100 000 population during the study period.<sup>14</sup>

3  
4 Adult in-hospital cardiac arrest (IHCA) rates and outcomes vary widely, and estimates are based  
5 on limited data from voluntary registries. In the United States, data from the American Heart  
6 Association (AHA) Get With The Guidelines<sup>®</sup>-Resuscitation program estimated 209 000  
7 hospitalized adults are treated for cardiac arrest annually, with a survival rate of 24.8%.<sup>15</sup> The  
8 United Kingdom National Cardiac Arrest Audit found an incidence of IHCA of 1.5 per 1000  
9 admissions across 144 acute-care hospitals, with an unadjusted survival rate of 18.4%.<sup>16</sup>

10

## 11 **[h1]BLS and Automated External Defibrillators**

12

13 Accurate recognition of cardiac arrest, prompt initiation of high-quality CPR, early defibrillation,  
14 and rapid activation of EMS are key interventions associated with improved outcome from adult  
15 cardiac arrest.<sup>17-25</sup> The 2015 Consensus on Science considered several topics related to the role  
16 of emergency dispatchers to improve recognition of, and bystander response to, OHCA. Given  
17 the global burden of lives lost and years of life lost from cardiac arrest, studies to address the  
18 knowledge gaps associated with cardiac arrest recognition, dispatch-assisted CPR, and public  
19 access defibrillation programs are considered to be of high impact and priority and are quite  
20 feasible.

21

1 ***[h2]Training/Protocols for Dispatcher-Assisted Recognition of Cardiac Arrest and***  
2 ***Interventions to Promote Bystander CPR***

3  
4 Bystander CPR improves survival rates by 2- to 4-fold.<sup>19-22,25</sup> The 2015 Consensus on Science  
5 noted a strong association between telephone-assisted CPR instruction and the provision of  
6 bystander CPR.<sup>26,27</sup> Specifically, unconsciousness and abnormal (agonal) breathing are the key  
7 diagnostic features to enable dispatchers to recognize cardiac arrest. Enhancing dispatcher  
8 recognition of cardiac arrest is likely to be one of the most cost-effective solutions to improving  
9 survival. Dispatcher training and protocol configuration can assist with cardiac arrest  
10 recognition, but further research is required to define the optimal system and processes. Specific  
11 knowledge gaps include the influence of a dispatcher’s background (clinical/nonclinical), the  
12 impact of and solutions to language barriers, and optimal questions/instructional sequence to  
13 provide to callers. Future studies should report the diagnostic performance characteristics (eg,  
14 sensitivity, specificity), time to key events (recognition of cardiac arrest, initiation of bystander  
15 CPR, time of first responder/EMS arrival), and relevant clinical outcomes (survival to hospital  
16 discharge and favorable neurologic outcome).

17  
18 ***[h2]Conventional vs Compression-Only CPR During Dispatch-Assisted CPR***

19  
20 The 2015 Consensus on Science recommends that dispatchers provide direction for chest  
21 compression-only CPR for adults with suspected OHCA.<sup>26,27</sup> These recommendations were  
22 drawn largely from randomized and observational studies of adults with cardiac arrest from a  
23 presumed *cardiac* cause. A recent systematic review supported that use of telephone-assisted  
24 continuous chest compressions by bystanders, compared with interruption of compressions for



1 rescue breathing, improved survival to discharge for nonasphyxial OHCA.<sup>28</sup> The 2015 CoSTR  
2 noted a paucity of evidence on the most effective strategy (conventional or compression-only  
3 CPR by bystanders) when the cause of arrest is believed to be noncardiac. Specific knowledge  
4 gaps include the optimal dispatch-assisted CPR recommendations for OHCA victims  
5 (compression-only or conventional CPR) and, if conventional CPR is to be given, the optimal  
6 instruction sequence for coaching callers. If conventional CPR is advised, the number of chest  
7 compressions to provide before ventilations are introduced also needs to be determined.

8

## 9 *[h2]System Configuration for Public Access Defibrillation and Automated External*

### 10 *Defibrillator Program Characteristics*

11

12 Defibrillation within the first few minutes after cardiac arrest can lead to survival rates of 50% to  
13 70%.<sup>22</sup> Systems enabling the public to have access to defibrillation have been developed and are  
14 associated with improved survival.<sup>23</sup> Training first responders (volunteers, police, fire services)  
15 to use an automated external defibrillator (AED) can also reduce time to defibrillation and  
16 positively impact survival.<sup>24,25</sup> While evidence supports the clinical effectiveness of these  
17 approaches, there is relatively little evidence informing the optimal system configuration or the  
18 most cost-effective approach. Specific knowledge gaps include effect of strategies to improve  
19 public access defibrillation usage and where best to place AEDs for public access. The role of  
20 digital/social media tools and applications to enhance volunteers' and first responders' ability to  
21 deliver timely defibrillation appear promising and warrant further evaluation.<sup>29,30</sup>

22

## 1 **[h2]CPR Quality**

2

3 High-quality chest compressions are associated with better patient outcomes. Evidence from  
4 observational studies supported the 2015 CoSTR recommendations on optimal chest  
5 compression depth and rate, minimizing interruptions in chest compressions, and avoidance of  
6 leaning.<sup>26,27</sup> The various components of chest compressions were noted, however, to be  
7 interrelated. For example, faster chest compression rates are associated with lower chest  
8 compression depth and greater leaning.<sup>31</sup> Further research is needed to identify which of the  
9 individual components of chest compressions should be prioritized when delivering CPR. At  
10 present, CPR feedback and prompt devices, which measure the quality of CPR delivered, are the  
11 most frequently used systems.<sup>32</sup> Some studies have suggested that optimizing CPR to achieve  
12 specific physiologic endpoints such as diastolic blood pressure, end-tidal carbon dioxide  
13 (ETCO<sub>2</sub>), or cerebral oximetry may be beneficial.<sup>33-35</sup> These studies highlight a key knowledge  
14 gap—identifying the best target(s) for optimizing CPR performance.

15

## 16 **[h1]Advanced Life Support**

17

18 The success of ALS interventions is dependent on the early recognition of cardiac arrest and use  
19 of BLS/AED treatments for cardiac arrest. ALS interventions that interfere with provision of  
20 high-quality chest compressions may reduce the likelihood of return of spontaneous circulation  
21 (ROSC). Postresuscitation care is increasingly recognized as a critical factor in outcome for  
22 patients achieving ROSC after cardiac arrest.

23

## 1 *[h2]Tracheal Intubation During CPR*

2

3 The 2015 Consensus on Science recommended use of bag-mask ventilations, a supraglottic  
4 airway, or an advanced airway for initial airway management during CPR and noted that  
5 selection and efficacy of an airway intervention depends on rescuer skills.<sup>36,37</sup> The role of  
6 tracheal intubation during CPR for both IHCA and OHCA remains uncertain with the possibility  
7 of both benefit and harm. A recent retrospective review from the Get With The Guidelines-  
8 Resuscitation registry found that for IHCA, tracheal intubation anytime during the first 15  
9 minutes of resuscitation, compared with no intubation during that minute, was associated with  
10 decreased survival to hospital discharge.<sup>38</sup> In contrast, another large retrospective series showed  
11 an inverse relationship between time to intubation during IHCA and survival to discharge and  
12 favorable neurologic outcome.<sup>39</sup> Specific knowledge gaps include the type and duration of  
13 training required for performing advanced airway management during CPR.

14

15 Because most patients receive more than 1 type of airway intervention during CPR, the study of  
16 the effect of tracheal intubation on outcome is difficult.<sup>40</sup> Two large RCTs are in progress to  
17 compare tracheal intubation with supraglottic airway devices during OHCA and should further  
18 inform our understanding of the role of an advanced airway during CPR.<sup>41,42</sup> The 2 trials use  
19 different devices and end points: AIRWAYS-2 (ISRCTN08256118), being conducted in the  
20 United Kingdom, is using the i-gel supraglottic airway and modified Rankin Scale score at  
21 hospital discharge, while the Pragmatic Airway Resuscitation Trial (PART) (NCT02419573), in  
22 progress in North America, is using the laryngeal tube and 72-hour survival.

23

## 1 *[h2]Vasopressors During Cardiac Arrest*

2

3 For many decades, the injection of epinephrine for cardiac arrest has been considered a standard  
4 of care during ALS. Given the limitations of observational studies and, therefore, uncertainty  
5 about the benefit or harm with regard to long-term survival, the 2015 Consensus on Science  
6 suggested that epinephrine continue to be administered to patients in cardiac arrest by using  
7 existing dosing and timing strategies.<sup>36,37</sup>

8

9 It is important to note that there have been no placebo-controlled prospective trials with adequate  
10 power to assess the effect of epinephrine on long-term outcome after cardiac arrest. An  
11 underpowered RCT of epinephrine vs placebo for OHCA documented a higher rate of ROSC  
12 without improving survival to discharge or neurologic outcome,<sup>43</sup> but in a meta-analysis of large  
13 observational studies, epinephrine has been associated with no change in survival to discharge  
14 and worse long-term neurologic outcomes even though ROSC rates were higher.<sup>44</sup> In a  
15 propensity-matched analysis of the Get With The Guidelines-Resuscitation registry published  
16 since the 2015 Consensus on Science, the administration of epinephrine within the first 2  
17 minutes after the first defibrillation to patients with IHCA and an initially shockable rhythm was  
18 associated with decreased odds of survival (odds ratio, 0.70; 95% confidence interval, 0.59–0.82;  
19  $P<0.001$ ).<sup>45</sup> In contrast, a recent retrospective review of multiple databases suggested that  
20 decreased time to vasopressor administration was associated with favorable neurologic  
21 outcome.<sup>46</sup>

22

1 In addition to the ongoing question of epinephrine’s benefit, specific knowledge gaps include the  
2 dose and timing of epinephrine administration during cardiac arrest. An ongoing randomized  
3 study of epinephrine (adrenaline) vs placebo for OHCA in the United Kingdom (Pre-hospital  
4 Assessment of the Role of Adrenaline: Measuring the Effectiveness of Drug administration In  
5 Cardiac arrest [PARAMEDIC-2]: The Adrenaline Trial, ISRCTN73485024)<sup>47</sup> will further our  
6 understanding of the role of epinephrine in outcome from cardiac arrest. Until high-quality, well-  
7 powered trials are completed comparing epinephrine with placebo, studies of dose response and  
8 timing of epinephrine are of lower priority except as a third arm embedded in an epinephrine-vs-  
9 placebo trial.

10

## 11 *[h2]Physiologic Targets During Post–Cardiac Arrest Care*

12

13 Approximately 60% of adult patients admitted to an intensive care unit (ICU) after either OHCA  
14 or IHCA die before hospital discharge.<sup>48-50</sup> Reducing post–cardiac arrest hospital mortality to  
15 50% would save 1 additional life for every 10 patients admitted to an ICU after resuscitation  
16 from cardiac arrest. Improving the neurologic outcome of those who survive is also a key metric  
17 of success.

18

19 When considering the 2015 Consensus on Science recommendations related to physiologic  
20 targets during post–cardiac arrest care,<sup>36,37</sup> there are multiple knowledge gaps related to what  
21 deviations from normal or baseline physiologic values are harmful or, perhaps, protective in  
22 post–cardiac arrest patients, and what is the timing, magnitude, and duration of these deviations  
23 that impact outcome. Specific parameters that have been most widely studied include mean

1 arterial pressure, partial pressure of oxygen and carbon dioxide in arterial blood (PaO<sub>2</sub> and  
2 PaCO<sub>2</sub>), central venous oxygen saturation (ScvO<sub>2</sub>), temperature, and serum glucose. With the  
3 exception of temperature, values significantly below and above the normal range have been  
4 associated with worse outcomes in retrospective observational studies, but prospective clinical  
5 trials comparing specific target ranges have been underpowered or are yet to be completed. In  
6 contrast, the currently recommended target temperature range of 32°C to 36°C for comatose  
7 adult cardiac arrest survivors is based on prospective randomized clinical trials demonstrating  
8 improved outcomes compared with no temperature control.<sup>51,52</sup> However, a follow-up trial  
9 detected no difference in outcomes with a target temperature of 33°C vs 36°C, although the time  
10 to reach target temperature was not reported.<sup>53</sup> A recent ILCOR advisory statement incorporated  
11 these findings as an update to the systematic review performed for the 2015 CoSTR  
12 recommendations.<sup>54</sup>

13  
14 There is ongoing interest to determine if targeted temperature management (TTM) provides  
15 equivalent outcomes to a strategy of “controlled normothermia” with strict avoidance of fever.  
16 The TTM2 study (NCT02908308), a follow-up to the TTM study comparing 33°C to 36°C  
17 targets,<sup>53</sup> will be an international, multicenter, parallel-group, noncommercial, randomized,  
18 superiority trial in which a target temperature of 33°C will be compared with standard care with  
19 early treatment of fever (37.8°C or greater) for adults who remain unconscious after cardiac  
20 arrest.<sup>55</sup> Likewise, the optimal target temperature for postarrest management is being evaluated  
21 by the Finding the Optimal Cooling temperature After Out-of-Hospital Cardiac Arrest  
22 (FROST) study, comparing target temperatures of 32°C, 33°C, and 34°C after ROSC for adults  
23 with OHCA and initial shockable rhythm.<sup>56</sup>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

It is challenging to design a prospective clinical trial examining a goal-directed strategy that optimizes only 1 physiologic parameter. One study currently under way in Belgium is the Neuroprotective Goal Directed Hemodynamic Optimization in Post-cardiac Arrest Patients (NEUROPROTECT) trial (NCT02541591).<sup>57</sup> NEUROPROTECT is an adult prospective randomized study comparing a hemodynamic optimization strategy that includes mean arterial pressure 85 to 100 mm Hg and ScvO<sub>2</sub> of 65% to 75% to a control group with a mean arterial pressure goal of greater than 65 mm Hg and no ScvO<sub>2</sub> goal.

Many experts believe the most successful approach will be a personalized precision strategy that individually optimizes each patient’s key physiologic parameters based on prearrest status, severity of injury, response to injury, and response to therapeutic interventions. However, which physiologic parameters are the most significant, as well as the time frame within which optimization is beneficial, remain to be determined. Moreover, more sophisticated monitoring modalities may be required, such as comparison between cerebral oxygen delivery and demand in real time at the bedside. The other significant challenge to this line of investigation is delays in physiologic optimization during early (pre-ICU) post–cardiac arrest care when monitoring capabilities are limited. However, if successful, the impact of optimizing immediate post–cardiac arrest physiology could be significant.

***[h2]Emergency Angiography in Comatose Patients After ROSC***

1 There have been numerous observational studies showing the consistent benefit of emergency  
2 angiography and primary percutaneous coronary intervention in comatose patients with ST-  
3 segment elevation myocardial infarction who have achieved ROSC, and a meta-analysis  
4 supporting use for patients with a mixed etiology of OHCA.<sup>58</sup> The most recent systematic review  
5 completed as part of the 2015 Consensus on Science also confirmed the benefit found in post-  
6 ROSC non-ST-segment elevation patients, but the studies were not consistent or explicit in the  
7 criteria for patient selection.<sup>59,60</sup> The evidence for those patients with non-ST-segment elevation  
8 on postarrest electrocardiogram (ECG) is more limited, and the decision to undertake emergent  
9 catheterization was infrequently documented, but it was commonly made at the discretion of the  
10 treating team. A number of factors could have influenced the decision, including age, duration of  
11 CPR, hemodynamic instability, presenting cardiac rhythm, neurologic status upon hospital  
12 arrival, and perceived likelihood of cardiac etiology.

13  
14 Specific knowledge gaps include determination of which subgroups of patients with non-ST-  
15 segment elevation are the most likely to benefit and which patients may not benefit at all. While  
16 there are no published randomized studies on emergency angiography and primary percutaneous  
17 coronary intervention after ROSC, multiple prospective trials are in progress. The Coronary  
18 Angiography After Cardiac Arrest (COACT) study currently in progress in the Netherlands is a  
19 prospective randomized study of immediate vs delayed coronary angiography for post-ROSC  
20 patients with non-ST-segment elevation, with a primary outcome of 90-day survival.<sup>61</sup> The Pilot  
21 Randomized Clinical Trial of Early Coronary Angiography Versus No Early Coronary  
22 Angiography for Post-Cardiac Arrest Patients Without ECG ST Segment Elevation (PEARL)  
23 trial is a US multicenter study that will assess the safety and efficacy of early (less than 120



1 minutes) angiography in adults with suspected cardiac etiology of cardiac arrest and without ST-  
2 segment elevation on ECG after ROSC.<sup>62</sup> In Germany, the Immediate Unselected Coronary  
3 Angiography Versus Delayed Triage in Survivors of Out-of-hospital Cardiac Arrest Without ST-  
4 segment Elevation (TOMAHAWK) study will also address the timing of angiography, with a  
5 primary outcome of 30-day survival.<sup>63</sup> The Scandinavian Direct or Subacute Coronary  
6 Angiography for Out-of-Hospital Cardiac Arrest (DISCO) trial will compare the safety and  
7 efficacy of these 2 approaches.<sup>64</sup> The study called A Randomised tRial of Expedited transfer to a  
8 cardiac arrest center for non-ST elevation ventricular fibrillation out-of-hospital cardiac arrest  
9 (ARREST) will explore the effectiveness of transfer to a cardiac arrest center and immediate PCI  
10 in patients without ST-segment elevation on postarrest ECG.<sup>65</sup> The National Institutes of Health-  
11 funded ACCESS trial will compare early cardiac catheterization and treatment for patients with  
12 ventricular fibrillation/ventricular tachycardia OHCA with cardiac catheterization lab activation  
13 only after ICU admission and consideration by the cardiology and treatment teams.<sup>66</sup> These  
14 studies will help to determine if emergency angiography should be performed in all comatose  
15 patients after ROSC or whether there should be clearer criteria for selection.

16

## 17 *[h2]Neuroprognostication in Comatose Survivors of Cardiac Arrest*

18

19 Determining the prognosis for recovery for patients after cardiac arrest is important for planning  
20 and coordinating postresuscitation care. In some settings, life-sustaining treatments are limited  
21 when the treating team believes favorable functional recovery is not possible<sup>67</sup>; the use of  
22 postarrest hypothermia added another layer of complexity.<sup>68,69</sup> The 2015 Consensus on Science  
23 identified multiple methods of patient assessment for estimating prognosis: clinical examination,

1 neurophysiologic studies, blood biomarkers, and imaging studies.<sup>36,37</sup> There were specific  
2 knowledge gaps regarding prognostic tests as well as global concerns related to the design of  
3 prognosis studies. First, understanding how sedative drugs and neuromuscular blocking drugs  
4 alter prognostic testing requires prospective studies on drug pharmacokinetics in post–cardiac  
5 arrest patients, especially those treated with temperatures below normal.<sup>70</sup> Next, studies are  
6 needed to investigate the reproducibility of clinical signs and inter-rater reliability of tests and  
7 examinations used to predict outcome in comatose postarrest patients.<sup>71</sup> Finally, blood tests have  
8 used a variety of laboratory standards and techniques, making comparisons between studies  
9 difficult. Studies to standardize handling and assays are needed.<sup>72</sup>

10

11 Future studies of prognostic testing should eliminate some of the design limitations of prior  
12 studies. In most prognostication studies, treating teams were not blinded to the results of tests,  
13 and the results were used as a criterion for limitation or suspension of life-sustaining treatment.  
14 This lack of blinding may bias results toward overestimating the accuracy of tests for predicting  
15 poor outcomes or death (“self-fulfilling prophecy”). Many studies of computed tomography (CT)  
16 and magnetic resonance imaging (MRI) included very selected populations who were referred  
17 for imaging. Prospective studies in unselected patient cohorts resuscitated from cardiac arrest are  
18 needed to determine the actual prognostic accuracy of imaging studies. Most studies examined  
19 the performance of a single variable for predicting prognosis, but clinical practice uses multiple  
20 modalities to assess an individual patient.<sup>73</sup> More studies are needed on the accuracy and  
21 predictive value of prognostic tests when used in combination.<sup>74</sup> Similarly, studies to determine  
22 the incremental value of a given test, when other data about the patient are already known, would  
23 help clinicians design the most efficient evaluation rather than performing every test on every

1 patient.<sup>75</sup> Many tests provide complex results, and the criterion for an “abnormal” or “poor”  
2 result varies widely between studies. For example, definitions of reactive and nonreactive  
3 electroencephalogram (EEG)-based predictors are inconsistent among prognostication studies.  
4 Standardized reporting of EEG, MRI, and CT scan data would improve the ability to compare  
5 cohorts.<sup>76,77</sup>

6

## 7 **[h1]Pediatric Life Support**

8

9 The global incidence of pediatric cardiac arrest is unknown; however, the premature death of a  
10 child leads to a greater proportion of life-years lost compared with an adult. The World Health  
11 Organization’s Global Health Observatory data estimated that 5.9 million children under the age  
12 of 5 died in calendar year 2015.<sup>78</sup> While primary heart disease is an uncommon cause of death in  
13 children, cardiac arrest represents the final common pathway for mortality from infection,  
14 trauma, and congenital anomalies.

15

16 Data from the ROC Epistry and the CARES registry estimate that 7000 children (aged 1–18  
17 years) experience EMS-assessed nontraumatic OHCA per year in North America.<sup>10</sup> Of the 78%  
18 who are treated by EMS, the survival to hospital discharge is 13.2% (ROC) to 11.4%  
19 (CARES).<sup>79</sup> This translates to approximately 6000 pediatric lives lost per year, with a high rate  
20 of neurologic morbidity in survivors. Within North America, the outcomes from pediatric cardiac  
21 arrest are variable by region, with survival to hospital discharge ranging from 2.6% to 14.7%.<sup>80</sup>  
22 The incidence of pediatric IHCA has been estimated to be approximately 1 per 1000 hospital  
23 discharges, based on reports from administrative databases.<sup>81</sup> Survival to discharge for pediatric

1 IHCA has been increasing, with recent estimated rates of 43% to 45%.<sup>81,82</sup> According to the  
2 Centers for Disease Control’s 2014 estimate of approximately 3 million overnight  
3 hospitalizations for patients under age 17, pediatric IHCA accounts for another 1300 deaths per  
4 year in the United States.<sup>83</sup>

## 6 ***[h2]Optimal Sequence of Interventions for Pediatric BLS (C-A-B vs A-B-C)***

7  
8 Prompt initiation of high-quality CPR is associated with increased rates of survival in both  
9 pediatric and adult patients.<sup>19,79</sup> The 2015 Consensus on Science<sup>84,85</sup> noted that time to first chest  
10 compression is decreased by the use of C-A-B vs A-B-C, based on manikin and simulation-based  
11 studies.<sup>86-88</sup> Specific knowledge gaps include whether suspected cardiac arrest etiology should  
12 influence the resuscitation sequence, but there are no published human studies comparing the  
13 outcomes of children in cardiac arrest who are treated with C-A-B vs A-B-C. Because of the lack  
14 of high-quality evidence, the CoSTR did not provide a treatment recommendation<sup>84,85</sup>;  
15 accordingly, the current pediatric BLS guidelines differ among ILCOR’s resuscitation councils.  
16 For instance, the European Resuscitation Council recommends the use of the A-B-C approach  
17 for children,<sup>89</sup> while the AHA supports the use of the C-A-B approach for both children and  
18 adults.<sup>90,91</sup> This “natural experiment” provides an opportunity to collect observational data,  
19 although controlling for other resuscitation and postresuscitation factors, as well as differences in  
20 healthcare systems, would prove challenging.

21

## 22 ***[h2]Goal-Directed Physiologic Targets During CPR***

23

1 CPR quality monitoring is recommended during adult and pediatric CPR.<sup>92</sup> Real-time visual  
2 feedback improves compliance with recommended chest compression depth,<sup>93</sup> which is  
3 associated with improved outcome from pediatric cardiac arrest.<sup>94</sup> ETCO<sub>2</sub> and arterial blood  
4 pressure are 2 physiologic parameters that can be used to guide chest compression performance,  
5 with pediatric animal studies showing that systemic perfusion pressure or diastolic blood  
6 pressure perform better than ETCO<sub>2</sub> to discriminate survivors from nonsurvivors.<sup>34,35</sup> In  
7 particular, using hemodynamic-directed CPR improves short-term survival from cardiac arrest in  
8 a pediatric animal model.<sup>35</sup> The 2015 Consensus on Science was unable to recommend targets  
9 for ETCO<sub>2</sub> or diastolic blood pressure because of lack of evidence.<sup>84,85</sup> Specific knowledge gaps  
10 include numeric physiologic parameters for ETCO<sub>2</sub> and diastolic blood pressure associated with  
11 high-quality pediatric CPR, as well as with improved outcomes.

12

## 13 *[h2]Vasopressors During Cardiac Arrest*

14

15 The efficacy of vasopressor administration during adult OHCA has been questioned due to high-  
16 quality randomized trials and meta-analyses.<sup>95,96</sup> There are no human studies comparing  
17 epinephrine or combination of vasopressors with controls for pediatric OHCA, although a small  
18 case series suggested that the use of epinephrine could be harmful in sudden, witnessed arrest.<sup>97</sup>  
19 The 2015 Consensus on Science recommended standard doses of epinephrine for pediatric  
20 cardiac arrest and noted a paucity of supporting evidence.<sup>84,85</sup>

21

22 For pediatric IHCA, the use of high-dose epinephrine has been shown to be harmful compared  
23 with standard-dose epinephrine,<sup>98</sup> but there are no placebo-controlled studies. A recent

1 retrospective registry study of time to epinephrine in pediatric IHCA for patients with  
2 nonshockable rhythms showed that delay to epinephrine administration was associated with  
3 reduced ROSC, survival to discharge, and favorable neurologic outcome.<sup>99</sup> Use of vasopressin as  
4 a rescue therapy for pediatric ICU patients who are refractory to epinephrine has been  
5 reported.<sup>100</sup> Specific knowledge gaps include the question of harm vs benefit of vasopressors for  
6 pediatric cardiac arrest. In addition, for selected circumstances (eg, pulmonary hypertension,  
7 myocarditis), the role of vasopressors vs extracorporeal CPR rescue should be further  
8 investigated.

9

## 10 *[h2]Defibrillation Energy Doses*

11

12 The 2015 Consensus on Science recommended that the energy dose for defibrillation of pediatric  
13 ventricular fibrillation/pulseless ventricular tachycardia range from 2 to 4 J/kg, monophasic or  
14 biphasic, in the absence of strong evidence for any specific or subsequent doses.<sup>84,85</sup> The  
15 available evidence is largely observational and influenced by confounders such as the duration of  
16 the arrhythmia, mixed analysis of primary and secondary ventricular fibrillation, and different  
17 energy waveforms being used for defibrillation.<sup>101,102</sup> As with the sequence of actions for  
18 pediatric BLS, the resuscitation council–specific guidelines differ. The AHA guidelines currently  
19 recommend an initial dose of 2 J/kg, followed by a second shock of 4 J/kg if needed.<sup>90</sup> In  
20 contrast, the European Resuscitation Council guidelines recommend initial and subsequent doses  
21 of 4 J/kg.<sup>89</sup> Specific knowledge gaps include the risk/benefits of specific or titrated energy doses  
22 in both primary and secondary shockable rhythms.

23

## 1 *[h2]TTM After IHCA*

2

3 Two RCTs on the use of TTM after pediatric cardiac arrest have now been published, comparing  
4 therapeutic hypothermia to normothermia in OHCA and IHCA.<sup>94,103</sup> The Therapeutic  
5 Hypothermia After Pediatric Cardiac Arrest (THAPCA) trials each included 2 cohorts: one group  
6 received 48 hours of therapeutic hypothermia at a target temperature of 33°C followed by 72  
7 hours of normothermia, while the other group was managed with normothermia for 5 days at a  
8 target temperature of 36.8°C. The results of both trials showed no statistical difference in the  
9 prespecified primary outcome of survival with good functional status at 1 year. At the time of  
10 publication of the 2015 Consensus on Science, only the OHCA study results were available.<sup>84,85</sup>

11

12 Limitations of the out-of-hospital THAPCA trial included the fact that it was underpowered to  
13 detect clinically important differences in outcome. Time to initiation of therapy was 5.9 hours,  
14 and time to achieve target temperature was not reported. The in-hospital THAPCA study was  
15 terminated for futility before the predefined enrollment. Despite the rigorous trial design,  
16 differences of opinion persist about the implications of the results for clinical practice, indicating  
17 a need for additional studies.<sup>95,96</sup> Specific knowledge gaps include the optimal target temperature  
18 and the duration of TTM after ROSC.

19

## 20 *[h3]Education, Implementation, and Teams*

21

22 The Utstein “Formula for Survival” emphasizes that cardiac arrest outcome is influenced by  
23 more than just our collective scientific understanding.<sup>104</sup> The 3 multiplicands in the formula

1 “Science,” “Education,” and “Implementation” interact together to affect survival rates after  
2 cardiac arrest. It is recognized that considerable variability still exists in the survival of cardiac  
3 arrest victims in both the out-of-hospital and in-hospital settings, and there continues to be a  
4 critical need to improve educational quality and efficiency as well as address systems of care.  
5 Research into optimizing resuscitation training and addressing implementation-related issues will  
6 be essential to improve the outcomes of cardiac arrest victims.

## 8 *[h2]Retraining Intervals for BLS and ALS*

9  
10 It is well documented that CPR skills can decay within weeks to months after traditional single-  
11 encounter courses and well before the usual retraining intervals of 1 to 2 years.<sup>105,106</sup> The  
12 evidence related to optimal retraining intervals for resuscitation education is limited in both  
13 quantity and quality. Educational psychology literature has suggested a pedagogic benefit to the  
14 “spacing effect” on learning: educational encounters spaced over time result in more efficient  
15 learning and improved learning retention.<sup>107</sup> Future research should ensure that the outcome  
16 measures used have established validity for their use because it is difficult to make generalizable  
17 conclusions without that evidence. Specific knowledge gaps include the effectiveness and  
18 efficiency of more frequent refreshers as well as the optimal amount of time to achieve specific  
19 educational objectives. Other key aspects of refresher training include feasibility, learner  
20 preference, self-efficacy,<sup>108</sup> and cost-effectiveness, especially the option of completing refresher  
21 training within the work flow of the clinical environment.

22



## 1 *[h2]Leadership and Team Training*

2

3 The 2015 Consensus on Science recommends team and leadership training as part of ALS  
4 training for healthcare providers despite low-quality evidence because of the potential benefit,  
5 lack of harm associated with the training, and high level of acceptability to learners and  
6 faculty.<sup>105,106</sup> The Education, Implementation, and Teams Task Force viewed these important  
7 considerations as more significant than the potential increased costs associated with team and  
8 leadership training. Specific knowledge gaps include the impact of team and leadership training  
9 on patient outcomes, long-term retention, and application of knowledge and skills in the clinical  
10 and simulated-learning environments. Furthermore, it would be useful to conduct a formal cost  
11 analysis of the additional training required for adequate teamwork and leadership skills and to  
12 assess the optimal methods to teach them.

13

## 14 *[h2]Social Media Strategies*

15

16 The 2015 Consensus on Science considered the potential impact of the use of social  
17 media/digital technologies as a mechanism for mobilizing citizen bystanders as cardiac arrest  
18 first responders.<sup>105,106</sup> A recent RCT conducted in Sweden showed improved bystander CPR  
19 rates by using a mobile phone positioning system for locating and recruiting CPR-trained lay  
20 responders, although survival was not increased, possibly due to lack of statistical power.<sup>29</sup>  
21 Because early bystander CPR has been shown to improve survival in OHCA, further research is  
22 required to better understand the role of social media in bystander recruitment  
23 (“crowdsourcing”).<sup>109</sup> In addition, there is a need for further research into the potential for the

1 audio, video, and motion-sensing capabilities of digital technologies to be exploited for real-time  
2 resuscitation events and education.<sup>110</sup>

3  
4 As identified by the recent AHA Scientific Statement, digital strategies, such as the use of  
5 mobile devices, social media, and crowdsourcing, are important to achieving the 2020 impact  
6 goals; however, “there is a need for rigorous research...to ensure safety and effectiveness.”<sup>30</sup>

## 8 *[h2]Impact of Cardiac Arrest Centers on Outcomes*

9  
10 There are no RCTs specifically testing the relationship between specialized cardiac arrest centers  
11 and patient outcomes. There is a growing body of evidence from observational studies showing a  
12 positive relationship between cardiac arrest case volume/hospital capabilities and survival  
13 outcomes.<sup>111-113</sup> Consistent with other emergent conditions such as trauma<sup>114</sup> and acute  
14 myocardial infarction,<sup>115</sup> the concept of cardiac arrest centers that deliver evidence-based  
15 postresuscitation care is intuitively appealing. However, as identified in the 2010<sup>116</sup> and  
16 2015<sup>105,106</sup> CoSTR, there are multiple specific knowledge gaps: the particular treatments to be  
17 provided by a cardiac arrest center, the safe patient transport interval (time taken to travel from  
18 scene to hospital), the optimal mode of transport (eg, ground ambulance, helicopter), the role of  
19 secondary transport (transfer from a receiving hospital to a cardiac arrest center), and the cost-  
20 effectiveness of cardiac arrest centers. It is uncertain if sufficient clinical equipoise exists for an  
21 RCT to be conducted.

22

1 ***[h2]Implementation of Resuscitation Guidelines***

2

3 The 2015 Consensus on Science recommended the implementation of resuscitation guidelines by  
4 organizations that provide care for cardiac arrest patients despite the absence of any strong  
5 evidence.<sup>105,106</sup> This so-called discordant recommendation was made in recognition of the likely  
6 benefits relative to harm of a consistent “evidence-based” response by rescuers to a life-  
7 threatening event. Despite the international effort that underpins the development of resuscitation  
8 guidelines, it has been shown that the uptake of resuscitation guidelines into practice at the local  
9 level can be delayed for a variety of reasons.<sup>117-120</sup> Specific knowledge gaps include the most  
10 effective (and cost-effective) strategies for dissemination and implementation of resuscitation  
11 guidelines.

12

13 ***[h3]First Aid***

14

15 *First aid* is defined as the helping behaviors and initial care provided for an acute illness or  
16 injury. The goals of first aid are to preserve life, alleviate suffering, prevent further illness or  
17 injury, and promote recovery. First aid can be initiated by anyone in any situation and is the  
18 overarching title for all resuscitation and associated life-preserving procedures.<sup>121,122</sup> First aid can  
19 therefore be considered to be the primary preventive precardiac arrest strategy to avert the need  
20 for resuscitation, as well as the post–cardiac arrest management that prevents further harm or  
21 rearrest. As with other domains, there is a serious lack of evidence-based science to support or  
22 refute the current first aid recommendations.

23

1 ***[h2]Recovery Position***

2

3 The 2015 Consensus on Science suggested the use of a lateral side-lying recovery position for  
4 persons with medical conditions who are unresponsive but breathing normally.<sup>121,122</sup> Specific  
5 knowledge gaps include determination of the optimal recovery position for both adults and  
6 children. Most studies have used discomfort in healthy volunteers as the outcome measure; only  
7 a single study has looked at aspiration in relation to position,<sup>123</sup> and no studies have looked at  
8 mortality. Further complicating the topic are anecdotal reports of resuscitation being delayed  
9 while the victim is placed in the recovery position and mistaking agonal gasps for normal  
10 respiratory effort.<sup>124</sup>

11

12 Specific knowledge gaps include situations where use of a recovery position may be beneficial.  
13 For example, placing a patient with a generalized seizure into a lateral recumbent recovery  
14 position may be associated with a lower incidence of sudden death in epilepsy.<sup>125</sup> Another recent  
15 study showed a lower incidence of hospital admissions among children with loss of  
16 consciousness primarily due to seizures and vasovagal syncope who were placed in a recovery  
17 position.<sup>126</sup> The use of a recovery position for acute drug and alcohol ingestions also deserves  
18 future study.

19

20 ***[h2]Use of Oxygen for Conditions Other Than Chest Pain/Cardiac Arrest***

21

22 The 2015 Consensus on Science focused on the use of oxygen for dyspnea and decompression  
23 injury.<sup>121,122</sup> A single large observational study evaluated oxygen use in suspected decompression

1 sickness,<sup>127</sup> and smaller studies examined oxygen use in patients with advanced cancer and  
2 dyspnea combined with hypoxemia.<sup>128-131</sup> Specific knowledge gaps include the use of oxygen by  
3 first aid providers for conditions such as suspected carbon monoxide poisoning, inhalation  
4 injury, seizures, or most other illnesses and injuries.

5

## 6 ***[h2]Control of Bleeding***

7

8 The ability to control severe, life-threatening bleeding is considered a primary first aid  
9 intervention and one that was the focus of a recent public health initiative in the United States  
10 (“Stop the Bleed” Campaign).<sup>132</sup> The 2015 Consensus on Science suggested the use of  
11 tourniquets for severe external limb bleeding that could not be controlled by direct pressure.<sup>121,122</sup>  
12 Research into the use of tourniquets comes primarily from the military setting in the form of case  
13 series or animal studies.<sup>123-140</sup> Civilian sector studies are limited and consist chiefly of small  
14 observational studies.<sup>141,142</sup>

15

16 There are no direct comparative studies looking at tourniquets vs hemostatic dressings for  
17 extremity wounds with massive bleeding or use of a second tourniquet when initial tourniquet  
18 application fails. There are no studies specifically comparing improvised tourniquets that might  
19 be used by a first aid provider for outcomes of control of bleeding. Research looking at  
20 hemostatic dressings is almost exclusively from animal studies. Further research comparing  
21 different methods of hemorrhage control are needed because it would be unethical to perform  
22 studies by using a control group that did not receive treatment. Specific knowledge gaps include

1 the optimal educational techniques for first aid providers and lay public members in the proper  
2 application of tourniquets, both commercial and improvised.

3

#### 4 ***[h2]Treatment of Mild Hypoglycemia***

5

6 The 2015 Consensus on Science suggested that first aid providers administer glucose tablets or  
7 other forms of dietary sugars for symptomatic patients suspected of being hypoglycemic.<sup>121,122</sup>

8 There are no studies looking at buccal or sublingual absorption of glucose gel or paste to  
9 determine a comparable dose equivalent to oral glucose tablet or liquid glucose administration.

10 Studies that look at buccal or sublingual administration suggest that the dose was swallowed  
11 rather than absorbed through mucosa.<sup>143</sup> Preliminary studies show comparable improvement in

12 moderate to severe hypoglycemia in children with malaria when treated with repeat doses of  
13 sublingual glucose as compared with intravenous glucose infusion.<sup>144</sup> Specific knowledge gaps

14 include the safety and efficacy of glucagon use by first aid responders for hypoglycemia, despite  
15 the existence in many states in the United States of protocols allowing the use of injectable  
16 glucagon by trained first aid providers or family members.

17

#### 18 ***[h2]Cervical Spine Motion Restriction***

19

20 While it is recognized that spinal immobilization has not been shown to prevent neurologic  
21 injury, there continue to be questions about the appropriate first aid response for patients at high

22 risk for spinal injury. Few studies have evaluated the effectiveness of manual stabilization  
23 techniques such as head-squeeze or trap-squeeze.<sup>145,146</sup> Specific knowledge gaps include

1 interventions that may be of benefit when providing first aid to a spine-injured/high-risk spinal  
2 injury patient, such as verbal prompts or manual stabilization, while awaiting arrival of advanced  
3 care providers.

4

## 5 **[h1]Summary**

6

7 The 2015 CoSTR used a rigorous process to evaluate the available evidence. Despite tremendous  
8 progress in the field of resuscitation medicine, multiple knowledge gaps remain. This Consensus  
9 Statement has outlined the major topic areas and attempted to prioritize the gaps in knowledge,  
10 based on potential impact on outcomes and feasibility to perform.

11

12

1

2 **Acknowledgments**

3 The writing group acknowledges the contribution of the ILCOR Task Force members who  
4 participated in the survey process, as well as the dedication and support of the AHA Emergency  
5 Cardiovascular Care staff.

6

7 **Sources of Funding**

8 The authors report no external funding for this manuscript.

9

10 **Disclosures**



## 1 [h1]References

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

1. O'Connor D, Higgins JPT, Green S. Defining the review questions and developing criteria for including studies. In: *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0. The Cochrane Collaboration. 2011. <http://handbook.cochrane.org/>. Accessed April 21, 2017.
2. International Liaison Committee on Resuscitation (ILCOR). ILCOR scientific evidence evaluation and review system. American Heart Association, American Stroke Association SharePoint website. <https://volunteer.heart.org/apps/pico/Pages/default.aspx>. Accessed April 21, 2017.
3. Schünemann H, Brożek J, Guyatt G, Oxman A, eds. *GRADE Handbook*. 2013. <http://www.guidelinedevelopment.org/handbook/>. Accessed April 21, 2017.
4. GRADEpro guideline development tool. GRADEpro GDT website. <http://www.guidelinedevelopment.org/>. Accessed April 21, 2017.
5. Morley PT, Lang E, Aickin R, Billi JE, Eigel B, Ferrer JM, Finn JC, Gent LM, Griffin RE, Hazinski MF, Maconochie IK, Montgomery WH, Morrison LJ, Nadkarni VM, Nikolaou NI, Nolan JP, Perkins GD, Sayre MR, Travers AH, Wyllie J, Zideman DA. Part 2: evidence evaluation and management of conflicts of interest: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2015;132(suppl 1):S40–S50. doi: 10.1161/CIR.0000000000000271.
6. Morley PT, Lang E, Aickin R, Billi JE, Eigel B, Ferrer JM, Finn JC, Gent LM, Griffin RE, Hazinski MF, Maconochie IK, Montgomery WH, Morrison LJ, Nadkarni VM, Nikolaou NI, Nolan JP, Perkins GD, Sayre MR, Travers AH, Wyllie J, Zideman DA. Part 2: evidence evaluation and management of conflicts of interest: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2015;95:e33–e41. doi: 10.1016/j.resuscitation.2015.07.040.

- 1 7. Chronic diseases and health promotion: global status report on NCDs. World Health Organization  
2 website. [http://www.who.int/chp/ncd\\_global\\_status\\_report/en/](http://www.who.int/chp/ncd_global_status_report/en/). Accessed April 24, 2017.
- 3 8. Gräsner JT, Lefering R, Koster RW, Masterson S, Böttiger BW, Herlitz J, Wnent J, Tjelmeland  
4 IB, Ortiz FR, Maurer H, Baubin M, Mols P, Hadzibegovic I, Ioannides M, Skulec R, Wissenberg M, Salo  
5 A, Hubert H, Nikolaou NI, Loczi G, Svavarsdottir H, Semeraro F, Wright PJ, Clarens C, Pijls R, Cebula  
6 G, Correia VG, Cimpoesu D, Raffay V, Trenkler S, Markota A, Strömsöe A, Burkart R, Perkins GD,  
7 Bossaert LL; for the EuReCa ONE Collaborators. EuReCa ONE-27 Nations, ONE Europe, ONE  
8 Registry: a prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in  
9 Europe. *Resuscitation*. 2016;105:188–195. doi: 10.1016/j.resuscitation.2016.06.004.
- 10 9. Berdowski J, Berg RA, Tijssen JG, Koster RW. Global incidences of out-of-hospital cardiac  
11 arrest and survival rates: systematic review of 67 prospective studies. *Resuscitation*. 2010;81:1479–1487.  
12 doi: 10.1016/j.resuscitation.2010.08.006.
- 13 10. Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, de Ferranti SD, Floyd J,  
14 Fornage M, Gillespie C, Isasi CR, Jimenez MC, Jordan LC, Judd SE, Lackland D, Lichtman JH, Lisabeth  
15 L, Liu S, Longenecker CT, Mackey RH, Matsushita K, Mozaffarian D, Mussolino ME, Nasir K, Neumar  
16 RW, Palaniappan L, Pandey DK, Thiagarajan RR, Reeves MJ, Ritchey M, Rodriguez CJ, Roth GA,  
17 Rosamond WD, Sasson C, Towfighi A, Tsao CW, Turner MB, Virani SS, Voeks JH, Willey JZ, Wilkins  
18 JT, Wu JH, Alger HM, Wong SS, Muntner P; for the American Heart Association Statistics Committee  
19 and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2017 update: a report from the  
20 American Heart Association. *Circulation*. 2017;135:e146–e603. doi: 10.1161/CIR.0000000000000485.
- 21 11. Daya MR, Schmicker RH, Zive DM, Rea TD, Nichol G, Buick JE, Brooks S, Christenson J,  
22 MacPhee R, Craig A, Rittenberger JC, Davis DP, May S, Wigginton J, Wang H; for the Resuscitation  
23 Outcomes Consortium Investigators. Out-of-hospital cardiac arrest survival improving over time: results  
24 from the Resuscitation Outcomes Consortium (ROC). *Resuscitation*. 2015;91:108–115. doi:  
25 10.1016/j.resuscitation.2015.02.003.

- 1 12. Stecker EC, Reinier K, Marijon E, Narayanan K, Teodorescu C, Uy-Evanado A, Gunson K, Jui J,  
2 Chugh SS. Public health burden of sudden cardiac death in the United States. *Circ Arrhythm*  
3 *Electrophysiol.* 2014;7:212–217. doi: 10.1161/CIRCEP.113.001034.
- 4 13. Hawkes C, Booth S, Ji C, Brace-McDonnell SJ, Whittington A, Mapstone J, Cooke MW, Deakin  
5 CD, Gale CP, Fothergill R, Nolan JP, Rees N, Soar J, Siriwardena AN, Brown TP, Perkins GD; for the  
6 OHCAO collaborators. Epidemiology and outcomes from out-of-hospital cardiac arrests in England.  
7 *Resuscitation.* 2017;110:133–140. doi: 10.1016/j.resuscitation.2016.10.030.
- 8 14. Kitamura T, Iwami T, Kawamura T, Nitta M, Nagao K, Nonogi H, Yonemoto N, Kimura T; for  
9 the Japanese Circulation Society Resuscitation Science Study Group. Nationwide improvements in  
10 survival from out-of-hospital cardiac arrest in Japan. *Circulation.* 2012;126:2834–2843. doi:  
11 10.1161/CIRCULATIONAHA.112.109496.
- 12 15. Merchant RM, Yang L, Becker LB, Berg RA, Nadkarni V, Nichol G, Carr BG, Mitra N, Bradley  
13 SM, Abella BS, Groeneveld PW; for the American Heart Association National Registry of  
14 Cardiopulmonary Resuscitation Investigators. Incidence of treated cardiac arrest in hospitalized patients  
15 in the United States. *Crit Care Med.* 2011;39:2401–2406. doi: 10.1097/CCM.0b013e3182257459.
- 16 16. Nolan JP, Soar J, Smith GB, Gwinnutt C, Parrott F, Power S, Harrison DA, Nixon E, Rowan K;  
17 for the National Cardiac Arrest Audit. Incidence and outcome of in-hospital cardiac arrest in the United  
18 Kingdom National Cardiac Arrest Audit. *Resuscitation.* 2014;85:987–992. doi:  
19 10.1016/j.resuscitation.2014.04.002.
- 20 17. Nichol G, Cobb LA, Yin L, Maynard C, Olsufka M, Larsen J, McCoy AM, Sayre MR. Briefer  
21 activation time is associated with better outcomes after out-of-hospital cardiac arrest. *Resuscitation.*  
22 2016;107:139–144. doi: 10.1016/j.resuscitation.2016.06.040.
- 23 18. Berdowski J, Beekhuis F, Zwinderman AH, Tijssen JG, Koster RW. Importance of the first link:  
24 description and recognition of an out-of-hospital cardiac arrest in an emergency call. *Circulation.*  
25 2009;119:2096–2102. doi: 10.1161/CIRCULATIONAHA.108.768325.

- 1 19. Sasson C, Rogers MA, Dahl J, Kellermann AL. Predictors of survival from out-of-hospital  
2 cardiac arrest: a systematic review and meta-analysis. *Circ Cardiovasc Qual Outcomes*. 2010;3:63–81.  
3 doi: 10.1161/CIRCOUTCOMES.109.889576.
- 4 20. Rajan S, Wissenberg M, Folke F, Hansen SM, Gerds TA, Kragholm K, Hansen CM, Karlsson L,  
5 Lippert FK, Køber L, Gislason GH, Torp-Pedersen C. Association of bystander cardiopulmonary  
6 resuscitation and survival according to ambulance response times after out-of-hospital cardiac arrest.  
7 *Circulation*. 2016;134:2095–2104. doi: 10.1161/CIRCULATIONAHA.116.024400.
- 8 21. Girotra S, van Diepen S, Nallamothu BK, Carrel M, Vellano K, Anderson ML, McNally B,  
9 Abella BS, Sasson C, Chan PS; and the CARES Surveillance Group and the HeartRescue Project.  
10 Regional variation in out-of-hospital cardiac arrest survival in the United States. *Circulation*.  
11 2016;133:2159–2168. doi: 10.1161/CIRCULATIONAHA.115.018175.
- 12 22. Hansen CM, Kragholm K, Granger CB, Pearson DA, Tyson C, Monk L, Corbett C, Nelson RD,  
13 Dupre ME, Fosbøl EL, Strauss B, Fordyce CB, McNally B, Jollis JG. The role of bystanders, first  
14 responders, and emergency medical service providers in timely defibrillation and related outcomes after  
15 out-of-hospital cardiac arrest: results from a statewide registry. *Resuscitation*. 2015;96:303–309. doi:  
16 10.1016/j.resuscitation.2015.09.002.
- 17 23. Kitamura T, Kiyohara K, Sakai T, Matsuyama T, Hatakeyama T, Shimamoto T, Izawa J, Fujii T,  
18 Nishiyama C, Kawamura T, Iwami T. Public-access defibrillation and out-of-hospital cardiac arrest in  
19 Japan. *N Engl J Med*. 2016;375:1649–1659. doi: 10.1056/NEJMsa1600011.
- 20 24. Zijlstra JA, Stieglis R, Riedijk F, Smeekes M, van der Worp WE, Koster RW. Local lay rescuers  
21 with AEDs, alerted by text messages, contribute to early defibrillation in a Dutch out-of-hospital cardiac  
22 arrest dispatch system. *Resuscitation*. 2014;85:1444–1449. doi: 10.1016/j.resuscitation.2014.07.020.
- 23 25. Malta Hansen C, Kragholm K, Pearson DA, Tyson C, Monk L, Myers B, Nelson D, Dupre ME,  
24 Fosbol EL, Jollis JG, Strauss B, Anderson ML, McNally B, Granger CB. Association of bystander and  
25 first-responder intervention with survival after out-of-hospital cardiac arrest in North Carolina, 2010-  
26 2013. *JAMA*. 2015;314:255–264. doi: 10.1001/jama.2015.7938.

- 1 26. Perkins GD, Travers AH, Berg RA, Castren M, Considine J, Escalante R, Gazmuri RJ, Koster  
2 RW, Lim SH, Nation KJ, Olasveengen TM, Sakamoto T, Sayre MR, Sierra A, Smyth MA, Stanton D,  
3 Vaillancourt C; for the Basic Life Support Chapter Collaborators. Part 3: adult basic life support and  
4 automated external defibrillation: 2015 International Consensus on Cardiopulmonary Resuscitation and  
5 Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2015;95:e43–  
6 e69. doi: 10.1016/j.resuscitation.2015.07.041.
- 7 27. Travers AH, Perkins GD, Berg RA, Castren M, Considine J, Escalante R, Gazmuri RJ, Koster  
8 RW, Lim SH, Nation KJ, Olasveengen TM, Sakamoto T, Sayre MR, Sierra A, Smyth MA, Stanton D,  
9 Vaillancourt C; for the Basic Life Support Chapter Collaborators. Part 3: Adult basic life support and  
10 automated external defibrillation: 2015 International Consensus on Cardiopulmonary Resuscitation and  
11 Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2015;132(suppl  
12 1):S51–S83. doi: 10.1161/CIR.0000000000000272.
- 13 28. Zhan L, Yang LJ, Huang Y, He Q, Liu GJ. Continuous chest compression versus interrupted  
14 chest compression for cardiopulmonary resuscitation of non-asphyxial out-of-hospital cardiac arrest.  
15 *Cochrane Database Syst Rev*. 2017;3:CD010134. doi: 10.1002/14651858.CD010134.pub2.
- 16 29. Ringh M, Rosenqvist M, Hollenberg J, Jonsson M, Fredman D, Nordberg P, Järnbert-Pettersson  
17 H, Hasselqvist-Ax I, Riva G, Svensson L. Mobile-phone dispatch of laypersons for CPR in out-of-  
18 hospital cardiac arrest. *N Engl J Med*. 2015;372:2316–2325. doi: 10.1056/NEJMoa1406038.
- 19 30. Rumsfeld JS, Brooks SC, Aufderheide TP, Leary M, Bradley SM, Nkonde-Price C, Schwamm  
20 LH, Jessup M, Ferrer JM; for the American Heart Association Emergency Cardiovascular Care  
21 Committee; Council on Cardiopulmonary CC, Perioperative and Resuscitation; Council on Quality of  
22 Care and Outcomes Research; Council on Cardiovascular and Stroke Nursing; and Council on  
23 Epidemiology and Prevention. Use of mobile devices, social media, and crowdsourcing as digital  
24 strategies to improve emergency cardiovascular care: a scientific statement from the American Heart  
25 Association. *Circulation*. 2016;134:e87–e108. doi: 10.1161/CIR.0000000000000428.

- 1 31. Field RA, Soar J, Davies RP, Akhtar N, Perkins GD. The impact of chest compression rates on  
2 quality of chest compressions—a manikin study. *Resuscitation*. 2012;83:360–364. doi:  
3 10.1016/j.resuscitation.2011.07.012.
- 4 32. Yeung J, Meeks R, Edelson D, Gao F, Soar J, Perkins GD. The use of CPR feedback/prompt  
5 devices during training and CPR performance: a systematic review. *Resuscitation*. 2009;80:743–751. doi:  
6 10.1016/j.resuscitation.2009.04.012.
- 7 33. Sutton RM, French B, Meaney PA, Topjian AA, Parshuram CS, Edelson DP, Schexnayder S,  
8 Abella BS, Merchant RM, Bembea M, Berg RA, Nadkarni VM; for the American Heart Association's Get  
9 With The Guidelines–Resuscitation Investigators. Physiologic monitoring of CPR quality during adult  
10 cardiac arrest: a propensity-matched cohort study. *Resuscitation*. 2016;106:76–82. doi:  
11 10.1016/j.resuscitation.2016.06.018.
- 12 34. Morgan RW, French B, Kilbaugh TJ, Naim MY, Wolfe H, Bratinov G, Shoap W, Hsieh TC,  
13 Nadkarni VM, Berg RA, Sutton RM. A quantitative comparison of physiologic indicators of  
14 cardiopulmonary resuscitation quality: diastolic blood pressure versus end-tidal carbon dioxide.  
15 *Resuscitation*. 2016;104:6–11. doi: 10.1016/j.resuscitation.2016.04.004.
- 16 35. Morgan RW, Kilbaugh TJ, Shoap W, Bratinov G, Lin Y, Hsieh TC, Nadkarni VM, Berg RA,  
17 Sutton RM; for the Pediatric Cardiac Arrest Survival Outcomes PiCASO Laboratory Investigators. A  
18 hemodynamic-directed approach to pediatric cardiopulmonary resuscitation (HD-CPR) improves  
19 survival. *Resuscitation*. 2017;111:41–47. doi: 10.1016/j.resuscitation.2016.11.018.
- 20 36. Soar J, Callaway CW, Aibiki M, Bottiger BW, Brooks SC, Deakin CD, Donnino MW, Drajer S,  
21 Kloeck W, Morley PT, Morrison LJ, Neumar RW, Nicholson TC, Nolan JP, Okada K, O'Neil BJ, Paiva  
22 EF, Parr MJ, Wang TL, Witt J; for the Advanced Life Support Chapter Collaborators. Part 4: advanced  
23 life support: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency  
24 Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*. 2015;95:e71–e120. doi:  
25 10.1016/j.resuscitation.2015.07.042.

- 1 37. Callaway CW, Soar J, Aibiki M, Bottiger BW, Brooks SC, Deakin CD, Donnino MW, Drajer S,  
2 Kloeck W, Morley PT, Morrison LJ, Neumar RW, Nicholson TC, Nolan JP, Okada K, O'Neil BJ, Paiva  
3 EF, Parr MJ, Wang TL, Witt J; for the Advanced Life Support Chapter Collaborators. Part 4: advanced  
4 life support: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency  
5 Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2015;132(suppl 1):S84–  
6 S145. doi: 10.1161/CIR.0000000000000273.
- 7 38. Andersen LW, Granfeldt A, Callaway CW, Bradley SM, Soar J, Nolan JP, Kurth T, Donnino  
8 MW; for the American Heart Association's Get With The Guidelines–Resuscitation Investigators.  
9 Association between tracheal intubation during adult in-hospital cardiac arrest and survival. *JAMA*.  
10 2017;317:494–506. doi: 10.1001/jama.2016.20165.
- 11 39. Wang CH, Chen WJ, Chang WT, Tsai MS, Yu PH, Wu YW, Huang CH. The association  
12 between timing of tracheal intubation and outcomes of adult in-hospital cardiac arrest: a retrospective  
13 cohort study. *Resuscitation*. 2016;105:59–65. doi: 10.1016/j.resuscitation.2016.05.012.
- 14 40. Voss S, Rhys M, Coates D, Greenwood R, Nolan JP, Thomas M, Bengert J. How do paramedics  
15 manage the airway during out of hospital cardiac arrest? *Resuscitation*. 2014;85:1662–1666. doi:  
16 10.1016/j.resuscitation.2014.09.008.
- 17 41. Taylor J, Black S, Brett J, Kirby K, Nolan JP, Reeves BC, Robinson M, Rogers CA, Scott LJ,  
18 South A, Stokes EA, Thomas M, Voss S, Wordsworth S, Bengert JR. Design and implementation of the  
19 AIRWAYS-2 trial: a multi-centre cluster randomised controlled trial of the clinical and cost effectiveness  
20 of the i-gel supraglottic airway device versus tracheal intubation in the initial airway management of out  
21 of hospital cardiac arrest. *Resuscitation*. 2016;109:25–32. doi: 10.1016/j.resuscitation.2016.09.016.
- 22 42. Wang HE, Prince DK, Stephens SW, Herren H, Daya M, Richmond N, Carlson J, Warden C,  
23 Colella MR, Brienza A, Aufderheide TP, Idris AH, Schmicker R, May S, Nichol G. Design and  
24 implementation of the Resuscitation Outcomes Consortium Pragmatic Airway Resuscitation Trial  
25 (PART). *Resuscitation*. 2016;101:57–64. doi: 10.1016/j.resuscitation.2016.01.012.

- 1 43. Jacobs IG, Finn JC, Jelinek GA, Oxe r HF, Thompson PL. Effect of adrenaline on survival in out-  
2 of-hospital cardiac arrest: a randomised double-blind placebo-controlled trial. *Resuscitation*.  
3 2011;82:1138–1143. doi: 10.1016/j.resuscitation.2011.06.029.
- 4 44. Loomba RS, Nijhawan K, Aggarwal S, Arora RR. Increased return of spontaneous circulation at  
5 the expense of neurologic outcomes: is prehospital epinephrine for out-of-hospital cardiac arrest really  
6 worth it? *J Crit Care*. 2015;30:1376–1381. doi: 10.1016/j.jcrc.2015.08.016.
- 7 45. Andersen LW, Kurth T, Chase M, Berg KM, Cocchi MN, Callaway C, Donnino MW; for the  
8 American Heart Association’s Get With The Guidelines–Resuscitation Investigators. Early administration  
9 of epinephrine (adrenaline) in patients with cardiac arrest with initial shockable rhythm in hospital:  
10 propensity score matched analysis. *BMJ*. 2016;353:i1577. doi: 10.1136/bmj.i1577.
- 11 46. Hubble MW, Tyson C. Impact of early vasopressor administration on neurological outcomes after  
12 prolonged out-of-hospital cardiac arrest. *Prehosp Disaster Med*. 2017:1–8. doi:  
13 10.1017/S1049023X17000115.
- 14 47. Perkins GD, Quinn T, Deakin CD, Nolan JP, Lall R, Slowther AM, Cooke M, Lamb SE, Petrou  
15 S, Achana F, Finn J, Jacobs IG, Carson A, Smyth M, Han K, Byers S, Rees N, Whitfield R, Moore F,  
16 Fothergill R, Stallard N, Long J, Hennings S, Horton J, Kaye C, Gates S. Pre-hospital Assessment of the  
17 Role of Adrenaline: Measuring the Effectiveness of Drug administration In Cardiac arrest  
18 (PARAMEDIC-2): trial protocol. *Resuscitation*. 2016;108:75–81. doi:  
19 10.1016/j.resuscitation.2016.08.029.
- 20 48. Carr BG, Kahn JM, Merchant RM, Kramer AA, Neumar RW. Inter-hospital variability in post-  
21 cardiac arrest mortality. *Resuscitation*. 2009;80:30–34. doi: 10.1016/j.resuscitation.2008.09.001.
- 22 49. Nolan JP, Ferrando P, Soar J, Beng er J, Thomas M, Harrison DA, Perkins GD. Increasing  
23 survival after admission to UK critical care units following cardiopulmonary resuscitation. *Crit Care*.  
24 2016;20:219. doi: 10.1186/s13054-016-1390-6.
- 25 50. Straney LD, Bray JE, Finn J, Bernard S, Pilcher D. Trends in intensive care unit cardiac arrest  
26 admissions and mortality in Australia and New Zealand. *Crit Care Resusc*. 2014;16:104–111.



- 1 51. Bernard SA, Gray TW, Buist MD, Jones BM, Silvester W, Gutteridge G, Smith K. Treatment of  
2 comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med*.  
3 2002;346:557–563. doi: 10.1056/NEJMoa003289.
- 4 52. Hypothermia after Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the  
5 neurologic outcome after cardiac arrest. *N Engl J Med*. 2002;346:549–556. doi: 10.1056/NEJMoa012689.
- 6 53. Nielsen N, Wetterslev J, Cronberg T, Erlinge D, Gasche Y, Hassager C, Horn J, Hovdenes J,  
7 Kjaergaard J, Kuiper M, Pellis T, Stammet P, Wanscher M, Wise MP, Aneman A, Al-Subaie N,  
8 Boesgaard S, Bro-Jeppesen J, Brunetti I, Bugge JF, Hingston CD, Juffermans NP, Koopmans M, Kober  
9 L, Langorgen J, Lilja G, Moller JE, Rundgren M, Rylander C, Smid O, Werer C, Winkel P, Friberg H; for  
10 the TTM Trial Investigators. Targeted temperature management at 33°C versus 36°C after cardiac arrest.  
11 *N Engl J Med*. 2013;369:2197–2206. doi: 10.1056/NEJMoa1310519.
- 12 54. Donnino MW, Andersen LW, Berg KM, Reynolds JC, Nolan JP, Morley PT, Lang E, Cocchi  
13 MN, Xanthos T, Callaway CW, Soar J; for the ILCOR ALS Task Force. Temperature management after  
14 cardiac arrest: an advisory statement by the Advanced Life Support Task Force of the International  
15 Liaison Committee on Resuscitation and the American Heart Association Emergency Cardiovascular  
16 Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation.  
17 *Resuscitation*. 2016;98:97–104. doi: 10.1016/j.resuscitation.2015.09.396.
- 18 55. History. TTM2 Trial Group website. <http://ttm2trial.org/about-study/history>. Accessed April 14,  
19 2017.
- 20 56. Finding the Optimal Cooling temperature After Out-of-Hospital Cardiac Arrest (FROST).  
21 <https://clinicaltrials.gov/ct2/show/NCT02035839>. Accessed April 14, 2017.
- 22 57. Neuroprotective Goal Directed Hemodynamic Optimization in Post-cardiac Arrest Patients  
23 (NEUROPROTECT). <https://clinicaltrials.gov/ct2/show/NCT02541591>. Accessed April 24, 2017.
- 24 58. Larsen JM, Ravkilde J. Acute coronary angiography in patients resuscitated from out-of-hospital  
25 cardiac arrest—a systematic review and meta-analysis. *Resuscitation*. 2012;83:1427–1433. doi:  
26 10.1016/j.resuscitation.2012.08.337.

- 1 59. Welsford M, Nikolaou NI, Beygui F, Bossaert L, Ghaemmaghami C, Nonogi H, O'Connor RE,  
2 Pichel DR, Scott T, Walters DL, Woolfrey KG; for the Acute Coronary Syndrome Chapter Collaborators.  
3 Part 5: acute coronary syndromes: 2015 International Consensus on Cardiopulmonary Resuscitation and  
4 Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2015;132(suppl  
5 1):S146–S176. doi: 10.1161/CIR.0000000000000274.
- 6 60. Nikolaou NI, Welsford M, Beygui F, Bossaert L, Ghaemmaghami C, Nonogi H, O'Connor RE,  
7 Pichel DR, Scott T, Walters DL, Woolfrey KG; for the Acute Coronary Syndrome Chapter Collaborators.  
8 Part 5: acute coronary syndromes: 2015 International Consensus on Cardiopulmonary Resuscitation and  
9 Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*.  
10 2015;95:e121–e146. doi: 10.1016/j.resuscitation.2015.07.043.
- 11 61. Lemkes JS, Janssens GN, Straaten HM, Elbers PW, van der Hoeven NW, Tijssen JG, Otterspoor  
12 LC, Voskuil M, van der Heijden JJ, Meuwissen M, Rijpstra TA, Vlachojannis GJ, van der Vleugel RM,  
13 Nieman K, Jewbali LS, Bleeker GB, Baak R, Beishuizen B, Stoel MG, van der Harst P, Camaro C,  
14 Henriques JP, Vink MA, Gosselink MT, Bosker HA, Crijs HJ, van Royen N; for the COACT  
15 investigators. Coronary angiography after cardiac arrest: rationale and design of the COACT trial. *Am*  
16 *Heart J*. 2016;180:39–45. doi: 10.1016/j.ahj.2016.06.025.
- 17 62. Early Coronary Angiography Versus Delayed Coronary Angiography (PEARL).  
18 <https://clinicaltrials.gov/ct2/show/NCT02387398>. Accessed April 24, 2017.
- 19 63. Immediate Unselected Coronary Angiography Versus Delayed Triage in Survivors of Out-of-  
20 hospital Cardiac Arrest Without ST-segment Elevation (TOMAHAWK).  
21 <https://clinicaltrials.gov/show/NCT02750462>. Accessed April 15, 2017.
- 22 64. Direct or Subacute Coronary Angiography for Out-of-hospital Cardiac Arrest (DISCO).  
23 <https://clinicaltrials.gov/ct2/show/NCT02309151>. Accessed April 15, 2017.
- 24 65. Patterson T, Perkins GD, Joseph J, Wilson K, Van Dyck L, Robertson S, Nguyen H, McConkey  
25 H, Whitbread M, Fothergill R, Nevett J, Dalby M, Rakhit R, MacCarthy P, Perera D, Nolan JP, Redwood  
26 SR. A Randomised tRial of Expedited transfer to a cardiac arrest centre for non-ST elevation ventricular

- 1 fibrillation out-of-hospital cardiac arrest: the ARREST pilot randomised trial [published online February  
2 4, 2017]. *Resuscitation*. 2017. doi: 10.1016/j.resuscitation.2017.01.020.
- 3 66. Yannopoulos D, Aufderheide TP. The ACCESS trial—CCC. Grantome website.  
4 <http://grantome.com/grant/NIH/R01-HL133818-01>. Accessed April 21, 2017.
- 5 67. Elmer J, Torres C, Aufderheide TP, Austin MA, Callaway CW, Golan E, Herren H, Jasti J,  
6 Kudenchuk PJ, Scales DC, Stub D, Richardson DK, Zive DM; the Resuscitation Outcomes Consortium.  
7 Association of early withdrawal of life-sustaining therapy for perceived neurological prognosis with  
8 mortality after cardiac arrest. *Resuscitation*. 2016;102:127–135. doi: 10.1016/j.resuscitation.2016.01.016.
- 9 68. Perman SM, Kirkpatrick JN, Reitsma AM, Gaieski DF, Lau B, Smith TM, Leary M, Fuchs BD,  
10 Levine JM, Abella BS, Becker LB, Merchant RM. Timing of neuroprognostication in postcardiac arrest  
11 therapeutic hypothermia. *Crit Care Med*. 2012;40:719–724. doi: 10.1097/CCM.0b013e3182372f93.
- 12 69. Mulder M, Gibbs HG, Smith SW, Dhaliwal R, Scott NL, Sprenkle MD, Geocadin RG.  
13 Awakening and withdrawal of life-sustaining treatment in cardiac arrest survivors treated with therapeutic  
14 hypothermia. *Crit Care Med*. 2014;42:2493–2499. doi: 10.1097/CCM.0000000000000540.
- 15 70. Anderson KB, Poloyac SM, Kochanek PM, Empey PE. Effect of hypothermia and targeted  
16 temperature management on drug disposition and response following cardiac arrest: a comprehensive  
17 review of preclinical and clinical investigations. *Ther Hypothermia Temp Manag*. 2016;6:169–179. doi:  
18 10.1089/ther.2016.0003.
- 19 71. Wijdicks EF, Bamlet WR, Maramattom BV, Manno EM, McClelland RL. Validation of a new  
20 coma scale: the FOUR score. *Ann Neurol*. 2005;58:585–593. doi: 10.1002/ana.20611.
- 21 72. Rundgren M, Cronberg T, Friberg H, Isaksson A. Serum neuron specific enolase—impact of  
22 storage and measuring method. *BMC Res Notes*. 2014;7:726. doi: 10.1186/1756-0500-7-726.
- 23 73. Choi SP, Park KN, Park HK, Kim JY, Youn CS, Ahn KJ, Yim HW. Diffusion-weighted magnetic  
24 resonance imaging for predicting the clinical outcome of comatose survivors after cardiac arrest: a cohort  
25 study. *Crit Care*. 2010;14:R17. doi: 10.1186/cc8874.

- 1 74. Youn CS, Callaway CW, Rittenberger JC; the Post Cardiac Arrest Service. Combination of initial  
2 neurologic examination and continuous EEG to predict survival after cardiac arrest. *Resuscitation*.  
3 2015;94:73–79. doi: 10.1016/j.resuscitation.2015.06.016.
- 4 75. Bouwes A, Binnekade JM, Verbaan BW, Zandbergen EG, Koelman JH, Weinstein HC, Hijdra A,  
5 Horn J. Predictive value of neurological examination for early cortical responses to somatosensory  
6 evoked potentials in patients with postanoxic coma. *J Neurol*. 2012;259:537–541. doi: 10.1007/s00415-  
7 011-6224-5.
- 8 76. Hirsch LJ, LaRoche SM, Gaspard N, Gerard E, Svoronos A, Herman ST, Mani R, Arif H, Jette N,  
9 Minazad Y, Kerrigan JF, Vespa P, Hantus S, Claassen J, Young GB, So E, Kaplan PW, Nuwer MR,  
10 Fountain NB, Drislane FW. American Clinical Neurophysiology society's standardized critical care EEG  
11 terminology: 2012 version. *J Clin Neurophysiol*. 2013;30:1–27. doi: 10.1097/WNP.0b013e3182784729.
- 12 77. Hirsch KG, Mlynash M, Jansen S, Persoon S, Eyngorn I, Krasnokutsky MV, Wijman CA,  
13 Fischbein NJ. Prognostic value of a qualitative brain MRI scoring system after cardiac arrest. *J*  
14 *Neuroimaging*. 2015;25:430–437. doi: 10.1111/jon.12143.
- 15 78. Global Health Observatory (GHO) data: causes of death. World Health Organization website.  
16 [http://www.who.int/gho/mortality\\_burden\\_disease/causes\\_death/en/](http://www.who.int/gho/mortality_burden_disease/causes_death/en/). Accessed April 21, 2017.
- 17 79. Naim MY, Burke RV, McNally BF, Song L, Griffis HM, Berg RA, Vellano K, Markenson D,  
18 Bradley RN, Rossano JW. Association of bystander cardiopulmonary resuscitation with overall and  
19 neurologically favorable survival after pediatric out-of-hospital cardiac arrest in the United States: a  
20 report from the Cardiac Arrest Registry to Enhance Survival Surveillance Registry. *JAMA Pediatr*.  
21 2017;171:133–141. doi: 10.1001/jamapediatrics.2016.3643.
- 22 80. Fink EL, Prince DK, Kaltman JR, Atkins DL, Austin M, Warden C, Hutchison J, Daya M,  
23 Goldberg S, Herren H, Tijssen JA, Christenson J, Vaillancourt C, Miller R, Schmicker RH, Callaway  
24 CW; for the Resuscitation Outcomes Consortium. Unchanged pediatric out-of-hospital cardiac arrest  
25 incidence and survival rates with regional variation in North America. *Resuscitation*. 2016;107:121–128.  
26 doi: 10.1016/j.resuscitation.2016.07.244.

- 1 81. Martinez PA, Totapally BR. The epidemiology and outcomes of pediatric in-hospital  
2 cardiopulmonary arrest in the United States during 1997 to 2012. *Resuscitation*. 2016;105:177–181. doi:  
3 10.1016/j.resuscitation.2016.06.010.
- 4 82. Berg RA, Nadkarni VM, Clark AE, Moler F, Meert K, Harrison RE, Newth CJ, Sutton RM,  
5 Wessel DL, Berger JT, Carcillo J, Dalton H, Heidemann S, Shanley TP, Zuppa AF, Doctor A, Tamburro  
6 RF, Jenkins TL, Dean JM, Holubkov R, Pollack MM; Eunice Kennedy Shriver National Institute of Child  
7 Health and Human Development Collaborative Pediatric Critical Care Research Network. Incidence and  
8 outcomes of cardiopulmonary resuscitation in PICUs. *Crit Care Med*. 2016;44:798–808. doi:  
9 10.1097/CCM.0000000000001484.
- 10 83. Summary health statistics tables for the U.S. population: national health interview survey: table  
11 P-10. Centers for Disease Control and Prevention (CDC)/National Center for Health Statistics website.  
12 [www.cdc.gov/nchs/fastats/hospital.htm](http://www.cdc.gov/nchs/fastats/hospital.htm). Accessed April 21, 2017.
- 13 84. Maconochie IK, de Caen AR, Aickin R, Atkins DL, Biarent D, Guerguerian AM, Kleinman ME,  
14 Kloeck DA, Meaney PA, Nadkarni VM, Ng KC, Nuthall G, Reis AG, Shimizu N, Tibballs J, Pintos RV;  
15 for the Pediatric Basic Life Support and Pediatric Advanced Life Support Chapter Collaborators. Part 6:  
16 pediatric basic life support and pediatric advanced life support: 2015 International Consensus on  
17 Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment  
18 Recommendations. *Resuscitation*. 2015;95:e147–e168. doi: 10.1016/j.resuscitation.2015.07.044.
- 19 85. de Caen AR, Maconochie IK, Aickin R, Atkins DL, Biarent D, Guerguerian AM, Kleinman ME,  
20 Kloeck DA, Meaney PA, Nadkarni VM, Ng KC, Nuthall G, Reis AG, Shimizu N, Tibballs J, Veliz Pintos  
21 R; for the Pediatric Basic Life Support and Pediatric Advanced Life Support Chapter Collaborators. Part  
22 6: pediatric basic life support and pediatric advanced life support: 2015 International Consensus on  
23 Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment  
24 Recommendations. *Circulation*. 2015;132(suppl 1):S177–S203. doi: 10.1161/CIR.0000000000000275.
- 25 86. Lubrano R, Cecchetti C, Bellelli E, Gentile I, Loayza Levano H, Orsini F, Bertazzoni G, Messi G,  
26 Rugolotto S, Pirozzi N, Elli M. Comparison of times of intervention during pediatric CPR maneuvers

- 1 using ABC and CAB sequences: a randomized trial. *Resuscitation*. 2012;83:1473–1477. doi:  
2 10.1016/j.resuscitation.2012.04.011.
- 3 87. Marsch S, Tschan F, Semmer NK, Zobrist R, Hunziker PR, Hunziker S. ABC versus CAB for  
4 cardiopulmonary resuscitation: a prospective, randomized simulator-based trial. *Swiss Med Wkly*.  
5 2013;143:w13856. doi: 10.4414/smw.2013.13856.
- 6 88. Sekiguchi H, Kondo Y, Kukita I. Verification of changes in the time taken to initiate chest  
7 compressions according to modified basic life support guidelines. *Am J Emerg Med*. 2013;31:1248–1250.  
8 doi: 10.1016/j.ajem.2013.02.047.
- 9 89. Maconochie IK, Bingham R, Eich C, López-Herce J, Rodríguez-Núñez A, Rajka T, Van de  
10 Voorde P, Zideman DA, Biarent D; for the Paediatric life support section Collaborators. European  
11 Resuscitation Council Guidelines for Resuscitation 2015: section 6: paediatric life support. *Resuscitation*.  
12 2015;95:223–248. doi: 10.1016/j.resuscitation.2015.07.028.
- 13 90. Atkins DL, Berger S, Duff JP, Gonzales JC, Hunt EA, Joyner BL, Meaney PA, Niles DE,  
14 Samson RA, Schexnayder SM. Part 11: pediatric basic life support and cardiopulmonary resuscitation  
15 quality: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and  
16 Emergency Cardiovascular Care. *Circulation*. 2015;132(suppl 2):S519–S525. doi:  
17 10.1161/CIR.0000000000000265.
- 18 91. Kleinman ME, Brennan EE, Goldberger ZD, Swor RA, Terry M, Bobrow BJ, Gazmuri RJ,  
19 Travers AH, Rea T. Part 5: adult basic life support and cardiopulmonary resuscitation quality: 2015  
20 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency  
21 Cardiovascular Care. *Circulation*. 2015;132:S414–S435. doi: 10.1161/CIR.0000000000000259.
- 22 92. Rodríguez-Núñez A, López-Herce J, del Castillo J, Bellón JM; Iberian-American Paediatric  
23 Cardiac Arrest Study Network RIPEPCI. Shockable rhythms and defibrillation during in-hospital  
24 pediatric cardiac arrest. *Resuscitation*. 2014;85:387–391. doi: 10.1016/j.resuscitation.2013.11.015.
- 25 93. Meaney PA, Nadkarni VM, Atkins DL, Berg MD, Samson RA, Hazinski MF, Berg RA; for the  
26 American Heart Association National Registry of Cardiopulmonary Resuscitation Investigators. Effect of

1 defibrillation energy dose during in-hospital pediatric cardiac arrest. *Pediatrics*. 2011;127:e16–e23. doi:  
2 10.1542/peds.2010-1617.

3 94. Moler FW, Silverstein FS, Holubkov R, Slomine BS, Christensen JR, Nadkarni VM, Meert KL,  
4 Clark AE, Browning B, Pemberton VL, Page K, Shankaran S, Hutchison JS, Newth CJ, Bennett KS,  
5 Berger JT, Topjian A, Pineda JA, Koch JD, Schleien CL, Dalton HJ, Ofori-Amanfo G, Goodman DM,  
6 Fink EL, McQuillen P, Zimmerman JJ, Thomas NJ, van der Jagt EW, Porter MB, Meyer MT, Harrison R,  
7 Pham N, Schwarz AJ, Nowak JE, Alten J, Wheeler DS, Bhalala US, Lidsky K, Lloyd E, Mathur M, Shah  
8 S, Wu T, Theodorou AA, Sanders RC Jr, Dean JM; for the THAPCA Trial Investigators. Therapeutic  
9 hypothermia after out-of-hospital cardiac arrest in children. *N Engl J Med*. 2015;372:1898–1908. doi:  
10 10.1056/NEJMoa1411480.

11 95. Moler FW, Silverstein FS, Holubkov R, Slomine BS, Christensen JR, Nadkarni VM, Meert KL,  
12 Browning B, Pemberton VL, Page K, Gildea MR, Scholefield BR, Shankaran S, Hutchison JS, Berger JT,  
13 Ofori-Amanfo G, Newth CJ, Topjian A, Bennett KS, Koch JD, Pham N, Chanani NK, Pineda JA,  
14 Harrison R, Dalton HJ, Alten J, Schleien CL, Goodman DM, Zimmerman JJ, Bhalala US, Schwarz AJ,  
15 Porter MB, Shah S, Fink EL, McQuillen P, Wu T, Skellett S, Thomas NJ, Nowak JE, Baines PB,  
16 Pappachan J, Mathur M, Lloyd E, van der Jagt EW, Dobyms EL, Meyer MT, Sanders RC Jr, Clark AE,  
17 Dean JM; for the THAPCA Trial Investigators. Therapeutic hypothermia after in-hospital cardiac arrest in  
18 children. *N Engl J Med*. 2017;376:318–329. doi: 10.1056/NEJMoa1610493.

19 96. Bistriz JF, Horton LM, Smaldone A. Therapeutic hypothermia in children after cardiac arrest: a  
20 systematic review and meta-analysis. *Pediatr Emerg Care*. 2015;31:296–303. doi:  
21 10.1097/PEC.0000000000000407.

22 97. Vincent JL, Taccone FS. Difficulty interpreting the results of some trials: the case of therapeutic  
23 hypothermia after pediatric cardiac arrest. *Crit Care*. 2015;19:391. doi: 10.1186/s13054-015-1121-4.

24 98. Mentzelopoulos SD, Zakynthinos SG, Siempos I, Malachias S, Ulmer H, Wenzel V. Vasopressin  
25 for cardiac arrest: meta-analysis of randomized controlled trials. *Resuscitation*. 2012;83:32–39. doi:  
26 10.1016/j.resuscitation.2011.07.015.

- 1 99. Lin S, Callaway CW, Shah PS, Wagner JD, Beyene J, Ziegler CP, Morrison LJ. Adrenaline for  
2 out-of-hospital cardiac arrest resuscitation: a systematic review and meta-analysis of randomized  
3 controlled trials. *Resuscitation*. 2014;85:732–740. doi: 10.1016/j.resuscitation.2014.03.008.
- 4 100. Enright K, Turner C, Roberts P, Cheng N, Browne G. Primary cardiac arrest following sport or  
5 exertion in children presenting to an emergency department: chest compressions and early defibrillation  
6 can save lives, but is intravenous epinephrine always appropriate? *Pediatr Emerg Care*. 2012;28:336–  
7 339. doi: 10.1097/PEC.0b013e31824d8c78.
- 8 101. Perondi MB, Reis AG, Paiva EF, Nadkarni VM, Berg RA. A comparison of high-dose and  
9 standard-dose epinephrine in children with cardiac arrest. *N Engl J Med*. 2004;350:1722–1730. doi:  
10 10.1056/NEJMoa032440.
- 11 102. Andersen LW, Berg KM, Saindon BZ, Massaro JM, Raymond TT, Berg RA, Nadkarni VM,  
12 Donnino MW; for the American Heart Association Get With The Guidelines–Resuscitation Investigators.  
13 Time to epinephrine and survival after pediatric in-hospital cardiac arrest. *JAMA*. 2015;314:802–810. doi:  
14 10.1001/jama.2015.9678.
- 15 103. Carroll TG, Dimas VV, Raymond TT. Vasopressin rescue for in-pediatric intensive care unit  
16 cardiopulmonary arrest refractory to initial epinephrine dosing: a prospective feasibility pilot trial. *Pediatr*  
17 *Crit Care Med*. 2012;13:265–272. doi: 10.1097/PCC.0b013e31822f1569.
- 18 104. Søreide E, Morrison L, Hillman K, Monsieurs K, Sunde K, Zideman D, Eisenberg M, Sterz F,  
19 Nadkarni VM, Soar J, Nolan JP; for the Utstein Formula for Survival Collaborators. The formula for  
20 survival in resuscitation. *Resuscitation*. 2013;84:1487–1493. doi: 10.1016/j.resuscitation.2013.07.020.
- 21 105. Finn JC, Bhanji F, Lockey A, Monsieurs K, Frengley R, Iwami T, Lang E, Ma MH, Mancini ME,  
22 McNeil MA, Greif R, Billi JE, Nadkarni VM, Bigham B; for the Education, Implementation, and Teams  
23 Chapter Collaborators. Part 8: education, implementation, and teams: 2015 International Consensus on  
24 Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment  
25 Recommendations. *Resuscitation*. 2015;95:e203–e224. doi: 10.1016/j.resuscitation.2015.07.046.



- 1 106. Bhanji F, Finn JC, Lockey A, Monsieurs K, Frengley R, Iwami T, Lang E, Ma MH, Mancini ME,  
2 McNeil MA, Greif R, Billi JE, Nadkarni VM, Bigham B; for the Education, Implementation, and Teams  
3 Chapter Collaborators. Part 8: education, implementation, and teams: 2015 International Consensus on  
4 Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment  
5 Recommendations. *Circulation*. 2015;132(suppl 1):S242–S268. doi: 10.1161/CIR.0000000000000277.
- 6 107. Dail TK, Christina RW. Distribution of practice and metacognition in learning and long-term  
7 retention of a discrete motor task. *Res Q Exerc Sport*. 2004;75:148–155. doi:  
8 10.1080/02701367.2004.10609146.
- 9 108. Turner NM, Dierselhuis MP, Draaisma JM, ten Cate OT. The effect of the Advanced Paediatric  
10 Life Support course on perceived self-efficacy and use of resuscitation skills. *Resuscitation*. 2007;73:430–  
11 436. doi: 10.1016/j.resuscitation.2006.10.018.
- 12 109. Hasselqvist-Ax I, Riva G, Herlitz J, Rosenqvist M, Hollenberg J, Nordberg P, Ringh M, Jonsson  
13 M, Axelsson C, Lindqvist J, Karlsson T, Svensson L. Early cardiopulmonary resuscitation in out-of-  
14 hospital cardiac arrest. *N Engl J Med*. 2015;372:2307–2315. doi: 10.1056/NEJMoa1405796.
- 15 110. Kalz M, Lenssen N, Felzen M, Rossaint R, Tabuenca B, Specht M, Skorning M. Smartphone  
16 apps for cardiopulmonary resuscitation training and real incident support: a mixed-methods evaluation  
17 study. *J Med Internet Res*. 2014;16:e89. doi: 10.2196/jmir.2951.
- 18 111. Schober A, Sterz F, Laggner AN, Poppe M, Sulzgruber P, Lobmeyr E, Datler P, Keferböck M,  
19 Zeiner S, Nuernberger A, Eder B, Hinterholzer G, Mydza D, Enzelsberger B, Herlich K, Schuster R,  
20 Koeller E, Publig T, Smetana P, Scheibenpflug C, Christ G, Meyer B, Uray T. Admission of out-of-  
21 hospital cardiac arrest victims to a high volume cardiac arrest center is linked to improved outcome.  
22 *Resuscitation*. 2016;106:42–48. doi: 10.1016/j.resuscitation.2016.06.021.
- 23 112. Elmer J, Rittenberger JC, Coppler PJ, Guyette FX, Doshi AA, Callaway CW; the Pittsburgh Post-  
24 Cardiac Arrest Service. Long-term survival benefit from treatment at a specialty center after cardiac  
25 arrest. *Resuscitation*. 2016;108:48–53. doi: 10.1016/j.resuscitation.2016.09.008.

- 1 113. Tranberg T, Lippert FK, Christensen EF, Stengaard C, Hjort J, Lassen JF, Petersen F, Jensen JS,  
2 Back C, Jensen LO, Ravkilde J, Botker HE, Terkelsen CJ. Distance to invasive heart centre, performance  
3 of acute coronary angiography, and angioplasty and associated outcome in out-of-hospital cardiac arrest:  
4 a nationwide study [published online March 29, 2017]. *Eur Heart J*. 2017. doi: 10.1093/eurheartj/ehx104.
- 5 114. Minei JP, Fabian TC, Guffey DM, Newgard CD, Bulger EM, Brasel KJ, Sperry JL, MacDonald  
6 RD. Increased trauma center volume is associated with improved survival after severe injury: results of a  
7 Resuscitation Outcomes Consortium study. *Ann Surg*. 2014;260:456–464. doi:  
8 10.1097/SLA.0000000000000873.
- 9 115. Tung YC, Chang GM, Chien KL, Tu YK. The relationships among physician and hospital  
10 volume, processes, and outcomes of care for acute myocardial infarction. *Med Care*. 2014;52:519–527.  
11 doi: 10.1097/MLR.0000000000000132.
- 12 116. Soar J, Mancini ME, Bhanji F, Billi JE, Dennett J, Finn J, Ma MH, Perkins GD, Rodgers DL,  
13 Hazinski MF, Jacobs I, Morley PT; for the Education, Implementation, and Teams Chapter Collaborators.  
14 Part 12: education, implementation, and teams: 2010 International Consensus on Cardiopulmonary  
15 Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations.  
16 *Resuscitation*. 2010;81(suppl 1):e288–e330. doi: 10.1016/j.resuscitation.2010.08.030.
- 17 117. Bigham BL, Koprowicz K, Aufderheide TP, Davis DP, Donn S, Powell J, Suffoletto B, Nafziger  
18 S, Stouffer J, Idris A, Morrison LJ; and the ROC Investigators. Delayed prehospital implementation of the  
19 2005 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency  
20 Cardiac Care. *Prehosp Emerg Care*. 2010;14:355–360. doi: 10.3109/10903121003770639.
- 21 118. Kim YM, Lee SJ, Jo SJ, Park KN. Implementation of the guidelines for targeted temperature  
22 management after cardiac arrest: a longitudinal qualitative study of barriers and facilitators perceived by  
23 hospital resuscitation champions. *BMJ Open*. 2016;6:e009261. doi: 10.1136/bmjopen-2015-009261.
- 24 119. Dainty KN, Brooks SC, Morrison LJ. Are the 2010 guidelines on cardiopulmonary resuscitation  
25 lost in translation? A call for increased focus on implementation science. *Resuscitation*. 2013;84:422–  
26 425. doi: 10.1016/j.resuscitation.2012.08.336.

- 1 120. Morrison LJ, Brooks SC, Dainty KN, Dorian P, Needham DM, Ferguson ND, Rubenfeld GD,  
2 Slutsky AS, Wax RS, Zwarenstein M, Thorpe K, Zhan C, Scales DC; for the Strategies for Post-Arrest  
3 Care Network. Improving use of targeted temperature management after out-of-hospital cardiac arrest: a  
4 stepped wedge cluster randomized controlled trial. *Crit Care Med.* 2015;43:954–964. doi:  
5 10.1097/CCM.0000000000000864.
- 6 121. Singletary EM, Zideman DA, De Buck ED, Chang WT, Jensen JL, Swain JM, Woodin JA,  
7 Blanchard IE, Herrington RA, Pellegrino JL, Hood NA, Lojero-Wheatley LF, Markenson DS, Yang HJ;  
8 for the First Aid Chapter Collaborators. Part 9: first aid: 2015 International Consensus on First Aid  
9 Science With Treatment Recommendations. *Circulation.* 2015;132(suppl 1):S269–S311. doi:  
10 10.1161/CIR.0000000000000278.
- 11 122. Zideman DA, Singletary EM, De Buck ED, Chang WT, Jensen JL, Swain JM, Woodin JA,  
12 Blanchard IE, Herrington RA, Pellegrino JL, Hood NA, Lojero-Wheatley LF, Markenson DS, Yang HJ;  
13 for the First Aid Chapter Collaborators. Part 9: first aid: 2015 International Consensus on First Aid  
14 Science With Treatment Recommendations. *Resuscitation.* 2015;95:e225–e261. doi:  
15 10.1016/j.resuscitation.2015.07.047.
- 16 123. Adnet F, Borron SW, Finot MA, Minadeo J, Baud FJ. Relation of body position at the time of  
17 discovery with suspected aspiration pneumonia in poisoned comatose patients. *Crit Care Med.*  
18 1999;27:745–748.
- 19 124. Freire-Tellado M, Pavón-Prieto Mdel P, Fernández-López M, Navarro-Patón R. Does the  
20 recovery position threaten cardiac arrest victim's safety assessment? *Resuscitation.* 2016;105:e1. doi:  
21 10.1016/j.resuscitation.2016.01.040.
- 22 125. Seyal M, Bateman LM, Li CS. Impact of periictal interventions on respiratory dysfunction,  
23 postictal EEG suppression, and postictal immobility. *Epilepsia.* 2013;54:377–382. doi: 10.1111/j.1528-  
24 1167.2012.03691.x.
- 25 126. Jullian S, Desmarest M, Gonzalez L, Ballesteros Y, Martinez A, Moretti R, Rivas A, Lacroix L,  
26 Biver A, Lejay E, Kanagarajah L, Portillo N, Cricchiutti G, Stefani C, Da Dalt L, Spiri D, Van De Voorde

- 1 P, Titomanlio L. Recovery position significantly associated with a reduced admission rate of children  
2 with loss of consciousness. *Arch Dis Child*. 2016;101:521–526. doi: 10.1136/archdischild-2015-308857.
- 3 127. Longphre JM, Denoble PJ, Moon RE, Vann RD, Freiburger JJ. First aid normobaric oxygen for  
4 the treatment of recreational diving injuries. *Undersea Hyperb Med*. 2007;34:43–49.
- 5 128. Bruera E, de Stoutz N, Velasco-Leiva A, Schoeller T, Hanson J. Effects of oxygen on dyspnoea  
6 in hypoxaemic terminal-cancer patients. *Lancet*. 1993;342:13–14.
- 7 129. Uronis HE, Currow DC, McCrory DC, Samsa GP, Abernethy AP. Oxygen for relief of dyspnoea  
8 in mildly- or non-hypoxaemic patients with cancer: a systematic review and meta-analysis. *Br J Cancer*.  
9 2008;98:294–299. doi: 10.1038/sj.bjc.6604161.
- 10 130. Philip J, Gold M, Milner A, Di Iulio J, Miller B, Spruyt O. A randomized, double-blind,  
11 crossover trial of the effect of oxygen on dyspnea in patients with advanced cancer. *J Pain Symptom*  
12 *Manage*. 2006;32:541–550. doi: 10.1016/j.jpainsymman.2006.06.009.
- 13 131. Booth S, Kelly MJ, Cox NP, Adams L, Guz A. Does oxygen help dyspnea in patients with  
14 cancer? *Am J Respir Crit Care Med*. 1996;153:1515–1518. doi: 10.1164/ajrccm.153.5.8630595.
- 15 132. Help us stop bleeding. Stop the Bleeding Coalition website. <http://stopthebleedingcoalition.org>.  
16 Accessed April 21, 2017.
- 17 133. Beekley AC, Sebesta JA, Blackburne LH, Herbert GS, Kauvar DS, Baer DG, Walters TJ,  
18 Mullenix PS, Holcomb JB; members of the 31st Combat Support Hospital Research Group. Prehospital  
19 tourniquet use in Operation Iraqi Freedom: effect on hemorrhage control and outcomes. *J Trauma*.  
20 2008;64:S28–S37. doi: 10.1097/TA.0b013e318160937e.
- 21 134. Lakstein D, Blumenfeld A, Sokolov T, Lin G, Bssorai R, Lynn M, Ben-Abraham R. Tourniquets  
22 for hemorrhage control on the battlefield: a 4-year accumulated experience. *J Trauma*. 2003;54:S221–  
23 S225. doi: 10.1097/01.TA.0000047227.33395.49.
- 24 135. King DR, van der Wilden G, Kragh JF Jr, Blackburne LH. Forward assessment of 79 prehospital  
25 battlefield tourniquets used in the current war. *J Spec Oper Med*. 2012;12:33–38.

- 1 136. Kragh JF Jr, Littrel ML, Jones JA, Walters TJ, Baer DG, Wade CE, Holcomb JB. Battle casualty  
2 survival with emergency tourniquet use to stop limb bleeding. *J Emerg Med*. 2011;41:590–597. doi:  
3 10.1016/j.jemermed.2009.07.022.
- 4 137. Kragh JF Jr, Cooper A, Aden JK, Dubick MA, Baer DG, Wade CE, Blackburne LH. Survey of  
5 trauma registry data on tourniquet use in pediatric war casualties. *Pediatr Emerg Care*. 2012;28:1361–  
6 1365. doi: 10.1097/PEC.0b013e318276c260.
- 7 138. Tien HC, Jung V, Rizoli SB, Acharya SV, MacDonald JC. An evaluation of tactical combat  
8 casualty care interventions in a combat environment. *J Am Coll Surg*. 2008;207:174–178. doi:  
9 10.1016/j.jamcollsurg.2008.01.065.
- 10 139. Kragh JF Jr, Nam JJ, Berry KA, Mase VJ Jr, Aden JK III, Walters TJ, Dubick MA, Baer DG,  
11 Wade CE, Blackburne LH. Transfusion for shock in US military war casualties with and without  
12 tourniquet use. *Ann Emerg Med*. 2015;65:290–296. doi: 10.1016/j.annemergmed.2014.10.021.
- 13 140. Brodie S, Hodgetts TJ, Ollerton J, McLeod J, Lambert P, Mahoney P. Tourniquet use in combat  
14 trauma: UK military experience. *J R Army Med Corps*. 2007;153:310–313.
- 15 141. Passos E, Dingley B, Smith A, Engels PT, Ball CG, Faidi S, Nathens A, Tien H; for the Canadian  
16 Trauma Trials Collaborative. Tourniquet use for peripheral vascular injuries in the civilian setting. *Injury*.  
17 2014;45:573–577. doi: 10.1016/j.injury.2013.11.031.
- 18 142. Kue RC, Temin ES, Weiner SG, Gates J, Coleman MH, Fisher J, Dyer S. Tourniquet use in a  
19 civilian emergency medical services setting: a descriptive analysis of the boston EMS experience.  
20 *Prehosp Emerg Care*. 2015;19:399–404. doi: 10.3109/10903127.2014.995842.
- 21 143. Ganeshalingam R, O'Connor M. Evidence behind the WHO guidelines: hospital care for children:  
22 what is the efficacy of sublingual, oral and intravenous glucose in the treatment of hypoglycaemia? *J*  
23 *Trop Pediatr*. 2009;55:287–289. doi: 10.1093/tropej/fmp099.
- 24 144. Graz B, Dicko M, Willcox ML, Lambert B, Falquet J, Forster M, Giani S, Diakite C, Dembele  
25 EM, Diallo D, Barennes H. Sublingual sugar for hypoglycaemia in children with severe malaria: a pilot  
26 clinical study. *Malar J*. 2008;7:242. doi: 10.1186/1475-2875-7-242.


1 145. Shrier I, Boissy P, Lebel K, Boulay J, Segal E, Delaney JS, Vacon LC, Steele RJ. Cervical spine  
2 motion during transfer and stabilization techniques. *Prehosp Emerg Care*. 2015;19:116–125. doi:  
3 10.3109/10903127.2014.936634.

4 146. Purvis TA, Carlin B, Driscoll P. The definite risks and questionable benefits of liberal pre-  
5 hospital spinal immobilisation [published online January 26, 2017]. *Am J Emerg Med*. 2017. doi:  
6 10.1016/j.ajem.2017.01.045.

7

1 **Figures**

Universal Chain of Survival: **Knowledge Gaps Map**



2

3 **Figure. Universal Chain of Survival: knowledge gaps map.**

4 A-B-C indicates airway, breathing, circulation; ALS, advanced life support; BLS, basic life  
 5 support; CA, cardiac arrest; C-A-B, compressions, airway, breathing; CPR, cardiopulmonary  
 6 resuscitation; EIT, Education, Implementation, and Teams; IHCA, in-hospital cardiac arrest;  
 7 ILCOR, International Liaison Committee on Resuscitation; PAC, postarrest care; PAD, public  
 8 access defibrillation; PAI, prearrival instructions; Peds, Pediatric; ROSC, return of spontaneous  
 9 circulation.

10