

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/107748>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

A GAP THEOREM FOR FREE BOUNDARY MINIMAL SURFACES IN THE THREE-BALL

LUCAS AMBROZIO AND IVALDO NUNES

ABSTRACT. We show that, among free boundary minimal surfaces in the unit ball in the three-dimensional Euclidean space, the flat equatorial disk and the critical catenoid are characterised by a pinching condition on the length of their second fundamental form.

1. INTRODUCTION

Let B^3 denote the unit ball in \mathbb{R}^3 , centred at the origin. The critical points of the area functional on the set of compact surfaces in B^3 whose boundaries are free to move in ∂B are minimal surfaces that intersect ∂B orthogonally.

The simplest examples of free boundary minimal surfaces in B^3 are the flat equatorial disk and the critical catenoid, which will be described in details below. Fraser and Schoen [7] constructed embedded examples of genus zero with any number of boundary components by finding metrics on surfaces of this topological type that maximize the first Steklov eigenvalue with fixed boundary length. The gluing construction of Pacard, Folha and Zolotareva [16] yielded free boundary minimal surfaces in B^3 of genus zero or one, and with any number of boundary components greater than a large constant. The existence of examples with three boundary components and arbitrarily large genus has been established by Martin Li and Kapouleas [11] and Ketover [12]. Although expected, it is not clear yet whether a free boundary minimal surface in B^3 can be of arbitrary genus and have any number of boundary components.

There are classifications theorems that single out the flat equatorial disk and the critical catenoid among free boundary minimal surfaces in B^3 . Some of them will be reviewed in Section 2. In this work, we sought for a new characterisation of the flat equatorial disk and the critical catenoid in terms of the length of their second fundamental forms. It turns out that a pinching condition on this quantity, that takes into

Mathematics Subject Classification codes: 53A05, 53A10.

account the support function of the surface, is sufficient. The result we obtained is the following:

Theorem. *Let Σ^2 be a compact free boundary minimal surface in B^3 . Assume that, for all points x in Σ ,*

$$(1) \quad |A|^2(x)\langle x, N(x) \rangle^2 \leq 2,$$

where $N(x)$ denotes a unit normal vector at the point $x \in \Sigma$ and A denotes the second fundamental form of Σ . Then

- i) either $|A|^2\langle x, N \rangle^2 \equiv 0$ and Σ is a flat equatorial disk;*
- ii) or $|A|^2(p)\langle p, N(p) \rangle^2 = 2$ at some point $p \in \Sigma$ and Σ is a critical catenoid.*

The result above bears some resemblance with the following theorem, which answered a question raised by the work of Simons [18]:

Theorem (Chern-do Carmo-Kobayashi [4], Lawson [10]). *Let Σ^n be a closed minimal hypersurface in the unit sphere S^{n+1} . Assume that*

$$|A|^2 \leq n.$$

Then

- i) either $|A|^2 \equiv 0$ and Σ is an equator;*
- ii) or $|A|^2 \equiv n$ and Σ is a Clifford hypersurface (product of two round spheres of appropriate radii and dimensions).*

However, the proof of our theorem is quite different and some steps are true only in low dimensions. It would be interesting to know if our result can be generalised to higher ambient dimension and submanifold co-dimensions, providing a full analogy between the results of [4] and this free boundary setting.

Using some arguments presented in this paper, Haizhong Li and Changwei Xiong [13] proved an interesting analogous gap result for free boundary minimal surfaces in geodesic balls of the hyperbolic three-space and of the round three-dimensional hemisphere.

2. THE FLAT EQUATORIAL DISK AND THE CRITICAL CATENOID

The flat equatorial disks are defined as the intersection of the unit ball B^3 with a plane containing the origin.

The flat equatorial disks are the only totally geodesic free boundary minimal surfaces in B^3 . Nitsche [15] has shown that they are in fact the only immersed free boundary minimal surfaces in B^3 that are homeomorphic to a disk. An interesting generalisation of his result to higher codimensions was obtained by Fraser and Schoen in [8]. In a different direction, it is known that the flat equatorial disk has the least

possible area among free boundary minimal surfaces in B^3 (see for example [17], Proposition 3, or [9], Theorem 5.4). This characterisation was generalised by Brendle [1]. Finally, flat equatorial disks are the only free boundary minimal surfaces with Morse index equal to one (a more general statement can be found in [7], Theorem 3.1).

Critical catenoids, on the other hand, are the only non-flat free boundary minimal surfaces in B^3 that are invariant under rotations around a given axis. They can be analytically defined as the image of the conformal maps

$$\begin{aligned} X : (t, \theta) &\in [-t_0, t_0] \times \mathbb{R} \\ &\mapsto a_0 \cosh(t) \cos(\theta)e_1 + a_0 \cosh(t) \sin(\theta)e_2 + a_0 t e_3 \in \mathbb{R}^3 \end{aligned}$$

where $\{e_1, e_2, e_3\}$ is some orthonormal basis of \mathbb{R}^3 . The constant t_0 is the unique positive solution to the equation $t \sinh(t) = \cosh(t)$, while $a_0 = (t_0 \cosh(t_0))^{-1}$. These constants are chosen in such way that the map X is conformal and its image is a piece of a catenoid contained in B^3 , meeting ∂B orthogonally, and whose axis of symmetry is the line generated by the vector e_3 .

It has been conjectured that the critical catenoid is the only embedded free boundary minimal annulus in B^3 (for example, in [6], Conjecture 1.1). McGrath [14] showed that this conjecture is true under the additional assumption that the surface is symmetric with respect to reflections through three mutually orthogonal planes. Fraser and Schoen [7] proved that the critical catenoid is the only free boundary minimal annulus that is immersed in B^3 by its first Steklov eigenfunctions. It would be also very interesting to classify the critical catenoid only by its Morse index.

Regarding our main result, we observe that, in the parametrization X given above, the second fundamental form and the support function are given by

$$|A|^2 = \frac{2}{a_0^2 \cosh^4(t)} \quad \text{and} \quad \langle x, N \rangle^2 = a_0^2 \left(1 - \frac{t \sinh(t)}{\cosh(t)} \right)^2.$$

In particular, since $|t| \leq t_0$,

$$|A|^2 \langle x, N \rangle^2 = \frac{2}{\cosh^6(t)} (\cosh(t) - t \sinh(t))^2 \leq 2.$$

Notice that the maximum value 2 is attained at $t = 0$, that is, on the circle defined by the intersection of the given critical catenoid and the plane through the origin that is orthogonal to its axis of symmetry.

3. PROOF OF THE THEOREM

Throughout the proof, by *surface* we mean a smooth compact surface with boundary Σ^2 smoothly immersed in the unit ball B^3 in such way that its boundary $\partial\Sigma$ is mapped in the unit sphere ∂B . We assume Σ is endowed with the induced metric from the Euclidean space. Passing to the double oriented cover, if necessary, we always assume Σ is orientable. As the arguments involve local computations and considerations about the intrinsic geometry of Σ , for the sake of notational convenience we do not distinguish between Σ and its image under the immersion.

The first key observation is to relate the pinching condition (1) to a condition on the Hessian of the distance function of points on Σ to the origin.

Lemma 1. *Let Σ^2 be a free boundary minimal surface in B^3 . Let f be the function defined by*

$$f(x) = \frac{|x|^2}{2}, \quad x \in \Sigma.$$

Then

- i) $\nabla^\Sigma f(x) = x$ for all $x \in \partial\Sigma$.*
- ii) For each $x \in \Sigma$, the eigenvalues of $\text{Hess}_\Sigma f(x)$ are given by*

$$1 - \frac{|A|(x)}{\sqrt{2}} \langle x, N(x) \rangle \quad \text{and} \quad 1 + \frac{|A|(x)}{\sqrt{2}} \langle x, N(x) \rangle.$$

Proof. In order to prove *i)*, we first note that $\nabla^\Sigma f(x) = x^T$ for all $x \in \Sigma$, where $(\cdot)^T$ stands for the orthogonal projection onto $T_x\Sigma$. Since Σ meets ∂B orthogonally along $\partial\Sigma$, $x \in T_x\Sigma$ for all $x \in \partial\Sigma$ and the assertion follows.

Given $x \in \Sigma$, let N be a local unit normal vector field in a neighbourhood of x . Denote the Levi-Civita connections of \mathbb{R}^3 and Σ by D and ∇^Σ , respectively. For all $X, Y \in T_x\Sigma$ we have

$$\begin{aligned} \text{Hess}_\Sigma f(x)(X, Y) &= XY(f) - (\nabla_X^\Sigma Y)(f) \\ &= X \langle x, Y \rangle - \langle x, \nabla_X^\Sigma Y \rangle \\ &= \langle X, Y \rangle + \langle x, D_X Y \rangle - \langle x, \nabla_X^\Sigma Y \rangle \\ &= \langle X, Y \rangle - \langle A(X), Y \rangle \langle x, N(x) \rangle \\ &= \langle X - A(X) \langle x, N(x) \rangle, Y \rangle. \end{aligned}$$

Therefore the eigenvalues of $\text{Hess}_\Sigma f(x)$ are given by

$$1 - k_1 \langle x, N(x) \rangle \quad \text{and} \quad 1 - k_2 \langle x, N(x) \rangle,$$

where $k_1 \leq k_2$ are the principal curvatures of Σ at x . Since Σ is minimal, $k_1 = -|A|(x)/\sqrt{2}$ and $k_2 = |A|(x)/\sqrt{2}$. This finishes the proof of *ii*). \square

Under the assumptions of the Theorem, we then conclude that the function f defined in Lemma 1 must be convex. This imposes very strong restrictions on the geometry of the surface Σ .

Proposition 2. *Let Σ^2 be a compact free boundary minimal surface in B^3 . Define*

$$\mathcal{C} = \{p \in \Sigma; |p| = \min_{x \in \Sigma} |x|\}.$$

If $|A|^2 \langle x, N \rangle^2 \leq 2$ on Σ , then

- i) either \mathcal{C} contains a single point $p \in \Sigma \setminus \partial\Sigma$, in which case Σ must be a flat equatorial disk;*
- ii) or \mathcal{C} is a simple closed geodesic in $\Sigma \setminus \partial\Sigma$ and Σ is homeomorphic to an annulus.*

Proof. Let $f : \Sigma \rightarrow \mathbb{R}$ be defined as in Lemma 1. We claim that \mathcal{C} is a totally convex subset of Σ (recall that a subset A of a Riemannian manifold (M^n, g) is totally convex when any geodesic in M^n joining two points in A actually lies in A , see for example [2]). In fact, given $p, q \in \mathcal{C}$, let $\gamma : [0, 1] \rightarrow \Sigma$ be a geodesic such that $\gamma(0) = p$ and $\gamma(1) = q$. By item *ii*) in Lemma 1, the geometric condition $|A|^2 \langle x, N \rangle^2 \leq 2$ on Σ is equivalent to $\text{Hess}_\Sigma f \geq 0$ on Σ . Hence, $(f \circ \gamma)''(t) \geq 0$ for all $t \in [0, 1]$, that is, $f \circ \gamma$ is convex on $[0, 1]$. By the definition of \mathcal{C} , as $f \circ \gamma$ attains its minimum at $t = 0$ and $t = 1$ we conclude that $(f \circ \gamma)(t) \equiv \min_\Sigma f$. Therefore $\gamma([0, 1]) \subset \mathcal{C}$ and the claim follows.

Since Σ meets ∂B orthogonally, we have that $\nu(x) = x$ for all points x in $\partial\Sigma$, where ν denotes the unit outward pointing conormal of Σ along $\partial\Sigma$. Hence, the geodesic curvature k_g of $\partial\Sigma$ in Σ satisfies $k_g = \langle \nabla_T^\Sigma \nu, T \rangle = \langle T, T \rangle = 1$ (here T denotes the a unit vector tangent to $\partial\Sigma$). In particular, $\partial\Sigma$ is strictly convex in Σ . This implies that for all $p, q \in \Sigma$ there exists a minimising geodesic in Σ joining p to q . Thus, the totally convex set \mathcal{C} is connected. Also, \mathcal{C} is clearly contained in the interior of Σ , because $\min_{x \in \Sigma} |x| < 1$ and points in $\partial\Sigma$ belong to the unit sphere ∂B .

Now, let us suppose that \mathcal{C} contains only one point $p \in \Sigma \setminus \partial\Sigma$. Given $[\alpha] \in \pi_1(\Sigma, p)$, let us assume that $[\alpha]$ is a non-trivial homotopy class. Since $\partial\Sigma$ is strictly convex, we can find a non-trivial geodesic loop $\gamma : [0, 1] \rightarrow \Sigma$, $\gamma(0) = \gamma(1) = p$, representing the class $[\alpha]$. Since \mathcal{C} is totally convex, $\gamma([0, 1]) \subset \mathcal{C}$. But $\mathcal{C} = \{p\}$, which is a contradiction. Therefore $\pi_1(\Sigma, p)$ is trivial, that is, Σ is homeomorphic to a disk. By

Nitsche's Theorem [15], Σ is a flat equatorial disk.

Finally, suppose that \mathcal{C} does not consist of a single point. By Nitsche's result, Σ cannot be homeomorphic to a disk. Let $p \in \mathcal{C}$ and let $[\alpha] \in \pi_1(\Sigma, p)$ be a non-trivial homotopy class. As above, since $\partial\Sigma$ is strictly convex, we can find a non-trivial geodesic loop $\gamma : [0, 1] \rightarrow \Sigma$, $\gamma(0) = \gamma(1) = p$, representing $[\alpha]$. We claim that $\gamma'(0) = \gamma'(1)$, $\gamma([0, 1])$ is a simple curve (has no self-intersections) and $\mathcal{C} = \gamma([0, 1])$. In fact, if $\gamma'(0) \neq \gamma'(1)$, then, because \mathcal{C} is totally convex, it is possible to join points on γ near the break at p by minimising geodesics and find an open set $U \subset \mathcal{C}$, which is a contradiction since Σ is minimal. By a similar reasoning, $\gamma([0, 1])$ has to be a simple curve and $\mathcal{C} = \gamma([0, 1])$, so that the set \mathcal{C} is a simple closed geodesic. Since the same argument shows that any non-trivial geodesic loop based at p must be contained in \mathcal{C} , we conclude that $\pi_1(\Sigma, p) = \mathbb{Z}$. By the topological classification of compact orientable surfaces, Σ is homeomorphic to an annulus. \square

The following corollary is a direct consequence of the above proof.

Corollary 3. *Let Σ^2 be a compact free boundary minimal surface in B^3 . If $|A|^2\langle x, N \rangle^2 < 2$ on Σ , then Σ is a flat equatorial disk.*

Proof. Since $|A|^2\langle x, N \rangle^2 < 2$, by item *ii*) of Lemma 1 it follows that the function f defined in Proposition 1 is strictly convex, that is, $\text{Hess}_\Sigma f > 0$ on Σ . Clearly, f attains its minimum value on the interior of Σ . By the strict convexity of f , the set of minima contains a single point. Thus, Proposition 2 implies that Σ is a flat equatorial disk. \square

In order to finish the proof of our main result, it remains only to analyse the situation where the function $|A|^2\langle x, N \rangle^2$ attains the maximum value 2 at some point on Σ .

Proposition 4. *Let Σ^2 be a compact free boundary minimal surface in B^3 . If $|A|^2\langle x, N \rangle^2 \leq 2$ on Σ and $|A|^2(p)\langle p, N(p) \rangle^2 = 2$ at some point $p \in \Sigma$, then Σ is a critical catenoid.*

Proof. Let $f : \Sigma \rightarrow \mathbb{R}$ and \mathcal{C} be defined as in Lemma 1 and Proposition 2, respectively. Since $|A|^2(p)\langle p, N(p) \rangle^2 = 2$ at some point $p \in \Sigma$, the surface Σ cannot be homeomorphic to a disk, for in this case it would be totally geodesic as a consequence of Nitsche's Theorem [15]. Thus, by Proposition 2, Σ is homeomorphic to an annulus and \mathcal{C} is a simple closed geodesic, say $\gamma : [0, \ell] \rightarrow \Sigma$, parametrized by arc length, in $\Sigma \setminus \partial\Sigma$. In particular, $\inf_\Sigma f > 0$.

Let $R > 0$ be defined by $R^2 = 2 \inf_\Sigma f$ and let $S_R \subset \mathbb{R}^3$ be the sphere of radius R centred at the origin. We claim that γ is a great circle in S_R . In fact, by definition of R , we have that $\Sigma \subset \{x \in \mathbb{R}^3 : |x| \geq R\}$

and $\Sigma \cap S_R = \gamma([0, \ell])$. This implies that $T_{\gamma(s)}\Sigma = T_{\gamma(s)}S_R$ for all $s \in [0, \ell]$. Since γ is a geodesic of Σ , we then conclude that γ is also a geodesic of S_R , that is, γ is a great circle of S_R .

If e_3 is a unit vector in \mathbb{R}^3 that is orthogonal to the plane containing the great circle γ , then $\{\gamma'(s), e_3\}$ is an orthonormal basis of $T_{\gamma(s)}\Sigma$ for all $s \in [0, \ell]$.

Let $u(x) = \langle x \wedge N(x), e_3 \rangle$, $x \in \Sigma$, where \wedge denotes the cross product in \mathbb{R}^3 . The function u is the infinitesimal normal speed of the variation of Σ by rotations around the e_3 -axis. As such, u is a Jacobi function of Σ , that is,

$$\Delta_\Sigma u + |A|^2 u = 0 \quad \text{on } \Sigma, \quad \text{and} \quad \frac{\partial u}{\partial \nu} = u \quad \text{on } \partial \Sigma.$$

Moreover, u vanishes identically if and only if Σ is a surface of revolution around the e_3 -axis.

Clearly, the circle $\gamma([0, \ell])$ is contained in $u^{-1}(0)$, which immediately implies that $du(\gamma(s))\gamma'(s) = 0$ for all $s \in [0, \ell]$. On the other hand, as $\gamma'(s)$ is a principal direction of Σ at $\gamma(s)$, so it is the orthogonal direction $e_3 \in T_{\gamma(s)}\Sigma$. Thus, $dN_{\gamma(s)}e_3$ is parallel to e_3 and

$$du(\gamma(s))e_3 = \langle e_3 \wedge N(\gamma(s)), e_3 \rangle + \langle \gamma(s) \wedge dN_{\gamma(s)}e_3, e_3 \rangle = 0.$$

The above argument has shown that every point in the circle \mathcal{C} , which is contained in $u^{-1}(0)$, is a critical point of u . But this implies that u vanishes identically on Σ . In fact, should this not be true, the nodal set $u^{-1}(0)$ of the Jacobi function u would contain only isolated critical points, by a result due to S. Y. Cheng ([3], Theorem 2.5). And this would be a contradiction.

Thus, Σ is an free boundary minimal annulus of revolution around the e_3 -axis in B^3 . In other words, Σ is a critical catenoid. \square

Remark 5. As pointed out by one of the referees, the final step of the proof also follows from the solution to the Björling problem, which states that two minimal surfaces in \mathbb{R}^3 containing a regular real analytic curve and having the same unit normal vector fields along that curve must be the same (see, for example, [5], Chapter 3, Section 4). On the final steps of our proof, we have shown that, possibly after a rotation, the free boundary minimal surface Σ contains the great circle $\mathcal{C} = \{(x, y, 0); x^2 + y^2 = R\}$ and its normal unit vector field along \mathcal{C} is equal to $(x/R, y/R, 0)$. Since the same conditions are satisfied by an appropriately chosen vertical catenoid, both surfaces are equal. In particular, Σ is rotational and the result follows.

Acknowledgements: L. A. was supported by the ERC Start Grant PSC and LMCF 278940 and I. N. was supported by CNPq-Brazil and

FAPEMA/CNPq. Both authors are grateful to André Neves, Celso Viana, Rafael Montezuma, Fernando Codá Marques, Ben Sharp and Alessandro Carlotto for their kind interest in this work, and to the referees for their comments.

REFERENCES

- [1] S. Brendle, *A sharp bound for the area of minimal surfaces in the unit ball*, Geom. Funct. Anal. **22** (2012), no. 3, 621-626.
- [2] J. Cheeger and D. Gromoll, *On the structure of complete manifolds of nonnegative curvature*, Ann. of Math. (2) **96** (1972), 413-443.
- [3] S. Y. Cheng, *Eigenfunctions and Nodal Sets*, Comment. Math. Helvetici **51** (1976), 43-55.
- [4] S.S. Chern, M. do Carmo, S. Kobayashi, *Minimal submanifolds of a sphere with second fundamental form of constant length*, in Functional Analysis and Related Fields (Proc. Conf. for M. Stone, Univ. Chicago, Chicago, Ill., 1968), Springer, New York, 1970, pp. 59-75.
- [5] U. Dierkes, S. Hildebrandt, F. Sauvigny, *Minimal Surfaces*, Grundlehren der Mathematischen Wissenschaften (339), Springer, Heidelberg, 2010.
- [6] A. Fraser and M. Li, *Compactness of the space of embedded minimal surfaces with free boundary in three-manifolds with nonnegative Ricci curvature and convex boundary*, J. Differential Geom. **96** (2014), no. 2, 183-200.
- [7] A. Fraser and R. Schoen, *Sharp eigenvalue bounds and minimal surfaces in the ball*, Invent. Math. **203** (2016), no. 3, 823-890.
- [8] A. Fraser and R. Schoen, *Uniqueness theorems for free boundary minimal disks in space forms*, Int. Math. Res. Not. IMRN (2015), no. 17, 8268-8274.
- [9] A. Fraser and R. Schoen, *The first Steklov eigenvalue, conformal geometry, and minimal surfaces*, Adv. Math. **226** (2011), 4011-4030.
- [10] B. Lawson, *Local rigidity theorems for minimal hypersurfaces*, Ann. of Math. **89** (1969), 187-197.
- [11] M. Li and N. Kapouleas, *Free boundary minimal surfaces in the unit three-ball via desingularization of the critical catenoid and the equatorial disk*, arXiv:1709.08556.
- [12] D. Ketover, *Free boundary minimal surfaces of unbounded genus*, arXiv:1612.08691.
- [13] H. Li and C. Xiong, *A gap theorem for free boundary minimal surfaces in geodesic balls of hyperbolic space and hemisphere*, to appear in the J. Geom Anal, published on-line (2017).
- [14] P. McGrath, *A characterization of the critical catenoid*, arxiv:1603.04114.
- [15] J. C. C. Nitsche, *Stationary partitioning of convex bodies*, Arch. Rational Mech. Anal. **89** (1985), 1-19.
- [16] F. Pacard, A. Folha and T. Zolotareva, *Free boundary minimal surfaces in the unit 3-ball*, Manuscripta Math. **154** (2017), no. 3-4, 359-409.
- [17] A. Ros and E. Vergasta, *Stability for hypersurfaces of constant mean curvature with free boundary*, Geom. Dedicata **56** (1995), no. 1, 19-33.
- [18] J. Simons, *Minimal varieties in riemannian manifolds*, Ann. of Math. (2) **88** (1968) 62-105.

L. AMBROZIO: MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, UK.

E-mail address: `l.coelho-ambrozio@warwick.ac.uk`

I. NUNES: DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO
MARANHÃO, CAMPUS DO BACANGA, SÃO LUÍS, MA, BRASIL.

E-mail address: `ivaldo.nunes@ufma.br`