

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/108614>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

© 2018 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>.



Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

The step Sidorenko property and non-norming edge-transitive graphs*

Daniel Král'[†] Taísa L. Martins[‡] Péter Pál Pach[§]
Marcin Wrochna[¶]

Abstract

Sidorenko's Conjecture asserts that every bipartite graph H has the Sidorenko property, i.e., a quasirandom graph minimizes the density of H among all graphs with the same edge density. We study a stronger property, which requires that a quasirandom multipartite graph minimizes the density of H among all graphs with the same edge densities between its parts; this property is called the step Sidorenko property. We show that many bipartite graphs fail to have the step Sidorenko property and use our results to show the existence of a bipartite edge-transitive graph that is not weakly norming; this answers a question of Hatami [Israel J. Math. 175 (2010), 125–150].

*The work of the first three authors has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 648509). This publication reflects only its authors' view; the European Research Council Executive Agency is not responsible for any use that may be made of the information it contains.

[†]Faculty of Informatics, Masaryk University, Botanická 68A, 602 00 Brno, Czech Republic, and Mathematics Institute, DIMAP and Department of Computer Science, University of Warwick, Coventry CV4 7AL, UK. E-mail: dkral@fi.muni.cz. The first author was also supported by the Engineering and Physical Sciences Research Council Standard Grant number EP/M025365/1.

[‡]Mathematics Institute and DIMAP, University of Warwick, Coventry CV4 7AL, UK. E-mail: t.lopes-martins@warwick.ac.uk. This author was also supported by the CNPq Science Without Borders grant number 200932/2014-4.

[§]Department of Computer Science and DIMAP, University of Warwick, Coventry CV4 7AL, UK and Department of Computer Science and Information Theory, Budapest University of Technology and Economics, 1117 Budapest, Magyar tudósok körútja 2., Hungary. E-mail: ppp@cs.bme.hu. This author was also supported by the National Research, Development and Innovation Office NKFIH (Grant Nr. K124171).

[¶]Institute of Informatics, University of Warsaw, Poland. E-mail: m.wrochna@mimuw.edu.pl. This author was supported by the National Science Centre of Poland grant number 2016/21/N/ST6/00475 and by the Foundation for Polish Science (FNP) via the START stipend programme. The visit of this author to the University Warwick was supported by the Leverhulme Trust 2014 Philip Leverhulme Prize of the first author.

1 Introduction

Sidorenko's Conjecture is one of the most important open problems in extremal graph theory. A graph H has the *Sidorenko property* if a quasirandom graph minimizes the density of H among all graphs with the same edge density. The beautiful conjecture of Erdős and Simonovits [21] and of Sidorenko [20] asserts that every bipartite graph has the Sidorenko property (it is easy to see that non-bipartite graphs fail to have the property). In this paper, we consider a more general property, the step Sidorenko property, and explore the link between this property and weakly norming graphs to show the existence of a bipartite edge-transitive graph that is not weakly norming. This answers a question of Hatami [13] whether such graphs exist.

Sidorenko's Conjecture is one of the central problems in extremal combinatorics. Sidorenko [20] confirmed the conjecture for trees, cycles and bipartite graphs with one of the sides having at most three vertices; the case of paths is equivalent to the Blakley-Roy inequality for matrices, which was proven in [1]. Additional graphs, such as cubes and bipartite graphs with a vertex complete to the other part, were added to the list of graphs with the Sidorenko property by Conlon, Fox and Sudakov [3], by Hatami [13], and by Szegedy [24]. Recursively described classes of bipartite graphs that have the Sidorenko property were obtained by Conlon, Kim, Lee and Lee [4], by Kim, Lee and Lee [16], by Li and Szegedy [17] and by Szegedy [23]. In particular, Szegedy [23] has described a class of graphs called thick graphs, which are amenable to showing the Sidorenko property using the entropy method argument that he developed. More recently, Conlon and Lee [6] showed that bipartite graphs such that one of the parts has many vertices of maximum degree have the Sidorenko property. Sidorenko's Conjecture is also known to hold in the local sense [18, Proposition 16.27], i.e., a small modification of a quasirandom graph preserving its edge density does not decrease the number of copies of any bipartite graph. A stronger statement of this type, which comes with uniform quantitative bounds, has recently been proven by Fox and Wei [9].

Sidorenko's Conjecture is also related to other well-studied problems in graph theory. We would like to particularly mention the connection to quasirandom graphs. We say that a graph H is *forcing* if all minimizers of the density of H among graphs with the same edge density are quasirandom graphs. Note that if H is forcing, then H has the Sidorenko property. The classical result of Thomason [25], also see [2], says that the cycle of length four is forcing. This result was generalized by Chung, Graham and Wilson [2], who showed that every complete bipartite graph $K_{2,n}$ is forcing, and by Skokan and Thoma [22], who showed that all complete bipartite graphs are forcing. A characterization of forcing graphs was stated as a question by Skokan and Thoma [22] and conjectured by Conlon, Fox and Sudakov [3]: a graph H is forcing if and only if H is bipartite and contains a cycle.

Another graph theoretic notion related to Sidorenko’s Conjecture is that of common graphs. A graph H is *common* if a quasirandom graph minimizes the sum of densities of H and the complement of H . An old theorem of Goodman [12] says that the complete graph K_3 is common. The conjecture of Erdős that the complete graph K_4 is also common was disproved by an ingenious construction of Thomason [26]; counterexamples with a simpler structure were found by Franek and Rödl in [11]. Jagger, Šťovíček and Thomason [15] showed that no graph containing K_4 is common. On the other hand, it is known that the graph obtained from K_4 by removing an edge [10] is common and so is the wheel W_5 [14]. The classification of common graphs remains a wide open problem.

Our results are motivated by the relation of Sidorenko’s Conjecture to *weakly norming* graphs, which are of substantial interest in the theory of graph limits. Due to its technical nature, we defer the definition to Section 2. Intuitively, these are graphs H such that the density of H in other graphs defines a norm on the space of graphons (graph limits). Chapter 14.1 in Lovász’ book [18] gives an introduction to this notion. Every weakly norming graph has the Sidorenko property [13]. However, every weakly norming graph also has a stronger property [18, Proposition 14.13], which we call the *step Sidorenko property*. Informally speaking, a graph H has the step Sidorenko property if a multipartite quasirandom graph minimizes the density of H among all multipartite graphs with the same density inside and between its parts; we give a formal definition in Section 2. It is not hard to find a graph that has the Sidorenko property but not the step Sidorenko property; the cycle of length four with an added pendant edge is an example (see Section 2).

In this paper, we present techniques for showing that a bipartite graph fails to have the step Sidorenko property. Our techniques allow us to show that graphs as simple and symmetric as toroidal grids, i.e., Cartesian products of any number of cycles, do not have the step Sidorenko property. The only exceptions are hypercubes (and single cycles of even length), which were shown to be weakly norming by Hatami [13] (see also [18, Proposition 14.2] for a concise presentation). The fact that most of the toroidal grids are not weakly norming is surprising when contrasted with the result of Conlon and Lee [5] that the incidence graphs of regular polytopes are weakly norming. Since toroidal grids $C_n \square C_n$ are edge-transitive, this answers in the negative a question of Hatami [13] whether all edge-transitive bipartite graphs are weakly norming.

2 Preliminaries

In this section, we introduce the notation that is used throughout the paper. In general, we follow standard graph theory notation. All graphs considered in this paper are simple and without loops. We sometimes consider graphs with vertices and edges assigned non-negative weights; when this is the case, we refer

to such a graph as a *weighted graph*. The order of a graph G , i.e., its number of vertices, will be denoted by $|G|$ and the size of a graph G , i.e., its number of edges, by $\|G\|$. If v and w are two vertices of G , then $\text{dist}(v, w)$ is the distance between v and w , i.e., the number of edges of the shortest path from v to w . The *Cartesian product* of graphs G_1, \dots, G_k , denoted $G_1 \square \dots \square G_k$, is the graph with vertex set equal to the Cartesian product of the vertex sets of G_1, \dots, G_k , where two vertices (u_1, \dots, u_k) and (v_1, \dots, v_k) are adjacent if there exists $1 \leq i_0 \leq k$ such that $u_{i_0}v_{i_0}$ is an edge of G_{i_0} and $u_i = v_i$ for all $i \neq i_0$.

In the rest of this section, we introduce notation related to graph homomorphisms and present notions from the theory of graph limits that we need for our exposition. We also formally define the Sidorenko property, the step Sidorenko property and weakly norming graphs.

2.1 Graph homomorphisms

A *homomorphism* from a graph H to a graph G is a mapping f from $V(H)$ to $V(G)$ such that if vv' is an edge of H , then $f(v)f(v')$ is an edge of G . If f is a homomorphism from H to G , $|f^{-1}(X)|$ for $X \subseteq V(G)$ denotes the number of vertices of H mapped to a vertex in X and $|f^{-1}(X)|$ for $X \subseteq E(G)$ denotes the number of edges mapped to an edge in X ; for simplicity, we write $|f^{-1}(x)|$ instead of $|f^{-1}(\{x\})|$.

We will need to consider homomorphisms extending a partial mapping between vertices of H and G and we now introduce notation that will be helpful in this setting. We write $H(v_1, \dots, v_k)$ for a graph H with k distinguished vertices v_1, \dots, v_k . If $H(v_1, \dots, v_k)$ and $G(v'_1, \dots, v'_k)$ are two graphs with k distinguished vertices, then a homomorphism from $H(v_1, \dots, v_k)$ to $G(v'_1, \dots, v'_k)$ is a homomorphism from H to G that maps v_i to v'_i for $i = 1, \dots, k$.

We will also consider homomorphisms to graphs with vertex and edge weights. As given earlier, a *weighted graph* is a graph G where each vertex and each edge of G is assigned a non-negative weight; the mapping w from $V(G) \cup E(G)$ assigning the weights will be referred to as a *weight function* of G . The *weight of a homomorphism* f from H to a weighted graph G , denoted $w(f)$, is defined as

$$\prod_{v \in V(H)} w(f(v)) \prod_{vv' \in E(H)} w(f(v)f(v')) = \prod_{v \in V(G)} w(v)^{|f^{-1}(v)|} \prod_{e \in E(G)} w(e)^{|f^{-1}(e)|}.$$

We will often speak about the sum of the weights of homomorphisms from a graph $H(v_1, \dots, v_k)$ to a weighted graph $G(v'_1, \dots, v'_k)$; this sum will be denoted by $\text{hom}(H(v_1, \dots, v_k), G(v'_1, \dots, v'_k))$ and we will understand it to be zero if no such homomorphism exists.

We also use the just introduced notation for graphs with distinguished vertices when talking about blow-ups of graphs. A *k-blow-up* of a graph $H(v)$ is the graph obtained from H by replacing the vertex v with k new vertices, which we refer

to as *clones* of v . The vertices different from v preserve their adjacencies, the clones of v form an independent set and each of them is adjacent precisely to the neighbors of v . Observe that if H is a weighted graph, then if the edges of the k -blow-up of $H(v)$ have the same weight as in H , the vertices of the k -blow-up except for the clones have the same weights as in H and each clone has weight equal to $1/k$ of the weight of v , then the sum of the weights of homomorphisms from G to H and the sum of the weights of homomorphisms from G to the k -blow-up are the same for every graph G .

2.2 Graph limits

The theory of graph limits offers analytic tools to study large graphs. We present here only those notions that we need further, and refer the reader to the monograph of Lovász [18] on the subject for a comprehensive introduction to the theory.

Let $t(H, G)$ be the normalized number of homomorphisms from a graph H to a graph G , i.e., $t(H, G) = \text{hom}(H, G) / |V(G)|^{|V(H)|}$ where G in $\text{hom}(H, G)$ is understood to have all the vertex and edge weights equal to one. A sequence $(G_n)_{n \in \mathbb{N}}$ of graphs is *convergent* if the sequence $t(H, G_n)$ converges for every graph H . A convergent sequence of graphs can be represented by an analytic object called a graphon. A *graphon* is a (Borel) measurable symmetric function W from $[0, 1]^2$ to $[0, 1]$, i.e., $W(x, y) = W(y, x)$ for all $(x, y) \in [0, 1]^2$. One can think (although very imprecisely) of a graphon as a continuous version of the adjacency matrix of a graph. Led by this intuition, we can define the density of a graph H in a graphon W as

$$t(H, W) = \int_{[0,1]^{V(H)}} \prod_{vv' \in E(H)} W(x_v, x_{v'}) \, dx^{V(H)} .$$

Note that the definition of $t(H, W)$ does not require W to be non-negative and we can define $t(H, f)$ in the same way for any bounded measurable function $f : [0, 1]^2 \rightarrow \mathbb{R}$.

We say that a graphon W is a *limit* of a convergent sequence $(G_n)_{n \in \mathbb{N}}$ of graphs if $t(H, W)$ is the limit density of $t(H, G_n)$ for every graph H . It is not hard to show that for every graphon W , there exists a convergent sequence of graphs such that W is its limit. The converse statement is also true as shown by Lovász and Szegedy [19], i.e., for every convergent sequence of graphs, there exists a graphon that is its limit.

The density $t(K_2, W)$ of K_2 is equal to the L_1 -norm of a graphon W as a function from $[0, 1]^2$. This leads to the question which graphs H can be used to define a norm on the space of measurable functions on $[0, 1]^2$ or, more restrictively, on the space of graphons. That is, we say that a graph H is *weakly norming* if the function $\|W\|_H = t(H, W)^{1/\|H\|}$ is a norm on the space of graphons, i.e., $\|W\|_H = 0$ if and only if W is equal to zero almost-everywhere and the triangle

inequality $\|W_1 + W_2\|_H \leq \|W_1\|_H + \|W_2\|_H$ holds for any two graphons W_1 and W_2 . Observe that H is weakly norming if and only if $\| |f| \|_H$ is a norm on the set of all bounded symmetric functions f from $[0, 1]^2$ to \mathbb{R} (if we required that $\|f\|_H$, without the absolute value, is a norm on all such functions, we would get the slightly stronger notion of *norming graphs*).

It is not hard to show that every weakly norming graph must be bipartite. Hatami [13] showed stronger statements as corollaries of his characterization of weakly norming graphs as those satisfying a certain Hölder type inequality. First, every weakly norming graph H must be biregular, i.e., all vertices in the same part of its bipartition have the same degree. Second, every subgraph H' of a connected weakly norming graph H must satisfy that

$$\frac{\|H'\|}{|H'| - 1} \leq \frac{\|H\|}{|H| - 1} .$$

Weakly norming graphs include complete bipartite graphs (in particular, stars), even cycles and hypercubes [13]; later, Conlon and Lee [5] presented a large class of weakly norming graphs, which they refer to as reflection graphs.

Every weighted graph G with a weight function w that assigns edges weights between 0 and 1 can be associated with a graphon W_G as follows. Each vertex v of G is associated with a measurable set J_v with measure $w(v)/w(V(G))$ in such a way that the sets J_v , $v \in V(G)$, form a partition of the interval $[0, 1]$; $w(V(G))$ denotes the sum of the weights of the vertices of G . For $x \in J_v$ and $y \in J_{v'}$, we set $W(x, y) = w(vv')$ if $vv' \in E(G)$ and $W(x, y) = 0$ otherwise (in particular, we set $W(x, y) = 0$ if $v = v'$). It is not hard to observe that $\text{hom}(H, G)$ is equal to $t(H, W_G) \cdot w(V(G))^{|H|}$; in particular, if the sum of the weights of vertices of G is one, then $\text{hom}(H, G) = t(H, W_G)$. This correspondence will allow us to study weakly norming graphs in terms of weighted homomorphisms.

2.3 Step Sidorenko property

We now use the language of graph limits to describe the Sidorenko property and to formally define the step Sidorenko property. A graph H has the *Sidorenko property* if

$$t(K_2, W)^{\|H\|} \leq t(H, W) \tag{1}$$

for every graphon W . The left hand side can also be written as $t(H, U_p)$, where $U_p \equiv p$ is the constant graphon with the same edge density $p = t(K_2, W)$ as W . A graph H is *forcing* if it has the Sidorenko property and (1) holds with equality only if W is equal to some $p \in [0, 1]$ almost everywhere. As we have presented earlier, Sidorenko's Conjecture asserts that every bipartite graph has the Sidorenko property and the Forcing Conjecture asserts that every bipartite graph with a cycle is forcing.

Let \mathcal{P} be a partition of the interval $[0, 1]$ into finitely many non-null measurable sets. We now define the *stepping operator*. If W is a graphon, then the graphon $W^{\mathcal{P}}$ is defined for $(x, y) \in [0, 1]^2$ as the ‘step-wise average’:

$$W^{\mathcal{P}}(x, y) = \frac{1}{|J||J'|} \int_{J \times J'} W(s, t) \, ds \, dt$$

where J and J' are the unique parts from \mathcal{P} such that $x \in J$ and $y \in J'$, and $|X|$ denotes the measure of a measurable subset $X \subseteq [0, 1]$. Note that the graphon $W^{\mathcal{P}}$ is constant on $J \times J'$ for any $J, J' \in \mathcal{P}$, i.e., the graphon $W^{\mathcal{P}}$ is a step graphon.

Let \mathcal{P}_0 be the partition with a single part being the interval $[0, 1]$ itself. A graph H has the Sidorenko property if and only if $t(H, W^{\mathcal{P}_0}) \leq t(H, W)$ for every graphon W . This motivates the following definition. A graph H has the *step Sidorenko property* if and only if

$$t(H, W^{\mathcal{P}}) \leq t(H, W)$$

for every graphon W and every partition \mathcal{P} of $[0, 1]$ into finitely many non-null measurable sets. Since all weakly norming graphs [18, Proposition 14.13] have the step Sidorenko property, it follows that complete bipartite graphs, even cycles, hypercubes and more generally reflection graphs defined by Conlon and Lee [5] all have the step Sidorenko property.

The definition of the step Sidorenko property yields that every graph that has the step Sidorenko property also has the Sidorenko property. However, the converse is not true in general as we now demonstrate. Let C_4^+ be the 5-vertex graph obtained from a cycle of length four by adding a single vertex adjacent to one of the vertices of the cycle. The graph C_4^+ has the Sidorenko property because, e.g., it is a bipartite graph with a vertex complete to the other part [3]. On the other hand, C_4^+ does not have the step Sidorenko property. Consider the partition $\mathcal{P} = \{[0, \frac{2}{5}), [\frac{2}{5}, 1]\}$ and the graphon W that is defined as follows (the symmetric cases of (x, y) are omitted).

$$W(x, y) = \begin{cases} 0.9 & \text{if } (x, y) \in [0, \frac{1}{5}) \times [0, \frac{1}{5}), \\ 0.85 & \text{if } (x, y) \in [0, \frac{1}{5}) \times [\frac{1}{5}, \frac{2}{5}), \\ 0.2 & \text{if } (x, y) \in [0, \frac{1}{5}) \times [\frac{2}{5}, 1], \\ 1 & \text{if } (x, y) \in [\frac{1}{5}, \frac{2}{5}) \times [\frac{1}{5}, \frac{2}{5}), \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$

A straightforward computation yields that

$$\begin{aligned} t(C_4^+, W) &\simeq 0.007453 \text{ and} \\ t(C_4^+, W^{\mathcal{P}}) &\simeq 0.007508 > t(C_4^+, W). \end{aligned}$$

Hence, the graph C_4^+ does not have the step Sidorenko property.

3 Grids

In this section, we demonstrate our techniques from Section 4 in a less general setting. We believe that this makes our presentation more accessible.

Intuitively, we consider a graph G with distinguished vertices u_0, u_1, u_2 such that u_0u_1 and u_0u_2 are edges. The idea is to blow-up u_0 into two copies and slightly perturb weights only on edges corresponding to u_0u_1 and u_0u_2 , increasing weights of edges for one copy and decreasing it for the other proportionally to a parameter α , resulting in a weighted graph G_α . A partition \mathcal{P} on the corresponding graphon W_α is then defined so that the stepping operator averages out this perturbation, returning to the original graph: $W_\alpha^{\mathcal{P}} = W_G$. The difference in homomorphism densities $t(H, W_\alpha^{\mathcal{P}}) - t(H, W_\alpha)$ is then analyzed in the limit of small perturbations α : first order changes (those linear in α) cancel out. Second order changes result in a condition that can be expressed fairly concisely as positive semidefiniteness of a matrix whose entries count certain constrained homomorphisms.

The more powerful setting in Section 4 uses essentially the same idea, only blowing up more vertices, resulting in a larger matrix and allowing us to further constraint the homomorphisms we have to count. We turn to choosing the starting weighted graph G and interpreting these counts in later corollaries.

Theorem 1. *Let H be a graph and let G be a weighted graph with three distinguished vertices u_0, u_1 and u_2 such that u_0u_1 and u_0u_2 are edges. For $i, j \in \{1, 2\}$, let M_{ij} be the sum of the weights of homomorphisms from $H(v_0, v_1, v_2)$ to $G(u_0, u_i, u_j)$ summed over all choices of vertices v_0, v_1 and v_2 in H such that v_0v_1 and v_0v_2 are edges, i.e.,*

$$M_{ij} = \sum_{v_0v_1, v_0v_2 \in E(H)} \text{hom}(H(v_0, v_1, v_2), G(u_0, u_i, u_j)).$$

If the (2×2) -matrix M is not positive semidefinite, i.e., $M_{11}M_{22} < M_{12}^2$, then H does not have the step Sidorenko property.

Proof. Let w be the weight function of G . We assume that the sum of the weights of vertices of G is one (if needed, we multiply the weights of all vertices by the same constant). Consider the step graphon W_G associated with the weighted graph G . Let J_u be the measurable set corresponding to a vertex u of G and set $\mathcal{P} = \{J_u, u \in V(G)\}$.

Suppose that the matrix M associated with G is not positive semidefinite and fix a vector $a = (a_1, a_2)^T$ such that $a^T M a < 0$. We next define a weighted graph G_α with a parameter $\alpha \geq 0$ as follows. The graph G_α is a 2-blow-up of $G(u_0)$; let u_0^+ and u_0^- be the clones of u_0 . Each of the clones u_0^+ and u_0^- has weight $w(u_0)/2$. The weight of the edge $u_0^+u_i$ is $w(u_0u_i)(1 + \alpha a_i)$ and the weight of the edge $u_0^-u_i$ is $w(u_0u_i)(1 - \alpha a_i)$, $i = 1, 2$. The remaining vertices and edges have

weights equal to their counterparts in G . Let W_α be the step graphon associated with the weighted graph G_α such that the set corresponding to a vertex $u \neq u_0$ is J_u and the sets corresponding to the vertices u_0^+ and u_0^- are subsets of J_{u_0} . Observe that $W_G = W_\alpha$ for $\alpha = 0$ and that $W_G = W_\alpha^P$ for any α .

Our aim is to show that $t(H, W_\alpha) < t(H, W_G)$ for some $\alpha \in (0, 1)$. To do so, we analyze the density $t(H, W_\alpha)$ as a function of α . Note that $t(H, W_\alpha)$ is actually a polynomial in α . We next wish to determine the coefficients c_1 and c_2 such that

$$t(H, W_\alpha) = t(H, W_G) + c_1\alpha + c_2\alpha^2 + O(\alpha^3). \quad (2)$$

The coefficient c_1 can be determined as follows:

$$c_1 = \sum_{v_0v_1 \in E(H)} a_1 \operatorname{hom}(H(v_0, v_1), G_0(u_0^+, u_1)) - a_1 \operatorname{hom}(H(v_0, v_1), G_0(u_0^-, u_1)) + \\ a_2 \operatorname{hom}(H(v_0, v_1), G_0(u_0^+, u_2)) - a_2 \operatorname{hom}(H(v_0, v_1), G_0(u_0^-, u_2)).$$

Since $\operatorname{hom}(H(v_0, v_1), G_0(u_0^+, u_i)) = \operatorname{hom}(H(v_0, v_1), G_0(u_0^-, u_i))$ for all edges $v_0v_1 \in E(G)$ and all $i \in \{1, 2\}$, we conclude that $c_1 = 0$.

We next analyze the coefficient c_2 . In this case, we need to count homomorphisms mapping two edges, say v_0v_1 and $v'_0v'_1$, of H to edges $u_0^+u_i$ and to $u_0^-u_i$ of G_0 , $i = 1, 2$. If $v_0 \neq v'_0$, then the contributions of the homomorphisms mapping the edge v_0v_1 to $u_0^+u_i$ and $u_0^-u_i$ have opposite signs and cancel out. Hence, we obtain the following formula for c_2 :

$$c_2 = \sum_{v_0v_1, v_0v_2 \in E(H)} \sum_{i,j=1}^2 a_i a_j (\operatorname{hom}(H(v_0, v_1, v_2), G_0(u_0^+, u_i, u_j)) + \\ \operatorname{hom}(H(v_0, v_1, v_2), G_0(u_0^-, u_i, u_j))).$$

The definition of the matrix M now yields that

$$c_2 = \sum_{i,j=1}^2 a_i a_j \cdot M_{ij} = a^T M a < 0.$$

Since $c_1 = 0$ and $c_2 < 0$, we conclude using $W_G = W_\alpha^P$ and (2) that $t(H, W_\alpha) < t(H, W_G)$ for small enough $\alpha > 0$. It follows that the graph H does not have the step Sidorenko property. \square

The setting of Theorem 1 is sufficient to prove that the only two-dimensional toroidal grid that is weakly norming is $C_4 \square C_4$ (note that the toroidal grids $C_\ell \square C_\ell$ with ℓ odd are not Sidorenko, and hence also not weakly norming, because they are not bipartite).

We apply Theorem 1 with $G = H = C_\ell \square C_\ell$. The identity homomorphism contributes to the off-diagonal entry of the matrix from Theorem 1 while the




Figure 1: Notation used in the proof of Corollary 2. The edges b_1 , b_2 , b_3 and b_4 are drawn bold.

homomorphisms contributing to the diagonal entries have to “fold” two edges onto one. We choose weights in the target grid in such a way that the contribution of the former homomorphisms becomes smaller, which makes the matrix not to be positive semidefinite.

Corollary 2. *Let $\ell \geq 6$ be an even integer. The Cartesian product $C_\ell \square C_\ell$ does not have the step Sidorenko property.*

Proof. Fix $\ell \geq 6$ and let G and H be both equal to the graph $C_\ell \square C_\ell$; we denote the vertices of G and H by (i, j) , $0 \leq i, j \leq \ell - 1$, in such a way that two vertices are adjacent if they agree in one of the coordinates and differ by one in the other (all computations with the entries are computed modulo ℓ throughout the proof). Let u_0 be the vertex $(0, 0)$, u_1 the vertex $(1, 0)$ and u_2 the vertex $(0, 1)$. Further, let b_1 be the edge $(1, 0)(1, -1)$, b_2 the edge $(1, 0)(2, 0)$, b_3 the edge $(0, 1)(-1, 1)$ and b_4 the edge $(0, 1)(0, 2)$ (see Figure 1).

We next define the weights of the vertices and the edges of G ; to do so, we use a parameter $\gamma \in \mathbb{N}$, which will be fixed later. The weight $w(v)$ of a vertex v is $\gamma^{\text{dist}(u_0, v)}$ for $v \neq u_0, u_1, u_2$, $w(u_0) = \gamma^{-3}$ and $w(u_i) = \gamma^{\text{dist}(u_0, u_i) - 3} = \gamma^{-2}$, $i = 1, 2$. The weights of all edges of G are equal to one except for the edges b_1, b_2, b_3 and b_4 that have weight $\gamma^{-1/4}$.

We wish to apply Theorem 1 with the graphs H and G , and the distinguished vertices u_0, u_1 and u_2 . Instead of verifying that the matrix M from the statement of Theorem 1 is not positive semidefinite, we consider the matrix M such that

$$M_{ij} = \sum_{v_1, v_2 \in N_H(u_0)} \text{hom}(H(u_0, v_1, v_2), G(u_0, u_i, u_j)) .$$

Since H is vertex-transitive, the considered matrix M is positive semidefinite if and only if the matrix from the statement of Theorem 1 is. Observe that $M_{1,1} = M_{2,2}$ and $M_{1,2} = M_{2,1}$.

Consider a homomorphism f from $H(u_0, v_1, v_2)$ to $G(u_0, u_i, u_j)$ for some $i, j \in \{1, 2\}$. Observe that the weight of the homomorphism f is equal to

$$\sum_{\gamma \in V(H)} \text{dist}(u_0, f(v)) - 3|f^{-1}(\{u_0, u_1, u_2\})| - \frac{1}{4}|f^{-1}(\{b_1, b_2, b_3, b_4\})|$$

Note that if f is the identity, then the weight of f is equal to γ^W where

$$W = \sum_{v \in V(H)} \text{dist}(u_0, v) - 10.$$

Since the identity is a homomorphism from $H(u_0, u_i, u_j)$ to $G(u_0, u_i, u_j)$ for $i \neq j$, it follows that the entries $M_{1,2}$ and $M_{2,1}$ are of order $\Omega(\gamma^W)$, as functions of γ . We next show that both $M_{1,1}$ and $M_{2,2}$ are of order $o(\gamma^W)$. Since $M_{1,1} = M_{2,2}$, it is enough to argue that that $M_{1,1} = o(\gamma^W)$.

We show that every homomorphism f from $H(u_0, v_1, v_2)$ to $G(u_0, u_1, u_1)$ has weight at most $\gamma^{W-\frac{1}{2}}$; this will imply that $M_{1,1} = o(\gamma^W)$. Fix a homomorphism f from $H(u_0, v_1, v_2)$ to $G(u_0, u_1, u_1)$ with weight at least γ^W . By symmetry, we may assume that $v_1 = (1, 0)$ and $v_2 \in \{(-1, 0), (0, 1)\}$. Note that $|f^{-1}(\{u_0, u_1, u_2\})| \geq 3$. Since f is a homomorphism, any shortest path from u_0 to v is mapped by f to a walk of at most length $\text{dist}(u_0, v)$ from $f(u_0) = u_0$ to $f(v)$, it follows that $\text{dist}(u_0, f(v)) \leq \text{dist}(u_0, v)$ for every vertex v . Also observe that the parities of $\text{dist}(u_0, f(v))$ and $\text{dist}(u_0, v)$ are the same since the graph $G = H$ is bipartite. Since the weight of f is at least γ^W , the following holds: $|f^{-1}(\{u_0, u_1, u_2\})| = 3$, $\text{dist}(u_0, f(v)) = \text{dist}(u_0, v)$ for every vertex v of H and $|f^{-1}(\{b_1, b_2, b_3, b_4\})| \leq 4$. Since $|f^{-1}(\{u_0, u_1, u_2\})| = 3$, no vertex other than u_0, v_1 and v_2 is mapped by f to any of u_0, u_1 and u_2 ; in particular, no vertex is mapped to u_2 .

To finish the proof, we distinguish two cases based on whether $v_2 = (-1, 0)$ or $v_2 = (0, 1)$. We start with analyzing the case $v_2 = (-1, 0)$. Let $i \in \{1, 2\}$ and let v be a neighbor of v_i different from $(0, 0)$ and $v_i + v_i$. If $f(v) = (1, 1)$ or $f(v) = (2, 0)$, then the common neighbor of $(0, 0)$ and v different from v_i must be mapped to u_1 or u_2 , which is impossible. Hence, $f(v) = (1, -1)$. Since the choice of i and v was arbitrary, it follows that all the four edges $(1, 0)(1, 1)$, $(1, 0)(1, -1)$, $(-1, 0)(-1, 1)$ and $(-1, 0)(-1, -1)$ are mapped to the edge b_1 ; in particular, no other edge is mapped to b_1 or b_2 . This implies that the vertex $(2, 0)$ is mapped by f to $(1, 1)$. It follows that the vertex $(2, 1)$, which is a common neighbor of $(1, 1)$ and $(2, 0)$, must be mapped to the unique common neighbor $u_1 = (1, 0)$ of the vertices $f((1, 1)) = (1, -1)$ and $f((2, 0)) = (1, 1)$, which is impossible. This finishes the analysis of the case $v_2 = (-1, 0)$.

It remains to analyze the case that $v_2 = (0, 1)$. If the vertex $(1, -1)$ was mapped to $(2, 0)$ or $(1, 1)$, then the vertex $(0, -1)$, which is a common neighbor of

$(1, -1)$ and $(0, 0)$, would have to be mapped to $(1, 0)$ or $(0, 1)$, which is impossible. Hence, the vertex $(1, -1)$ is mapped by f to itself and the vertex $(0, -1)$ is also mapped to itself. Since swapping coordinates is a symmetry mapping v_1 and v_2 between each other, a symmetric argument yields that the vertex $(-1, 0)$ is mapped to $(0, -1)$.

Next, if the vertex $(2, 0)$ was mapped to the vertex $(1, 1)$, then the vertex $(2, -1)$, which is a common neighbor of $(2, 0)$ and $(1, -1)$, would have to be mapped to $(1, 0)$, which is impossible. It follows that the vertex $(2, 0)$ must be mapped to $(2, 0)$ or $(1, -1)$. We conclude that the edge b_1 is mapped to itself and the edge b_2 to either b_1 or b_2 . A symmetric argument yields that the edge b_3 is mapped to b_1 and the edge b_4 to b_1 or b_2 . In particular, no other edges of G are mapped to any of the edges b_1, b_2, b_3 and b_4 . This implies that the vertex $(1, 1)$ is mapped by f to itself. Consequently, the vertex $(2, 0)$ is also mapped to itself (otherwise, the vertex $(2, 1)$ would have to be mapped to $(1, 0)$).

We now prove the following statement for $r = 1, \dots, \ell/2 - 1$ by induction on r : all the vertices $(r, 1)$, $(r, -1)$ and $(r + 1, 0)$ are mapped by f to themselves. We have already established this statement for $r = 1$, so it remains to present the induction step. Fix $r = 2, \dots, \ell/2 - 1$ and assume that all the vertices $(r - 1, 1)$, $(r - 1, -1)$ and $(r, 0)$ are mapped to themselves. The vertex $(r, 1)$, which is a common neighbor of $(r - 1, 1)$ and $(r, 0)$, must be mapped to a common neighbor of $(r - 1, 1)$ and $(r, 0)$ at the distance $r + 1$ from $(0, 0)$. However, the only such vertex is $(r, 1)$. A symmetric argument yields that the vertex $(r, -1)$ is mapped to itself. Since the vertex $(r + 1, 0)$ must be mapped to a neighbor of $(r, 0)$ at distance $r + 1$ from $(0, 0)$, it can only be mapped to one of the vertices $(r, 1)$, $(r + 1, 0)$ and $(r, -1)$. By symmetry, it is enough to exclude that it is mapped to $(r, 1)$. If this was the case, then the vertex $(r + 1, -1)$, which is a common neighbor of $(r, -1)$ and $(r + 1, 0)$, must be mapped to $(r, 0)$, which is impossible. Hence, the vertex $(r + 1, 0)$ is mapped to itself, concluding the proof of the statement.

We have just shown that the vertex $(\ell/2, 0) = (-\ell/2, 0)$ is mapped to itself; earlier, we have shown that the vertex $(-1, 0)$ is mapped to $(0, -1)$. However, the path $(-1, 0)(-2, 0) \cdots (-\ell/2, 0)$ must be mapped by f to a walk with at most $\ell/2$ vertices but there is no such walk between the vertices $(0, -1)$ and $(-\ell/2, 0)$. Hence, there is no homomorphism from $H(u_0, v_1, v_2)$ to $G(u_0, u_1, u_1)$ with weight at least γ^W . \square

4 General Condition

We now present our general technique for establishing that certain graphs do not have the step Sidorenko property. One difference is that instead of considering only two neighbors of a distinguished vertex u_0 , we can choose any number of neighbors u_1, \dots, u_k , giving a larger matrix. More importantly, we are able to restrict homomorphisms considered in the statement to only those that map the

neighborhood of each u_i bijectively (to the neighborhood of the image of u_i , or a chosen subset of it).

The proof extends the arguments presented in the proof of Theorem 1. The main new idea is that by blowing up u_i , and appropriately choosing weights on copies of the edges to its neighbors, we can obtain an expression that is counting homomorphisms to the original graph, but with a weight that is an arbitrary function of how many neighbors of u_i map to each neighbor of the image of u_i . We choose this function to ensure that exactly one neighbor of u_i (or exactly zero) must map to each neighbor of its image.

Theorem 3. *Let H be a graph and let G be a weighted graph with $k + 1$ distinguished vertices u_0, u_1, \dots, u_k such that u_0u_1, \dots, u_0u_k are edges and u_1, \dots, u_k form an independent set. Further, let U_i , $i = 1, \dots, k$, be a subset of neighbors of u_i containing u_0 , and let M be the $(k \times k)$ -matrix such that the entry M_{ij} is the sum of the weights of homomorphisms from $H(v_0, v_1, v_2)$ to $G(u_0, u_i, u_j)$, where the sum runs over all choices of vertices v_0, v_1 and v_2 in H , such that the neighbors of v_1 are one-to-one mapped to U_i and the neighbors of v_2 to U_j . If the matrix M is not positive semidefinite, then H does not have the step Sidorenko property.*

Proof. Suppose that the matrix M is not positive semidefinite and fix a vector a such that $a^T M a < 0$. Let w be the weight function of G . As in the proof of Theorem 1, we assume that the sum of the weights of vertices of G is one. Similarly, we assume that the weight of each edge is at most $1/2$ (if needed, we can multiply the weights of all edges by the same constant).

We next define a weighted graph $G_{\varepsilon, \alpha}$, which is parameterized by $\varepsilon > 0$ and $\alpha \in \mathbb{R}$. The structure of the graph is independent of ε and α and is the following. Let n be the number of vertices of H . We consider the 3-blow-up of a vertex u_0 and $(n^{|U_i|-1} + 1)$ -blow-up of a vertex u_i . The three clones of u_0 will be denoted by u'_0, u_0^+ and u_0^- ; one of the $n^{|U_i|-1} + 1$ clones of u_i will be denoted by u'_i and the remaining ones by $u_{i, j_1, \dots, j_{|U_i|-1}}$ where $1 \leq j_1, \dots, j_{|U_i|-1} \leq n$. We next remove every edge going from the vertex $u_{i, j_1, \dots, j_{|U_i|-1}}$ to a vertex outside the set U_i that is not u_0^+ or u_0^- , i.e., the vertex $u_{i, j_1, \dots, j_{|U_i|-1}}$ is adjacent to u_0^+, u_0^- and the vertices of $U_i \setminus \{u_0\}$.

The weight of the vertex u'_0 is $(1 - 2\varepsilon)w(u_0)$ and the weight of each of the vertices u_0^+ and u_0^- is $\varepsilon w(u_0)$. The weight of the vertex u'_i is $(1 - n^{|U_i|-1}\varepsilon)w(u_i)$ and the weight of each of the vertices $u_{i, j_1, \dots, j_{|U_i|-1}}$ is $\varepsilon w(u_i)$. The remaining vertices of $G_{\varepsilon, \alpha}$ have the same weights as in G .

Before defining the weights of the edges, we define an auxiliary matrix B . The matrix B has n rows and n columns and $B_{ij} = 2^{(i-1)(j-1)}$. Note that B is a Vandermonde matrix. Since the matrix B is invertible, there exists a vector b such that $Bb = (0, 1, 0, \dots, 0)^T$. The weight of the edge between u_0^+ and $u_{i, j_1, \dots, j_{|U_i|-1}}$

is equal to

$$w(u_0 u_i) \left(1 + a_i \alpha \prod_{m=1}^{|U_i|-1} b_{j_m} \right),$$

and the weight of the edge between u_0^- and $u_{i,j_1,\dots,j_{|U_i|-1}}$ is equal to

$$w(u_0 u_i) \left(1 - a_i \alpha \prod_{m=1}^{|U_i|-1} b_{j_m} \right).$$

The weights of the edges incident with u'_0 and the remaining edges incident with u_0^+ and u_0^- are equal to the weights of their counterparts in G . Fix $i \in \{1, \dots, k\}$ and let $z_1, \dots, z_{|U_i|-1}$ be the vertices of U_i different from u_0 . The weight of the edge between the vertices $u_{i,j_1,\dots,j_{|U_i|-1}}$ and z_m is equal to $2^{j_m-1} w(u_i z_m)$. The weights of the edges incident with the vertex u'_i are the same as the weights of their counterparts in G . We have just defined the weights of all edges incident with at least one clone. The weights of the remaining edges are the same as in G .

We analyze $t(H, W_{\varepsilon,\alpha})$ as a function of α for $\alpha, \varepsilon \in (0, 1)$. In particular, we will show that

$$t(H, G_{\varepsilon,\alpha}) = t(H, G_{\varepsilon,0}) + c_\varepsilon \varepsilon^3 \alpha^2 + O(\varepsilon^4 \alpha^2) \quad (3)$$

for a coefficient c_ε , which we will estimate. Since the coefficient c_ε depends on ε , it is important to emphasize that the constants hidden in big O notation in (3) are independent of ε and α , i.e., the equality (3) represents that there exists $K > 0$, which is independent of ε , and a coefficient c_ε for every $\varepsilon \in (0, 1)$ such that the value of $t(H, G_{\varepsilon,\alpha})$ differs from $t(H, G_{\varepsilon,0}) + c_\varepsilon \varepsilon^3 \alpha^2$ by at most $K \varepsilon^4 \alpha^2$ for every $\alpha \in (0, 1)$.

We now proceed with analyzing the function $t(H, W_{\varepsilon,\alpha})$. As in the proof of Theorem 1, we observe that $t(H, W_{\varepsilon,\alpha})$ is a polynomial in α and the linear terms in α cancel out by pairing homomorphisms using u_0^+ and those using u_0^- . Hence, only quadratic and higher order terms remain. To estimate c_ε , we need to consider the terms corresponding to homomorphisms mapping exactly three vertices of H to the vertices of $G_{\varepsilon,\alpha}$ with weight ε and these vertices must induce a 2-edge path with the middle vertex mapped to u_0^+ or to u_0^- (the contribution of other homomorphisms cancels out by pairing those using u_0^+ and those using u_0^- , similarly as in the proof of Theorem 1). We arrive at the following identity.

$$\begin{aligned} c_\varepsilon \varepsilon^3 = & \sum_{v_0 v_1, v_0 v_2 \in E(H)} \sum_{i, i'=1}^k \sum_{j \in [n]^{|U_i|-1}} \sum_{j' \in [n]^{|U_{i'}|-1}} a_i a_{i'} \prod_{m=1}^{|U_i|-1} b_{j_m} \prod_{m=1}^{|U_{i'}|-1} b_{j'_m} \times \\ & \left(\text{hom}(H(v_0, v_1, v_2), G_{\varepsilon,0}(u_0^+, u_{i,j_1,\dots,j_{|U_i|-1}}, u_{i',j'_1,\dots,j'_{|U_{i'}|-1})) \right) + \\ & \left(\text{hom}(H(v_0, v_1, v_2), G_{\varepsilon,0}(u_0^-, u_{i,j_1,\dots,j_{|U_i|-1}}, u_{i',j'_1,\dots,j'_{|U_{i'}|-1})) \right) \end{aligned}$$

It follows that

$$\lim_{\varepsilon \rightarrow 0} c_\varepsilon = \sum_{\substack{v_0 v_1 \\ v_0 v_2} \in E(H)} \sum_{i, i'=1}^k \sum_h 2a_i a_{i'} w(h) \sum_{j \in [n]^{|U_i|-1}} \sum_{j' \in [n]^{|U_{i'}|-1}} \prod_{m=1}^{|U_i|-1} b_{j_m} 2^{(j_m-1)h(v_1 \leftrightarrow z_m)} \prod_{m=1}^{|U_{i'}|-1} b_{j'_m} 2^{(j'_m-1)h(v_2 \leftrightarrow z'_m)}$$

where the sum is taken over all homomorphisms h from H to G such that $h(v_0) = u_0$, $h(v_1) = u_i$ and $h(v_2) = u_{i'}$, and $w(h)$ denotes the weight of the homomorphism h , $h(v_1 \leftrightarrow z_m)$ denotes the number of neighbors of v_1 mapped to $z_m \in U_i$ and $h(v_2 \leftrightarrow z'_m)$ denotes the number of neighbors of v_2 mapped to $z'_m \in U_{i'}$. Observe that b was chosen so that the expression

$$\sum_{j_1, \dots, j_{|U_i|-1}=1}^n \prod_{m=1}^{|U_i|-1} b_{j_m} 2^{(j_m-1)h(v_1 \leftrightarrow z_m)} = \prod_{m=1}^{|U_i|-1} \sum_{j_m=1}^n b_{j_m} 2^{(j_m-1)h(v_1 \leftrightarrow z_m)}$$

is one if $h(v_1 \leftrightarrow z_m) = 1$ and it is zero otherwise. Hence, it follows that

$$\lim_{\varepsilon \rightarrow 0} c_\varepsilon = \sum_{v_0 v_1, v_0 v_2 \in E(H)} \sum_{i, i'=1}^k \sum_h a_i a_{i'} w(h)$$

where the sum is taken over homomorphisms h from H to G such that $h(v_0) = u_0$, $h(v_1) = u_i$, $h(v_2) = u_{i'}$, all neighbors of v_1 are one-to-one mapped to U_i and all neighbors of v_2 are one-to-one mapped to $U_{i'}$. The definition of the matrix M now implies that

$$\lim_{\varepsilon \rightarrow 0} c_\varepsilon = \sum_{i, i'=1}^k M_{ii'} a_i a_{i'} = a^T M a < 0. \quad (4)$$

The expressions (3) and (4) imply that there exist $\varepsilon > 0$ and $\alpha > 0$ such that $t(H, G_{\varepsilon, \alpha}) < t(H, G_{\varepsilon, 0})$. Fix such ε and α for the rest of the proof.

Consider the graphons W_0 and W_α associated with the weighted graphs $G_{\varepsilon, 0}$ and $G_{\varepsilon, \alpha}$, respectively. Let J_u be the measurable set corresponding to the vertex u of $G_{\varepsilon, 0}$; we can assume that the measurable set corresponding to the vertex u of $G_{\varepsilon, \alpha}$ is also J_u . Let \mathcal{P} be the partition of $[0, 1]$ formed by $J_{u_0^+} \cup J_{u_0^-}$ and J_u , $u \neq u_0^+, u_0^-$. Observe that $W_0 = W_\alpha^{\mathcal{P}}$. Since $t(H, W_0) = t(H, G_{\varepsilon, 0})$ and $t(H, W_\alpha) = t(H, G_{\varepsilon, \alpha})$, we conclude that the graph H does not have the step Sidorenko property. \square

Theorem 3 yields immediately the following corollary, which in particular rules out many non-biregular graphs to have the step Sidorenko property. Note that the assumptions of the corollary are easy to verify.

Corollary 4. *Let H be a graph and \mathcal{D}_H the set of degrees of its vertices. Further let M be the matrix with rows and columns indexed by the elements of \mathcal{D}_H such that the entry $M_{dd'}$ is equal to the number of 2-edge paths from a vertex of degree d to a vertex of degree d' in H . If the matrix M is not positive semidefinite, then H does not have the step Sidorenko property.*

Proof. We can assume without loss of generality that H is bipartite; if not, H does not even have the Sidorenko property. Let $n = |H|$, let $d_1 < \dots < d_k$ be the degrees of vertices of H , i.e., $\mathcal{D}_H = \{d_1, \dots, d_k\}$, and let $D = d_1 + \dots + d_k$. We next construct a weighted bipartite graph G_ε with weights depending on a parameter $\varepsilon > 0$. One part of G_ε has $k + 1$ vertices, which are denoted by u_1, \dots, u_{k+1} , and the other part has $D - k + 1$ vertices. One of the vertices of the second part is denoted by u_0 and the remaining $D - k$ vertices are split into disjoint sets U_1, \dots, U_k such that $|U_i| = d_i - 1$, $i = 1, \dots, k$. The vertices u_0 and u_{k+1} have weight one, each of the vertices u_i has weight $\varepsilon^{\frac{1}{|U_i|}}$ and each vertex contained in a set U_i has weight $\varepsilon^{\frac{1}{|U_i|}} / (|U_i| - 1)!$, $i = 1, \dots, k$. The weights of all edges of G_ε are equal to one.

We will apply Theorem 3 with the weighted graph G_ε , vertices u_0, \dots, u_k and sets $U_1 \cup \{u_0\}, \dots, U_k \cup \{u_0\}$. Let M_ε be the matrix from the statement of Theorem 3 for the graph G_ε . Fix $i, j \in \{1, \dots, k\}$ and a 2-edge path $v_1 v_0 v_2$ such that the degree of v_1 is d_i and the degree of v_2 is d_j . Let h be a mapping such that $h(v_0) = u_0$, $h(v_1) = u_i$ and $h(v_2) = u_j$. The mapping h can be extended to $(|U_i| - 1)! (|U_j| - 1)!$ homomorphisms from H to G such that

- the neighbors of v_1 are one-to-one mapped to $U_i \cup \{u_0\}$,
- the neighbors of v_2 are one-to-one mapped to $U_j \cup \{u_0\}$, and
- all other vertices of H are mapped to u_0 or to u_{k+1} .

Each such homomorphism has weight $\frac{\varepsilon^2}{(|U_i| - 1)! (|U_j| - 1)!}$, i.e., their total weight is ε^2 . Any other extensions of h to a homomorphism from H to G such that the neighbors of v_1 are one-to-one mapped to $U_i \cup \{u_0\}$ and the neighbors of v_2 to $U_j \cup \{u_0\}$ has weight at most ε^{2+1/d_k} . We conclude that the entry of the matrix M_ε in the i -th row and the j -th column is equal to $M_{ij} \varepsilon^2 + O(\varepsilon^{2+1/d_k})$. It follows that there exists $\varepsilon > 0$ such that the matrix M_ε is not positive semidefinite. Theorem 3 now yields that H does not have the step Sidorenko property. \square

The weights of vertices and edges of the graph G in Theorem 3 can be set to lower the weight of specific homomorphisms, as we did in Corollary 2. We first formalize the ideas used there, so that we can focus on just the existence of very restricted homomorphisms, without counting or weights.

Lemma 5. *Let H be a vertex-transitive graph. Let u_0, u_1 and u_2 be (distinct) distinguished vertices in H such that $u_0 u_1$ and $u_0 u_2$ are edges. Suppose that*

for each distinct neighbors v_1 and v_2 of u_0 , there is no homomorphism f from $H(u_0, v_1, v_2)$ to $H(u_0, u_1, u_1)$ that simultaneously satisfies the following:

- neighbors of v_i are one-to-one mapped to neighbors of u_1 for $i = 1, 2$,
- distances from u_0 are preserved, i.e., $\text{dist}(v, u_0) = \text{dist}(f(v), u_0)$ for each $v \in V(H)$, and
- no vertex other than u_0, v_1 and v_2 is mapped to any of u_0, u_1 and u_2 .

Then H does not have the step Sidorenko property.

Proof. We start with constructing a weighted graph G_γ where the weights depend on a parameter $\gamma \in \mathbb{N}$. The graph G_γ is obtained from H by setting $w(v) := \gamma^{\text{dist}(u_0, v)-1}$ for $v \in \{u_0, u_1, u_2\}$ and $w(v) := \gamma^{\text{dist}(u_0, v)}$ for each vertex $v \neq u_0, u_1, u_2$. The weights of all edges of G_γ are one. We apply Theorem 3 to H and G_γ with the distinguished vertices u_0, u_1 and u_2 . Since H is vertex-transitive, we will analyze the matrix M such that M_{ij} is the sum of weights of homomorphisms from $H(u_0, v_1, v_2)$ to $G_\gamma(u_0, u_i, u_j)$ such that the neighbors of v_1 and v_2 are mapped one-to-one to the neighbors of u_i and u_j , respectively, where the sum runs over all choices of v_1 and v_2 in H . Note that the matrix from the statement Theorem 3 is the considered matrix M with each entry multiplied by $|G|$, in particular, it is enough to show that the considered matrix M is not positive semidefinite for some γ .

Let $W := \sum_{v \in V(H)} \text{dist}(v, u_0) - 3$. We show that $M_{1,1} = o(\gamma^W)$, $M_{1,2} = M_{2,1} = \Omega(\gamma^W)$ and $M_{2,2} = O(\gamma^W)$ (as functions of the parameter γ). Hence, if γ is large enough, the matrix M is not positive semidefinite and H does not have the step Sidorenko property by Theorem 3.

By the definition, the entry $M_{1,2}$ contains a summand corresponding to the identity homomorphism from $H(u_0, v_1, v_2)$ to $G_\gamma(u_0, u_1, u_2)$; the weight of this summand is exactly γ^W . It follows $M_{1,2} = M_{2,1} = \Omega(\gamma^W)$.

Consider a homomorphism f contributing to the sum defining the entry $M_{i,i}$ for $i \in \{1, 2\}$. Observe that f satisfies $|f^{-1}(\{u_0, u_1, u_2\})| \geq 3$ (at least the three vertices u_0, v_1 and v_2 are mapped to u_0 and u_i) and $\text{dist}(u_0, f(v)) \leq \text{dist}(u_0, v)$ for every vertex v (a shortest walk from u_0 to v is mapped by f to a walk of at most the same length from u_0 to $f(v)$). Hence, it holds that $w(f(v)) \leq w(v)$ for every vertex v , and the equality holds for all vertices v if and only if $\text{dist}(u_0, f(v)) = \text{dist}(u_0, v)$ for every vertex v of H and $|f^{-1}(\{u_0, u_1, u_2\})| = 3$. In particular, the equality does not hold for any homomorphism f contributing to the sum defining the entry $M_{1,1}$. It follows that each summand in the sum defining the entry $M_{1,1}$ is of order $O(\gamma^{W-1})$ and each summand in the sum defining the entry $M_{2,2}$ is of order $O(\gamma^W)$. Since the number of the summands is independent of γ , we conclude that $M_{1,1} = o(\gamma^W)$ and $M_{2,2} = O(\gamma^W)$. \square

We conclude the paper with applying Lemma 5 to show that all multidimensional grids other than hypercubes are not weakly norming.

Corollary 6. *Let $k \geq 2$. The Cartesian product $C_{\ell_1} \square \cdots \square C_{\ell_k}$ has the step Sidorenko property if and only if the length of each cycle in the product is four, i.e., $\ell_1 = \cdots = \ell_k = 4$.*

Proof. Let $H = C_{\ell_1} \square \cdots \square C_{\ell_k}$. By symmetry, we can assume that ℓ_1 is the largest and ℓ_2 is the smallest among ℓ_1, \dots, ℓ_k . If $\ell_1 = \cdots = \ell_k = 4$, the graph H is isomorphic to the $2k$ -dimensional hypercube graph, which is weakly norming, see [13] and [18, Proposition 14.2]; this implies that H has the step Sidorenko property [18, Proposition 14.13]. If ℓ_i is odd for some i , then the graph H is not bipartite, which implies that it fails to even have the Sidorenko property. Hence, we can assume that all ℓ_i are even and $\ell_1 > 4$.

We will view the vertices of H as the elements of $\mathbb{Z}_{\ell_1} \times \cdots \times \mathbb{Z}_{\ell_k}$ and perform all computations involving the i -th coordinate modulo ℓ_i . Let e_i be the i -th unit vector. Note that two vertices of H are adjacent if their difference is equal to e_i or $-e_i$ for some $i = 1, \dots, k$. Also observe that if v is a vertex of H and $\ell_i > 4$, then v is the only common neighbor of $v + e_i$ and $v - e_i$.

We apply Lemma 5 with $u_0 = (0, \dots, 0)$ and $u_i = e_i$ for $i = 1, 2$. Suppose that for some distinct vertices v_1 and v_2 , there is a homomorphism f from $H(u_0, v_1, v_2)$ to $H(u_0, e_1, e_1)$ contradicting the assumption of Lemma 5, i.e.,

- (i) the neighbors of v_i are one-to-one mapped to neighbors of e_1 , for $i = 1, 2$,
- (ii) $\text{dist}(u_0, v) = \text{dist}(u_0, f(v))$ for each $v \in V(H)$, and
- (iii) no vertex other than u_0, v_1 and v_2 is mapped to any of the vertices u_0, e_1 and e_2 .

We will show that the existence of such a homomorphism f leads to a contradiction. By symmetry, we can assume that $v_1 = e_{i_1}$ for some i_1 and either $v_2 = -e_{i_1}$ or $v_2 = e_{i_2}$ for some $i_2 \neq i_1$.

Note that the neighbors of v_1 are one-to-one mapped to the neighbors of e_1 , and let i' be such that $f(e_{i_1} + e_{i'}) = e_1 + e_1$. If $i' \neq i_1$, both common neighbors of u_0 and $e_{i_1} + e_{i'}$, which are e_{i_1} and $e_{i'}$, must be mapped to the unique common neighbor of u_0 and $e_1 + e_1$, which is the vertex e_1 (note that $\ell_1 > 4$). However, this would contradict (iii). Hence, $i' = i_1$, i.e., $f(v_1 + v_1) = f(e_{i_1} + e_{i_1}) = e_1 + e_1$. It follows that there exists a bijection π between $\{\pm e_{i'} \mid i' \neq i_1\}$ and $\{\pm e_{j'} \mid j' \neq 1\}$ such that $f(e_{i_1} + e) = e_1 + \pi(e)$ for $e \in \{\pm e_{i'} \mid i' \neq i_1\}$. Observe that a symmetric argument to the one that we have just presented yields that $f(v_2 + v_2) = e_1 + e_1$.

To exclude the case that $v_2 = -e_{i_1}$, let $e = \pi^{-1}(e_2)$, i.e., $f(e_{i_1} + e) = e_1 + e_2$. Note that $e \neq \pm e_{i_1}$. It follows that the vertex e , which is a common neighbor of u_0 and $e_{i_1} + e$, must be mapped to a common neighbor of u_0 and $e_1 + e_2$, i.e., either to e_1 or to e_2 . The first case would contradict (iii), hence e is mapped to e_2 , meaning $v_2 = e$. We conclude that $v_2 = e_{i_2}$ for some $i_2 \neq i_1$ and that $f(e_{i_1} + e_{i_2}) = e_1 + e_2$.

Suppose that $\ell_2 = 4$ and recall that $f(v_2 + v_2) = e_1 + e_1$. If additionally $\ell_{i_2} = 4$, then $-e_{i_2}$, which is a common neighbor of u_0 and $e_{i_2} + e_{i_2}$, must be mapped to the unique common neighbor of u_0 and $e_1 + e_1$, i.e., to the vertex e_1 ; this is impossible by (iii). Hence, $\ell_{i_2} \neq 4$.

Let us call two vertices v and v' *close* if they have at least two common neighbors. Observe that two close distinct neighbors v and v' of e_{i_1} must be mapped to close neighbors of e_1 ; otherwise, all common neighbors of v and v' would be mapped to e_{i_1} , contradicting (iii). Since the neighborhood of e_{i_1} is one-to-one mapped to the neighborhood of e_1 and the number of pairs of close neighbors of e_{i_1} is the same as the number of pairs of close neighbors of e_1 , it follows that pairs of close neighbors of e_{i_1} are one-to-one mapped to pairs of close neighbors of e_1 and pairs of non-close neighbors of e_{i_1} are one-to-one mapped to pairs of non-close neighbors of e_1 . Since $\ell_{i_2} \neq 4$, the neighbors $e_{i_1} + e_{i_2}$ and $e_{i_1} - e_{i_2}$ of e_{i_1} are not close. On the other hand, since $\ell_2 = 4$, the vertex $f(e_{i_1} + e_{i_2}) = e_1 + e_2$ has a common neighbor other than e_1 with each neighbor of e_1 . In particular, $f(e_{i_1} + e_{i_2})$ and $f(e_{i_1} - e_{i_2})$ are close, which is impossible. We conclude that $\ell_2 \neq 4$. Since ℓ_2 is the smallest among ℓ_1, \dots, ℓ_k , it follows that each ℓ_i is at least six.

As the final step of the proof of the corollary, we prove the following statement for $r = 1, \dots, \ell_{i_1}/2$ by induction on r :

$$\begin{aligned} f((r-1)e_{i_1}) &= (r-1)e_1, & f(re_{i_1}) &= re_1, \text{ and} \\ f(re_{i_1} + e) &= re_1 + \pi(e) \text{ for } e \in \{\pm e_{i'} \mid i' \neq i_1\}. \end{aligned} \quad (5)$$

The case $r = 1$ follows from the definition of i_1 and π . We assume that the above statement holds for r and prove it for $r + 1 \leq \ell_{i_1}/2$. We first show that $f((r+1)e_{i_1}) = (r+1)e_1$. Note that $f(re_{i_1} + e_{i_1})$ cannot be $re_1 - e_1$ by (ii). If $f(re_{i_1} + e_{i_1})$ is $re_1 + e_j$ for some $j \neq 1$, then the common neighbor $re_{i_1} + e_{i_1} + \pi^{-1}(-e_j)$ of $re_{i_1} + e_{i_1}$ and $re_{i_1} + \pi^{-1}(-e_j)$ must be mapped to the unique common neighbor of $re_1 + e_j$ and $re_1 - e_j$, which is re_1 , contradicting (ii). An analogous argument excludes that $f(re_{i_1} + e_{i_1})$ is $re_1 - e_j$ for some $j \neq 1$. Since the vertex $f((r+1)e_{i_1})$ must be a neighbor of $f(re_{i_1}) = re_1$, it follows that $f((r+1)e_{i_1}) = (r+1)e_1$.

We next analyze $f((r+1)e_{i_1} + e)$ for $e \neq \pm e_{i_1}$. Since the vertex $(r+1)e_{i_1} + e = re_{i_1} + e_{i_1} + e$ is a common neighbor of $re_{i_1} + e_{i_1}$ and $re_{i_1} + e$, it must be mapped to a common neighbor of $re_1 + e_1$ and $re_1 + \pi(e)$, i.e., to re_1 or $re_1 + e_1 + \pi(e)$. Since the former is excluded by (ii), it follows that $f((r+1)e_{i_1} + e) = (r+1)e_1 + \pi(e)$. This concludes the proof of (5).

The statement (5) implies that $f(\ell_{i_1}/2 \cdot e_{i_1}) = \ell_{i_1}/2 \cdot e_1$, in particular $\ell_{i_1} \geq \ell_1$ by (ii). Since the path $u_0, -e_{i_1}, -2e_{i_1}, \dots, -\ell_{i_1}/2 \cdot e_{i_1}$ must be mapped to a path from u_0 to $f(-\ell_{i_1}/2 \cdot e_{i_1}) = f(\ell_{i_1}/2 \cdot e_{i_1}) = \ell_{i_1}/2 \cdot e_1$ and the vertices of the path must be mapped to vertices at distances $0, 1, \dots, \ell_{i_1}/2$ from u_0 by (ii), the path can be mapped only to the path $u_0, e_1, 2e_1, \dots, \ell_{i_1}/2 \cdot e_1$ or, if $\ell_1 = \ell_{i_1}$, to the path $u_0, -e_1, -2e_1, \dots, -\ell_{i_1}/2 \cdot e_1$. The former case is impossible since $-e_{i_1}$ cannot

be mapped to e_1 by (iii). It follows that $\ell_1 = \ell_{i_1}$ and $f(-e_{i_1}) = -e_1$. Hence, the vertex $e_{i_2} - e_{i_1} \neq u_0$, which is a common neighbor of e_{i_2} and $-e_{i_1}$, must be mapped to the unique common neighbor of $f(e_{i_2}) = e_1$ and $f(-e_{i_1}) = -e_1$, which is u_0 . However, this contradicts (iii). We conclude there is no homomorphism f satisfying (i)–(iii). Lemma 5 now implies that H does not have the step Sidorenko property. \square

5 Conclusion

Corollary 2 and Corollary 6 give an infinite class of edge-transitive graphs that are not weakly norming, which answers in the negative a question of Hatami [13]. Conlon and Lee [5, Conjecture 6.3] present a large class of weakly norming graphs, which they call reflection graphs, and conjecture that a bipartite graph is weakly norming if and only if it is edge-transitive under a subgroup of its automorphism group (generated by so called ‘cut involutions’). In particular, this would imply that all weakly norming graphs are edge-transitive.

Since every weakly norming graph has the step Sidorenko property, it is natural to ask whether the converse is true for connected graphs, i.e., whether every connected graph with the step Sidorenko property is weakly norming. This question has been very recently answered in the affirmative by Doležal et al. [7] who showed the following: a connected graph G is weakly norming if and only if it has the step Sidorenko property.

Finally, it is natural to wonder about the Forcing Conjecture in the setting of the step Sidorenko property. Let us say that a graph H has the *step forcing property* if and only if

$$t(H, W^{\mathcal{P}}) \leq t(H, W)$$

for every graphon W and every partition \mathcal{P} of $[0, 1]$ into finitely many non-null measurable sets and the equality holds if and only if $W^{\mathcal{P}}$ and W are equal almost everywhere. It can be shown that all even cycles have the step forcing property (while an ad hoc argument can be given, this also follows from [7, Theorem 3.14]). Graphs with the step forcing property are related to the proof of the existence of graphons via weak* limits given by Doležal and Hladký [8]; in particular, if H has the step forcing property, minimizing the entropy of W in the arguments given in [8] can be replaced by maximizing $t(H, W)$.

Acknowledgements

The first author would like to thank Karel Král and László Miklós Lovász for discussions on the step Sidorenko property and the step forcing property.

References

- [1] G. R. Blakley and P. Roy: *A Hölder type inequality for symmetric matrices with nonnegative entries*, Proc. Amer. Math. Soc. **16** (1965), 1244–1245.
- [2] F. R. K. Chung, R. L. Graham and R. M. Wilson: *Quasi-random graphs*, Combinatorica **9** (1989), 345–362.
- [3] D. Conlon, J. Fox and B. Sudakov: *An approximate version of Sidorenko’s conjecture*, Geom. Funct. Anal. **20** (2010), 1354–1366.
- [4] D. Conlon, J. H. Kim, C. Lee and J. Lee: *Some advances on sidorenko’s conjecture*, to appear in J. Lond. Math. Soc. .
- [5] D. Conlon and J. Lee: *Finite reflection groups and graph norms*, Adv. Math. **315** (2017), 130–165.
- [6] D. Conlon and J. Lee: *Sidorenko’s conjecture for blow-ups*, preprint arXiv:1809.01259 (2018).
- [7] M. Doležal, J. Grebík, J. Hladký and V. Rocha, I. Rozhoň: *Cut distance identifying graphon parameters over weak* limits*, preprint arXiv:1809.03797 (2018).
- [8] M. Doležal and J. Hladký: *Cut-norm and entropy minimization over weak* limits*, preprint arXiv:1701.09160 (2017).
- [9] J. Fox and F. Wei: *On the local approach to Sidorenko’s conjecture*, in: Electronic Notes in Discrete Mathematics **61** (2017), 459–465 (Proceedings of EuroComb’17).
- [10] F. Franek and V. Rödl: *Ramsey problem on multiplicities of complete subgraphs in nearly quasirandom graphs*, Graphs Combin. **8** (1992), 299–308.
- [11] F. Franek and V. Rödl: *2-colorings of complete graphs with a small number of monochromatic K_4 subgraphs*, Discrete Math. **114** (1993), 199–203.
- [12] A. W. Goodman: *On sets of acquaintances and strangers at any party*, Amer. Math. Monthly **66** (1959), 778–783.
- [13] H. Hatami: *Graph norms and Sidorenko’s conjecture*, Israel J. Math. **175** (2010), 125–150.
- [14] H. Hatami, J. Hladký, D. Král’, S. Norine and A. Razborov: *Non-three-colourable common graphs exist*, Combin. Probab. Comput. **21** (2012), 734–742.

- [15] C. Jagger, P. Štoviček and A. Thomason: *Multiplicities of subgraphs*, *Combinatorica* **16** (1996), 123–141.
- [16] J. H. Kim, C. Lee and J. Lee: *Two approaches to Sidorenko’s conjecture*, *Trans. Amer. Math. Soc.* **368** (2016), 5057–5074.
- [17] J. L. X. Li and B. Szegedy: *On the logarithmic calculus and Sidorenko’s conjecture*, to appear in *Combinatorica* .
- [18] L. Lovász: *Large Networks and Graph Limits*, *Colloquium Publications*, volume 60.
- [19] L. Lovász and B. Szegedy: *Limits of dense graph sequences*, *J. Combin. Theory Ser. B* **96** (2006), 933–957.
- [20] A. Sidorenko: *A correlation inequality for bipartite graphs*, *Graphs Combin.* **9** (1993), 201–204.
- [21] M. Simonovits: *Extremal graph problems, degenerate extremal problems, and supersaturated graphs*, in: *Progress in graph theory* (Waterloo, Ont., 1982), 419–437.
- [22] J. Skokan and L. Thoma: *Bipartite subgraphs and quasi-randomness*, *Graphs Combin.* **20** (2004), 255–262.
- [23] B. Szegedy: *An information theoretic approach to sidorenko’s conjecture*, preprint arXiv:1406.6738 (2014).
- [24] B. Szegedy: *On Sidorenko’s conjecture for determinants and Gaussian Markov random fields*, preprint arXiv:1701.03632 (2017).
- [25] A. Thomason: *Pseudorandom graphs*, in: *Random graphs ’85* (Poznań, 1985), *North-Holland Math. Stud.*, volume 144, 307–331.
- [26] A. Thomason: *A disproof of a conjecture of Erdős in Ramsey theory*, *J. London Math. Soc.* **39** (1989), 246–255.