

**Manuscript version: Author's Accepted Manuscript**

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

**Persistent WRAP URL:**

<http://wrap.warwick.ac.uk/115187>

**How to cite:**

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk).

1 **Investigating the potential for call combinations in a life-long vocal learner.**

2

3 Sarah L. Walsh<sup>a</sup>, Simon W. Townsend<sup>b,c</sup>, Kate Morgan<sup>a</sup> & Amanda R. Ridley<sup>a</sup>

4

5 <sup>a</sup>School of Biological Sciences, The University of Western Australia, Crawley, WA 6009, Australia;

6 <sup>b</sup>Department of Comparative Linguistics, University of Zurich, Zurich, 8032, Switzerland; <sup>c</sup>Department of

7 Psychology, University of Warwick, Coventry CV47AL, United Kingdom.

8

9 **ABSTRACT**

10

11 The ability for humans to create seemingly infinite meaning from a finite set of sounds has

12 likely been a critical component in our success as a species, allowing the unbounded

13 communication of information. Syntax, the combining of meaningful sounds into phrases, is

14 one of the primary features of language that enables this extensive expressivity. The

15 evolutionary history of syntax, however, remains largely debated, and it is only very recently

16 that comparative data for syntax in animals has been revealed. Here, we provide further

17 evidence for a structural basis of potential syntactic-like call combinations in the vocal

18 communication system of a group-living songbird. Acoustic analyses indicate that Western

19 Australian magpies (*Cracticus tibicen dorsalis*) structurally combine generic alarm calls with

20 acoustically distinct alert calls to produce an alarm-alert sequence. These results are distinct

21 from previous examples of call combinations as, to our knowledge, evidence for this capacity

22 is yet to be demonstrated in a non-human species that is capable of vocal learning throughout

23 life. These findings offer prospects for experimental investigation into the presence and

24 function of magpie call combinations, extending our understanding of animal vocal

25 complexity.

26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

*Keywords: call combinations, syntax, animal vocal communication, language evolution, Western Australian magpie, Cracticus tibicen dorsalis.*

## **1. Introduction**

The immensely generative nature of language has long been thought to be a key factor distinguishing humans from other animals (Collier et al., 2014; Hurford, 2011). The property of language that enables extensive combinatorial power is known as double articulation and refers to the combining of sounds into words (phonology), and the grammatical combining of meaningful words into phrases (syntax, Hurford, 2011). While there is no doubt humans are capable of vast combinatorial power (Hurford, 2012), recent comparative research has begun to draw parallels between language’s combinatoriality and potential equivalents within non-human animal vocal communication systems (Arnold & Zuberbühler, 2006; Engesser et al., 2016; Ouattara et al., 2009; Suzuki et al., 2016; Townsend et al., 2018).

Many non-human animals, such as some cetacean and avian species for example, have been found capable of stringing sounds together into larger sequences, that, at least superficially, are reminiscent of the phonological layer of human language (Engesser et al., 2015; Hauser et al., 2002; Suzuki et al., 2006; Tyack, 1983). Moreover, a growing body of work has revealed interesting potential homologous and analogous examples of call combinations in animals which can be feasibly compared with simple syntactic structures in language (Arnold & Zuberbühler, 2006; Ouattara et al., 2009; Townsend et al., 2018).

50 The combination of context- specific calls into a meaningful sequence has been suggested in  
51 two different studies in related monkey taxa (Arnold & Zuberbühler, 2006; Ouattara et al.,  
52 2009). Putty-nosed monkeys (*Cercopithecus nictitans*) combine two predator-specific alert  
53 calls into a larger meaningful sequence (Arnold & Zuberbühler, 2006). However, despite the  
54 discrete calls each being elicited upon discovery of the presence of a specific predator, the  
55 sequence evokes group movement regardless of predatory context. It is, therefore, uncertain  
56 whether this call sequence is indicative of compositional or instead combinatorial syntax,  
57 where the meaning of the resultant sequence (i.e. ‘Let’s go’) is not a product of the  
58 combination of the alarm calls’ meanings (Collier et al., 2014; Hurford, 2012).

59

60 Perhaps more convincing evidence of rudimentary compositional syntax in primates is shown  
61 in a study on male Campbell’s monkeys (*Cercopithecus campbelli campbelli*), where the use  
62 of an acoustically invariant suffix modifies the meaning of predator-specific alarm calls in a  
63 predictable way (from more specific to more general, Ouattara et al., 2009). Specifically, this  
64 has been argued to be akin to abstract meaning operators in language, such as “like”, that  
65 generalises the meaning of words (“monkey” to “monkey-like”) and is therefore suggestive  
66 of basic compositional syntax (Collier et al., 2014; see also Townsend et al., 2018).

67

68 Recently, additional evidence of simple syntax has been seen in more distantly related  
69 species. The combining of warning and recruitment calls in the Japanese great tit (*Parus*  
70 *minor*) elicits a compound response consisting of a combination of the behaviours prompted  
71 by each individual call (Suzuki et al., 2016). Critically, when artificially reversing the  
72 sequence, the authors find a change in behavioural response, which is argued to indicate the  
73 sequence is perceived as a single meaningful unit (Suzuki et al., 2016). Similarly, alert and

74 recruitment calls in the cooperatively breeding southern pied babbler (*Turdoides bicolor*) are  
75 combined into a sequence eliciting mobbing behaviour from conspecifics in the presence of a  
76 territorial threat (Engesser et al., 2016). Experiments demonstrated that the individual calls  
77 have an independent meaning that are distinct from, but contribute to, the meaning of the  
78 mobbing sequence and thus are strongly indicative of rudimentary syntax (Engesser et al.,  
79 2016; see Suzuki et al., 2018, Townsend et al., 2018 and Zuidema & de Boer, 2018). These  
80 studies have laid the foundation for further research into basic syntactic abilities in non-  
81 human animals.

82

83 In this study, we build on emerging evidence of avian syntactic abilities by investigating the  
84 occurrence of call combinations in the Western Australian magpie. Magpies in south-western  
85 Australia are highly territorial, and cooperatively defend a year-round territory (Ashton et al.,  
86 2018; Farabaugh & Brown, 1988; Hughes et al., 1996). All group members participate in  
87 territorial defence behaviours throughout the year, including vocalizations, inter-group  
88 fighting, and aggressive posturing (Farabaugh & Brown, 1988; Kaplan, 2004; Kaplan, 2006;  
89 Kaplan et al., 2009). Both sexes have a similarly well-developed song and extensive vocal  
90 repertoire (Kaplan, 2005; Suthers et al., 2011; Walsh, 2017). Males do not use song or any  
91 other specific vocalization for courtship and thus share the same basic repertoire with females  
92 (Kaplan, 2005).

93

94 The vocal repertoire of adult Western Australian magpies (hereafter magpies) can be  
95 separated into short alarm calls, distress or alert calls, begging, feeding grunts, and carols or  
96 warbling calls (Baker, 2009; Kaplan, 2004; Silvestri, 2017; Walsh, 2017). Previous research  
97 has revealed the potential for context specificity in a variety of magpie vocalizations, such as

98 carols and some alarm calls (Baker, 2009; Kaplan, 2008). For example, carolling calls may  
99 signal territoriality and willingness to participate in defence of the group territory (Baker,  
100 2009). Furthermore, there is evidence that some alarm calls may be predator specific in  
101 nature (Kaplan, 2005; Kaplan, 2006; Kaplan, 2008; Kaplan et al., 2009; Kaplan & Rogers,  
102 2013). In light of the context-specific nature of magpie calls (Kaplan, 2008), and due to the  
103 clear capabilities for flexibility in call structure (Kaplan, 2000), we reasoned rudimentary  
104 syntactic structuring might also exist in the vocal system of magpies. While basic  
105 combinatoriality in magpie vocal communication has been alluded to in past research  
106 (Kaplan, 2005) to date, systematic investigations have not been conducted.

107

108 Moreover, to our knowledge, syntactic capabilities have not yet been indicated in a life-long  
109 vocal learner other than humans. Magpies have been shown to be capable of flexibly  
110 interspersing mimicry into their own song, indicating a high plasticity of vocal learning and  
111 high retention of learned material throughout life (Brown & Farabaugh, 1991; Kaplan, 2000;  
112 Kaplan, 2005). The fact that magpies can be considered as open-ended vocal learners is  
113 particularly interesting because theory suggests that the transition from a non-compositional  
114 system to a compositional one, relates to a constraint in the number of available calls  
115 (Nowack, 2000). As this would seemingly not be a constraint for a life-long vocal learner, if  
116 syntactic capability is revealed in magpies it could provide additional insight into the  
117 potential evolutionary forces promoting the progression of syntax.

118

119 One instrumental first step prior to unambiguously identifying a syntactic-like combination in  
120 non-human animal communication systems involves isolating naturally-occurring,  
121 stereotyped, call combinations and then demonstrating that the calls comprising the

122 combination and the same calls produced alone are acoustically equivalent (*sensu* Hedwig et  
123 al., 2014), such that the combination can be feasibly be considered as composed of  
124 independently-occurring calls (see Hedwig et al., 2015, Engesser et al., 2016 & Suzuki et al.,  
125 2016).

126

127 Observational work has indicated that magpies do indeed produce such a sequence in the  
128 form of an “alarm-alert sequence” appearing to be comprised of two independently occurring  
129 calls, a generic alarm call and an alert vocalisation (Figure 1), and used in a variety of threat-  
130 related contexts. Furthermore, field observations suggest the call may convey information to  
131 receivers with individuals responding predominantly with vigilance-type behaviours such as  
132 scanning, vigilance, mobbing a predator, and movement directly following the call (Walsh,  
133 2017). Regarding the individual call types, observations indicated alarm calls produced  
134 predominantly during the approach of humans, dogs and general disturbance (Silvestri, 2017;  
135 Walsh, 2017). The alert call, however, coincides with the presence of a potential predatory  
136 threat, often eliciting response behaviours such as vigilance, scanning or carolling  
137 vocalisations from other group members (Silvestri, 2017; Walsh, 2017).

138

139 The fact that two independently occurring calls (alarm and alert calls) are concatenated into a  
140 single structure (alarm-alert sequence) is suggestive that the call might represent a syntactic-  
141 like structure (Hurford, 2011), providing a solid basis for the investigation into this  
142 combinatorial ability (Townsend et al., 2018). Here, we provide the first step in  
143 demonstrating that magpies may be capable of producing call combinations by quantifying  
144 the extent to which the calls comprising the combination are (a) acoustically distinct units  
145 and (b) acoustically equivalent to those produced alone.

146

## 147 **2. Methods**

148

### 149 *2.1. Study site and animals*

150

151 Fourteen groups of free-living magpies were chosen for this study. The groups were situated  
152 in the urban Perth suburbs of Guildford (31°89'S, 115°96'E) and Crawley (31°98'S,  
153 115°81'E), within Western Australia. Many individuals within these groups have been ringed  
154 for identification and are habituated to human presence, therefore allowing observation and  
155 vocal recording at a close distance (Ashton et al., 2018; Edwards et al., 2015; Mirville et al.,  
156 2016).

157

### 158 *2.2. Natural observations and acoustic recording*

159

160 We recorded naturally occurring vocalizations and the context in which they were given,  
161 while following groups in the field to compile a database of independent calls and potential  
162 call combinations. Vocalizations were recorded at a 44.1 kHz sampling rate using a Roland  
163 R-05 digital wave/mp3 recorder coupled to a Rode NTG2 directional condenser microphone  
164 encased in a Rode Blimp Suspension Windshield. Data collection was conducted during  
165 observations sessions (4-6 hours/session) on 27 days between June and August 2017. Data  
166 was primarily collected in the morning to coincide with the higher activity levels of magpies  
167 (Edwards et al., 2015). Recording focused on adult magpies, with no restriction based on sex.  
168

169 We obtained over 500 recordings of distinct magpie vocalizations, which were extracted from

170 ~150 hours spent following groups in the field. The recordings we obtained were added to a  
171 sound database that already contained over 2000 recordings (~12 hours of raw material)  
172 collected by the University of Western Australia's (UWA) magpie research group from 2014  
173 to 2017. Within the database, the independent vocalizations were categorised into the  
174 following call types based primarily on auditory and observational features; begging, feeding  
175 grunt, alarm, alert, carol, chorus (more than two individuals carolling at the same time), and  
176 mimicry. Within each call type, calls were then further categorised into similar acoustic units  
177 based on visual (spectrographic) features. The following contextual details were logged  
178 where possible for each recording; caller identity and behaviour immediately prior  
179 to/following vocalization, group members present, stimulus or disturbance, and group  
180 response (vocal and behavioural).

181

182 Periods of silence have been used as a method of classifying boundaries between acoustic  
183 units in numerous previous studies, including research on birdsong in zebra finches  
184 (*Taeniopygia guttata*, Franz & Goller, 2002), killer whale (*Orcinus orca*) calls (Shapiro,  
185 2011), and gorilla (*Gorilla beringei beringei* & *G. gorilla gorilla*) close calls (Hedwig et al.,  
186 2014). Previous research on magpie song production indicated duration of mini-breaths  
187 between song syllables to be roughly < 0.3 seconds (Suthers et al., 2011). As this study  
188 focussed on alarm and alert calls rather than song production, we have slightly increased the  
189 period of silence between calls to reflect the observed trend in periods of silence in call series  
190 (independent calls produced consecutively) versus call combinations (two or more calls  
191 produced in concatenation). As such, in this study, we defined calls as either discrete calls  
192 (singular independent units separated by a period of silence equal to or greater than 0.5  
193 seconds) or potential call combinations (independent units separated by a period of silence

194 less than 0.5 seconds). This is based on visual inspection of spectrograms of individual calls,  
195 discrete call series, and potential call combinations.

196

197 A subset database was created focussing on the calls that, from preliminary field observation  
198 and acoustic investigation, were (a) given frequently and (b) flexibly combined into a larger  
199 sequence (see Figure 2 for examples). The subset database included the number of recordings  
200 obtained for each call type from each individual from each group, and their respective  
201 observational contexts. From the subset database, one type of alarm-alert sequence (Figure 1)  
202 was chosen for acoustic analysis. This sequence was chosen based on the number of different  
203 groups recorded producing this vocalization (N=11) and the high frequency of occurrence of  
204 this sequence (and its comprising calls) detected during observational research.

205

### 206 2.3. *Acoustic analysis*

207

208 Calls were assessed for quality (where high quality = high signal to background noise ratio),  
209 and the percentage of the call demonstrating clear, tonal structures using Audacity 2.1.2  
210 (Audacity Team, 2016). Due to a limitation in the number of good-quality recordings of each  
211 discrete call type in all test groups, only 26 recordings of the alarm, 9 recordings of the alert  
212 and 19 recordings of the alert-alarm sequence, were used for analysis (our study site is  
213 located within an urban matrix and so many recordings were compromised by background  
214 noise). Spectrographic cross-correlation (SPCC) analyses were run to compare the recordings  
215 within each call-type to determine if the calls were correlated. Analyses were then undertaken  
216 to determine if the two calls are distinct in structure, and to compare the recordings of  
217 discrete calls with the respective sequence component to determine similarity in structure.

218 SPCC compares the structural features in time-frequency spectrograms of two calls or  
219 sounds, taking a peak correlation value as a measure of similarity and generating a matrix of  
220 similarity values for each pair of calls in the analysis (Cortopassi & Bradbury, 2012).

221

222 To verify that alarm and alert calls are independent units that are structurally distinct from  
223 each other, both the discrete calls and their respective sequence components were cross-  
224 correlated with the other call type using SPCC. A procedure based on the Mantel test was  
225 then used to test the null hypothesis that the calls grouped into the alarm or alert categories  
226 were no more similar to other calls within their own group than to calls within the other call-  
227 type group. As such, the SPCC matrix was compared with a second ‘hypothesis’ matrix  
228 which contained a binary code, whereby ‘1’ represented same-group comparisons and ‘0’  
229 represented different-group comparisons in equivalent positions (‘group’ is defined as the  
230 recording being either an alarm or alert call, Schnell et al., 1985). A p-value less than the  
231 alpha (set at 0.05) indicates the two calls are structurally distinct.

232

233 SPCC and Mantel testing were then used to test the null hypothesis that the discrete calls  
234 were no more similar to each other than to their respective component in the call sequence,  
235 using the same protocol as described above (Schnell et al., 1985). A p-value > 0.05 indicates  
236 the structure of the discrete call is similar to the structure of its respective sequence  
237 component, therefore confirming the two discrete calls are structurally combined into a  
238 sequence.

239

240 All SPCC analyses were conducted using Batch Correlation Analysis in Raven (Bioacoustics  
241 Research Program, Cornell University, Ithaca, 2017) using normalized Hamming window

242 spectrographic correlation analysis on peak values, with a band pass filter from 500Hz to 22,  
243 000Hz.

244

245

### 246 **3. Results**

247

248 Spectrographic cross-correlation analyses revealed distinct call structure, with calls correlated  
249 within each call-type. The mean correlation coefficient was consistent across cross-  
250 comparisons of alarm calls (SPCC:  $\bar{x} = 0.30$ ,  $n = 325$ ), alert calls (SPCC:  $\bar{x} = 0.34$ ,  $n = 36$ ),  
251 and alarm-alert sequences (SPCC:  $\bar{x} = 0.34$ ,  $n = 171$ ).

252

253 Mantel tests comparing correlation coefficients for alarm and alert call-type comparisons  
254 were highly significant (10,000 permutations,  $r = 0.284$ ,  $p < 0.0001$ ), indicating the two calls  
255 (alert and alarm) are structurally distinct from one another (Table 1).

256

257 Pairwise comparison analysis indicated the two discrete calls were correlated with their  
258 respective sequence counter-parts. Mantel tests comparing correlation coefficients from  
259 pairwise comparisons of each of the discrete calls with their respective cut-sequence  
260 components were non-significant for both alarm (10,000 permutations,  $r = 0.022$ ,  $p = 0.488$ )  
261 and alert calls (10,000 permutations,  $r = 0.019$ ,  $p = 0.706$ ) (Figure 3, Table 1). Therefore,  
262 recordings of the discrete calls were no more similar to one another than to their respective  
263 component cut from the sequence.

264

### 265 **4. Discussion**

266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289

Using acoustic analyses, we investigated a potential call combination in the Western Australian magpie. Our results indicate that the alarm and alert calls are both acoustically distinct from one another but similar in structure to their respective sequence counterparts. Furthermore, spectrographic correlation analyses revealed similarity across recordings from multiple individuals, of the alarm, alert and sequence calls. These results provide evidence that the alarm and alert calls are distinct, highly repeatable calls that occur across all study groups and, at least at the structural level, are combined into an alarm-alert sequence.

The extent to which the detected combination of alarm and alert calls in the Western Australian magpie represents a syntactic-like structure, however, requires follow up experimental verification. Such experiments are key to unpacking the function of the independent calls and the combination via assessing receiver response in the absence of the stimuli that naturally elicit the vocalisations (*see* Suzuki et al., 2016 & Engesser et al., 2016).

Furthermore, although initial evidence is provided here for one potential call combination, preliminary observational data indicate magpies may be capable of flexibly creating combinations larger than a two-call construction, which has, to our knowledge, yet to be demonstrated in a non-human animal communication system (see Figure 2 e-h for potential examples, Russell & Townsend, 2017). Hence, follow up work is critical to assess the extent to which (a) magpies use call combinations, (b) whether these combinations are functionally relevant and meaningful to receivers, and (c) whether these combinations are indeed indicative of compositional syntax, or rather combinatoriality.

290 Our finding of a candidate call combination in magpies (if validated under future  
291 experimental investigation), in addition to previous research indicating similar rudimentary  
292 compositionality in primates (Ouattara et al., 2009) and other bird species (Engesser et al.,  
293 2016; Suzuki et al., 2016), further suggests that the capability for syntactic-like structuring in  
294 vocal communication is not confined to humans, and may instead be phylogenetically  
295 widespread (Collier et al., 2014). Such research can aid in bridging the gap between human  
296 and animal communication, and potentially shed light on the evolutionary development of  
297 complex vocal communication systems (Russell & Townsend, 2017).

298

299 It has been hypothesized that a transition to syntax may have evolved as a result of natural  
300 selection favouring the combining of meaningful calls when the number of relevant events  
301 needing to be communicated exceeds the number of calls available (Nowack et al., 2000).  
302 Thereafter, simple two-word or sign combinations may have occurred as a means of  
303 increasing lexicon size (Russell & Townsend, 2017). Interestingly, previous and ongoing  
304 work suggests magpies' already have an extensive vocal repertoire facilitated in part by their  
305 capacity to vocally imitate (Brown & Farabaugh, 1997; Kaplan, 2005; Suthers et al., 2011).  
306 Why magpies would need to combine calls to communicate new events if they could instead  
307 incorporate a new call/sound into their repertoire is therefore unclear. A bias towards  
308 combinatoriality could, in principle, be driven by working memory constraints on the part of  
309 the receiver: deriving meaning compositionally places less load on working memory than  
310 learning and storing a new call type (Nowack et al., 2000). Further work investigating the  
311 processing of calls will help shed light on the pressures driving combinatoriality in magpies,  
312 which in turn has the potential to shed light on the evolution of human syntax given that  
313 humans are also open-ended vocal production learners.

314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337

Investigating the presence of combinatoriality in animals is a critical first step to understanding the origins and phylogenetic spread of this key linguistic ability. Our results demonstrating combinatorial structures in Western Australian magpies contributes to recent work showing that avian species are capable of producing and processing call combinations (Engesser et al., 2016; Suzuki et al., 2016), in turn indicating that the phylogenetic spread of this capacity is greater than previously thought. Critically, our research provides important initial evidence of the presence of call combinations in a vocal learning species (excluding *Homo sapiens*), which has interesting implications for theories explaining the evolution of combinatoriality more generally. These results set the foundation for further experimental investigations into syntactic-like structures in this species – a vital step in furthering our understanding of this largely enigmatic field (Suzuki et al., 2018; Townsend et al., 2018).

### **Acknowledgements**

This project was funded by the School of Biological Sciences and a University Collaboration Award from the University of Western Australia and an Australian Research Council Discovery Grant (DP140101921) to ARR. SWT was funded by the Swiss National Science Foundation (Grant number: PP00P3\_163850).

### **References**

Arnold, K., & Zuberbühler, K. (2006). Semantic combinations in primate calls. *Nature*, *441*, 303. doi: 10.1038/441303a

338 Ashton, B., Ridley, A., Edwards, E., & Thornton, A. (2018). Cognitive performance is linked  
339 to group size and affects fitness in Australian magpies. *Nature*, 554(7692), 364-367. doi:  
340 10.1038/nature25503  
341

342 Audacity Team. (2016). Audacity (R): Free Audio Editor and Recorder (Version 2.1.2)  
343 [Computer Program]. Retrieved from <http://audacity.sourceforge.net/>  
344

345 Baker, M. C. (2009). Information Content in Chorus Song of the Group-Living Australian  
346 Magpie (*Cracticus tibicen dorsalis*) in Western Australia. *Ethology*, 115, 227-238. doi:  
347 10.1111/j.1439-0310.2008.01606.x  
348

349 Bioacoustics Research Program. (2017). Raven Pro: Interactive Sound Analysis Software  
350 (Version 1.5) [Computer Software]. Ithaca, NY: The Cornell Lab of Ornithology. Retrieved  
351 from [www.birds.cornell.edu/raven](http://www.birds.cornell.edu/raven)  
352

353 Brown, E., & Farabaugh, S. (1991). Song sharing in a group-living songbird, the Australian  
354 magpie, *Gymnorhina tibicen*, III: Sex Specificity and individual specificity of vocal part in  
355 communal chorus and duet songs. *Behaviour*, 118, 244-274.  
356

357 Brown, E., & Farabaugh, S. (1997). What birds with complex relationships can tell us about  
358 vocal learning: vocal sharing in avian groups. *Social influences on vocal development*, 98127.  
359

360 Collier, K., Bickle, B., van Schaik, C. P., Manser, M. B., & Townsend, S. W. (2014).  
361 Language evolution: syntax before phonology? *Proceedings of the Royal Society B*,

362 281(1788), 20140263. doi: 10.1098/rspb.2014.0263

363

364 Cortopassi, K. A., & Bradbury, J. W. (2012). The Comparison of Harmonically Rich Sounds  
365 using Spectrographic Cross-Correlation and Principal Coordinates Analysis. *Bioacoustics:*

366 *The International Journal of Animal Sound and its Recording*, 11(2), 89-127. doi:

367 10.1080/09524622.2000.9753454

368

369 Edwards, E., Mitchell, N., & Ridley, A. (2015). The impact of high temperatures on foraging  
370 behaviour and body condition in the Western Australian Magpie *Cracticus tibicen dorsalis*.

371 *Ostrich*, 86(1-2), 137-144. doi: 10.2989/00306525.2015.1034219

372

373 Engesser, S., Crane, J. M., Savage, J. L., Russell, A. F., & Townsend, S. W. (2015).

374 Experimental Evidence for Phonemic Contrasts in a Nonhuman Vocal System. *PLoS Biol*,

375 13(6), e1002171. doi: 10.1371/journal.pbio.1002171

376

377 Engesser, S., Ridley, A., & Townsend, S. W. (2016). Meaningful call combinations and

378 compositional syntax in the southern pied babbler. *Proceedings of the National Academy of*

379 *Sciences*, 113(21), 5976-5981. doi: 10.1073/pnas.1600970113

380

381 Farabaugh, S., & Brown, E. (1988). Song sharing in a group living songbird, the Australian  
382 magpie, *Gymnorhina tibicen*. Part I. Vocal sharing within and among social groups.

383 *Behaviour*, 104, 1-29.

384

385 Franz, M., & Goller, F. (2002). Respiratory units of motor production and song imitation in

386 the zebra finch. *Journal of Neurobiology*, 51(2), 129-141. doi: 10.1002/neu.10043  
387

388 Hauser, M. D., Chomsky, N., & Fitch, W. T. (2002). The Faculty of Language: What Is It,  
389 Who Has It, and How Did It Evolve? *Science*, 298(5598), 1569-1579.  
390

391 Hedwig, D., Hammerschmidt, K., Mundry, R., Robbins, M., & Boesch, C. (2014). Acoustic  
392 structure and variation in mountain and western gorilla close calls: a syntactic approach.  
393 *Behaviour*, 151, 1091-1120. doi: 10.1163/1568539X-00003175  
394

395 Hedwig, D., Mundry, R., Robbins, M., & Boesch, C. (2015). Contextual correlates of  
396 syntactic variation in mountain and western gorilla close-distance vocalizations: Indications  
397 for lexical or phonological syntax? *Animal Cognition*, 18, 423-435. doi: 10.1007/s10071-  
398 0140812-6  
399

400 Hughes, J., Hesp, J., Kallioinen, R., Kempster, M., Lange, C., Hedstrom, K., Mather, P.,  
401 Robinson, A., & Wellbourn, M. (1996). Differences in Social Behaviour Between  
402 Populations of the Australian Magpie *Gymnorhina tibicen*. *EMU*, 96, 65-70.  
403

404 Hurford, J. (2011). The evolution of human communication and language. In P. D'Ettorre &  
405 H. D. P (Eds.), *Sociobiology communication: and interdisciplinary perspective*. Oxford, UK:  
406 Oxford University Press.  
407

408 Hurford, J. (2012). *The Origins of Grammar: Language in the Light of Evolution II*. Oxford,  
409 UK: Oxford University Press.

410

411 Kaplan, G. (2000). Song structure and function of mimicry in the Australian magpie  
412 (*Gymnorhina tibicen*) compared to the lyrebird (*Menura* ssp.). *Int J Comp Psych*, 12, 219241.

413

414 Kaplan, G. (2004). *Australian Magpie: Biology and Behaviour of an Unusual Songbird*.  
415 Collingwood, Victoria: CSIRO Pub.

416

417 Kaplan, G. (2005). Vocal behaviour of Australian Magpies (*Gymnorhina tibicen*): A Study of  
418 Vocal Development, Song Learning, Communication and Mimicry in the Australian Magpie.  
419 PhD Thesis, University of Queensland.

420

421 Kaplan, G. (2006). Alarm calls, communication and cognition in Australian magpies  
422 (*Gymnorhina tibicen*). *Act Zool Sinica*, 52, 614-617.

423

424 Kaplan, G. (2008). Alarm calls and referentiality in Australian magpies: Between midbrain  
425 and forebrain, can a case be made for complex cognition? *Brain Research Bulletin*, 76,  
426 253263. doi: 10.1016/j.brainresbull.2008.02.006

427

428 Kaplan, G., Johnson, G., Koboroff, A., & Rogers, L. J. (2009). Alarm calls of the Australian  
429 Magpie (*Gymnorhina tibicen*): Predators Elicit Complex Vocal Responses and Mobbing  
430 Behaviour. *The Open Ornithology Journal*, 2, 7-16. doi: 10.2174/1874453200902010007

431

432 Kaplan, G., & Rogers, L. J. (2013). Stability of referential signalling across time and  
433 locations: testing alarm calls of Australian magpies (*Gymnorhina tibicen*) in urban and rural

434 Australia and in Fiji. *PeerJ*, e112. doi: 10.7717/peerj.112

435

436 Mirville, M., Kelley, J., & Ridley, A. (2016). Group size and associative learning in the  
437 Australian magpie (*Cracticus tibicen dorsalis*). *Behavioural ecology and sociobiology*, 70(3),  
438 417-427. doi: 10.1007/s00265-016-2062-x

439

440 Nowack, M., Plotkin, J., & Jansen, V. (2000). The evolution of syntactic communication.  
441 *Nature*, 404, 495-498. doi: 10.1038/35006635

442

443 Ouattara, K., Lemasson, A., & Zuberbühler, K. (2009). Campbell's Monkeys use Affixation  
444 to Alter Call Meaning. *PLoS ONE*, 4(11), e7808. doi: 10.1371/journal.pone.0007808

445

446 Russell, A. F., & Townsend, S. W. (2017). Communication: Animal Steps on the Road to  
447 Syntax? *Current Biology*, 27, R753-R755. doi: 10.1016/j.cub.2017.06.066.

448

449 Schnell, G. D., Watt, D. J., & Douglas, M. E. (1985). Statistical comparison of proximity  
450 matrices: applications in animal behaviour. *Animal Behaviour*, 33, 239-253.

451

452 Shapiro, A. D., Tyack, P. L., Seneff, S. (2011). Comparing call-based versus subunit-based  
453 methods for categorizing Norwegian killer whale, *Orcinus orca*, vocalizations. *Animal*  
454 *Behaviour*, 81(2), 377-386.

455

456 Silvestri, A. (2017) The discrimination of alarm callers based on reliability in Western  
457 Australian magpies (*Cracticus tibicen dorsalis*). Honours' Thesis, University of Western

458 Australia.

459

460 Suthers, R. A., Wild, J. M., & Kaplan, G. (2011). Mechanisms of song production in the  
461 Australian magpie. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol*, 197(1), 45-49.  
462 doi: 10.1007/s00359-010-0585-6.

463

464 Suzuki, R., Buck, J. R., & Tyack, P. L. (2006). Information entropy of humpback whale  
465 songs. *Journal of the Acoustical Society of America*, 119(3), 1849-1866. doi:  
466 10.1121/1.2161827

467

468 Suzuki, T. N., Wheatcroft, D., & Griesser, M. (2016). Experimental evidence for  
469 compositional syntax in bird calls. *Nature Communications*, 7, 10986. doi:  
470 10.1038/ncomms10986

471

472 Suzuki, T. N., Wheatcroft, D., & Griesser, M. (2018). Call combinations in birds and the  
473 evolution of compositional syntax. *PLoS Biol*, 16 (8): e2006532

474

475 Townsend, S. W., Engesser, S., Stoll, S., Zuberbühler, K., Bickel, B. (2018).  
476 Compositionality in animals and humans. *PLoS Biol*, 16.

477

478 Tyack, P. L. (1983). Differential response of humpback whales, *Megaptera noveangliae*, to  
479 playback of song or social sounds. *Behavioral Ecology and Sociobiology*, 13, 49-55.

480

481 Walsh, Sarah. (2017). Evidence for potential compositional syntax in an Australian songbird:

482 a step toward understanding the evolution of language. Honours' Thesis, University of  
483 Western Australia.

484

485 Zuidema, W & de Boer, B. (2018). The evolution of combinatorial structure in language.  
486 *Current Opinion in Behavioural Sciences*, 21: 138-144.

487

## 488 **TABLE**

489

490 **Table 1. Comparisons of different calls types using Mantel test.**

Comparison	Mean Correlation	r	P
Alarm vs Sequence-Cut Alarm	0.315	0.022	0.488
Alert vs Sequence-Cut Alert	0.355	0.019	0.706
Alarm vs Alert	0.269	0.284	<0.0001*

491 Mantel tests were based on correlation coefficients obtained from SPCC on discrete calls versus respective  
492 sequence components and alarm call versus alert call. 18 recordings of each sequence-cut call, 26 recordings of  
493 alarm calls, and 9 recordings of alert calls were used in analysis. Asterisks indicates significant difference in call  
494 comparison whereby  $p < 0.05$ .

495

## 496 **FIGURE LEGENDS**

497

498 **Figure 1.** Spectrograms of the independent (a) alarm and (b) alert calls, and the (c) sequence  
499 they comprise which was used for acoustic analysis. Calls are separated into independent  
500 units (indicated by black lines) based on their occurrence with a period of at least 0.5 second  
501 of silence in between calls.

502

503 **Figure 2.** Spectrograms of magpie alarm (a & b), alert (c & d) calls and potential call  
504 combinations (e-h). Calls are separated into independent units (indicated by black lines)  
505 based on their occurrence with a period of at least 0.5 second of silence in between calls.

506

507 **Figure 3.** Distribution of correlation coefficients obtained from SPCC on pairwise  
508 comparison of each alarm and alert call-types. Calls were correlated within each call-type.  
509 Boxes show the medians and interquartile ranges, and whiskers show the ranges (n = number  
510 of comparisons).