

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/128725>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

ON PERTURBATIONS OF THE SCHWARZSCHILD ANTI-DE SITTER SPACES OF POSITIVE MASS

LUCAS C. AMBROZIO

ABSTRACT. In this paper we prove the Penrose inequality for metrics that are small perturbations of the Schwarzschild anti-de Sitter metrics of positive mass. We use the existence of a global foliation by weakly stable constant mean curvature spheres and the monotonicity of the Hawking mass.

1. INTRODUCTION

Let (M^3, g) be a complete Riemannian three-manifold, possibly with boundary, with exactly one end. Assume that the complement of a compact subset of M is diffeomorphic to \mathbb{R}^3 minus a ball. Roughly speaking, (M, g) is called *asymptotically hyperbolic* when the metric g decays sufficiently fast to the hyperbolic metric when expressed in the spherical coordinates induced by this chart. If an asymptotically hyperbolic manifold (M, g) has scalar curvature $R \geq -6$ and its boundary ∂M is empty or has mean curvature $H \leq 2$, there is a well-defined geometric invariant called the *total mass* of (M, g) , a non-negative number m that is zero if and only if (M, g) is isometric to the hyperbolic space. This is the content of the Positive Mass Theorem in the asymptotically hyperbolic setting, proved with spinorial methods by X. Wang [24] when ∂M is empty and by P. Chruściel and M. Herzlich [6] under weaker asymptotic conditions.

If an asymptotically hyperbolic manifold (M, g) has scalar curvature $R \geq -6$ and its boundary ∂M is a *connected minimal surface* that is *outermost*, i.e., if there are no other closed minimal surfaces in M , then it is conjectured that the total mass m of (M, g) is related to the area of ∂M by the following inequality:

$$(1) \quad \left(\frac{|\partial M|}{16\pi} \right)^{\frac{1}{2}} + 4 \left(\frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}} \leq m.$$

This statement is known as the *Penrose Conjecture* in the asymptotically hyperbolic setting. We refer the reader to the surveys [3] and [17], where he will find a comprehensive discussion on these types of inequalities in various settings and also on the physics behind these conjectures. In the original

The author was supported by FAPERJ and CNPq-Brasil.

asymptotically flat setting, the Penrose Conjecture was proved with different techniques by G. Huisken and T. Ilmanen [14] and by H. Bray [2].

The Penrose Conjecture contains also a rigidity statement. There are important models, known as the *Schwarzschild anti-de Sitter spaces of positive mass*, that satisfy the equality in (1). They are obtained as spherically symmetric metrics g_m on $M = [0, +\infty) \times S^2$ with constant scalar curvature -6 , where the parameter m is a positive real number that coincides with the total mass of (M, g_m) above discussed (see Section 2 for more details). The Penrose Conjecture also asserts that the Schwarzschild anti-de Sitter spaces of positive mass are the unique asymptotically hyperbolic manifolds, with scalar curvature $R \geq -6$ and an outermost minimal boundary, that satisfy the equality in (1).

A special feature of the Schwarzschild anti-de Sitter spaces of positive mass is that they are foliated by constant mean curvature spheres which are *weakly stable*. This kind of foliation of an asymptotically hyperbolic manifold is interesting, among other reasons, because of a monotonicity result observed by H. Bray in [1]: if $R \geq -6$, the so-called Hawking mass functional, defined for closed surfaces Σ in (M, g) by

$$(2) \quad m_H(\Sigma) := \sqrt{\frac{|\Sigma|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma} (H^2 - 4) d\Sigma \right),$$

is monotone non-decreasing along these foliations in the direction of increasing area of the leaves.

If ∂M is a minimal surface in (M, g) , its Hawking mass is exactly the left-hand side of (1). This suggests an approach to prove the Penrose inequality (1), at least for the class of asymptotically hyperbolic manifolds (M, g) , with $R \geq -6$ and an outermost minimal boundary, admitting this kind of foliation starting at ∂M and sweeping out all M : Inequality (1) would follow if the Hawking mass of the leaves near the infinity converges to the total mass of (M, g) .

There are asymptotically hyperbolic manifolds admitting unique foliations by weakly stable CMC spheres *near the infinity*. Results in this direction were proved by R. Rigger [23] and then by A. Neves and G. Tian in a quite general setting (see [21] and [22]). See also [19] for other results of this nature in asymptotically hyperbolic settings. For the original asymptotically flat setting, see the pioneering work of G. Huisken and S.T. Yau [15].

In the present paper, we consider metrics on $M = [0, +\infty) \times S^2$ that are small *global* perturbations of the Schwarzschild anti-de Sitter metric g_m in a suitable sense and show that the foliation constructed in [22] outside a compact set can be extended up to ∂M . Following the argument outlined above, we then show the Penrose inequality is true for these asymptotically hyperbolic manifolds.

In order to state our results more precisely, let us introduce some terminology. Given $m > 0$, the perturbations of g_m we consider belong to the space $\mathcal{M}(M, m)$ of metrics g of class C^3 on $M := \{p = (s, x) \in [0, +\infty) \times S^2\}$ such that ∂M is a minimal surface in (M, g) and

$$d(g, g_m) := \sup_{p \in M} \left(\sum_{i=0}^3 \exp(4s) \|(\nabla^m)^i (g - g_m)\|_{g_m}(p) \right) < +\infty,$$

see Section 2 for more details. All these metrics are asymptotically hyperbolic with total mass m in the sense of [24].

Our main results can be summarized as follows (see Theorem 7 and Theorem 8):

Theorem. *Let $m > 0$. There exists $\epsilon > 0$ such that for every metric $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ the following statements hold:*

- i) There exists a foliation $\{\Sigma_t\}_{t \in [0, +\infty)}$ of (M, g) by weakly stable constant mean curvature spheres such that $\Sigma_0 = \partial M$ is an outermost minimal surface.*
- ii) (The Penrose inequality). If the scalar curvature of g is greater than or equal to -6 , then*

$$\left(\frac{|\partial M|}{16\pi} \right)^{\frac{1}{2}} + 4 \left(\frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}} \leq \lim_{t \rightarrow +\infty} m_H(\Sigma_t) = m,$$

with equality if and only if (M, g) is isometric to the Schwarzschild anti-de Sitter space of mass m .

The paper is organized as follows. In Section 2, we describe the geometric properties of the models and define the class of perturbations of (M, g_m) we are going to work with. In Section 3, we construct foliations of compact regions of M by weakly stable CMC spheres which begin at the minimal boundary. In Section 4, we recall A. Neves and G. Tian's result in [22] about existence and uniqueness of foliations near the infinity of certain asymptotically hyperbolic manifolds by weakly stable CMC spheres, stating a version adapted to the space of metrics we consider. In Section 5, we calculate the limit of the Hawking mass of the leaves of that foliation. In Section 6, we argue that the foliations constructed before match on their overlap if the perturbation is small enough and show that the obtained foliation has the properties described in the above theorem. In Section 7, we prove the Penrose inequality for these small perturbations following the monotonicity argument outlined above.

We remark that in the asymptotically hyperbolic setting there is also another form of the Penrose Conjecture where the boundary corresponds to some $H = 2$ surface (see [3] and [17]). In the end of the paper we briefly discuss it and explain the modifications of the previous results that establish it for small perturbations (see Theorem 9 and Theorem 10).

The approach to the Penrose Conjecture involving the monotonicity of the Hawking mass for a family of surfaces that interpolates the outermost boundary and the infinity was originally suggested in the asymptotically flat setting by R. Geroch [10], who proposed the inverse mean curvature flow to produce such family. This program was successfully implemented by G. Huisken and T. Ilmanen [14]. In the asymptotically hyperbolic setting, however, there are serious difficulties in using this approach, see the work of A. Neves [20].

In the asymptotically hyperbolic setting, the Penrose Conjecture in its full generality is still an open problem. Using Bray's monotonicity as in [1], J. Corvino, A. Gerek, M. Greenberg and B. Krummel [7] proved the conjecture for asymptotically hyperbolic manifolds that are isometric to a Schwarzschild anti-de Sitter space of positive mass outside a compact set under restrictive hypotheses on the behavior of its isoperimetric surfaces, among them the assumption that the coordinate spheres near infinity are isoperimetric surfaces. This hypothesis has been proved to be always verified for compact perturbations of the models by O. Chodosh [5], whose recent work has also showed that isoperimetric surfaces exists for all sufficiently large volumes in asymptotically hyperbolic manifolds with scalar curvature $R \geq -6$. Finally, F. Girão and L. L. de Lima proved both cases of the conjecture and the higher dimensional analogues of them for another class of asymptotically hyperbolic manifolds, those that are certain graphical hypersurfaces of the hyperbolic space (see [11] and [12]). For related work on Penrose type inequalities in asymptotically hyperbolic settings, we refer the reader to [4], [8] and [16].

As pointed out by one of the referees, our result may be also conceived as a "local" Penrose inequality and may be related to the dynamical stability of the Schwarzschild anti-de Sitter spaces of positive mass as solutions to the Einstein vacuum equations with negative cosmological constant (for further information on this subject, see [9] and [13]).

Acknowledgements. I am grateful to my Ph.D advisor at IMPA, Fernando Codá Marques, for his constant encouragement and advice. I would like to thank the referees for their many useful comments. I was supported by FAPERJ and CNPq-Brasil.

2. THE SCHWARZSCHILD ANTI-DE SITTER SPACES AND ITS PERTURBATIONS

Let m be a real number. Let $\rho_m : (r_0, +\infty) \rightarrow \mathbb{R}$ be the function given by $\rho_m(r) = \sqrt{1 + r^2 - 2m/r}$, where $r_0 = r_0(m)$ is the unique positive zero of ρ_m (if $m > 0$) or 0 (if $m \leq 0$). Let (S^2, g_0) be the standard round sphere of constant Gaussian curvature 1 and let dS^2 denote its volume element.

We call *Schwarzschild anti-de Sitter space of mass m* the metric completion of the Riemannian manifold $((r_0, +\infty) \times S^2, g_m)$, where using the

natural r coordinate the metric g_m is written as

$$g_m = \frac{dr^2}{\rho_m^2(r)} + r^2 g_0 = \frac{dr^2}{1 + r^2 - \frac{2m}{r}} + r^2 g_0.$$

Although the expression of g_m in this coordinate system becomes singular at r_0 , it can be proved that when $m > 0$ the metric g_m extends to a (smooth) Riemannian metric on $M = [r_0, \infty) \times S^2$. On the other hand a smooth extension is not possible if $m < 0$. Notice also that if we let the parameter m to be zero we recover the hyperbolic space (this can be easily seen by performing the coordinate change $r = \sinh s$).

We will call *coordinate spheres* the surfaces $S_r = \{r\} \times S^2 \subset M$. The following proposition describes the geometry of (M, g_m) and of its coordinate spheres.

Proposition 1. (*Geometry of the Schwarzschild anti-de Sitter space of mass m*)

i) *The Ricci curvature of g_m is given by*

$$\text{Ric}_m = \left(-2 - \frac{2m}{r^3}\right) \frac{1}{\rho_m^2(r)} dr^2 + \left(-2 + \frac{m}{r^3}\right) r^2 g_0.$$

ii) *The scalar curvature of g_m is constant and equal to -6 .*

iii) *The coordinate spheres S_r are totally umbilic surfaces with constant mean curvature given by*

$$H_m(r) = \frac{2}{r} \sqrt{1 + r^2 - \frac{2m}{r}}.$$

iv) *The Hawking mass of all coordinate spheres is m .*

v) *The Jacobi operator of the coordinate sphere S_r is given by*

$$L_r = \frac{1}{r^2} \left(\Delta_0 + \left(2 - \frac{6m}{r} \right) \right),$$

where Δ_0 is the Laplacian operator of the round sphere (S^2, g_0) .

Proof. A calculation in coordinates. □

Notice that when $m > 0$ the Jacobi operator of S_r is invertible except for $r = 3m$. In any case, it is invertible when restricted to the space of zero mean value functions. However, a degeneration occurs when r goes to infinity: up to normalization, it becomes $\Delta_0 + 2$, which is no longer invertible in this restricted space.

From now on we assume $m > 0$. Let s be the function that gives the distance of a point of (M, g_m) to ∂M . Using $s \in [0, +\infty)$ as coordinate, one can write

$$(3) \quad g_m = \frac{dr^2}{1 + r^2 - \frac{2m}{r}} + r^2 g_0 = ds^2 + \sinh^2(s) v_m(s) g_0,$$

where v_m is a positive function defined on $[0, +\infty)$ that has the following expansion as s goes to infinity:

$$v_m(s) = 1 + \frac{2m}{3 \sinh^3 s} + O(\exp(-5s)).$$

Although we have explicit formulas as in Proposition 1 only when we use the r coordinate, it will be more convenient to use the s coordinate. We will then consider g_m to be defined on $M = [0, +\infty) \times S^2$ by formula (3) above, and as a small abuse of notation we use s both for the first coordinate of a point $p \in M$ and for the function $r \in (r_0, +\infty) \mapsto s(r) \in (0, +\infty)$ that gives the coordinate change described above. It is worth noting that, as a function of s , the r coordinate expands as $r = \sinh(s)(1 + O(\exp(-3s)))$ as s goes to infinity. In particular, for example, the mean curvature of the coordinate spheres $S_s := S_{r(s)}$ behaves as

$$H_m(s) = 2 \frac{\cosh s}{\sinh s} - \frac{2m}{\sinh^3 s} + O(\exp(-5s)) \quad \text{as } s \text{ goes to infinity.}$$

To conclude our description of the Schwarzschild anti-de Sitter space, it may be useful to remark that the double of M , $\hat{M} = (-\infty, +\infty) \times S^2$, can be endowed with a (smooth) Riemannian metric whose restriction to $M = [0, +\infty) \times S^2$ is the metric g_m and such that the involution $i : (s, x) \in \hat{M} \mapsto (-s, x) \in \hat{M}$ is an isometry fixing $\{0\} \times S^2$, a surface identified with the totally geodesic boundary of (M, g_m) . We still denote this metric by g_m .

Now we define the class of metrics on $M = [0, +\infty) \times S^2$ we are going to work with.

Definition. Given $m > 0$, let $\mathcal{M}(M, m)$ be the set of metrics g of class C^3 on $M = [0, +\infty) \times S^2$ such that

- a) $\partial M = \{0\} \times S^2$ is a minimal surface in (M, g) ; and
- b) There exists a constant $C > 0$ such that for every $p = (s, x) \in M$,

$$(\|g - g_m\| + \|\nabla^m g\| + \|(\nabla^m)^2 g\| + \|(\nabla^m)^3 g\|)(p) \leq C \exp(-4s).$$

Here the norm is calculated with respect to the metric g_m and ∇^m denotes the Levi-Civita connection of g_m .

The space $\mathcal{M}(M, m)$ has a distance function

$$d(g_1, g_2) := \sup_{p \in M} \left(\sum_{i=0}^3 \exp(4s) \|(\nabla^m)^i (g_1 - g_2)\|_{g_m}(p) \right).$$

We remark that each metric in $\mathcal{M}(M, m)$ is asymptotically hyperbolic with total mass m , according to the definitions of [24] and [6]. The control of g up to the third derivative is needed in order to apply the results of [22].

Observe that we do not assume a priori that ∂M is outermost.

Given $g \in \mathcal{M}(M, m)$ one can calculate the expansions of its Ricci tensor, its scalar curvature and the mean curvature of the coordinate spheres in (M, g) as follows: one adds terms of order $O(\exp(-4s))$ to the expansion of

the corresponding quantities of (M, g_m) in the s coordinate.

We finish this section by discussing the geometry of surfaces in (M, g) , $g \in \mathcal{M}(M, m)$.

In this paper, we consider closed surfaces $\Sigma \subset M = [0, +\infty) \times S^2$ such that $M \setminus \Sigma$ has two connected components, one of them containing ∂M . We define the inner radius and the outer radius of such Σ to be

$$\underline{s} := \min\{s(x); x \in \Sigma\} \quad \text{and} \quad \bar{s} := \max\{s(x); x \in \Sigma\},$$

respectively. We also use the convention that the unit normal vector N points toward the unbounded component of $M \setminus \Sigma$ and that the mean curvature is the trace of the second fundamental form A given by $A(X, Y) := g(\nabla_X N, Y)$ for every pair of vectors X, Y tangent to Σ .

A constant mean curvature surface Σ in (M, g) is called weakly stable when its Jacobi operator,

$$L_\Sigma := \Delta_\Sigma + Ric(N, N) + |A|^2,$$

is such that $-\int_\Sigma L_\Sigma(\phi)\phi d\Sigma \geq 0$ for all functions $\phi \in C^\infty(\Sigma)$ with $\int_\Sigma \phi d\Sigma = 0$. This analytic definition is equivalent to the geometric one that the second variation of the area of Σ under volume preserving variations is non-negative.

Throughout this paper, we will frequently consider surfaces in (M, g) that are graphical over a coordinate sphere S_s . Given $u \in C^2(S^2)$, we write

$$S_s(u) := \{(s + u(x), x) \in M; x \in S^2\}.$$

We denote by $H(s, u, g)$ the mean curvature of $S_s(u)$ in (M, g) , which we consider to be a real function defined on S^2 .

Remark 1. In the Section 3, it will be important to understand how $H(s, u, g)$ depends on the variables s, u and g . One way to analyze this is the following. Choose $(x^0 = s, x^1, x^2)$ local coordinates on M , where $x = (x^1, x^2)$ are local coordinates on S^2 . The local parametrization $x \mapsto (x^0 + u(x^1, x^2), x^1, x^2)$ of $S_s(u)$ gives explicit tangent vectors to $S_s(u) \subset (M, g)$, whose coordinates depends only on $\partial_i u(x)$ for $i = 1, 2$. From this, we conclude that the coordinates of the normal field N to $S_s(u)$ in (M, g) are smooth functions of $\partial_i u(x)$ and $g_{AB}(x^0 + u(x^1, x^2), x^1, x^2)$ for all $i, j = 1, 2$, $A, B = 0, 1, 2$. By the definition of the mean curvature, one then verify that $H(s, g, u)(x)$ is a smooth function of $\partial_i \partial_j u(x)$, $\partial_i u(x)$, $g_{AB}(x^0 + u(x), x^1, x^2)$ and $\partial_A g_{BC}(x^0 + u(x), x^1, x^2)$ for all $i, j = 1, 2$, $A, B, C = 0, 1, 2$. From this rough analysis, for example, we see that taking derivatives of $H(s, u, g)$ with respect to the s and u variables requires the functions g_{AB} to be twice differentiable.

3. FOLIATION OF COMPACT REGIONS

For metrics $g \in \mathcal{M}(M, m)$ that are close enough to g_m , we use the implicit function theorem to construct a family of weakly stable CMC spheres on compact regions of (M, g) .

For the rest of the paper, we fix some $\alpha \in (0, 1)$.

Theorem 2. *Let $m > 0$. Given $S > 0$, there exists $\eta_0 > 0$ such that for all $\eta \in (0, \eta_0)$ there exists $\epsilon > 0$ with the following property:*

For every $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ and for every $s \in [0, S]$ there exists a unique function $u(s, g) \in C^{2,\alpha}(S^2)$ with $\|u(s, g)\|_{C^{2,\alpha}} < \eta$ and $\int_{S^2} u(s, g) dS^2 = 0$ such that the surface

$$\Sigma_s(g) := S_s(u(s, g)) = \{(s + u(s, g)(x), x) \in M; x \in S^2\}$$

has constant mean curvature with respect to the metric g .

Moreover, for every $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$, $\Sigma_0(g) = \partial M$ and the family $\{\Sigma_s(g)\}_{s \in [0, S]}$ gives a foliation of a compact region of (M, g) by weakly stable CMC spheres, with positive mean curvature if $s \in (0, S]$.

Finally, when $S > s(3m)$, given any constant $\kappa > 0$ and any compact interval $I \subset (s(3m), S]$, it is possible to choose the ϵ above in such way that for every $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ the mean curvature $H_g(s)$ of $\Sigma_s(g)$ in (M, g) is monotone decreasing on I and satisfies $|H_g(s) - H_m(s)| < \kappa$ for every $s \in I$.

Proof. For a minor technical reason, choose $\theta > 0$. In order to consider the geometry of graphs of small functions over ∂M , we have to extend each metric in $\mathcal{M}(M, m)$ to a fixed neighborhood of M in $\hat{M} = (-\infty, +\infty) \times S^2$. It is always possible to extend $g \in \mathcal{M}(M, m)$ as a C^3 metric \tilde{g} on $\tilde{M} := [-\theta, +\infty) \times S^2$ in such way that $\|\tilde{g} - g_m\|_{C^3(\tilde{M})} \leq \lambda \|g - g_m\|_{C^3(M)}$ for some fixed $\lambda > 1$. In particular, if $d(g, g_m) < \epsilon$ then $\|\tilde{g} - g_m\|_{C^3(\tilde{M})} < \lambda\epsilon$. For simplicity we refer to this extension of g using the same letter.

On the round unit sphere (S^2, g_0) , consider the Banach spaces

$$E = \{u \in C^{2,\alpha}(S^2); \int_{S^2} u dS^2 = 0\} \text{ and } F = \{u \in C^{0,\alpha}(S^2); \int_{S^2} u dS^2 = 0\},$$

endowed with their standard norms.

Given $\eta_0 \in (0, \theta)$, for every $s \in [0, S]$ and every $u \in E$ with $\|u\|_{C^{2,\alpha}} < \eta_0$ we can consider the surfaces

$$S_s(u) = \{(s + u(x), x) \in \tilde{M}; x \in S^2\}.$$

Let $H(s, u, g)$ be the mean curvature of $S_s(u)$ with respect to a metric $g \in \mathcal{M}(M, m)$. It defines a C^1 map from $[0, S] \times (B(0, \eta_0) \subset E) \times \mathcal{M}(M, m)$ to $C^{0,\alpha}(M)$, see Remark 1 in the end of Section 2.

Let $\Phi : [0, S] \times \mathcal{M}(M, m) \times (B(0, \eta_0) \subset E) \rightarrow F$ be the C^1 map given by

$$\Phi(s, g, u) = H(s, u, g) - \frac{1}{4\pi} \int_{S^2} H(s, u, g) dS^2.$$

Observe that $\Phi(s, g, u) = 0$ if and only if $S_s(u)$ is a CMC surface in (\tilde{M}, g) . In particular, $\Phi(s, g_m, 0) = 0$ for all $s \in [0, S]$.

We claim that, for every $s \in [0, S]$, $D\Phi_{(s, g_m, 0)}$ is an isomorphism when restricted to E . In fact, for every $v \in E$, the family $t \mapsto S_s(tv)$ is a normal

variation of the coordinate sphere S_s in (M, g_m) with speed v . Therefore, if L_s is the Jacobi operator of S_s with respect to g_m , we have

$$\frac{d}{dt}\Big|_{t=0} \Phi(s, g_m, tv) = -L_s(v) + \frac{1}{4\pi} \int_{S^2} L_s(v) dS^2 = -L_s(v).$$

The last equality follows because $L_s(v) = (1/r^2)(\Delta_0 + (2 - 6m/r))(v)$ (see Proposition 1, considering the coordinate change $r = r(s)$) and v has zero mean value. Since $m > 0$, $\Delta_0 + (2 - 6m/r)$ is an invertible operator from E to F for all $s \in [0, S]$ and the claim follows.

Therefore we can apply the implicit function theorem: there exists (a possibly smaller) $\eta_0 > 0$ such that for all $\eta \in (0, \eta_0)$, there exists a small ball B around (s, g_m) in $[0, S] \times \mathcal{M}(M, m)$ and a C^1 function $(\tilde{s}, g) \in B \mapsto u(\tilde{s}, g) \in B(0, \eta) \subset E$ such that $u(s, g_m) = 0$ and $u(\tilde{s}, g)$ is uniquely defined in $B(0, \eta)$ by the equation $\Phi(\tilde{s}, g, u(\tilde{s}, g)) = 0$ for all $(\tilde{s}, g) \in B$.

Since u is a C^1 function of s and g , by compactness we can choose η_0 in such way that for all $\eta \in (0, \eta_0)$ there exists $\epsilon > 0$ such that u is uniquely defined on $[0, S] \times \{g \in \mathcal{M}(M, m); d(g, g_m) < \epsilon\}$ and takes values on $B(0, \eta) \subset E$. Thus for each metric $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ we have constructed a family

$$\{\Sigma_s(g)\}_{s \in [0, S]} := \{S_s(u(s, g))\}_{s \in [0, S]}$$

of CMC spheres in (\tilde{M}, g) , where $u(s, g) \in E$ has norm $\|u(s, g)\|_{C^{2,\alpha}} < \eta$. Notice that the surfaces $\Sigma_s(g)$ depend smoothly on s and g , and also that $\{\Sigma_s(g_m)\}$ is precisely the foliation of M by coordinate spheres.

Since $\partial M = S_0(0)$ is minimal for all metrics $g \in \mathcal{M}(M, m)$, the uniqueness of the function u above constructed implies that $u(0, g) = 0$, i.e., $\Sigma_0(g) = \partial M$ for every $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$.

In order to prove that $\{\Sigma_s(g)\}$ is a foliation of some region of M , we have to analyze the sign of its lapse function, that is, its normal speed. Since for g_m the constructed family $\{\Sigma_s(g_m)\}$ is a foliation, its lapse function is positive on $[0, S]$, hence the lapse function of $\{\Sigma_s(g)\}_{s \in [0, S]}$ with respect to all $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ is also positive, at least when we choose a possibly smaller ϵ . Since each of these families starts at $\partial M = \{0\} \times S^2 \subset \tilde{M}$, the families $\{\Sigma_s(g)\}_{s \in [0, S]}$ foliate a compact region of M .

To see that $\Sigma_g(s)$ has positive mean curvature in (M, g) for all $s \in (0, S]$, observe that this is true for $\Sigma_s(g_m)$ in (M, g_m) and also that, by Proposition 1, $H'_m(0) = -(1/r_0^2)(2 - 6m/r_0) > 0$. Moreover, $H_g(s) := H(s, u(s, g), g)$ is a C^1 function of s and g . Therefore by continuity we can arrange ϵ in such way that for every $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ the surface $\Sigma_s(g)$ has positive mean curvature in (M, g) for all $s \in (0, S]$.

Now we argue that the leaves are weakly stable. In fact, since $m > 0$, for every $s \in [0, S]$ the Jacobi operator L_s of $\Sigma_s(g_m)$ in (M, g_m) satisfies

$$-\int_{\Sigma_s(g_m)} L_s(\phi) \phi d\Sigma_s(g_m) = \int_{S^2} |\nabla_0 \phi|^2 - \left(2 - \frac{6m}{r(s)}\right) \phi^2 dS^2 \geq \frac{6m}{r(S)} \int_{S^2} \phi^2 dS^2$$

for every $\phi \in C^\infty(\Sigma_s(g_m))$ with $\int \phi d\Sigma_s(g_m) = 0$. Arguing by contradiction we then conclude that for a possibly smaller ϵ there exists a constant $c > 0$ such that for every $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ and every $s \in [0, S]$ the Jacobi operator $L_{(s,g)}$ of the surface $\Sigma_s(g)$ in (M, g) is such that

$$-\int_{\Sigma_s(g)} L_{(s,g)}(\phi)\phi d\Sigma_s(g) \geq c \int_{\Sigma_s(g)} \phi^2 d\Sigma_s(g) \geq 0$$

for every $\phi \in C^\infty(\Sigma_s(g))$ with $\int \phi d\Sigma_s(g) = 0$, i.e., $\Sigma_s(g)$ is weakly stable.

The last statement of the theorem also follows by continuity, since $H'_m(s) < 0$ on $(s(3m), +\infty)$. \square

Notice that in the previous result the mean curvature of the graph $\Sigma_s(g) = S_s(u(s, g))$ is not prescribed and can be different from $H_m(s)$. Having in mind the matching argument of Section 6, we finish this section showing the existence and uniqueness of a graph over the coordinate sphere S_s with prescribed mean curvature $H_m(s)$ in (M, g) , at least when $d(g, g_m)$ is sufficiently small and s is large enough. More precisely, we have:

Theorem 3. *Let $m > 0$. Given a compact interval $I \subset (s(3m), +\infty)$, there exists $\eta_0 > 0$ such that for all $\eta \in (0, \eta_0)$ there exists $\epsilon > 0$ with the following property:*

For every $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ and for every $s \in I$ there exists a unique function $h(s, g) \in C^{2,\alpha}(S^2)$ with $\|h(s, g)\|_{C^{2,\alpha}} < \eta$ such that the surface

$$S_s(h(s, g)) = \{(s + h(s, g)(x), x) \in M; x \in S^2\}$$

has constant mean curvature $H_m(s)$ in (M, g) .

Moreover, the family $\{S_s(h(s, g))\}_{s \in I}$ gives a foliation of a compact region of (M, g) by weakly stable CMC spheres.

Proof. Following the notations of Theorem 2, given $\eta_0 > 0$ sufficiently small we consider the C^1 map $\Psi : I \times \mathcal{M}(M, m) \times (B(0, \eta_0) \subset C^{2,\alpha}(S^2)) \rightarrow C^{0,\alpha}(S^2)$ given by $\Psi(s, g, u) = H(s, u, g) - H_m(s)$. Notice that $\Psi(s, g_m, 0) = 0$ for all $s \in I$.

The linearisation of Ψ at $(s, g_m, 0)$ is such that, for every $v \in C^{2,\alpha}(S^2)$,

$$D\Psi_{(s,g_m,0)}(0, 0, v) = -L_s(v) = -\frac{1}{r^2}(\Delta_0 + (2 - \frac{6m}{r})),$$

where we use the coordinate $r = r(s)$, see Proposition 1. Since $I \subset (s(3m), +\infty)$, the map $v \in C^{2,\alpha}(S^2) \mapsto L_s(v) \in C^{0,\alpha}(S^2)$ is an isomorphism for all $s \in I$. Hence, we can apply the implicit function theorem and the result follows by the same arguments presented in Theorem 2. \square

4. FOLIATION NEAR THE INFINITY

The next theorem is the version of the existence and uniqueness theorem of A. Neves and G. Tian [22] adapted to the asymptotically hyperbolic

manifolds (M, g) where g belongs to the space of metrics $\mathcal{M}(M, m)$ (we refer the reader to Theorem 2.2 and the proof of Theorem 8.2 in [22]).

Theorem 4 ([22]). *Let $m > 0$. Given $\epsilon_0 > 0$, there exists $\delta > 0$, $C > 0$ and $\underline{s}_0 > s(3m)$ with the following properties:*

- 1) *Given $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon_0$, for all $\ell \in (2, 2 + \delta)$, there exists a unique sphere $\Sigma_\ell = \Sigma_\ell(g) \subset M$ such that*
 - a) *$M \setminus \Sigma_\ell$ has two connected components, one of them containing ∂M ;*
 - b) *Σ_ℓ is a weakly stable constant mean curvature sphere in (M, g) with mean curvature $H = \ell$; and*
 - c) *The inner radius \underline{s}_ℓ and the outer radius \bar{s}_ℓ of Σ_ℓ satisfy*

$$\underline{s}_\ell \geq \underline{s}_0 \quad \text{and} \quad \bar{s}_\ell - \underline{s}_\ell \leq 1.$$

- 2) *The family $\{\Sigma_\ell\}_{\ell \in (2, 2 + \delta)}$ gives a smooth foliation of the complement of a compact set of M and $\lim_{\ell \rightarrow 2} \underline{s}_\ell = +\infty$.*
- 3) *Given $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon_0$ and $\ell \in (2, 2 + \delta)$, if for the above surface Σ_ℓ in (M, g) we define \hat{s}_ℓ by the equality*

$$|\Sigma_\ell| = 4\pi \sinh^2 \hat{s}_\ell,$$

then:

- a) *If we set $w_\ell(p) = s(p) - \hat{s}_\ell$ for $p \in \Sigma_\ell$, then*

$$\sup_{\Sigma_\ell} |w_\ell| \leq C \exp(-\underline{s}_\ell) \quad \text{and} \quad \int_{\Sigma_\ell} |\partial_s^\top|^2 d\Sigma_\ell \leq C \exp(-2\underline{s}_\ell).$$

b)

$$\int_{\Sigma_\ell} |\mathring{A}_\ell|^2 d\Sigma_\ell \leq C \exp(-4\underline{s}_\ell).$$

- c) *There exists a function $f \in C^2(S^2)$ with $\|f\|_{C^2} \leq C$ such that*

$$\Sigma_\ell = S_{\hat{s}_\ell}(f) = \{(\hat{s}_\ell + f(x), x) \in M; x \in S^2\}.$$

Remark 2. Using the terminology of [22], the metrics $g \in \mathcal{M}(M, m)$ satisfy condition (H) with same constants r_1, C_2 and C_3 , and the constant C_1 depends only on $d(g, g_m)$. Therefore, if we fix the constant $C_4 = 1$, all constants appearing in Neves and Tian's theorem depend only on the distance between the metric $g \in \mathcal{M}(M, m)$ and g_m . This uniform dependence on $d(g, g_m)$ is crucial to the matching argument.

Remark 3. Theorem 4 is proved by the continuity method (Theorem 8.2 in [22]). A fundamental step is to prove that a weakly stable CMC sphere in (M, g) with mean curvature ℓ close enough to 2 and large enough inner radius is graphical over *some* coordinate sphere (this is the content of item 3) of the above theorem) and that its Jacobi operator is invertible (see Proposition 8.1 in [22]). To prove the *existence* of the foliation, following the argument of [22], page 92, given $g \in \mathcal{M}(M, g)$ with $d(g, g_m) < \epsilon_0$, consider the family of metrics $g_t = (1 - t)g_m + tg \in \mathcal{M}(M, g)$, $t \in [0, 1]$. The rough idea is then

to prove that for each $s \in (s(3m), +\infty)$ large enough the set A of $t \in [0, 1]$ such that there exists a weakly stable CMC sphere $\Sigma \subset (M, g_t)$ with mean curvature $H_m(s)$ and large enough inner radius is open and closed.

The set A clearly contains $t = 0$. The proof that A is open follows the reasoning of the proof of Theorem 3: given t in this set, since the Jacobi operator of Σ in (M, g) is invertible and since Σ is a graph over a coordinate sphere, the implicit function theorem implies the existence of small graphs over the *same* coordinate sphere which have the *same* mean curvature with respect to nearby g_t metrics. The proof that the set A is closed uses the uniform estimates on the norm of the function that gives the graph (see item 3) of the above theorem).

Since the CMC sphere in (M, g_m) with mean curvature $H_m(s)$ is precisely the coordinate sphere S_s , we conclude the following. Given $\epsilon_0 > 0$, let δ, \bar{s}_0 and C be given by Theorem 4. Let $I \subset (\bar{s}_0, +\infty)$ be a compact interval such that $H_m(s) \in (2, 2 + \delta)$ for all $s \in I$. Then, the corresponding $\eta_0 > 0$ given by Theorem 3 is such that for every $\eta \in (0, \eta_0)$ there exists $\epsilon \in (0, \epsilon_0)$ such that for every $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ the surfaces of the foliation $\{\Sigma_\ell(g)\}$ described in Theorem 4 with mean curvature $\ell = H_m(s) \in (2, 2 + \delta)$ for $s \in I$ are exactly the surfaces obtained in Theorem 3.

Observe that the above argument does not make any reference to the *uniqueness* statement of Theorem 4. And since it contains all the information needed for the matching argument, we will use only it in the proof of Theorem 7 (see Section 6).

5. LIMIT OF THE HAWKING MASS

Let g be a metric in $\mathcal{M}(M, m)$ with scalar curvature $R \geq -6$. Recall that the Hawking mass of a closed surface Σ in (M, g) is

$$m_H(\Sigma) = \sqrt{\frac{|\Sigma|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma} (H^2 - 4) d\Sigma \right).$$

We want to calculate the limit of the right hand side of the above expression for the weakly stable CMC spheres Σ_ℓ in (M, g) given by Theorem 4 as they approach the infinity. In order to do this, we need the following consequence of Gauss equation (see [21]).

Lemma 5. *Given $g \in \mathcal{M}(M, m)$, let $\{\Sigma_t\}_{t>t_0}$ be a family of constant mean curvature spheres in (M, g) such that $\underline{s}_t \rightarrow +\infty$ as t goes to infinity. Then*

$$(H_t^2 - 4)|\Sigma_t| = 16\pi - \int_{\Sigma_t} \frac{8m - 12m|\partial_s^\top|^2}{\sinh^3 s} d\Sigma_t + 2 \int_{\Sigma_t} |\dot{A}_t|^2 d\Sigma_t + |\Sigma_t| O(\exp(-4\underline{s}_t)).$$

Proof. Let K_t be the Gaussian curvature of Σ_t . The Gauss equation for Σ_t in (M, g) can be written as

$$\begin{aligned} (4) \quad 2K_t &= R - 2\text{Ric}(N_t, N_t) + H_t^2 - |A_t|^2 \\ &= (R + 6) - 2(\text{Ric}(N_t, N_t) + 2) + \frac{H_t^2 - 4}{2} - |\dot{A}_t|^2. \end{aligned}$$

By Proposition 1, for metrics $g \in \mathcal{M}(M, m)$, if $\{\partial_s, e_1, e_2\}$ is a g_m -orthonormal referential, then

$$\begin{aligned} Ric(\partial_s, \partial_s) &= -2 - \frac{2m}{\sinh^3 s} + O(\exp(-4s)), \\ Ric(e_i, e_j) &= \left(-2 + \frac{m}{\sinh^3 s}\right) \delta_{ij} + O(\exp(-4s)), \\ Ric(\partial_s, e_i) &= O(\exp(-4s)), \quad \text{and} \\ R + 6 &= O(\exp(-4s)). \end{aligned}$$

Considering the g_m -orthogonal decomposition $N_t = a\partial_s + X$, it follows that

$$(5) \quad 4K_t = (H_t^2 - 4) + \left(\frac{8m - 12m|X|_{g_m}^2}{\sinh^3 s}\right) - 2|\dot{A}_t|^2 + O(\exp(-4s)).$$

Observe that if ν is the unit normal of a coordinate sphere in (M, g) , then $\partial_s = \nu + W$, where $g(\nu, W) = O(\exp(-4s))$ and $|W|_g = O(\exp(-4s))$. Hence, the g -orthogonal decomposition $N_t = b\nu + Y$ is such that $|X|_{g_m}^2 = |Y|_g^2 + O(\exp(-4s))$. On the other hand, if $\nu = bN_t + \nu^\top$ is the g -orthonormal decomposition of ν corresponding to the tangent space of Σ_t and its g -normal N_t , we have $|Y|_g^2 = |\nu^\top|_g^2$. Therefore one can change $|X|_{g_m}^2$ by $|\partial_s^\top|_g^2$ in (5).

Since Σ_t is a sphere of constant mean curvature, the lemma follows after integration of (5). \square

Proposition 6. *Let $m > 0$. Given $g \in \mathcal{M}(M, m)$, the family $\{\Sigma_\ell\}_{\ell \in (2, 2+\delta)}$ in (M, g) given by Theorem 4 is such that*

$$\lim_{\underline{s}_\ell \rightarrow +\infty} \sqrt{\frac{|\Sigma_\ell|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma_\ell} (H_\ell^2 - 4) d\Sigma_\ell\right) = m.$$

Proof. Let \hat{s}_ℓ be defined by $|\Sigma_\ell| = 4\pi \sinh^2 \hat{s}_\ell$. We use the information given by Theorem 4, item 3), and calculate all expansions as \underline{s}_ℓ goes to infinity.

Lemma 5 and the estimates of Theorem 4, item 3) on the behavior of $|\partial_s^\top|$ and $|\dot{A}_\ell|$ implies

$$(6) \quad \begin{aligned} &\sqrt{\frac{|\Sigma_\ell|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma_\ell} (H_\ell^2 - 4) d\Sigma_\ell\right) = \\ &= \frac{|\Sigma_\ell|^{1/2}}{8\pi^{3/2}} \left(\int_{\Sigma_\ell} \frac{m}{\sinh^3 s} d\Sigma_\ell + |\Sigma_\ell| O(\exp(-4\underline{s}_\ell)) + O(\exp(-4\underline{s}_\ell))\right). \end{aligned}$$

In order to analyze (6), recall that Σ_ℓ is a graph over the coordinate sphere $S_{\hat{s}_\ell}$ given by a function f whose C^2 norm is bounded for all $\ell \in (2, 2+\delta)$ by the same constant $C > 0$. Therefore $|\underline{s}_\ell - \hat{s}_\ell| = |\min_{x \in S^2} \{f(x)\}| \leq C$ for all ℓ , which implies

$$(7) \quad |\Sigma_\ell| = 4\pi \sinh^2 \hat{s}_\ell = O(\exp(2\underline{s}_\ell)).$$

On the other hand, for every $p = (s, x) \in \Sigma_\ell$, since $|w_\ell(p)| \leq C \exp(-\underline{s}_\ell)$,

$$\frac{\sinh \hat{s}_\ell}{\sinh s} = \frac{\sinh(s - w_\ell(p))}{\sinh s} = \cosh w_\ell(p) - \frac{\cosh s}{\sinh s} \sinh w_\ell(p) = 1 + O(\exp(-\underline{s}_\ell)).$$

Therefore

$$(8) \quad \int_{\Sigma_\ell} \frac{|\Sigma_\ell|^{1/2}}{\sinh^3(s)} d\Sigma_\ell = \frac{(4\pi)^{3/2}}{|\Sigma_\ell|} \int_{\Sigma_\ell} \left(\frac{\sinh \hat{s}}{\sinh s} \right)^3 d\Sigma_\ell = 8\pi^{3/2}(1 + O(\exp(-\underline{s}_\ell))).$$

The proposition follows from (6), (7) and (8). \square

6. THE GLOBAL FOLIATION

We now argue that when a metric $g \in \mathcal{M}(M, m)$ is sufficiently close to g_m the foliations of (M, g) constructed in Sections 3 and 4 match on their overlap. It is possible to argue using the uniqueness of Neves and Tian's foliation, but we follow a different reasoning using the implicit function theorem, see Remark 3 in Section 4.

Theorem 7. *Let $m > 0$. There exists $\epsilon > 0$ with the following property:*

If $g \in \mathcal{M}(M, m)$ is such that $d(g, g_m) < \epsilon$, then there exists a foliation $\{\Sigma_t\}_{t \in [0, +\infty)}$ of M such that:

- i) Each Σ_t is a weakly stable CMC sphere in (M, g) , with positive mean curvature when $t > 0$;*
- ii) $\Sigma_0 = \partial M$ is an outermost minimal surface in (M, g) ; and*
- iii) $\lim_{t \rightarrow +\infty} \sqrt{\frac{|\Sigma_t|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma_t} (H_t^2 - 4) d\Sigma_t \right) = m$.*

Proof. Given an arbitrary $\epsilon_0 > 0$, let $\delta > 0$, $C > 0$ and $\underline{s}_0 > s(3m)$ be given by Theorem 4. Recall that $H_m(s)$ is a smooth monotone decreasing function on the interval $(s(3m), +\infty)$ and converges to 2 as s goes to infinity. Let $s_0 > \underline{s}_0$ be such that $H_m(s_0) < 2 + \delta$.

Let $S > s_0 + 1$. Given this choice of S , we can choose $\eta < 1$ and $\epsilon \in (0, \epsilon_0)$ sufficiently small in such way that Theorem 2 holds for S and Theorem 3 holds for the interval $[S - 1, S]$ for every metric $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$.

In particular, we can assume that for every metric $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ and for every $s \in [S - 1, S]$ the surface $\Sigma_\ell = \Sigma_\ell(g)$ in (M, g) described in Theorem 4 with $\ell = H_m(s)$ is given by

$$\Sigma_\ell = S_s(h),$$

where h is the unique function in $C^{2,\alpha}(S^2)$ with norm $< \eta$ such that its graph over S_s has constant mean curvature $H_m(s)$, see the remarks after Theorem 4.

Given $\kappa \in (0, \eta)$, let $[c, d]$ be the image of the interval $[S - 3\kappa/4, S - \kappa/4]$ under the map $H_m(s)$. We can moreover assume that $\epsilon \in (0, \epsilon_0)$ is sufficiently small in such way that for every $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$

the foliation $\{\Sigma_s^1(g)\}_{s \in [0, S]}$ constructed in Theorem 2 has in particular the following properties:

- a) For every $s \in [S - 1, S]$, there exists a function $u(s, g) \in C^{2, \alpha}(S^2)$ with $\|u(s, g)\|_{C^{2, \alpha}} < \eta/2$ such that

$$\Sigma_s^1(g) = S_s(u(s, g)).$$

- b) The mean curvature $H_g(s)$ of $\Sigma_s^1(g)$ in (M, g) is a decreasing function on the interval $[S - 1, S]$ with

$$|H_g(s) - H_m(s)| < (d - c)/4$$

for all $s \in [S - 1, S]$. Hence, there exists an interval $(a, b) \subset [S - 3\kappa/4, S - \kappa/4]$ such that for every $s \in (a, b)$ there exists a unique $\tilde{s} \in (S - 3\kappa/4, S - \kappa/4)$ with $H_g(s) = H_m(\tilde{s})$. This defines \tilde{s} as a function of s , which is of class C^1 since $H_g(s)$ is a C^1 function of s .

We now prove that the theorem is true for this choice of ϵ .

In fact, fix some $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$. By item a) and b) above, given $s \in (a, b)$, if we define the function $\tilde{h} = s - \tilde{s}(s) + u(s, g) \in C^{2, \alpha}(S^2)$, then

$$\|\tilde{h}\|_{C^{2, \alpha}} \leq |s - \tilde{s}(s)| + \|u\|_{C^{2, \alpha}} < \kappa/2 + \eta/2 < \eta$$

and the graph

$$S_{\tilde{s}(s)}(\tilde{h}) = S_s(u(s, g)) = \Sigma_s^1(g)$$

has constant mean curvature $H_g(s) = H_m(\tilde{s}(s))$. These are the conditions that uniquely characterize the function that gives Σ_ℓ in (M, g) with $\ell = H_m(\tilde{s}(s))$ as a graph over $S_{\tilde{s}(s)}$. Therefore $\Sigma_s^1(g) = \Sigma_\ell(g)$ where $\ell = H_m(\tilde{s}(s))$ for all $s \in (a, b)$.

This proves that the foliations $\{\Sigma_s^1(g)\}$ and $\{\Sigma_\ell(g)\}$ given by Theorems 2 and 4, respectively, match on their overlap. The obtained foliation, which we reparametrize as $\{\Sigma_t\}_{t \in [0, +\infty)}$, is a foliation of (M, g) by weakly stable CMC spheres, starting at the minimal $\Sigma_0 = \partial M$, such that each Σ_t has positive mean curvature for $t > 0$ and such that

$$\lim_{t \rightarrow +\infty} \sqrt{\frac{|\Sigma_t|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma_t} (H_t^2 - 4) d\Sigma_t \right) = m,$$

see Theorem 2, Theorem 4 and Proposition 6. It remains only to prove that ∂M is an outermost minimal surface in (M, g) . This is a consequence of the Maximum Principle, since we proved $M \setminus \partial M$ is foliated by surfaces with positive mean curvature. \square

7. THE PENROSE INEQUALITY FOR PERTURBATIONS OF THE SCHWARZSCHILD ANTI-DE SITTER SPACES OF POSITIVE MASS

Using the foliation by weakly stable CMC spheres constructed above on (M, g) , $g \in \mathcal{M}(M, m)$ sufficiently close to g_m , and the remark of H. Bray

that the Hawking mass is monotone along such families (see [1]), we prove the Penrose inequality for this class of asymptotically hyperbolic manifolds.

Theorem 8. *Given $m > 0$, let $\epsilon > 0$ be given by Theorem 7. If $g \in \mathcal{M}(M, m)$ with $d(g, g_m) < \epsilon$ has scalar curvature $R \geq -6$, then*

$$(9) \quad \left(\frac{|\partial M|}{16\pi} \right)^{\frac{1}{2}} + 4 \left(\frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}} \leq m.$$

Moreover, equality holds if and only if (M, g) is isometric to (M, g_m) .

Proof. Let $\{\Sigma_t\}_{t \geq 0}$ be the foliation of (M, g) by weakly stable CMC spheres constructed in Theorem 7.

By assumption, g has scalar curvature $R \geq -6$. Then the Hawking mass of Σ_t ,

$$m_H(\Sigma_t) = \sqrt{\frac{|\Sigma_t|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma_t} (H_t^2 - 4) d\Sigma_t \right),$$

is monotone non-decreasing in t . In fact, we can be more precise:

Claim: $m'_H(\Sigma_t) \geq 0$. Moreover, $m'_H(\Sigma_t)$ is zero for $t > 0$ if and only if Σ_t satisfies the following properties:

- a) R is constant and equal to -6 along Σ_t ;
- b) Σ_t is totally umbilic; and
- c) Σ_t has constant Gaussian curvature.

The proof of the claim goes as follows. Choose a parametrization of this foliation by some function $F : [0, +\infty) \times S^2 \rightarrow M$ such that for each $t \in [0, +\infty)$, $F_t : S^2 \rightarrow M$ is a parametrization of Σ_t and $\partial_t F$ does not vanish. Let ρ_t be the lapse function, i.e., $\rho_t = g(N_t, \partial_t F)$ where N_t is the unit normal of the surface Σ_t pointing to infinity. Since we have a foliation, $\rho_t > 0$ on Σ_t for all t . We also define the mean value $\bar{\rho}_t = \int \rho_t d\Sigma_t / |\Sigma_t|$.

By the first variation formula of the Hawking mass (see, for example, [18]), we have

$$(16\pi)^{\frac{3}{2}} m'_H(\Sigma_t) = 2|\Sigma_t|^{\frac{1}{2}} \int_{\Sigma_t} (\Delta_t H_t + Q_t H_t) \rho_t d\Sigma_t,$$

where

$$Q_t = \frac{1}{2}(R + 6) + \left(\frac{4\pi}{|\Sigma_t|} - K_t \right) + \frac{1}{2} \left(|A_t|^2 - \frac{1}{2|\Sigma_t|} \int_{\Sigma_t} H_t^2 d\Sigma_t \right).$$

Since H_t is constant for each t ,

$$\begin{aligned} (16\pi)^{\frac{3}{2}} m'_H(\Sigma_t) &= 2|\Sigma_t|^{\frac{1}{2}} H_t \int_{\Sigma_t} Q_t \rho_t d\Sigma_t \\ &= 2|\Sigma_t|^{\frac{1}{2}} H_t \int_{\Sigma_t} (\Delta_t + Q_t)(\rho_t - \bar{\rho}_t) d\Sigma_t + 2|\Sigma_t|^{\frac{1}{2}} H_t \bar{\rho}_t \int_{\Sigma_t} Q_t d\Sigma_t \\ &= 2|\Sigma_t|^{\frac{1}{2}} H_t \int_{\Sigma_t} L_t(\rho_t - \bar{\rho}_t) d\Sigma_t + 2|\Sigma_t|^{\frac{1}{2}} H_t \bar{\rho}_t \int_{\Sigma_t} Q_t d\Sigma_t. \end{aligned}$$

In the last line, we used the Gauss equation (4) to see that the operator $\Delta_t + Q_t$ and the Jacobi operator $L_t = \Delta_t + Ric(N_t, N_t) + |A_t|^2$ of Σ_t differ by a constant.

Since $\partial M = \Sigma_0$ is minimal, the derivative of $m_H(\Sigma_t)$ is zero for $t = 0$. When $t > 0$, Σ_t has positive mean curvature. Observe also that $\int_{\Sigma_t} Q_t d\Sigma_t \geq 0$, since $R \geq -6$ and Σ_t is a sphere. Therefore

$$(16\pi)^{\frac{3}{2}} m'_H(\Sigma_t) \geq 2|\Sigma_t|^{\frac{1}{2}} H_t \int_{\Sigma_t} L_t(\rho_t - \bar{\rho}_t) d\Sigma_t.$$

Now we use the weak stability of Σ_t . Since H_t is constant for each t , $L_t(\rho_t) = -\partial_t H_t$ is also constant on Σ_t . Hence the stability inequality gives

$$0 \leq - \int_{\Sigma_t} L_t(\rho_t - \bar{\rho}_t)(\rho_t - \bar{\rho}_t) d\Sigma_t = \int_{\Sigma_t} L_t(\bar{\rho}_t)(\rho_t - \bar{\rho}_t) d\Sigma_t = \bar{\rho}_t \int_{\Sigma_t} L_t(\rho_t - \bar{\rho}_t) d\Sigma_t.$$

This implies that $m'_H(\Sigma_t)$ is non-negative. If $m'_H(\Sigma_t) = 0$ for $t > 0$, then

$$\int_{\Sigma_t} Q_t d\Sigma_t = 0 \quad \text{and} \quad \int_{\Sigma_t} L_t(\rho_t - \bar{\rho}_t)(\rho_t - \bar{\rho}_t) d\Sigma_t = 0.$$

The first equality implies that Σ_t satisfies a) and b). By the weak stability, the second equality implies that $L_t(\rho_t - \bar{\rho}_t)$ is constant, for it must be orthogonal to every function on Σ_t with zero mean value. Since $L_t(\rho_t)$ is constant, this implies that $L_t(\bar{\rho}_t)$ is also constant. Then c) follows from a), b) and the Gauss equation (4).

Once we proved the claim, inequality (9) follows immediately:

$$\left(\frac{|\partial M|}{16\pi} \right)^{\frac{1}{2}} + 4 \left(\frac{|\partial M|}{16\pi} \right)^{\frac{3}{2}} = m_H(\Sigma_0) \leq \lim_{t \rightarrow +\infty} m_H(\Sigma_t) = m.$$

Now we analyze the equality. In this case, $m_H(\Sigma_t)$ must be constant and equal to m . By the second part of the claim, each Σ_t satisfies a), b) and c) for all $t > 0$. Then, possibly after a change of the parametrization $F : [0, +\infty) \times S^2 \rightarrow M$, F^*g is a metric on $M = [0, +\infty) \times S^2$ that can be written in the form $ds^2 + V^2(s)g_0$, has constant scalar curvature $R = -6$, and is such that all slices $\{s\} \times S^2$ have Hawking mass m . These conditions uniquely characterize the metric g_m . And this finishes the proof. \square

Another Penrose inequality. In Proposition 1, we saw that the mean curvature of the coordinate spheres in the Schwarzschild anti-de Sitter spaces of mass $m > 0$ is given by the function $H_m(r) = (2/r)\sqrt{1 + r^2 - 2m/r}$. Observe that $H_m(2m) = 2$ and that $H_m(r) > 2$ for all $r > 2m$. In particular, the Maximum Principle implies that there are no other closed surfaces with constant mean curvature 2 in $([2m, +\infty) \times S^2, g_m)$.

Let (M, g) be an asymptotically hyperbolic three-manifold with connected boundary ∂M . Assume that (M, g) has scalar curvature $R \geq -6$ and that ∂M is an outermost $H = 2$ surface, meaning that there are no closed surfaces in M with constant mean curvature $H = 2$ other than ∂M . In this setting, the

Penrose Conjecture is that the area of ∂M and the total mass m of (M, g) are related by the inequality

$$\left(\frac{|\partial M|}{16\pi}\right)^{\frac{1}{2}} \leq m,$$

and that equality holds if and only if (M, g) is isometric to the piece of the Schwarzschild anti-de Sitter space of mass m outside the region bounded by the coordinate sphere of mean curvature 2.

Given $m > 0$ and $M_2 = [s(2m), +\infty) \times S^2$, we analogously define the space $\mathcal{M}(M_2, m)$ of C^3 metrics g on M_2 such that $\partial M_2 = \{s(2m)\} \times S^2$ has constant mean curvature 2 in (M_2, g) and

$$d(g, g_m) = \sup_{p \in M_2} \left(\sum_{i=0}^3 \exp(4s) \|(\nabla^m)^i (g - g_m)\|_{g_m}(p) \right) < +\infty$$

The analogous versions of Theorem 7 and Theorem 8 follows immediately by the same arguments:

Theorem 9. *Let $m > 0$. There exists $\epsilon > 0$ with the following property:*

If $g \in \mathcal{M}(M_2, m)$ is such that $d(g, g_m) < \epsilon$, then there exists a foliation $\{\Sigma_t\}_{t \in [0, +\infty)}$ of M_2 such that:

- i) Each Σ_t is a weakly stable CMC sphere in (M_2, g) , with mean curvature $H_t > 2$ when $t > 0$;*
- ii) $\Sigma_0 = \partial M_2$ is an outermost $H = 2$ surface in (M_2, g) ; and*
- iii) $\lim_{t \rightarrow +\infty} \sqrt{\frac{|\Sigma_t|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma_t} (H_t^2 - 4) d\Sigma_t\right) = m$.*

Theorem 10. *Given $m > 0$, let $\epsilon > 0$ be given by Theorem 9. If $g \in \mathcal{M}(M_2, m)$ with $d(g, g_m) < \epsilon$ has scalar curvature $R \geq -6$, then*

$$\left(\frac{|\partial M_2|}{16\pi}\right)^{\frac{1}{2}} \leq m.$$

Moreover, equality holds if and only if (M_2, g) is isometric to (M_2, g_m) .

REFERENCES

- [1] H. Bray, *The Penrose inequality in general relativity and volume comparison theorems involving scalar curvature*, Thesis, Stanford University (1997).
- [2] H. Bray, *Proof of the Riemannian Penrose inequality using the positive mass theorem*, J. Differential Geom. 59 (2001), no. 2, 177-267.
- [3] H. Bray and P. Chruściel, *The Penrose inequality*, The Einstein equations and the large scale behavior of gravitational fields, Birkhäuser, Basel (2004), 39-70.
- [4] S. Brendle and O. Chodosh, *A volume comparison theorem for asymptotically hyperbolic manifolds*, Commun. Math. Phys, vol 332 (2014), no 2, 839-846.
- [5] O. Chodosh, *Large isoperimetric regions in asymptotically hyperbolic manifolds*, arXiv:1403.6108.
- [6] P. Chruściel and M. Herzlich, *The mass of asymptotically hyperbolic Riemannian manifolds*, Pacific J. Math. 212 (2003), no. 2, 231-264.
- [7] J. Corvino, A. Gerek, M. Greenberg and B. Krummel, *On isoperimetric surfaces in general relativity*, Pac. J. Math. 231 (2007), no. 1, 63-84.

- [8] M. Dahl, R. Gicquard and A. Sakovich, *Penrose type inequalities for asymptotically hyperbolic graphs*, Ann. Henri Poincaré 14 (2013), 1135-1168.
- [9] P. Figueras, K. Murata and H. S. Reall, *Black hole instabilities and local Penrose inequalities*, Class. Quantum Grav. 28 (2011), 225030 (30pp).
- [10] R. Geroch, *Energy extraction*, Ann. New York Acad. Sci. 224 (1973), 108-117.
- [11] F. Girão and L. L. de Lima, *Positive mass and Penrose type inequalities for asymptotically hyperbolic hypersurfaces*, arXiv:1201.4991.
- [12] F. Girão and L. L. de Lima, *An Alexandrov-Fenchel-type inequality in hyperbolic space with an application to a Penrose inequality*, arXiv:1209.0438.
- [13] S. Hollands and R. M. Wald, *Stability of black holes and black branes*, Commun. Math. Phys. 321 (2013), no. 3, 629-680.
- [14] G. Huisken and T. Ilmanen, *The inverse mean curvature flow and the Riemannian Penrose inequality*, J. Differential Geom. 59 (2001), no. 3, 353-437.
- [15] G. Huisken and S.T. Yau, *Definition of center of mass for isolated physical systems and unique foliations by stable spheres with constant mean curvature*, Invent. Math. 124 (1996), 281-311.
- [16] D. Lee and A. Neves, *The Penrose inequality for asymptotically locally hyperbolic spaces with nonpositive mass*, arXiv: 1310.3002.
- [17] M. Mars, *Present status of the Penrose inequality*, Classical and Quantum Gravity 26 (2009), no. 19, 193001, 59 pp.
- [18] D. Maximo and I. Nunes, *Hawking mass and local rigidity of minimal two-spheres in three-manifolds*, Comm. Anal. Geom. 21 (2013), no. 2, 409-432.
- [19] R. Mazzeo and F. Pacard, *Constant curvature foliations in asymptotically hyperbolic spaces*, Rev. Mat. Iberoam. 27 (2011), no. 1, 303-333.
- [20] A. Neves, *Insufficient convergence of inverse mean curvature flow on asymptotically hyperbolic manifolds*, J. Differential Geom. 84 (2010), no. 1, 191-229.
- [21] A. Neves and G. Tian, *Existence and uniqueness of constant mean curvature foliation of asymptotically hyperbolic 3-manifolds*, Geom. Funct. Anal. 19 (2009), no. 3, 910-942.
- [22] A. Neves and G. Tian, *Existence and uniqueness of constant mean curvature foliation of asymptotically hyperbolic 3-manifolds. II*, J. Reine Angew. Math. 641 (2010), 69-93.
- [23] R. Rigger, *The foliation of asymptotically hyperbolic manifolds by surfaces of constant mean curvature (including the evolution equations and estimates)*, Manuscripta Math. 113 (2004), 403-421.
- [24] X. Wang, *The mass of asymptotically hyperbolic manifolds*, J. Differential Geom. 57 (2001), no. 2, 273-299.