

Manuscript version: Published Version

The version presented in WRAP is the published version (Version of Record).

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/136896>

How to cite:

The repository item page linked to above, will contain details on accessing citation guidance from the publisher.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

Algebra & Number Theory

Volume 14
2020
No. 9

On asymptotic Fermat over \mathbb{Z}_p -extensions of \mathbb{Q}

Nuno Freitas, Alain Kraus and Samir Siksek



On asymptotic Fermat over \mathbb{Z}_p -extensions of \mathbb{Q}

Nuno Freitas, Alain Kraus and Samir Siksek

Let p be a prime and let $\mathbb{Q}_{n,p}$ denote the n -th layer of the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} . We prove the effective asymptotic FLT over $\mathbb{Q}_{n,p}$ for all $n \geq 1$ and all primes $p \geq 5$ that are non-Wieferich, i.e., $2^{p-1} \not\equiv 1 \pmod{p^2}$. The effectivity in our result builds on recent work of Thorne proving modularity of elliptic curves over $\mathbb{Q}_{n,p}$.

1. Introduction

Let F be a totally real number field. The *asymptotic Fermat's last theorem over F* is the statement that there exists a constant B_F , depending only on F , such that, for all primes $\ell > B_F$, the only solutions to the equation $x^\ell + y^\ell + z^\ell = 0$, with $x, y, z \in F$ are the trivial ones satisfying $xyz = 0$. If B_F is effectively computable, we refer to this as the *effective asymptotic Fermat's last theorem over F* . Let p be a prime, n a positive integer and write $\mathbb{Q}_{n,p}$ for the n -th layer of the cyclotomic \mathbb{Z}_p -extension. In [Freitas et al. 2020], the authors established the following theorem.

Theorem 1. *The effective asymptotic Fermat's last theorem holds over each layer $\mathbb{Q}_{n,2}$ of the cyclotomic \mathbb{Z}_2 -extension.*

The proof of Theorem 1 relies heavily on class field theory and the theory of 2-extensions, and the method depends crucially on the fact that 2 is totally ramified in $\mathbb{Q}_{n,2}$. We establish the following.

Theorem 2. *Let $p \geq 5$ be a prime. Suppose p is non-Wieferich, i.e., $2^{p-1} \not\equiv 1 \pmod{p^2}$. The effective asymptotic Fermat's last theorem holds over each layer $\mathbb{Q}_{n,p}$ of the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} .*

We remark that the only Wieferich primes currently known are 1093 and 3511. It is fascinating to observe that these primes originally arose in connection with historical attempts at proving Fermat's last theorem. Indeed Wieferich [1909] showed that if $2^{p-1} \not\equiv 1 \pmod{p^2}$ then the first case of Fermat's last theorem holds for exponent p .

In contrast to Theorem 1, the proof of Theorem 2 makes use of a criterion (Theorem 3 below) established in [Freitas and Siksek 2015] for asymptotic FLT in terms of solutions to a certain S -unit equation. The proof of that criterion builds on many deep results including modularity lifting theorems due to Breuil, Diamond, Gee, Kisin, and others, and Merel's uniform boundedness theorem, and exploits the strategy of Frey, Serre, Ribet, Wiles and Taylor, utilized in Wiles' proof of Fermat's last theorem [1995]. We use elementary

Freitas is supported by a Ramón y Cajal fellowship (RYC-2017-22262). Siksek is supported by EPSRC grant "Moduli of Elliptic curves and Classical Diophantine Problems" (EP/S031537/1).

MSC2020: primary 11D41; secondary 11R23.

Keywords: Fermat, unit equation, \mathbb{Z}_p -extensions.

arguments to study these S -unit equations in $\mathbb{Q}_{n,p}$ and this study, together with the S -unit criterion, quickly yields Theorem 2. The effectivity in Theorem 2 builds on the following great theorem due to Thorne [2019].

Theorem (Thorne). *Elliptic curves over $\mathbb{Q}_{n,p}$ are modular.*

2. An S -unit criterion for asymptotic FLT

The following criterion for asymptotic FLT is a special case of [Freitas and Siksek 2015, Theorem 3].

Theorem 3. *Let F be a totally real number field. Suppose the Eichler–Shimura conjecture over F holds. Assume that 2 is inert in F and write $\mathfrak{q} = 2\mathcal{O}_F$ for the prime ideal above 2. Let $S = \{\mathfrak{q}\}$ and write \mathcal{O}_S^\times for the group of S -units in F . Suppose every solution (λ, μ) to the S -unit equation*

$$\lambda + \mu = 1, \quad \lambda, \mu \in \mathcal{O}_S^\times \tag{2-1}$$

satisfies both of the following conditions:

$$\max\{|\text{ord}_{\mathfrak{q}}(\lambda)|, |\text{ord}_{\mathfrak{q}}(\mu)|\} \leq 4, \quad \text{ord}_{\mathfrak{q}}(\lambda\mu) \equiv 1 \pmod{3}. \tag{2-2}$$

Then the asymptotic Fermat’s last theorem holds over F . Moreover, if all elliptic curves over F with full 2-torsion are modular, then the effective asymptotic Fermat’s last theorem holds over F .

For a discussion of the Eichler–Shimura conjecture see [Freitas and Siksek 2015, Section 2.4], but for the purpose of this paper we note that the conjecture is known to hold for all totally real fields of odd degree. In particular, it holds for $\mathbb{Q}_{n,p}$ for all odd p .

To apply Theorem 3 to $F = \mathbb{Q}_{n,p}$ we need to know for which p is 2 inert in F . The answer is given by the following lemma, which for $n = 1$ is [Washington 1997, Exercise 2.4].

Lemma 2.1. *Let $p \geq 3, q$ be distinct primes. Then q is inert in $\mathbb{Q}_{n,p}$ if and only if $q^{p-1} \not\equiv 1 \pmod{p^2}$.*

Proof. Let $L = \mathbb{Q}(\zeta_{p^{n+1}})$ and $F = \mathbb{Q}_{n,p}$. Write σ_q and τ_q for the Frobenius elements corresponding to q in $\text{Gal}(L/\mathbb{Q})$ and $\text{Gal}(F/\mathbb{Q})$. The prime q is inert in F precisely when τ_q has order p^n . The natural surjection $\text{Gal}(L/\mathbb{Q}) \rightarrow \text{Gal}(F/\mathbb{Q})$ sends σ_q to τ_q and its kernel has order $p - 1$. Thus q is inert in F if and only if the order of σ_q is divisible by p^n , which is equivalent to σ_q^{p-1} having order p^n . There is a canonical isomorphism $\text{Gal}(L/\mathbb{Q}) \rightarrow (\mathbb{Z}/p^{n+1}\mathbb{Z})^\times$ sending σ_q to $q + p^{n+1}\mathbb{Z}$. Thus q is inert in F if and only if $q^{p-1} + p^{n+1}\mathbb{Z}$ has order p^n . This is equivalent to $q^{p-1} \not\equiv 1 \pmod{p^2}$. □

3. Proof of Theorem 2

Lemma 3.1. *Let \mathfrak{p} be the unique prime above p in $F = \mathbb{Q}_{n,p}$. Let $\lambda \in \mathcal{O}_F$. Then $\lambda \equiv \text{Norm}_{F/\mathbb{Q}}(\lambda) \pmod{\mathfrak{p}}$.*

Proof. As p is totally ramified in F , we know that the residue field $\mathcal{O}_F/\mathfrak{p}$ is \mathbb{F}_p . Thus there is some $a \in \mathbb{Z}$ such that $\lambda \equiv a \pmod{\mathfrak{p}}$. Let $\sigma \in G = \text{Gal}(F/\mathbb{Q})$. Since $\mathfrak{p}^\sigma = \mathfrak{p}$, we have $\lambda^\sigma \equiv a \pmod{\mathfrak{p}}$. Hence

$$\text{Norm}_{F/\mathbb{Q}}(\lambda) = \prod_{\sigma \in G} \lambda^\sigma \equiv a^{\#G} \pmod{\mathfrak{p}}.$$

However $\#G = p^n$ so $\text{Norm}_{F/\mathbb{Q}}(\lambda) \equiv a \equiv \lambda \pmod{\mathfrak{p}}$. □

Lemma 3.2. *Let $p \neq 3$ be a rational prime. Let $F = \mathbb{Q}_{n,p}$. Then the unit equation*

$$\lambda + \mu = 1, \quad \lambda, \mu \in \mathcal{O}_F^\times \tag{3-1}$$

has no solutions.

Proof. Let (λ, μ) be a solution to (3-1). By Lemma 3.1, $\lambda \equiv \pm 1 \pmod{p}$ and $\mu \equiv \pm 1 \pmod{p}$. Thus $\pm 1 \pm 1 \equiv \lambda + \mu = 1 \pmod{p}$. This is impossible as $p \neq 3$. \square

Remark 3.3. Lemma 3.2 is false for $p = 3$. Indeed, let $p = 3$ and $n = 1$. Then $F = \mathbb{Q}_{1,3} = \mathbb{Q}(\theta)$, where θ satisfies $\theta^3 - 6\theta^2 + 9\theta - 3 = 0$. The unit equation has solution $\lambda = 2 - \theta$ and $\mu = -1 + \theta$. In fact, the unit equation solver of the computer algebra system Magma [Bosma et al. 1997] gives a total of 18 solutions.

Lemma 3.4. *Let $p \geq 5$ be a rational prime. Let $F = \mathbb{Q}_{n,p}$. Suppose 2 is inert in F and write $\mathfrak{q} = 2\mathcal{O}_F$ for the unique prime above 2. Let $S = \{\mathfrak{q}\}$ and write \mathcal{O}_S^\times for the group of S -units. Then every solution to the S -unit equation (2-1) satisfies one of the following:*

- (i) $\text{ord}_{\mathfrak{q}}(\lambda) = 1, \text{ord}_{\mathfrak{q}}(\mu) = 0$;
- (ii) $\text{ord}_{\mathfrak{q}}(\lambda) = 0, \text{ord}_{\mathfrak{q}}(\mu) = 1$;
- (iii) $\text{ord}_{\mathfrak{q}}(\lambda) = \text{ord}_{\mathfrak{q}}(\mu) = -1$.

Proof. Write $n_\lambda = \text{ord}_{\mathfrak{q}}(\lambda)$ and $n_\mu = \text{ord}_{\mathfrak{q}}(\mu)$. Suppose first $n_\lambda \geq 2$. Then $n_\mu = 0$ and so $\mu \in \mathcal{O}_F^\times$. Moreover, as $4 \mid \lambda$, we have $\mu \equiv 1 \pmod{4}$ and so $\mu^\sigma \equiv 1 \pmod{4}$ for all $\sigma \in G = \text{Gal}(F/\mathbb{Q})$. Hence $\text{Norm}_{F/\mathbb{Q}}(\mu) = \prod \mu^\sigma \equiv 1 \pmod{4}$. But $\text{Norm}_{F/\mathbb{Q}}(\mu) = \pm 1$, thus $\text{Norm}_{F/\mathbb{Q}}(\mu) = 1$. As before, denote the unique prime above p by \mathfrak{p} . By Lemma 3.1 we have $\mu \equiv 1 \pmod{\mathfrak{p}}$. Hence \mathfrak{p} divides $1 - \mu = \lambda$ giving a contradiction.

Thus $n_\lambda \leq 1$. Next suppose $n_\lambda \leq -2$. Then $n_\lambda = n_\mu$. Let $\lambda' = 1/\lambda$ and $\mu' = -\mu/\lambda$. Then (λ', μ') is a solution to the S -unit equation satisfying $n_{\lambda'} \geq 2$, giving a contradiction by the previous case. Hence $-1 \leq n_\lambda \leq 1$ and by symmetry $-1 \leq n_\mu \leq 1$. From Lemma 3.2 either $n_\lambda \neq 0$ or $n_\mu \neq 0$. Thus one of (i), (ii), (iii) must hold. \square

Remark 3.5. Possibilities (i), (ii), (iii) cannot be eliminated because of the solutions $(2, -1)$, $(-1, 2)$ and $(\frac{1}{2}, \frac{1}{2})$ to the S -unit equation.

Proof of Theorem 2. We suppose $p \geq 5$ and non-Wieferich. It follows from Lemma 2.1 that 2 is inert in $F = \mathbb{Q}_{n,p}$. Write $\mathfrak{q} = 2\mathcal{O}_F$. By Lemma 3.4 all solutions (λ, μ) to the S -unit equation (2-1) satisfy (2-2). We now apply Theorem 3. As elliptic curves over $\mathbb{Q}_{n,p}$ are modular thanks to Thorne’s theorem, we conclude that the effective Fermat’s last theorem holds over $\mathbb{Q}_{n,p}$. \square

Remark 3.6. The proof of Theorem 2 for $p = 3$ and for the Wieferich primes seems out of reach at present. There are solutions to the unit equation in $\mathbb{Q}_{1,3}$ (as indicated in Remark 3.3), and therefore in $\mathbb{Q}_{n,3}$ for all n , and these solutions violate the criterion of Theorem 3. For p a Wieferich prime, 2 splits in $\mathbb{Q}_{n,p}$ into at least p prime ideals and we would need to consider the S -unit equation (2-1) with S the set of primes above 2. It appears difficult to treat the S -unit equation in infinite families of number fields where $\#S \geq 2$ (see [Freitas et al. 2020, Theorem 7] and its proof).

4. A generalization

In fact, the proof of Theorem 2 establishes the following more general theorem.

Theorem 4. *Let F be a totally real number field and $p \geq 5$ be a rational prime. Suppose that the following conditions are satisfied.*

- (a) F is a p -extension of \mathbb{Q} (i.e., F/\mathbb{Q} is a Galois extension of degree p^n for some $n \geq 1$).
- (b) p is totally ramified in F .
- (c) 2 is inert in F .

Then the asymptotic Fermat's last theorem holds for F .

Example 4.1. A quick search on the L-Functions and Modular Forms Database [LMFDB 2020] yields 153 fields of degree 5 satisfying conditions of the theorem with $p = 5$. The one with smallest discriminant is $\mathbb{Q}_{1,5}$. The one with the next smallest discriminant is $F = \mathbb{Q}(\theta)$ where $\theta^5 - 110\theta^3 - 605\theta^2 - 990\theta - 451 = 0$. The discriminant of F is $5^8 \cdot 11^4$. It is therefore not contained in any \mathbb{Z}_p -extension of \mathbb{Q} .

References

- [Bosma et al. 1997] W. Bosma, J. Cannon, and C. Playoust, “The Magma algebra system, I: The user language”, *J. Symbolic Comput.* **24**:3-4 (1997), 235–265. See also Magma computational algebra system. MR Zbl
- [Freitas and Siksek 2015] N. Freitas and S. Siksek, “The asymptotic Fermat's last theorem for five-sixths of real quadratic fields”, *Compos. Math.* **151**:8 (2015), 1395–1415. MR Zbl
- [Freitas et al. 2020] N. Freitas, A. Kraus, and S. Siksek, “Class field theory, Diophantine analysis and the asymptotic Fermat's last theorem”, *Adv. Math.* **363** (2020), art. id. 106964. MR Zbl
- [LMFDB 2020] The LMFDB Collaboration, “The L -functions and modular forms database”, 2020, <http://www.lmfdb.org>.
- [Thorne 2019] J. A. Thorne, “Elliptic curves over \mathbb{Q}_∞ are modular”, *J. Eur. Math. Soc.* **21**:7 (2019), 1943–1948. MR Zbl
- [Washington 1997] L. C. Washington, *Introduction to cyclotomic fields*, 2nd ed., Grad. Texts in Math. **83**, Springer, 1997. MR Zbl
- [Wieferich 1909] A. Wieferich, “Zum letzten Fermatschen Theorem”, *J. Reine Angew. Math.* **136** (1909), 293–302. MR Zbl
- [Wiles 1995] A. Wiles, “Modular elliptic curves and Fermat's last theorem”, *Ann. of Math. (2)* **141**:3 (1995), 443–551. MR Zbl

Communicated by Andrew Granville

Received 2020-04-02 Accepted 2020-05-11

nunobfreitas@gmail.com

Department de Matemàtiques i Informàtica, Universitat de Barcelona, Barcelona, Spain

alain.kraus@imj-prg.fr

Institut de Mathématiques de Jussieu - Paris Rive Gauche, Sorbonne Université, UMR 7586 CNRS - Paris Diderot, Paris, France

s.siksek@warwick.ac.uk

Mathematics Institute, University of Warwick, Coventry, United Kingdom

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Jason P. Bell	University of Waterloo, Canada	Susan Montgomery	University of Southern California, USA
Bhargav Bhatt	University of Michigan, USA	Martin Olsson	University of California, Berkeley, USA
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
Frank Calegari	University of Chicago, USA	Jonathan Pila	University of Oxford, UK
Antoine Chambert-Loir	Université Paris-Diderot, France	Irena Peeva	Cornell University, USA
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Michael Rapoport	Universität Bonn, Germany
Samit Dasgupta	Duke University, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Michael Singer	North Carolina State University, USA
Sergey Fomin	University of Michigan, USA	Christopher Skinner	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Wee Teck Gan	National University of Singapore	Shunsuke Takagi	University of Tokyo, Japan
Andrew Granville	Université de Montréal, Canada	Pham Huu Tiep	University of Arizona, USA
Ben J. Green	University of Oxford, UK	Ravi Vakil	Stanford University, USA
Joseph Gubeladze	San Francisco State University, USA	Michel van den Bergh	Hasselt University, Belgium
Christopher Hacon	University of Utah, USA	Akshay Venkatesh	Institute for Advanced Study, USA
Roger Heath-Brown	Oxford University, UK	Marie-France Vignéras	Université Paris VII, France
János Kollár	Princeton University, USA	Melanie Matchett Wood	University of California, Berkeley, USA
Michael J. Larsen	Indiana University Bloomington, USA	Shou-Wu Zhang	Princeton University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne		

PRODUCTION

production@msp.org
Silvio Levy, Scientific Editor

See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2020 is US \$415/year for the electronic version, and \$620/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 14 No. 9 2020

The Brauer group of the moduli stack of elliptic curves BENJAMIN ANTIEAU and LENNART MEIER	2295
Modular forms from Noether–Lefschetz theory FRANÇOIS GREER	2335
Quadratic Chabauty for (bi)elliptic curves and Kim’s conjecture FRANCESCA BIANCHI	2369
The Prasad conjectures for GSp_4 and PGSp_4 HENGFEI LU	2417
Invertible functions on nonarchimedean symmetric spaces ERNST-ULRICH GEKELER	2481
On a cohomological generalization of the Shafarevich conjecture for K3 surfaces TEPPEI TAKAMATSU	2505
Iterated local cohomology groups and Lyubeznik numbers for determinantal rings ANDRÁS C. LÓRINCZ and CLAUDIU RAICU	2533
On asymptotic Fermat over \mathbb{Z}_p -extensions of \mathbb{Q} NUNO FREITAS, ALAIN KRAUS and SAMIR SIKSEK	2571



1937-0652(2020)14:9;1-T