



NHC-catalyzed enantioselective synthesis of β -trifluoromethyl- β -hydroxyamides

Alyn T. Davies, Mark D. Greenhalgh, Alexandra M. Z. Slawin and Andrew D. Smith*

Letter

Open Access

Address:
EaStCHEM, School of Chemistry, University of St Andrews, North
Haugh, St Andrews, Fife KY16 9ST, United Kingdom

Email:
Andrew D. Smith* - ads10@st-andrews.ac.uk

* Corresponding author

Keywords:
enantioselective catalysis; formal [2 + 2] cycloaddition; N-heterocyclic
carbene; ring opening; trifluoromethyl group

Beilstein J. Org. Chem. **2020**, *16*, 1572–1578.
doi:10.3762/bjoc.16.129

Received: 09 March 2020
Accepted: 05 June 2020
Published: 30 June 2020

This article is part of the thematic issue "Organo-fluorine chemistry V".

Guest Editor: D. O'Hagan

© 2020 Davies et al.; licensee Beilstein-Institut.
License and terms: see end of document.

Abstract


The N-heterocyclic carbene (NHC)-catalyzed formal [2 + 2] cycloaddition between α -aroyloxyaldehydes and trifluoroacetophenones, followed by ring opening with an amine or a reducing agent is described. The resulting β -trifluoromethyl- β -hydroxyamide and alcohol products are produced with reasonable diastereocontrol (typically $\approx 70:30$ dr) and excellent enantioselectivity, and they can be isolated in moderate to good yield as a single diastereoisomer.

Introduction


The trifluoromethyl unit holds a prominent and privileged position within organic chemistry [1-6]. The incorporation of this motif is widely employed within the pharmaceutical, agrochemical, and materials industries as it can be used strategically to increase lipophilicity as well as to enhance metabolic stability and binding selectivity [7-9]. In this context, the generation of effective and practical methodologies capable of the enantioselective incorporation of the trifluoromethyl unit into organic molecules has been developed widely [10-12], with a series of elegant organocatalytic strategies utilized toward these aims [13]. One general strategy for the organocatalytic construction of stereogenic trifluoromethyl centers is through enantioselective addition of enolates or their equivalents to prochiral tri-

fluoromethyl ketones (Figure 1A). Within this area, a common catalytic approach has utilized aliphatic ketones as enolate equivalents using prolinamide, cinchona, or hybrid catalysts that proceed via enamine intermediates (Figure 1B) [14-19]. The state of the art within this area has been recently demonstrated by Dixon and co-workers, who showed that bifunctional BIMP catalysts could promote the enantioselective addition of typically recalcitrant aryl ketones to trifluoromethyl ketones (Figure 1C) [20].

Over the last twenty years, NHCs have been widely exploited as highly efficient organocatalysts that have found use in numerous applications and were the subject of many extensive



reviews [21–26]. Among the most common reactive intermediates generated using NHCs, the azolium enolate has been widely used [27,28]. Methods to generate azolium enolates from a number of precursors have been reported, including the use of ketenes [29,30], α -functionalized aldehydes [31–35], enals [36–38], aryl esters [39–42], or aldehydes [43–45] in the presence of an oxidant. As representative examples of the use of azolium enolates in reactions with trifluoroacetophenone derivatives, Ye and co-workers have shown that using disubstituted ketenes as azolium enolate precursors and NHC precatalyst **1** allowed access to trifluoromethyl-substituted β -lactone products in a high yield, diastereo-, and enantioselectivity (Figure 2A) [46]. Chi and co-workers have reported limited examples of oxidative [2 + 2] cycloadditions using hydrocinnamaldehyde as an azolium enolate precursor and the NHC precatalyst **2**, giving β -lactone products in high enantioselectivity but requiring superstoichiometric quantities of quinone as an oxidant (Figure 2B) [47]. In our previous work, we have developed α -aryloxyaldehydes as reactive, bench-stable precursors of azolium enolates [48–53], which can be synthesized in a single step from the desired aldehyde [54]. We have previously applied these precursors in the NHC-catalyzed formal [2 + 2]



cycloadditions between α -aryloxyaldehydes and perfluoroalkyl-substituted ketones using NHC precatalyst **3** to produce polyfluorinated oxetanes and β -perfluoroalkyl- β -hydroxyamides after derivatization [55]. In this paper, this formal [2 + 2] protocol is extended to the use of trifluoroacetophenones to develop an enantioselective route to β -trifluoromethyl- β -hydroxy carboxylic acid derivatives (Figure 2C).

Results and Discussion

Optimization


Optimization of the NHC-catalyzed formal [2 + 2] cycloaddition using the α -aryloxyaldehyde **4** and trifluoromethylacetophenone (**5**) as reactants began using the NHC precatalyst **3**, triethylamine as the base, and THF as the solvent (Table 1, entry 1). A moderate conversion (48%, as determined by NMR analysis) to the desired β -lactone product **6** as a 70:30 mixture of diastereoisomers was observed, which served as a basis for further optimization. Variation of the base showed caesium carbonate to be significantly more effective than the organic bases examined, giving 88% conversion to product **6** (Table 1, entries 2 and 3). Trialling a number of alternative solvents, including dichloromethane, diethyl ether, and toluene showed no improvement upon the yield obtained with THF (Table 1, entries 4–6). Unfortunately, attempts to isolate the desired β -lactone product **6** were unsuccessful, with **6** being unstable to chromatographic purification under a variety of different conditions. As such, following the NHC-catalyzed formal [2 + 2]

cycloaddition in THF, a subsequent ring opening step with allylamine was investigated. Although the diastereomeric ratio of the resultant crude reaction mixture was 75:25 after chromatographic purification, as seen by NMR analysis, the corresponding β -trifluoromethyl- β -hydroxyamide **7** was isolated as a single diastereoisomer in 57% yield and >99:1 er (Table 1, entry 7).

Scope and limitations


Having optimized this process with a model system, further work probed the scope and limitations of this process through a sequential variation of the nucleophile used for the ring opening procedure, aryl and heteroaryl substitution of the trifluoromethylacetophenone, as well as variation of the α -aryloxyaldehyde. Using α -aryloxyaldehyde **4** and trifluoromethylacetophenone (**5**) as reactants, the variation of the nucleophile showed that a range of amine-based nucleophiles was readily tolerated in this process (Scheme 1). The use of benzylamine, pyrrolidine, and ammonia all gave the corresponding β -trifluoromethyl- β -hydroxyamide product in \approx 75:25 dr, with purification giving **8–10**, respectively, as a single diastereoisomer in 42–55% yield and excellent enantioselectivity. Using methanol as a nucleophile gave the corresponding ester, however, this product proved unstable to purification. Generation of the ester in situ, followed by subsequent reduction with LiAlH_4 , gave the diol **11** in 48% isolated yield as a single diastereoisomer in 96:4 er.

Table 1: Optimization of the NHC-catalyzed formal [2 + 2] cycloaddition.^a



entry	solvent	base	dr ^b	yield (%)	er ^c
1	THF	Et_3N	70:30	48 ^d	–
2	THF	$i\text{Pr}_2\text{NEt}$	70:30	50 ^d	–
3	THF	Cs_2CO_3	75:25	88 ^d	–
4	CH_2Cl_2	Cs_2CO_3	80:20	61 ^d	–
5	Et_2O	Cs_2CO_3	70:30	58 ^d	–
6	toluene	Cs_2CO_3	70:30	55 ^d	–
7	THF	Cs_2CO_3	75:25	57 ^e	>99:1

^aThe data contained in this Table has previously been published as part of our previous communication in this area [55]. ^bDetermined by ^1H NMR spectroscopic analysis of the crude reaction product mixture. ^cer of the major diastereoisomer (>95:5 dr), determined by GC analysis using a chiral support. ^dCombined yield of both diastereoisomers of **6**, determined by the analysis of the crude ^1H NMR spectra with reference to 2,5-dimethylfuran as an internal standard. ^eIsolated yield of product **7** (>95:5 dr).




Scheme 1: Reaction scope with respect to the nucleophile. ^aIsolated yield of the product in >95:5 dr. ^bDetermined by ¹H NMR spectroscopic analysis of the crude reaction product mixture. ^cDetermined by HPLC analysis or GC analysis using a chiral support.


Subsequent variation of the electronic nature of the aromatic substituent within the trifluoroacetophenone component, followed by a ring opening with allylamine, showed that the introduction of electron-withdrawing groups (positive Hammett sigma constants) [56], such as *p*-bromo, *p*-fluoro, and *p*-trifluoromethyl groups, respectively, were well tolerated, giving a consistent diastereoselectivity of $\approx 75:25$ dr (Scheme 2). Following chromatographic purification, the desired products **12–14**, respectively, were isolated in 59–71% yield as single diastereoisomers with excellent enantioselectivity (95:5 to >99:1 er). The relative and absolute configuration of (2*S*,3*S*)- β -trifluoromethyl- β -hydroxyamide **12** was confirmed by single crystal X-ray crystallographic analysis, with all other products in this series assigned by analogy to **12** [57]. The introduction of an electron-donating *p*-tolyl substituent was also tolerated in the system, providing the corresponding β -trifluoromethyl- β -hydroxyamide **15** in a moderate yield of 46%, although the er of **15** could not be determined by either GC or HPLC analysis. The trifluoroacetophenone derivatives with stronger electron-donating *p*-methoxy and *p*-dimethylamino substituents proved unreactive, consistent with the expected lower electrophilicity. A heterocyclic substituent could also be incorporated using this methodology, with thiophene-substituted β -trifluoromethyl- β -hydroxyamide **16** being produced in 80:20 dr, with purification

giving **16** in 51% yield as a single diastereoisomer and 96:4 er. The substrate scope was further investigated by variation of the α -aryloxyaldehyde component of the reaction. Functionalized α -aryloxyaldehydes containing a C2-benzyl and a C2-benzyl-oxy derivative were tested, producing the corresponding β -trifluoromethyl- β -hydroxyamides **17** and **18**, respectively, in $\approx 70:30$ dr after a ring opening with allylamine. Purification gave **17** and **18**, respectively, in a moderate yield as single diastereoisomers and with a good enantioselectivity.

The mechanism of this NHC redox process is believed to proceed through the following mechanism (Scheme 3): After deprotonation of the triazolium salt precatalyst **3** [58], reversible addition of the free NHC **I** to the aldehyde leads to adduct **II** [59]. A subsequent deprotonation allows access to Breslow intermediate **III**, which can eliminate *para*-nitrobenzoate to leave azolium enol **IV**. Deprotonation gives azolium enolate intermediate **V**, which undergoes a formal [2 + 2] cycloaddition with the trifluoroacetophenone derivative to give **VI** [60]. Elimination of the NHC catalyst completes the catalytic cycle and provides the corresponding β -lactone product. Subsequent ring opening with a suitable nucleophilic amine leads to the isolable β -trifluoromethyl- β -hydroxyamide (Scheme 3).



Scheme 2: Reaction scope with respect to the trifluoroacetophenone derivative and α -aryloxyaldehyde. ^aIsolated yield of the product in >95:5 dr. ^bDetermined by ¹H NMR spectroscopic analysis of the crude reaction product mixture. ^cer of the major diastereoisomer (>95:5 dr), determined by HPLC analysis or GC analysis using a chiral support. ^dMolecular representation of the X-ray crystal structure of **12**. The unit cell of **12** contained two molecules of **12**, with only one shown for clarity.



Scheme 3: Proposed mechanism.

Conclusion

In this paper, we showed that azolium enolates generated from α -aroyloxyaldehydes can undergo NHC-catalyzed formal [2 + 2] cycloadditions with trifluoroacetophenone derivatives. Although the β -lactone products proved unstable to chromatographic purification, ring opening with amine nucleophiles allowed access to β -trifluoromethyl- β -hydroxyamides in moderate to good yield as single diastereoisomers in excellent er following purification [61].

Supporting Information

Supporting Information File 1

Experimental procedures, product characterization data (mp, NMR, IR, HRMS, $[\alpha]_D$, HPLC), and spectra (^1H , ^{13}C , and ^{19}F NMR, HPLC).

[<https://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-16-129-S1.pdf>]

Supporting Information File 2

Crystallographic details for **12**.

[<https://www.beilstein-journals.org/bjoc/content/supplementary/1860-5397-16-129-S2.cif>]

Acknowledgements

We thank the EPSRC UK National Mass Spectrometry Facility at Swansea University.

Funding

We thank the European Research Council under the European Union's Seventh Framework Program (FP7/2007-2013) ERC grant agreement no. 279850 (A. T. D.)

ORCID® iDs

Mark D. Greenhalgh - <https://orcid.org/0000-0002-4176-7633>

Alexandra M. Z. Slawin - <https://orcid.org/0000-0002-9527-6418>

Andrew D. Smith - <https://orcid.org/0000-0002-2104-7313>

References

- Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320–330. doi:10.1039/b610213c
- Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881–1886. doi:10.1126/science.1131943
- Studer, A. *Angew. Chem., Int. Ed.* **2012**, *51*, 8950–8958. doi:10.1002/anie.201202624
- Tomashenko, O. A.; Grushin, V. V. *Chem. Rev.* **2011**, *111*, 4475–4521. doi:10.1021/cr1004293
- Dilman, A. D.; Levin, V. V. *Eur. J. Org. Chem.* **2011**, 831–841. doi:10.1002/ejoc.2011001558
- Liu, H.; Gu, Z.; Jiang, X. *Adv. Synth. Catal.* **2013**, *355*, 617–626. doi:10.1002/adsc.201200764
- Hagmann, W. K. *J. Med. Chem.* **2008**, *51*, 4359–4369. doi:10.1021/jm800219f
- Sani, M.; Volonterio, A.; Zanda, M. *ChemMedChem* **2007**, *2*, 1693–1700. doi:10.1002/cmdc.200700156
- Léger, S.; Bayly, C. I.; Black, W. C.; Desmarais, S.; Falguyret, J.-P.; Massé, F.; Percival, M. D.; Truchon, J.-F. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 4328–4332. doi:10.1016/j.bmcl.2007.05.024
- Nie, J.; Guo, H.-C.; Cahard, D.; Ma, J.-A. *Chem. Rev.* **2011**, *111*, 455–529. doi:10.1021/cr100166a
- Shibata, N.; Matsnev, A.; Cahard, D. *Beilstein J. Org. Chem.* **2010**, *6*, No. 65. doi:10.3762/bjoc.6.65
- Ma, J.-A.; Cahard, D. *Chem. Rev.* **2004**, *104*, 6119–6146. doi:10.1021/cr030143e
- Valero, G.; Companyó, X.; Rios, R. *Chem. – Eur. J.* **2011**, *17*, 2018–2037. doi:10.1002/chem.201001546
See for a review.
- Yang, W.; Cui, Y.-M.; Zhou, W.; Li, L.; Yang, K.-F.; Zheng, Z.-J.; Lu, Y.; Xu, L.-W. *Synlett* **2014**, *25*, 1461–1465. doi:10.1055/s-0033-1341281
- Vlasserou, I.; Sfetsa, M.; Gerokonstantis, D.-T.; Kokotos, C. G.; Moutevelis-Minakakis, P. *Tetrahedron* **2018**, *74*, 2338–2349. doi:10.1016/j.tet.2018.03.054
- Duangdee, N.; Harnying, W.; Rulli, G.; Neudörfel, J.-M.; Gröger, H.; Berkessel, A. *J. Am. Chem. Soc.* **2012**, *134*, 11196–11205. doi:10.1021/ja302511t
- Hara, N.; Tamura, R.; Funahashi, Y.; Nakamura, S. *Org. Lett.* **2011**, *13*, 1662–1665. doi:10.1021/ol2001039
- Kokotos, C. G. *J. Org. Chem.* **2012**, *77*, 1131–1135. doi:10.1021/jo2020104
- Qui, L.-H.; Shen, Z.-X.; Shi, C.-Q.; Liu, Y.-H.; Zhang, Y.-W. *Chin. J. Chem.* **2005**, *23*, 584–588. doi:10.1002/cjoc.200590584
- Thomson, C. J.; Barber, D. M.; Dixon, D. J. *Angew. Chem., Int. Ed.* **2020**, *59*, 5359–5364. doi:10.1002/anie.201916129
- Moore, J. L.; Rovis, T. *Top. Curr. Chem.* **2010**, *291*, 77–144.
- Marion, N.; Díez-González, S.; Nolan, S. P. *Angew. Chem., Int. Ed.* **2007**, *46*, 2988–3000. doi:10.1002/anie.200603380
- Enders, D.; Niemeier, O.; Henseler, A. *Chem. Rev.* **2007**, *107*, 5606–5655. doi:10.1021/cr068372z
- Hopkinson, M. N.; Richter, C.; Schedler, M.; Glorius, F. *Nature* **2014**, *510*, 485–496. doi:10.1038/nature13384
- Ryan, S. J.; Candish, L.; Lupton, D. W. *Chem. Soc. Rev.* **2013**, *42*, 4906–4917. doi:10.1039/c3cs35522e
- Flanigan, D. M.; Romanov-Michaílidis, F.; White, N. A.; Rovis, T. *Chem. Rev.* **2015**, *115*, 9307–9387. doi:10.1021/acs.chemrev.5b00060
- Vora, H. U.; Wheeler, P.; Rovis, T. *Adv. Synth. Catal.* **2012**, *354*, 1617–1639. doi:10.1002/adsc.201200031
- Douglas, J.; Churchill, G.; Smith, A. D. *Synthesis* **2012**, *44*, 2295–2309. doi:10.1055/s-0031-1289788
- Zhang, Y.-R.; He, L.; Wu, X.; Shao, P.-L.; Ye, S. *Org. Lett.* **2008**, *10*, 277–280. doi:10.1021/ol702759b
- Duguet, N.; Campbell, C. D.; Slawin, A. M. Z.; Smith, A. D. *Org. Biomol. Chem.* **2008**, *6*, 1108–1113. doi:10.1039/b800857b
- He, M.; Uc, G. J.; Bode, J. W. *J. Am. Chem. Soc.* **2006**, *128*, 15088–15089. doi:10.1021/ja066380r
- He, M.; Beahm, B. J.; Bode, J. W. *Org. Lett.* **2008**, *10*, 3817–3820. doi:10.1021/ol801502h
- Kobayashi, S.; Kinoshita, T.; Uehara, H.; Sudo, T.; Ryu, I. *Org. Lett.* **2009**, *11*, 3934–3937. doi:10.1021/ol901544q

34. Yang, L.; Wang, F.; Chua, P. J.; Lv, Y.; Zhong, L.-J.; Zhong, G. *Org. Lett.* **2012**, *14*, 2894–2897. doi:10.1021/ol301175z
35. Jian, T.-Y.; Sun, L.-H.; Ye, S. *Chem. Commun.* **2012**, *48*, 10907–10909. doi:10.1039/c2cc35273g
36. Lv, H.; Chen, X.-Y.; Sun, L.-h.; Ye, S. *J. Org. Chem.* **2010**, *75*, 6973–6976. doi:10.1021/jo101318u
37. Jian, T.-Y.; Shao, P.-L.; Ye, S. *Chem. Commun.* **2011**, *47*, 2381–2383. doi:10.1039/c0cc04839a
38. Chen, X.; Fang, X.; Chi, Y. R. *Chem. Sci.* **2013**, *4*, 2613–2618. doi:10.1039/c3sc50666e
39. Hao, L.; Du, Y.; Lv, H.; Chen, X.; Jiang, H.; Shao, Y.; Chi, Y. R. *Org. Lett.* **2012**, *14*, 2154–2157. doi:10.1021/ol300676w
40. Lee, A.; Younai, A.; Price, C. K.; Izquierdo, J.; Mishra, R. K.; Scheidt, K. A. *J. Am. Chem. Soc.* **2014**, *136*, 10589–10592. doi:10.1021/ja505880r
41. Cheng, J.-T.; Chen, X.-Y.; Ye, S. *Org. Biomol. Chem.* **2015**, *13*, 1313–1316. doi:10.1039/c4ob02330g
42. Lee, A.; Scheidt, K. A. *Chem. Commun.* **2015**, *51*, 3407–3410. doi:10.1039/c4cc09590a
43. Zhao, X.; Ruhl, K. E.; Rovis, T. *Angew. Chem., Int. Ed.* **2012**, *51*, 12330–12333. doi:10.1002/anie.201206490
44. Dong, X.; Yang, W.; Hu, W.; Sun, J. *Angew. Chem., Int. Ed.* **2015**, *54*, 660–663. doi:10.1002/anie.201409961
45. Lin, L.; Yang, Y.; Wang, M.; Lai, L.; Guo, Y.; Wang, R. *Chem. Commun.* **2015**, *51*, 8134–8137. doi:10.1039/c5cc01587a
46. Wang, X.-N.; Shao, P.-L.; Lv, H.; Ye, S. *Org. Lett.* **2009**, *11*, 4029–4031. doi:10.1021/ol901290z
47. Mo, J.; Yang, R.; Chen, X.; Tiwari, B.; Chi, Y. R. *Org. Lett.* **2013**, *15*, 50–53. doi:10.1021/ol303035r
48. Ling, K. B.; Smith, A. D. *Chem. Commun.* **2011**, *47*, 373–375. doi:10.1039/c0cc02456b
49. Davies, A. T.; Taylor, J. E.; Douglas, J.; Collett, C. J.; Morrill, L. C.; Fallan, C.; Slawin, A. M. Z.; Churchill, G.; Smith, A. D. *J. Org. Chem.* **2013**, *78*, 9243–9257. doi:10.1021/jo401433q
50. Taylor, J. E.; Daniels, D. S. B.; Smith, A. D. *Org. Lett.* **2013**, *15*, 6058–6061. doi:10.1021/ol402955f
51. Davies, A. T.; Pickett, P. M.; Slawin, A. M. Z.; Smith, A. D. *ACS Catal.* **2014**, *4*, 2696–2700. doi:10.1021/cs500667g
52. Attaba, N.; Taylor, J. E.; Slawin, A. M. Z.; Smith, A. D. *J. Org. Chem.* **2015**, *80*, 9728–9739. doi:10.1021/acs.joc.5b01820
53. Kerr, R. W. F.; Greenhalgh, M. D.; Slawin, A. M. Z.; Arnold, P. L.; Smith, A. D. *Tetrahedron: Asymmetry* **2017**, *28*, 125–134. doi:10.1016/j.tetasy.2016.10.012
54. Uyanik, M.; Suzuki, D.; Yasui, T.; Ishihara, K. *Angew. Chem., Int. Ed.* **2011**, *50*, 5331–5334. doi:10.1002/anie.201101522
55. Davies, A. T.; Slawin, A. M. Z.; Smith, A. D. *Chem. – Eur. J.* **2015**, *21*, 18944–18948. doi:10.1002/chem.201504256
56. McDaniel, D. H.; Brown, H. C. *J. Org. Chem.* **1958**, *23*, 420–427. doi:10.1021/jo01097a026
57. Crystallographic data for **12** has been deposited with the Cambridge Crystallographic Data Centre (ascension number: CCDC 1987497).
58. Massey, R. S.; Collett, C. J.; Lindsay, A. G.; Smith, A. D.; O'Donoghue, A. C. *J. Am. Chem. Soc.* **2012**, *134*, 20421–20432. doi:10.1021/ja308420c
59. Collett, C. J.; Massey, R. S.; Maguire, O. R.; Batsanov, A. S.; O'Donoghue, A. C.; Smith, A. D. *Chem. Sci.* **2013**, *4*, 1514–1522. doi:10.1039/c2sc22137c
60. Barrios Antúnez, D.-J.; Greenhalgh, M. D.; Brueckner, A. C.; Walden, D. M.; Elías-Rodríguez, P.; Roberts, P.; Young, B. G.; West, T. H.; Slawin, A. M. Z.; Ha-Yeon Cheong, P.; Smith, A. D. *Chem. Sci.* **2019**, *10*, 6162–6173. doi:10.1039/c9sc00390h
61. The research data underpinning this publication can be accessed at doi:<https://doi.org/10.17630/7bc9507c-e817-4da7-988a-02fcd1a6f4fa>.

License and Terms

This is an Open Access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>). Please note that the reuse, redistribution and reproduction in particular requires that the authors and source are credited.

The license is subject to the *Beilstein Journal of Organic Chemistry* terms and conditions: (<https://www.beilstein-journals.org/bjoc>)

The definitive version of this article is the electronic one which can be found at: doi:[10.3762/bjoc.16.129](https://doi.org/10.3762/bjoc.16.129)