

**Manuscript version: Author's Accepted Manuscript**

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

**Persistent WRAP URL:**

<http://wrap.warwick.ac.uk/138965>

**How to cite:**

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

© 2020 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>.



**Publisher's statement:**

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk).

## Optimising the duration of adjuvant trastuzumab in early breast cancer in the UK

Helena M Earl, Louise Hiller, Janet Dunn, Iain Macpherson, Daniel Rea, Luke Hughes-Davies, Karen McAdam, Peter Hall, Janine Mansi, Duncan Wheatley, Jean Abraham, Carlos Caldas, Sophie Gasson, Liz O'Riordan, Maggie Wilcox, David Miles, David Cameron, Andrew Wardley.

Adjuvant trastuzumab for patients with HER2 positive early breast cancer demonstrated significant improvements in both disease-free and overall survival with 12 months treatment (1, 2), and this was NICE approved in the UK in 2006. When the FinHer trial demonstrated similar results with nine weeks trastuzumab (3), there was significant interest in whether shorter durations might be as effective. Additional benefits for patients could be less toxicity, fewer hospital visits and a more rapid return to normal life, with considerable societal benefits of reduced costs. PERSEPHONE was the pragmatic UK duration trial funded by the NIHR HTA, which showed that 6 months adjuvant trastuzumab was non-inferior to 12 months with 4-year disease-free survival rate of 89.4% compared with 89.8% respectively (non-inferiority  $p = 0.01$ ) (4). Less toxicity was reported with 6 months particularly cardiac toxicity and there were cost savings over the first 2 years (5), which were maintained over an average patient's lifetime when extrapolated using an economic model. After the publication of these results in June 2019, the Optimal Duration of Adjuvant Trastuzumab Working Group was convened, comprising a diverse, multi-disciplinary membership. There were representatives from the PERSEPHONE Trial Management Group including patient advocates, the NCRI Breast Group, the Association of Cancer Physicians (ACP), the Royal College of Radiologists (RCR), and the Independent Cancer Patients' Voice (ICPV). By November 2019, both dual antibody treatment with trastuzumab and pertuzumab (<https://www.nice.org.uk/guidance/ta569>) and extended neratinib after single agent trastuzumab (<https://www.nice.org.uk/guidance/ta612>) had been approved by NICE only for those at high risk of recurrence. Therefore, single agent trastuzumab remained standard of care for those at lower risk of recurrence and recommendations were made for these patients.

With the aim of assessing current practice and implementation of the PERSEPHONE results, we surveyed breast oncologists in the UK, for their views on the recommendations from the Working Group for 6 months adjuvant trastuzumab. The following three questions were included in the survey:

1. *Do you agree with the following statement? "Patients with HER2-positive breast cancer who are receiving adjuvant single agent trastuzumab with chemotherapy (concurrent or sequential timing) should be considered for 6 months trastuzumab as standard."*
2. *Do you agree with the following statement? "Patients receiving adjuvant single agent trastuzumab and suffering severe toxicities including cardiac toxicity should be told that receiving only 6 months treatment will not result in significant loss of benefit from trastuzumab"*
3. *Following the results of the PERSEPHONE trial have you reduced trastuzumab duration for any of your patients?*

The survey was hosted by the University of Warwick and used the QUALTRICS online survey tool. It was sent to 330 members of the UK Breast Cancer Group (UKBCG) on 10<sup>th</sup> January 2020. The Warwick Clinical Trials Unit sent a reminder one month later to the principal investigators and recruiting consultants of the PERSEPHONE sites, the majority of whom were in the original mailing.

One hundred and seventeen out of 330 contacted, returned completed questionnaires (35%) from 77 sites. The majority were consultant oncologists (113/117; 97%), more than half of whom (65/117; 56%) were practising in Cancer Centres, and 47 (47/117; 40%) in Cancer Units. Two thirds of respondents (83/117; 71%) were PERSEPHONE investigators or recruiters.

### **Statement 1:**

More than three-quarters of respondents (91/117; 78%, see figure) agreed that for patients receiving single agent trastuzumab, 6 months should be considered as standard. Sixty-eight did not make any qualifying text comments (68/91; 75%), which represents more than half of all respondents (68/117; 58%). 23/91 (25%) of those who agreed with statement 1, qualified their response and 11 (48%) considered 6 months trastuzumab standard for patients with a lower risk of relapse (usually node negative). Of these 11 respondents, three also limited 6 months to patients with ER positive tumours, two to patients with T1 and one to T1b tumours. Five other comments related to the use of single agent paclitaxel for patients at low risk of relapse (APT (6)), and whether the PERSEPHONE results could be applied to these patients. One of these five also commented on the use of trastuzumab and pertuzumab in the neoadjuvant setting. Two comments mentioned the need for change in national guidelines before practice changes. Other comments included: (i) a requirement for longer follow-up; (ii) a question about the neratinib treatment pathway; (iii) shared decision-making with patients discussing risks and benefits; and (iv) two confirming their support.

Respondents who did not agree with the statement (26/117; 22%) shared comments in half (13/26: 50%). Three considered that higher risk patients should be excluded, and 3 felt that longer follow-up and an independent meta-analysis were required before any change in practice. Three respondents expressed concerns about low risk patients who had already de-escalated chemotherapy on the APT regimen (6), with one of these highlighting the predominant use of anthracyclines in the trial. Two respondents referred to PHARE (7) and HORG (8) trials, which had not demonstrated non-inferiority for 6 months. One respondent discussed the uncertainties of duration with patients, and one said that with the increase of neoadjuvant therapy there was no plan to de-escalate trastuzumab.

### **Statement 2:**

Nearly all respondents (114/117; 97%, see figure) agreed with the statement that reassurance should be given to patients who had to stop trastuzumab after 6 months because of severe

toxicities, that there would not be a significant loss of benefit from trastuzumab. Ten respondents made a comment (10/114: 9%), with four simply confirming their views. Two requested a definition of toxicity, and one suggested a minor rewording of the statement. One felt that whilst 12 months should remain the standard, patients who were frail, elderly or who had co-morbidities could be reduced to 6 months. One respondent reported that advice would depend on patients' risk profiles and one reported if toxicities were affecting QOL then 6 months' was reasonable. Three respondents who did not agree with the statement made no comments.

### **Statement 3:**

Just under half the respondents (53/117: 45%, see figure) said they had reduced trastuzumab for some of their patients since the PERSEPHONE results were published, and of these 25/53 (47%) added a comment. The most frequent (19/25: 76%) related to stopping after 6 months due to cardiac or other toxicity. Three respondents discussed 6 months treatment with patients, two in a selective way with low risk patients and one as routine. This last respondent also discussed stopping trastuzumab and pertuzumab after 6 months with a pathological complete response to neoadjuvant treatment. One respondent was giving 6 months in T1N0 patients with paclitaxel only chemotherapy (APT) (6), but expressed concern about reducing chemotherapy as well as duration of trastuzumab in these patients. One respondent excluded patients receiving neoadjuvant therapy and patients with >3 node positive disease from 6 months therapy. One had switched to 6 months in all lower risk patients including those receiving weekly taxol, and those with concerns about cardiotoxicity.

Just over half of respondents (64/117: 55%) reported not reducing trastuzumab duration, with 24/64 (37.5%) supplying comments. The most common reason was waiting for local/national guidelines to change (13/24: 54%, see figure). Three other respondents said that for low risk patients they had reduced chemotherapy to paclitaxel only (APT) (6), and for high risk patients escalated to dual antibodies. Two respondents said they would only reduce trastuzumab duration for toxicity, which one reported they were doing already. Two were not involved in decision-making for these patients. Other comments included: *'not an easy change to sell - not complete consensus with colleagues'*; *'not yet'*; *'have offered but none have accepted'*; *'considering reduction now there are published results'*.

### **Single agent taxane regimens**

Nine respondents commented at some point in the questionnaire on low risk patients who are receiving paclitaxel for 12 weeks with concurrent trastuzumab continued for 12 months (6). The number of patients receiving taxane-only chemotherapy within the PERSEPHONE trial is very small (35, 12 month patients and 38, 6 month patients) (9). Hence, it is impossible to make any recommendations based on such limited data. However since the trial results as a whole confirm non-inferiority for 6 months treatment it is reasonable to conclude that this can apply to all types of chemotherapy.

## Summary

The majority of respondents (78%) agreed that 6-months trastuzumab should be a standard option for patients with lower risk disease receiving single agent treatment. In Scotland the situation is different and dual therapy is not approved for high risk patients. Hence, we would advise that those in Scotland who elsewhere in the UK would be eligible for dual antibody therapy or extended neratinib, should continue with 12 months' trastuzumab. There was a clear overwhelming consensus (97%) that with severe toxicity, patients should be reassured that stopping at six months would not result in a significant loss of benefit from trastuzumab. Although the majority agreed with 6-months for patients with lower risk disease, it was notable that over half had not yet introduced this in their clinical practice. This was despite an interval of 19 months and 7 months, respectively, since initial presentation (10) and subsequent full publication (4) of the PERSEPHONE results. This is not unexpected given the well-documented barriers to de-escalation of cancer therapy (11). Although not specifically explored in our survey, it is likely that the results of PHARE (7) and the HORG (8) study may have led to uncertainty around the strength of the evidence provided by PERSEPHONE. However, it is also crucial to recognise that movement towards de-escalation of therapy is not determined solely by scientific data. Historical, economic, professional, and social factors may all favour entrenched behaviour, even in the face of robust evidence (12). Consistent with professional and organisational norms being powerful drivers of clinician behaviour, the most common reason given for not reducing trastuzumab duration was waiting for local or national guidelines to change.

The unprecedented crisis of the COVID-19 pandemic significantly increases the acute risks for cancer patients attending hospital for treatment. The UKBCG has issued prioritisation guidelines for breast cancer treatments (13). On the strength of the PERSEPHONE data, the UKBCG executive committee has advised that those at low risk of recurrence receiving single agent trastuzumab should stop at 6 months with immediate effect, since the acute risks of attending hospital clinics are significant and outweigh any minimal loss of long-term benefit. Many hospitals have implemented this prioritisation guidance.


**Figure legend:** Responses to the three statements within the survey

## References


1. Piccart-Gebhart MJ, Procter M, Leyland-Jones B, Goldhirsch A, Untch M, Smith I, et al. Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *The New England journal of medicine*. 2005;353(16):1659-72.
2. Romond EH, Perez EA, Bryant J, Suman VJ, Geyer CE, Jr., Davidson NE, et al. Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *The New England journal of medicine*. 2005;353(16):1673-84.

3. Joensuu H, Kellokumpu-Lehtinen PL, Bono P, Alanko T, Kataja V, Asola R, et al. Adjuvant docetaxel or vinorelbine with or without trastuzumab for breast cancer. *The New England journal of medicine*. 2006;354(8):809-20.
4. Earl HM, Hiller L, Vallier AL, Loi S, McAdam K, Hughes-Davies L, et al. 6 versus 12 months of adjuvant trastuzumab for HER2-positive early breast cancer (PERSEPHONE): 4-year disease-free survival results of a randomised phase 3 non-inferiority trial. *Lancet*. 2019;393(10191):2599-612.
5. Hulme C, Hall P, Shinkins B, Chehadah F, McCabe C, Hiller L, et al., editors. PERSEPHONE: 6 versus 12 months (m) of adjuvant trastuzumab in patients (pts) with HER2 positive (+) early breast cancer (EBC): Cost effectiveness analysis results. ESMO Munich, 2018. Oral Presentation, Early Breast Cancer Session. *Annals of Oncology*; Volume 29, Issue suppl\_8, October 2018.
6. Tolaney SM, Guo H, Pernas S, Barry WT, Dillon DA, Ritterhouse L, et al. Seven-Year Follow-Up Analysis of Adjuvant Paclitaxel and Trastuzumab Trial for Node-Negative, Human Epidermal Growth Factor Receptor 2-Positive Breast Cancer. *J Clin Oncol*. 2019;37(22):1868-75.
7. Pivot X, Romieu G, Debled M, Pierga JY, Kerbrat P, Bachelot T, et al. 6 months versus 12 months of adjuvant trastuzumab in early breast cancer (PHARE): final analysis of a multicentre, open-label, phase 3 randomised trial. *Lancet*. 2019;393(10191):2591-8.
8. Mavroudis D, Saloustros E, Malamos N, Kakolyris S, Boukovinas I, Papakotoulas P, et al. Six versus 12 months of adjuvant trastuzumab in combination with dose-dense chemotherapy for women with HER2-positive breast cancer: a multicenter randomized study by the Hellenic Oncology Research Group (HORG). *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO*. 2015;26(7):1333-40.
9. Earl HM, Hiller L, Vallier A-L, Loi S, McAdam K, Hughes-Davies L et al. Six versus 12 months' adjuvant trastuzumab in patients with HER2-positive early breast cancer: the PERSEPHONE non-inferiority Randomised Controlled Trial (RCT). *Health Technology Assessment Journal*. 2020; Accepted January 2020.
10. Earl HM, Hiller L, Vallier A-L, Loi S, Howe D, Higgins HB, et al. PERSEPHONE: 6 versus 12 months (m) of adjuvant trastuzumab in patients (pts) with HER2 positive (+) early breast cancer (EBC): Randomised phase 3 non-inferiority trial with definitive 4-year (yr) disease-free survival (DFS) results. *J Clin Oncol*. 2018;36 (suppl; 506 abstr). Oral presentation.
11. Norton WE, Chambers DA, Kramer BS. Conceptualizing De-Implementation in Cancer Care Delivery. *Journal of Clinical Oncology*. 2018;37(2):93-6.
12. Montini T, Graham ID. "Entrenched practices and other biases": unpacking the historical, economic, professional, and social resistance to de-implementation. *Implementation Science*. 2015;10(1):24.
13. UKBCG. Prioritisation Guidance for Systemic Breast Cancer Treatment during COVID-19. 2020, to UK Breast Cancer Group Members.

1) Patients with HER2-positive breast cancer who are receiving adjuvant single agent trastuzumab with chemotherapy (concurrent or sequential timing) should be considered for 6 months trastuzumab as standard.



2) Patients receiving adjuvant single agent trastuzumab and suffering severe toxicities including cardiac toxicity should be told that receiving only 6 months treatment will not result in significant loss of benefit from trastuzumab



3) Following the results of the Persephone trial have you reduced Herceptin duration for any of your patients?

