

**Original citation:**

Barnes, A. D., et al. Kinematical studies of the low-mass X-ray binary GR Mus (XB 1254-690). Monthly Notices of the Royal Astronomical Society, 380(3), pp. 1182-1190.

**Permanent WRAP url:**

<http://wrap.warwick.ac.uk/31364>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes the work of researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

The definitive version is available at [www.blackwell-synergy.com](http://www.blackwell-synergy.com)  
DOI: 10.1111/j.1365-2966.2007.12174.x

**A note on versions:**

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP url' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)



# Kinematical Studies of the Low Mass X-Ray Binary GR Mus (XB 1254–690)

A. D. Barnes<sup>1\*</sup>, J. Casares<sup>2</sup>, R. Cornelisse<sup>1</sup>, P. A. Charles<sup>1,3</sup>, D. Steeghs<sup>4,5</sup>,  
R.I. Hynes<sup>6</sup> and K. O’Brien<sup>7</sup>

<sup>1</sup>*School of Physics and Astronomy, University of Southampton, SO17 1BJ, UK*

<sup>2</sup>*Instituto de Astrofísica de Canarias, 38200 La Laguna, Tenerife, Spain*

<sup>3</sup>*South African Astronomical Observatory, PO Box 9, Observatory 7935, Cape, South Africa*

<sup>4</sup>*Harvard-Smithsonian Center for Astrophysics, MS-67, 60 Garden Street, Cambridge, MA 02138, USA*

<sup>5</sup>*Department of Physics, University of Warwick, Coventry, CV4 7AL, UK*

<sup>6</sup>*Department of Physics and Astronomy, 202 Nicholson Hall, Louisiana State University, Baton Rouge, LA 70803, USA*

<sup>7</sup>*European Southern Observatory, Casilla 19001, Santiago 19, Chile*

## ABSTRACT

We present simultaneous high-resolution optical spectroscopy and X-ray data of the X-ray binary system GR Mus (XB 1254–690), obtained over a full range of orbital phases. The X-ray observations are used to re-establish the orbital ephemeris for this source. The optical data includes the first spectroscopic detection of the donor star in this system, through the use of the Doppler Tomography technique on the Bowen fluorescence blend ( $\sim 4630\text{--}4650$  Å). In combination with an estimate for the orbital parameters of the compact object using the wings of the He II  $\lambda 4686$  emission line, dynamical mass constraints of  $1.20 \leq M_X/M_\odot \leq 2.64$  for the neutron star and  $0.45 \leq M_2/M_\odot \leq 0.85$  for the companion are derived.

**Key words:** binaries: close - stars: binaries: individual: GR Mus, XB 1254–690 - binaries: spectroscopic - X-rays: binaries.

## 1 INTRODUCTION

XB 1254–690 is a persistently bright, low-mass X-ray binary (LMXB). It was identified with a faint blue star (GR Mus,  $V = 19.1$ ), which exhibited the Bowen blend of N III and C III in emission (Griffiths et al. 1978). The object produced Type 1 X-ray bursts (Mason et al. 1980; Courvoisier et al. 1986), indicating the presence of a neutron star as the compact object. Dips of up to 95% of the 1–10 keV flux, with a recurrence period of  $P_X = 3.88 \pm 0.15$  hr and lasting  $\sim 0.8$  hr per cycle, were discovered in *EXOSAT* data by Courvoisier et al. (1986). The dips are caused by obscuration of the central source by a bulge on the outer edge of the accretion disc, implying a moderately high inclination.

From *V*-band observations of the optical counterpart, Motch et al. (1987) determined an optical ephemeris where minimum light occurs at  $T_{min} = \text{JD } 2,445,735.693 (4) + 0.163890 (9)$  days, equivalent to a period of  $P_{opt} = 3.9334 \pm 0.0002$  hr. This broad optical modulation is probably due to the changing visibility of the heated face of the secondary star in the system, with an additional contribution from the

X-ray heated bulge, although other explanations such as an asymmetrical disc or occultation of the disc by the companion cannot be ruled out. Within the errors, the optical and X-ray period measurements are consistent, with later X-ray observations indicating that the optical modulation provides a more accurate measurement of the orbital period (Smale et al. 2002; Levine & Corbet 2006). A short section of simultaneous X-ray and optical coverage showed that the optical minimum occurs  $\sim 0.16$  in phase after the centre of the X-ray dips (Motch et al. 1987).

Simple geometrical modelling is possible by considering the X-ray illuminated area of the companion star, disc radius, disc opening angle, and inclination at given distances which are compatible with the observed optical flux and the amplitude of the light curve. This leads to constraints on the source inclination of  $65^\circ\text{--}73^\circ$ , and on the distance of 8–15 kpc (Motch et al. 1987).

The first mass constraints for this system were based upon a unique relation between the orbital period and the mass of the companion, assuming that the companion is a low mass, zero-age main sequence star filling its Roche lobe (Warner 1976). Using this approximation, Courvoisier et al. (1986) estimate the mass of the companion to be in the region of  $\sim 0.45M_\odot$ .

\* E-mail: adb@astro.soton.ac.uk (ADB)

**Table 1.** Log of observations of GR Mus / XB 1254–690

HJD	Observatory	Total Int. Time
2453151.52	VLT	3.75 h
2453151.51	RXTE	4.75 h
2453152.52	VLT	4 h
2453152.69	RXTE	5.25 h
2453153.53	VLT	2 h

The first kinematic mass constraints were proposed by Cowley et al. (1988), who obtained intermediate resolution ( $\sim 3\text{\AA}$ ) spectra using the 4m telescope at the Cerro Tololo Inter-American Observatory. Assuming that velocities derived from the He II  $\lambda 4686$  emission line centroid represent the motion of the compact star and not, for example, the streaming of gas between the stars or out of the system, then approximate stellar masses can be derived using the mass function. The result is consistent with a low-mass main-sequence star ( $M \simeq 0.5\text{--}0.8 M_{\odot}$ ) and a canonical  $1.4 M_{\odot}$  neutron star.

In the remainder of this paper we will attempt to confirm and better constrain these mass estimates using high-resolution blue optical spectroscopy. We will measure the wings of the He II  $\lambda 4686$  emission line to gain an estimate for the compact object velocity (Section 3.3), and use the Bowen Fluorescence technique (Steeghs & Casares 2002) to measure the donor star velocity (Section 3.4). We also obtained simultaneous X-ray observations (see Section 3.1) in order to re-establish the ephemeris of Motch et al. (1987). This cannot be extended to the observations described herein because the propagated uncertainty in phase is now comparable to the orbital period itself.

## 2 OBSERVATIONS

We have obtained simultaneous optical and X-ray data for GR Mus / XB 1254–690. The journal of observations is given in Table 1.

### 2.1 Optical data

Phase-resolved spectroscopy of GR Mus was obtained from 26 to 28 May 2004 using the FORS 2 spectrograph mounted on the VLT/UT4 at ESO’s Paranal observatory. During each night we observed GR Mus for  $\simeq$  one full orbit, resulting in a total of 38 spectra with an integration time of 900s each. We used the 1400V volume-phased holographic grism, and a slit width of  $0.7''$ , giving a wavelength coverage of  $\lambda\lambda 4513\text{--}5814$  with a resolution of  $70 \text{ km s}^{-1}$  (FWHM). The seeing during these observations varied between 0.5 and 2.1 arcsec. The slit was orientated at a position angle of  $88.15^{\circ}$  to include a comparison star in order to calibrate slit losses. We observed the flux standard Feige 110 with the same instrumental set-up in order to correct for the instrumental response. He, Ne, Hg and Cd arc lamp exposures were taken during daytime for the wavelength calibration scale. We de-biased and flat-fielded all of the images and used optimal extraction techniques to maximise the signal-to-noise ratio of the extracted spectra (Horne 1986). The pixel-to-wavelength scale was derived through polynomial fits to a large number of identified reference lines resulting in a dispersion of  $0.64 \text{ \AA}$

$\text{pix}^{-1}$ . Velocity drifts due to instrumental flexure (found to be always  $< 5 \text{ km s}^{-1}$ ) were corrected by cross-correlation with the sky spectra.

### 2.2 X-ray data

Simultaneous X-ray data from XB 1254–690 were obtained using the Proportional Counter Array (PCA) on board the *Rossi X-Ray Timing Explorer (RXTE)*. The Proportional Counter Array (PCA; for a detailed description, see Jahoda et al. (1996)) onboard the *RXTE* satellite consists of an array of 5 co-aligned Proportional Counter Units (PCUs) that are sensitive to photons of energy 2 to 60 keV with a total collecting area of  $6500 \text{ cm}^2$ . We obtained 36 ks of total coverage during our optical observations. For our analysis we only used the data from the Standard 2 configuration, with a time resolution of 16 seconds.


## 3 ANALYSIS

### 3.1 Determination of orbital phase


The ephemeris of Motch et al. (1987) is now out of date because the propagated uncertainty in phase is over a factor of 2 greater than the orbital period. We therefore need to make use of our contemporaneous *RXTE* observations to set a new zero point using the X-ray dips (note that we still use the Motch et al. (1987) orbital period). This is not quite as simple as it would first appear since there are large variations in dip depth and structure that are apparent from observation to observation, as well as cycle to cycle (Courvoisier et al. 1986). Additionally, a note of caution must be sounded since in a number of earlier *RXTE* and *BeppoSAX* observations (Smale & Wachter 1999; Iaria et al. 2001), X-ray dipping was not observed at all, with an accompanying decrease in the mean optical variability of the source. With the combination of these effects, it was suggested that the vertical structure on the accretion disc edge, associated with the impact point of the accretion stream, had decreased in angular size at the time of these earlier observations. The cause of this intermittent reduction in opening angle is not understood. Our observations include just one deep dipping episode (Fig. 1), co-incident with our first night of optical observations.

XB 1254–690 typically exhibits ‘shoulders’ of dipping, i.e. regions both before and after the deep dipping in which there is a small decrease of intensity. Each shoulder of the dip lasts  $\sim 650\text{--}925$  seconds, during which the total count rate in the 2–25 keV band drops by  $\sim 15\%$  with no strong changes in the hardness ratio. The shoulder is followed by an interval of strongly variable deep dipping lasting  $\sim 1100$  seconds (Smale et al. 2002), with associated strong changes in the hardness ratio (Fig. 2). The hardening of the spectra in the dips is most likely to be due to absorption by cold matter.

Our observation of a dipping episode opens with the system already in the first shoulder phase, the onset of deep dipping occurring approximately 200–250 seconds into the observation (before a brief return of the source intensity to the non-dip level), and then continuing to capture the full




**Figure 1.** 2–20 keV X-ray lightcurve showing the deep dip used to calibrate a new ephemeris for the system. The dipping episode appears to begin shortly before JD 2453151.64, and occurs for the typical dip duration of  $\sim 1100$  seconds (Smale et al. 2002).



**Figure 2.** 2–4 keV (top) and 4–9 keV (centre) X-ray lightcurves displaying the deep dipping episode. The bottom panel displays hardness as a function of intensity. There is an anti-correlation between intensity and hardness ratio during the deep dipping episode, which begins shortly after 200 seconds on the above scale.

gress from deep dipping, into the shoulder and then returning to the continuum level. Note that the return of the source intensity to the non-dip level during this episode is not unusual, and the light curve can in fact be interpreted as superpositions of small micro-dips (e.g. Uno et al. 1997).

Unfortunately, the lack of information regarding the full ingress to deep dipping through the preceding shoulder introduces a degree of uncertainty into our estimate of the dip centre. For example we could potentially be observing only a small section of an extended deep dipping episode (e.g. Uno et al. 1997, report dips lasting for  $\sim 40$  minutes). If this is the case, the absence of the beginning of the dip will cause the estimate for dip centre to be incorrect. Nevertheless, we do have cause for optimism. The deep dipping which we observe is approximately 1100 seconds in length, equivalent to the typical period of a dipping episode in XB 1254–690 (see above). This is a strong indicator that we have full, or nearly-full, coverage of the deep dipping phase. We estimate



**Figure 3.** Continuum lightcurve of GR Mus, normalised to the flux level of the first observation in our dataset. Note that the errors on each measurement are smaller than the plotted points.

the dip centre by eye, taking the mid-point between the onset of and exit from deep dipping (which can be determined with the help of the associated hardness ratio changes, see Fig. 2). This occurs at  $T_{dip} = \text{JD } 2453151.647$  (3). From Motch et al. (1987) we know that the dip centre has an average orbital phase of 0.84 (with phase 0 occurring at optical minimum). We therefore estimate a new value for  $T_0$  of JD 2453151.509 (3).


There are problems with the reliability of an ephemeris generated from just a single dipping episode, not least because we do not have full coverage either side of the dip. It is therefore hard to estimate the dip centre with absolute confidence. However, we are encouraged by the fact that our dip lies within 0.05 phase of the 1999 dips (the propagated phase uncertainty from these dips is of the order  $\sim 0.6$ ) reported by in’t Zand et al. (2003). More importantly, the continuum lightcurve from the resulting ephemeris possesses the same phasing relation reported by Motch et al. (1987) (Fig. 3)<sup>1</sup>. In light of this evidence we are confident that we have identified the dip centre correctly and we will use this zero-point to calculate orbital phases throughout the remainder of this paper.

### 3.2 Optical emission spectrum

The continuum subtracted average of the 38 individual optical spectra, corresponding to an exposure time of nearly 10 hours, is given in Fig. 4. The overall spectrum resembles that of other LMXBs, with high excitation lines superposed on a blue continuum. This is typical of strongly X-ray heated accretion discs (e.g. van Paradijs & McClintock 1995).

The blue end of the optical spectrum in this system is dominated by the Bowen emission blend of C III and N III. He II  $\lambda 4686$  is also seen strongly in emission, in addition to weaker H $\beta$  and He II  $\lambda 5411$ . The absorption component longward of H $\beta$  is observed in a few other LMXBs

<sup>1</sup> Note that the large intrinsic scatter in the optical continuum lightcurve makes the generation of a new ephemeris solely from a relatively limited dataset such as ours rather difficult.




**Figure 4.** Average normalised spectrum of GR Mus. The most prominent emission lines are indicated, including the Bowen blend of C III and N III, He II and H $\beta$ . The emission line near 5577Å is a night sky feature.

such as X 1822–371 (Casares et al. 2003), XTE J2123–058 (Hynes et al. 2001), GX 339–4 (Buxton & Vennes 2003) and MM Ser (Hynes et al. 2004).

The narrow emission feature at 5577Å is [O I], a strong night-sky line useful for confirming the veracity of our wavelength calibration. The broad features between 4900–5300Å remain unidentified. The absence of these features in our comparison star lead us to believe that they are real. We also see some of these ‘bumps’ in other LMXBs (e.g. GX 9+9, X 1822–371, V801 Ara, V926 Sco), and perhaps they are blends of a number of different weak emission lines. Part of this could be He I  $\lambda$ 4922 and  $\lambda$ 5016, and also Fe II  $\lambda$ 4924,  $\lambda$ 5018,  $\lambda$ 5169 and  $\lambda$ 5297. There are also several O II and O III lines between 4891–5006Å which may explain the large bump at  $\sim$ 4930Å. Note also that the Bowen blend (and to a lesser extent the He II  $\lambda$ 4686 emission line) appears to be sitting upon a broad emission ‘hump’. This has been noted in other LMXBs (e.g. V801 Ara; Augusteijn et al. 1998), and attributed to a blend of Fe II emission lines (Schachter et al. 1989).

In Fig. 5, we display the Bowen blend and He II  $\lambda$ 4686 in the form of a trailed spectrogram. All of the individual spectra were first normalized to the continuum using a third-order polynomial fit, and then phase-folded into 15 bins. The spectra were then smoothed using a Gaussian filter with a FWHM of two pixels. We see a clear S-wave structure modulated on the orbital period in the He II  $\lambda$ 4686 emission. The blue-to-red crossing phase is  $\sim$ 0.6 and the semi-amplitude  $\sim$ 250 km s $^{-1}$ . This behaviour indicates a close association with the accretion disc or compact object. There is also an additional, weaker, component present. Again, this is modulated on the orbital period, approximately a quarter cycle out of phase with the more prominent component. There also appear to be faint traces of two anti-phased sharp components in the Bowen blend though the broad underlying component moves in phase with the accretion disc S-wave (maximum blue-shift at phase 0.25). The tomograms discussed in Section 3.4 will help to characterise these spectral trails more thoroughly.

The H $\beta$  line is shown in the right-hand panel in Fig. 5,



**Figure 6.** Velocity profile of the He II  $\lambda$ 4686 emission line displaying a complex core structure. The extreme blue wing is contaminated by the Bowen emission blend.


along with the broad emission feature at  $\sim$ 4930Å. The absorption component longward of H $\beta$  is present at all phases, and in common with the emission component displays an S-wave approximately in phase with the expected motion of the accretion disc. The broad feature at  $\sim$ 4930Å also displays a S-wave. The He II  $\lambda$ 5411 emission line is too weak in the individual spectra to produce a trailed spectrogram.

### 3.3 Motion of the compact object

Early attempts to determine any kinematic constraints on this system met with limited success. Motch et al. (1987) noted that the relative weakness of the emission lines in their spectra did not allow for the study of their variation with orbital phase. However, they did exclude the possibility of orbital motion of He II  $\lambda$ 4686 (which requires a source of highly ionising radiation, and is typically seen in accretion discs) in excess of 300 km s $^{-1}$ . Later observations by Cowley et al. (1988) did result in a measurement of the same He II line’s core velocity, which possessed a semi-amplitude of  $114 \pm 13$  km s $^{-1}$ . However, this is not a particularly reliable measurement of the compact object velocity. The line core can often be heavily contaminated by other emission sites (see Fig. 6), for example from the accretion stream or ‘hot-spots’ on the accretion disc itself.

Instead, the compact object velocity can be better estimated by examining the wings of the emission line. These wings find their origin in the high-velocity gas of the inner accretion disc, very close to the compact object itself. Presumably this gas will closely replicate the motion of the compact object, and it is far less likely that alternative emission sites will provide a strong contaminating effect. However, this method is not immune to contamination and we must be cautious in accepting the results of this analysis.

We have followed the double-Gaussian technique (Schneider & Young 1980) to estimate both the velocity semi-amplitude  $K_1$  and the systemic velocity  $\gamma$  of the compact object from He II  $\lambda$ 4686. We employed a double Gaussian bandpass with FWHM of 100 km s $^{-1}$  and Gaussian separations of 300–1700 km s $^{-1}$  in steps of 100 km s $^{-1}$ . Radial velocity curves of line sections with separations of 700–



**Figure 5.** Trailed spectra of the Bowen complex and He II  $\lambda 4686$  (left) and the  $H\beta$  region (right). The data are plotted twice for clarity. Note that the wavelength scales differ. There are clear multiple S-waves evident in the He II emission line,  $\sim 0.25$  out of phase with each other, and also in the Bowen complex. The absorption component redwards of  $H\beta$  is present at all phases.


$1400 \text{ km s}^{-1}$  yield consistent results with a median blue-to-red crossing phase of  $0.58 \pm 0.03$ , median velocity semi-amplitude  $K_1 = 130 \pm 16 \text{ km s}^{-1}$  and median systemic velocity  $\gamma = 173 \pm 12 \text{ km s}^{-1}$  (Fig. 7). The errors are conservatively taken from the maximum error obtained from sine wave fits to the results from each individual line section. Separations below  $700 \text{ km s}^{-1}$  suffer badly from contamination by the line core, whilst separations over  $1400 \text{ km s}^{-1}$  become corrupted by continuum noise and the Bowen emission blend. The small difference in phasing (0.08 orbits; slightly larger than the ephemeris uncertainty of  $\sim 0.02$  orbits) indicates that even the line wings may be contaminated to some extent by inhomogeneities in the disc. This contamination will also affect the measurements of  $K_1$  and  $\gamma$  to some extent. Nevertheless, this remains the best method for estimating  $K_1$  from our own and other currently available data.

Unfortunately, the He II  $\lambda 5411$  and  $H\beta$  emission lines, which are clearly seen in our averaged spectrum (Fig. 4), are too weak in the individual spectra for a similar analysis. The fitting procedures are simply dominated by noise.

### 3.4 Doppler tomography

We can use Doppler tomography (Marsh & Horne 1988) to measure the radial velocity amplitudes of any weak, narrow, emission lines that may be present. By mapping the observed data onto a velocity coordinate frame, Doppler tomography makes use of all data at once and can thus be used to search for features that are too weak to be distinguished in each individual spectrum. One effectively resolves the distribution of line emission in the co-rotating frame of the binary system, providing an excellent tool for identifying the origin and kinematics of the various emission components. It also allows us to map the emission line distribution from the broader He II and  $H\beta$  lines.

Secondary star emission is readily identifiable in Doppler tomograms since the solid body rotation of the Roche lobe is mapped to a corresponding Roche lobe area along the positive  $V_y$ -axis. For reference, in Figs. 8–11 we overplot the area of the Roche lobe and the centres of mass of the donor star (uppermost cross), full system (map centre) and the compact object (lower cross) assuming a mass ratio of  $\sim 0.4$  (equivalent to the mass ratio proposed by Cowley et al. 1988) and compact object velocity of  $130$




**Figure 7.** Diagnostic diagram for the He II  $\lambda 4686$  emission line wings, displaying the systemic velocity  $\gamma$  (top), phasing (centre) and radial velocity  $K_1$  (bottom).

$\text{km s}^{-1}$ . Additionally, we have overplotted the ballistic accretion stream for such a system. The circle, centred upon the compact object and with radius  $600 \text{ km s}^{-1}$ , indicates the common emission regions from a generic LMXB accretion disc. We employed a maximum entropy implementation of Doppler tomography whereby the data are fitted under the added constraint of maximizing the entropy (i.e. image smoothness) of the tomogram (Marsh & Horne 1988). This reduces the presence of noise artifacts in the recovered tomograms and allows for a simultaneous fit to a number of heavily blended lines.

In Fig. 8 we display a tomogram of the He II  $\lambda 4686$  emission line. This displays a clear enhancement in the lower left quadrant and an additional component close to the accretion stream. We do not observe the classic ring-like distribution of the accretion disc, presumably due to the strength of the line core features vastly overwhelming the high velocity inner disc emission. For this reason we cannot be fully confident in the results of the double Gaussian technique (see Section 3.3), though this remains at present the best method for estimating  $K_1$ .

The enhancement in the (lower) left quadrant is also seen in the tomograms of H $\beta$  (Fig. 9) and He II  $\lambda 5411$  (Fig. 10). Whilst this phenomenon is a common feature in the remarkable SW Sex-type systems (Hellier 2000), it has also been noted in some other LMXBs, notably XTE J2123–058 (Hynes et al. 2001), GX 9+9 (Cornelisse et al. 2006) and J0422+32 (Casares et al. 1995). However, in this case (and in the case of X 1636–536 and X 1735–444, Casares et al. 2006) the emission is not quite so localised in the lower-left quadrant, but is present in a broad band on the left-hand side of the maps. Rather than being an analogue to the highly magnetic SW Sex-type systems, it is likely that this emission is produced in an extended disc bulge. In most LMXBs, the He II emission is predominantly produced in a hotspot where the accretion stream impacts the disc (thus producing a spot in the upper left quadrant of a Doppler map, e.g. Pearson et al. 2006) or is more evenly distributed around the bulk of the accretion disc. If the emitting gas was symmetrically distributed around the primary, the Doppler image would reveal a circular distribution centred on  $V_x = 0$ ,




**Figure 8.** Doppler tomogram reconstructed from the He II  $\lambda 4686$  emission line using maximum entropy optimisation,  $\gamma = 185 \text{ km s}^{-1}$ . The area of the Roche lobe, the ballistic accretion stream and the centres of mass of the donor star (uppermost cross), system (map centre) and the compact object (lower cross) are plotted assuming a mass ratio of  $\sim 0.4$  and compact object velocity of  $130 \text{ km s}^{-1}$ . The circle is centred upon the compact object and possesses a radius of  $600 \text{ km s}^{-1}$ .

$V_y = -K_1$ . This would have been a useful way of confirming the  $K_1$  estimate derived from the wings of the He II  $\lambda 4686$  emission line.


Like He II  $\lambda 4686$ , the He II  $\lambda 5411$  tomogram also indicates the presence of some diffuse emission at velocities which partially correspond to the Roche lobe of the donor star. However, as this emission is diffuse in the Doppler map, extending over a broad range of velocities, it is difficult to directly associate this extended emission with the donor star. Fluorescent emission from the donor in this case should manifest itself as a tightly concentrated spot situated on the positive  $V_y$  axis.

In Fig. 11 we present a tomogram of the N III  $\lambda 4640$  emission line. The maximum entropy method of producing Doppler tomograms allows for a simultaneous fit to a number of different lines to produce one map. However, in this case the N III  $\lambda 4640$  line strongly dominates over any others in the Bowen blend. We see two regions of enhanced emission. The brighter of the two spots occurs at a combination of phasing and velocity where we would expect the donor star to be fluorescing. The small horizontal offset of  $V_x \approx 35 \pm 30 \text{ km s}^{-1}$  is consistent with the errors in our ephemeris ( $\pm 0.02$  in phase).


The second spot occurs at a much lower velocity, and is  $1\text{--}2 \sigma$  weaker (relative to the local noise level) than the first. It is difficult to provide a truly satisfactory explanation for its origin. Perhaps it traces some form of low velocity outflow (e.g. disc wind), though why it should be preferentially observed at an apparent phase of  $\sim 0.25$  remains a mystery



**Figure 9.** Doppler tomogram reconstructed from the  $H\beta$  emission line using maximum entropy optimisation,  $\gamma = 185 \text{ km s}^{-1}$ . The symbols are as for Fig. 8.



**Figure 11.** Doppler tomogram reconstructed from the  $N \text{ III } \lambda 4640$  emission line using maximum entropy optimisation,  $\gamma = 185 \text{ km s}^{-1}$ . The symbols are as for Fig. 8.




**Figure 10.** Doppler tomogram reconstructed from the  $\text{He II } \lambda 5411$  emission line using maximum entropy optimisation,  $\gamma = 185 \text{ km s}^{-1}$ . The symbols are as for Fig. 8.

(orbital phase can be considered to increase in a clockwise-fashion around a Doppler map).

If we presume that the bright spot in Fig. 11 is caused by the irradiated inner face of the donor star, this is an excellent tracer for the kinematics of the system. The precise location of the spot can be measured using a two-dimensional Gaussian fit (though bearing in mind that the actual shape of the spot is not expected to be Gaussian), providing a velocity of  $K_{\text{em}} = 245 \pm 30 \text{ km s}^{-1}$ . Using this value we can shift all of our spectra into the rest-frame of the (irradiated face of the) donor. A sharp  $N \text{ III } \lambda 4640$  component is clearly resolved (Fig. 12). However, since this is a measure of the motion of the irradiated inner face of the donor star, it only provides us with a lower-limit on the velocity of the centre of mass (see e.g. Steeghs & Casares 2002; Muñoz-Darias et al. 2005).

One final piece of information remains to be extracted from the Doppler tomograms; assigning the correct systemic velocity should produce the most sharply defined features in a Doppler map. We used the  $\text{He II}$  and  $N \text{ III}$  Doppler maps to verify the systemic velocity. The  $\chi^2$  value of the map was calculated for a range of  $\gamma$ , and the best fit in terms of minimal  $\chi^2$  was achieved for  $\gamma = 192 \pm 2 \text{ km s}^{-1}$  for the  $N \text{ III}$  map and  $\gamma = 185 \pm 2 \text{ km s}^{-1}$  for  $\text{He II } \lambda 4686$ . These are both consistent with, if slightly higher than, the results derived from the  $\text{He II } \lambda 4686$  wings in Section 3.3. The complex, asymmetric core profile of the  $\text{He II}$  emission line could be contaminating one wing more than the other, leading to this small offset. We heretofore adopt the value for  $\gamma$  derived from the  $\text{He II } \lambda 4686$  Doppler map. This should be more reliable than the  $N \text{ III}$  map since only a single, isolated emission line is involved in the fitting process, rather than the more complicated blend of Bowen lines.



**Figure 12.** Average spectrum created by combining all of the individual unshifted spectra from our observations (top), showing the Bowen emission blend and He II  $\lambda 4686$  region. Below this is shown the average spectrum when all of the individual spectra are shifted into the rest-frame of the donor star. A sharp N III  $\lambda 4640$  component is clearly resolved.

## 4 DISCUSSION


### 4.1 Systemic velocity

Whilst the systemic velocities inferred from both the wings of the He II  $\lambda 4686$  line and the Doppler tomograms are self-consistent ( $\sim 185 \text{ km s}^{-1}$ ), they are considerably higher than the earlier estimate by Cowley et al. (1988) ( $98 \pm 10 \text{ km s}^{-1}$ ). This measurement was obtained from rather low quality data, with poor spectral and temporal resolution. Additionally, they based their measurements on the motion of the He II  $\lambda 4686$  line core. There are a number of different emitting regions which contribute to this line, leading to a complex core structure (see e.g. Figs. 6, 8). A basic measurement of the line centre under the assumption that this will accurately trace the motion of the compact object will therefore be subject to large systematic errors.


In light of this, we believe that our estimate for the systemic velocity will be a better reflection of the true value. Although large, our measurements do seem reasonable in the context of the high galactic latitude ( $b = -6.42$ ) of GR Mus; the neutron star is likely to have received a substantial ‘kick’ out of the Galactic Plane at birth. A number of statistical studies on pulsar velocities have been carried out. These studies give a mean birth velocity of  $100\text{--}500 \text{ km s}^{-1}$ , with some possessing velocities of  $\geq 1000 \text{ km s}^{-1}$  (Wang et al. 2006).

### 4.2 Interpretation of the Bowen blend Doppler map

The N III Doppler map (Fig. 11) appears to display enhanced emission in the region of and at the phasing where we would expect to see the companion star Roche lobe, in addition to an abundance of noisy structure in the outer regions of the map. A useful way to confirm the validity of the Doppler map is to observe the behaviour of sharp components in our trailed spectrogram (Fig. 5). This is not trivial due to the



**Figure 13.** The right hand panel displays the observed trailed spectra of the He II  $\lambda 4686$  emission line and the left hand panel shows the ideal trailed spectrogram reconstructed from the Doppler tomogram shown in Fig. 8. Orbital phase is plotted along the  $x$ -axis.



**Figure 14.** The right hand panel displays the observed trailed spectra of the N III  $\lambda 4640$  emission line, part of the Bowen emission complex. Next to this is plotted the ideal trailed spectrogram reconstructed from the Doppler tomogram in Fig. 11. Orbital phase is plotted along the  $x$ -axis.

faintness of GR Mus compared to a system such as Sco X-1, which clearly features narrow Bowen lines moving in phase with the donor star (Steeeghs & Casares 2002).

We can reconstruct an ideal trailed spectrogram from the Doppler tomograms to test for systematic residuals in the maps. As a test, we first apply this technique to the He II  $\lambda 4686$  emission line. The two distinct components we see in the Doppler map (Fig. 8) are reproduced clearly in the predicted trailed spectrogram (left hand panel, (Fig. 13), and can be easily identified in the observed data. For the case of the N III  $\lambda 4640$  emission line (Fig. 14) we again see two components in the reconstructed spectrogram, though not quite so distinctly as for the case of He II. Comparing this idealised image to the real data it is indeed possible to detect faint traces of these two narrow, approximately anti-phased lines which produce the two distinct spots in Fig. 11. The second spot is therefore unlikely to be a systematic residual produced when creating the Doppler map, and its unusual phasing/positioning could perhaps therefore be attributed to a region which violates one of the axioms of Doppler Tomography, for example that all motion should be parallel to the orbital plane. Note that a violation of these axioms does not mean that the map is erroneous, but it does complicate the successful interpretation of the map.

**Table 2.** Derived system parameters for GR Mus / XB 1254–690

Parameter		Parameter	
$T_{dip}$ (HJD)	2453151.647 (3)	$\gamma$ (km s <sup>-1</sup> )	185 ± 2
$T_0$ (HJD)	2453151.509 (3)	$q$	0.32–0.43
$K_1$ (km s <sup>-1</sup> )	130 ± 16	$M_X(M_\odot)$	1.20–2.64
$K_{em}$ (km s <sup>-1</sup> )	245 ± 30	$M_2(M_\odot)$	0.45–0.85

Note: Inclination constrained to 65°–73°; Motch et al. (1987)

### 4.3 Refined mass limits

de Jong et al. (1996) estimated an opening (semi-) angle for the accretion disc in GR Mus of  $\sim 12^\circ$ , using a simple geometric model for the reprocessing of X-rays in LMXBs. It is important to note that this value is hostage to the implicit assumptions underlying these models, for example the geometric description of the disc and of the X-ray emitter. This will entail a systematic uncertainty that likely exceeds the internal model accuracy of less than a degree. Nevertheless, this value agrees well with Motch et al. (1987) who estimated an opening angle of  $\sim 9^\circ - 13^\circ$  in order to explain the observed amplitude of the orbital photometric modulation.

We can use this information to derive a mass estimate for the individual components by using the ‘K-correction’ algorithm developed by Muñoz-Darias et al. (2005). This measures the deviation between the reprocessed light centre of the fluorescence emission lines and the centre of mass of a Roche lobe filling star in a persistent LMXB, including the screening effects by a flared accretion disk. Using the fourth order polynomial fits given by Muñoz-Darias et al. (2005) for disc opening angles of 8°, 10°, 12° and 14° gives a total mass range for the neutron star of  $1.20 \leq M_X/M_\odot \leq 2.64$ . Since the neutron star mass and the mass ratio are related by the mass function<sup>2</sup>, we obtain  $0.32 \leq q \leq 0.43$ , and limits on the companion mass of  $0.45 \leq M_2/M_\odot \leq 0.85$ .

This correction is dependent upon the opening angle of the accretion disc, monotonically decreasing with opening angle such that the largest opening angle gives the lowest neutron star mass. Since the largest opening angle we have used exceeds the estimates of both Motch et al. (1987) and Muñoz-Darias et al. (2005), this gives a very conservative lower limit to the neutron star mass, and the true value is unlikely to be so low. Equally, the upper limit is derived using an opening angle lower than that suggested by Motch et al. (1987) and as such the neutron star is unlikely to be quite so massive.

Limiting ourselves to a disc opening angle of exactly  $12^\circ$  (de Jong et al. 1996), the mass constraints upon the neutron star tighten up to  $1.35 \leq M_X/M_\odot \leq 2.32$ , with  $0.33 \leq q \leq 0.41$  and  $0.49 \leq M_2/M_\odot \leq 0.79$ . These limits are not a vast improvement over the ones derived above using a range of opening angles from 8–14°. This indicates that the major source of uncertainty in this case is still incurred from the measurements of  $K_{em}$  and particularly of  $K_1$  (see Section 3.3), in addition to the system inclination angle. The choice of disc opening angle (within a reasonable range) has only a small effect in comparison. Nevertheless, we feel it is prudent not to restrict our final mass estimate by the assump-

tion of a disc opening angle of exactly  $12^\circ$ , preferring the more conservative range of 8–14°. The derived parameters for GR Mus are listed in Table 2.

### 4.4 Roche lobe size

We know that the companion must be filling its Roche lobe in order to permit persistent accretion. There is a well known and useful relationship between the fractional Roche lobe radius and the mass-ratio (Paczynski 1971) which can be combined with Kepler’s 3rd law to provide an estimate of the size of the Roche lobe as a function of period. In the case of this system (with a 3.9 hour period), the relationship may be expressed as:  $R_L = 0.58M_2^{1/3}$ .

We can thus derive the range of potential Roche lobe radii consistent with the mass range of the companion star that we have derived kinematically. This turns out to be  $0.44 \leq R_L/R_\odot \leq 0.55$ . Whilst at the lower end of our mass constraints the donor would be consistent with a main-sequence-like object as proposed by Motch et al. (1987), we cannot rule out the possibility that GR Mus harbours an over-massive evolved companion that has been stripped of its outer layers (e.g. King & Schenker 2002).

## 5 CONCLUSIONS

We have used the Bowen emission blend and Doppler tomography to trace for the first time the motion of the donor star in the persistent LMXB GR Mus. This has allowed us to derive kinematical mass limits for this system. In combination with estimates for the opening angle of the accretion disc, and new measurements of the He II  $\lambda 4686$  emission line (in an attempt to trace the motion of the compact object), we have derived tentative new mass constraints of  $1.20 \leq M_X/M_\odot \leq 2.64$  for the compact object and  $0.45 \leq M_2/M_\odot \leq 0.85$  for the companion star.

We cannot rule out the possibility that GR Mus harbours an over-massive evolved companion star, or more intriguingly an over-massive neutron star. Tighter constraints on the system parameters may therefore have important implications for our knowledge of the equation of state of nuclear matter (e.g. Cook et al. 1994) or indeed on the formation scenarios for typical LMXBs (e.g. Pfahl et al. 2003).

Future investigations should also concentrate on determining the origin of the unusual second spot in the N III Doppler map (Fig. 11). This feature is hard to explain, and perhaps could be caused by a violation of the axioms of Doppler Tomography. Related to this point, the broad emission band seen in the left hand side of the Doppler maps for He II and H $\beta$  (in this system and a number of others) is of uncertain origin. We believe this emission to be produced in an extended disc bulge, but physical modelling will be necessary to confirm this.

## ACKNOWLEDGMENTS

ADB was supported by a PPARC Research Studentship. DS acknowledges a Smithsonian Astrophysical Observatory Clay Fellowship as well as support through NASA GO grant

<sup>2</sup>  $f(M_X) = \frac{K_2^3 P_{orb}}{2\pi G} = \frac{M_X \sin^3 i}{(1+q)^2}$

NNG06GC05G. Based on observations made with ESO Telescopes at the Paranal Observatory under programme ID 073.D-0819. We gratefully acknowledge the use of the MOLLY and DOPPLER packages developed by T.R. Marsh.

## REFERENCES

- Augusteijn T., van der Hooft F., de Jong J.A., et al., 1998, *A&A*, 332, 561
- Buxton M., Vennes S., 2003, *MNRAS*, 342, 105
- Casares J., Marsh T.R., Charles P.A., et al., 1995, *MNRAS*, 274, 565
- Casares J., Steeghs D., Hynes R.I., Charles P.A., O'Brien K., 2003, *ApJ*, 590, 1041
- Casares J., Cornelisse R., Steeghs D., et al., 2006, *MNRAS*, submitted
- Cook G.B., Shapiro S.L., Teukolsky S.A., 1994, *ApJ*, 424, 823
- Cornelisse R., Steeghs D., Casares J., Charles P.A., Barnes A.D., et al., 2006, submitted to *MNRAS*
- Courvoisier T.J.L., Parmar A.N., Peacock A., Pakull M., 1986, *ApJ*, 309, 265
- Cowley A.P., Hutchings J.B., Crampton D., 1988, *ApJ*, 333, 906
- de Jong J., van Paradijs J., Augusteijn T., 1996, *A&A*, 314, 484
- Griffiths R.E., Gursky H., Schwartz D.A., Schwartz J., Bradt H., Doxsey R.E., Charles P.A., Thorstensen J.R., 1978, *Nature*, 276, 247
- Hellier C., 2000, *NewAR*, 44, 131
- Horne K., 1986, *PASP*, 98, 609
- Hynes R.I., Charles P.A., Haswell C.A., et al., 2001, *MNRAS*, 324, 180
- Hynes R.I., Charles P.A., van Zyl L., et al., 2004, *MNRAS*, 348, 100
- Iaria R., DiSalvo T., Burderi L., Rossa N.R., 2001, *ApJ*, 548, 883
- Jahoda K., Swank J.H., Stark M.J., et al. 1996, *Proc. SPIE*, 2808, 59
- King A.R., Schenker K., 2002, in 'The Physics of Cataclysmic Variables and Related Objects', ASP Conference Proceedings, Vol. 261. Edited by B. T. Gnsicke, K. Beuermann, and K. Reinsch., San Francisco: Astronomical Society of the Pacific, p.233
- Levine A.M., Corbet R., 2006, *ATel*, 940
- Mason K.O., Middleditch J., Nelson J.E., White N.E., 1980, *Nature*, 287, 516
- Marsh T. R., Horne K., 1988, *MNRAS*, 235, 269
- McClintock J.E., Canizares C.R., Tarter C.B., 1975, *ApJ*, 198, 641
- Motch C., Pedersen H., Beuermann K., Pakull M.W., Courvoisier T.J.L., 1987, *ApJ*, 313, 792
- Muñoz-Darias T., Casares J., Martínez-Pais I.G., 2005, *ApJ*, 635, 502
- Paczynski B., 1971, *ARA & A*, 9, 183
- Pearson K.J., Hynes R.I., Steeghs D., Jonker P.G., Haswell C.A., King A.R., O'Brien K., Nelemans G., Mndez M., 2006, *ApJ*, 648, 1169
- Pfahl E., Rappaport S., Podsiadlowski Ph., 2003, *ApJ*, 597, 1036
- Schachter J., Filippenko A.V., Kahn S.M., 1989, *ApJ*, 340, 1049
- Schneider D.P., Young P., 1980, *ApJ*, 238, 946
- Smale A.P., Wachter S., 1999, *ApJ*, 527, 341
- Smale A.P., Church M.J., Balucinska-Church M., 2002, *ApJ*, 581, 1286
- Smith D.A., Dhillon V.S., 1998, *MNRAS*, 301, 767
- Steeghs D., Casares J., 2002, *ApJ*, 568, 273
- Uno S., Mitsuda K., Aoki T., Makino F., 1997, *PASJ*, 49, 353
- van Paradijs J., McClintock J.E., 1995, *Optical and ultraviolet properties of X-ray binaries (X-Ray Binaries, Cambridge University Press, edited by Lewin, W.H.G., van Paradijs, J. & van der Heuvel, P.J. )*
- Wade R.A., Horne K., 1988, *ApJ*, 324, 411
- Wang C., Lai D., Han J.L., 2006, *ApJ*, 639, 1007
- Warner B., 1976, in *IAU Symposium 73, Structure and Evolution of Close Binary Systems*, ed. Eggleton P., Mitton S., Whelan J., (Dordrecht:Reidel), p. 85
- in't Zand J.J.M., Kuulkers E., Verbunt F., Heise J., Cornelisse R., 2003, *A&A*, 411, L487