



University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): MAURY BRAMSON, KRZYSZTOF BURDZY, AND WILFRID KENDALL

Article Title: SHY COUPLINGS, $CAT(0)$ SPACES, AND THE LION AND MAN

Year of publication: Forthcoming

Link to published article:

<http://projecteuclid.org/DPubS?service=UI&version=1.0&verb=Display&page=past&handle=euclid.aop>

Publisher statement: None

SHY COUPLINGS, CAT(0) SPACES, AND THE LION AND MAN

MAURY BRAMSON, KRZYSZTOF BURDZY, AND WILFRID KENDALL

ABSTRACT. Two random processes X and Y on a metric space are said to be ε -*shy coupled* if there is positive probability of them staying at least a positive distance ε apart from each other forever. Interest in the literature centres on nonexistence results subject to topological and geometric conditions; motivation arises from the desire to gain a better understanding of probabilistic coupling. Previous non-existence results for co-adapted shy coupling of reflected Brownian motion required convexity conditions; we remove these conditions by showing the non-existence of shy co-adapted couplings of reflecting Brownian motion in any bounded CAT(0) domain with boundary satisfying uniform exterior sphere and interior cone conditions, for example, simply-connected bounded planar domains with C^2 boundary.

The proof uses a Cameron–Martin–Girsanov argument, together with a continuity property of the Skorokhod transformation and properties of the intrinsic metric of the domain. To this end, a generalization of Gauss’ Lemma is established that shows differentiability of the intrinsic distance function for closures of CAT(0) domains with boundaries satisfying uniform exterior sphere and interior cone conditions. By this means, the shy coupling question is converted into a Lion and Man pursuit–evasion problem.

1. INTRODUCTION

1.1. Results and motivation. Benjamini, Burdzy, and Chen (2007) introduced the notion of *shy coupling*: a coupling of Brownian motions X and Y (more generally, of two random processes X and Y on a metric space) is said to be *shy* if there is an $\varepsilon > 0$ such that $\mathbb{P}[\text{dist}(X(t), Y(t)) \geq \varepsilon \text{ for all } t] > 0$. For example consider Brownian motion on the circle: if Y is produced from X by a non-trivial rotation then X and Y exhibit a shy coupling, since $\text{dist}(X, Y)$ is then constant. Interest in the existence or non-existence of such couplings arises from the study of couplings of reflected Brownian motions, which occur in various contexts. Benjamini et al. (2007) discussed existence and non-existence of shy couplings for Brownian motions on graphs and for reflected Brownian motions in domains (connected open subsets of Euclidean space) satisfying suitable boundary regularity conditions. They restricted attention to Markovian couplings and we will do essentially the same,

1991 *Mathematics Subject Classification.* 60J65.

Key words and phrases. CAT(0); co-adapted coupling; coupling; eikonal equation; first geodesic variation; Gauss’ lemma; greedy strategy; intrinsic metric; Lion and Man problem; Lipschitz domain; Markovian coupling; pursuit-evasion problem; reflected Brownian motion; Reshetnyak majorization; shy coupling; Skorokhod transformation; total curvature; uniform exterior sphere condition; uniform interior cone condition.

Research supported in part by NSF Grants CCF-0729537 and DMS-0906743, and by grant N N201 397137, MNiSW, Poland.

by restricting attention to co-adapted couplings. (This is only slightly more general, but is more convenient for expression in terms of stochastic calculus.) In particular the results in [Benjamini et al. \(2007\)](#) showed that no shy co-adapted couplings can exist for reflected Brownian motion in convex bounded planar domains with C^2 boundary satisfying a strict convexity condition (namely, that all line segments in the boundary are trivial). Their argument used a large deviations argument bearing some resemblance to methods from differential game theory. [Kendall \(2009\)](#) showed that neither differentiability nor strict convexity is required for the planar result, and also generalized the result to convex bounded domains in higher dimensions whose boundaries need no longer be smooth but still satisfy the regularity condition of triviality of all line segments on the boundary. These more recent results are based on direct proofs using ideas from stochastic control.

The work described below both generalizes the above results and also shows that failure of shyness is not confined to the case of convexity. We consider a bounded domain with boundary satisfying uniform exterior sphere and interior cone conditions and that satisfies a CAT(0) condition (see [Definition 4](#)) when furnished with the intrinsic metric, and we show that such domains cannot support shy co-adapted coupling of reflected Brownian motion. We do this by establishing a rather direct connection between (the non-existence of) Brownian shy co-adapted couplings and deterministic pursuit–evasion problems. As part of this process, we generalize Gauss’ lemma (on the differentiability of the distance function) to the case of closures of CAT(0) domains furnished with the intrinsic metric and satisfying uniform exterior sphere and interior cone conditions. It may not be evident to the reader exactly how the stochastic and undirected notion of Brownian motion can be connected to the deterministic and intentional notion of a pursuit–evasion problem, and it was not initially evident to us (though in retrospect this is latent in [Benjamini et al., 2007](#)), but nonetheless the connection is both immediate and useful.

The pursuit–evasion problem in question is a well-known problem concerning a Lion chasing a Man in a disk, both travelling at unit speed: Rado’s celebrated “Lion and Man” problem. Our shy coupling problem leads us to consider the generalization in which the Lion chases the Man in a bounded domain which is CAT(0) in its intrinsic metric. [Isaacs \(1965\)](#) is the classic reference for pursuit–evasion problems; [Nahin \(2007\)](#) provides an accessible exposition of the special case of the Lion and Man problem in the unit disk. [Littlewood \(1986, pp.114–117 in Bollobas’ extended edition\)](#) provides a brief description of the Lion and Man problem with an indication of its history, including a presentation of Besicovitch’s celebrated proof that in the disc the Man can evade the Lion indefinitely, even though the distance between Lion and Man may tend to zero. A generalization of discrete-time pursuit–evasion to bounded CAT(0) domains is dealt with in [Alexander, Bishop, and Ghrist \(2006\)](#); we summarize concepts from metric geometry and develop results required for the continuous-time variant in [Section 2](#), and it is here that we generalize the Gauss lemma to the case of closures of CAT(0) domains with sufficient boundary regularity ([Proposition 13](#)).

In particular, [Section 2](#) rigorously develops the geometric results required to reason with these concepts in the context of the intrinsic metric for the domain D (determined by lengths of paths restricted to lie within D). On a first reading one should feel free to note only the general ideas of

Section 2, and then to pass quickly on to the probabilistic arguments in the remaining sections of the paper.

In Section 3, we describe how continuous-time pursuit-evasion problems can be solved in CAT(0) domains. We obtain an upper bound for the time of ε -capture, expressed in terms of domain geometry.

Pursuit–evasion games involve control of the velocity of the pursuer so as to bring it arbitrarily close to the evader, regardless of what strategy may be adopted by the evader. In order to show non-existence of Brownian shy couplings, we investigate the possibility of bringing the Brownian pursuer (the *Brownian Lion*) arbitrarily close to the Brownian evader (the *Brownian Man*), regardless of how the Brownian motion of the Brownian Man is coupled to that of the Brownian Lion. The connection between coupling and deterministic Lion and Man problems is described in Section 4: a suitable pursuit strategy generates a vectorfield χ on the configuration manifold generated by the locations of Brownian Lion and Man. (More pedantically, it generates a section of the pullback of the tangent bundle of D to the configuration space of the pursuer and evader before capture.) If this pursuit strategy can be guaranteed to bring the Lion within $\varepsilon/2$ of Man by a bounded time t_c in the deterministic problem, then a Cameron–Martin–Girsanov argument together with a continuity property for the Skorokhod transformation shows that the Brownian Lion has a positive probability of getting within distance ε of the Brownian Man, whatever coupling strategy might be adopted by the Brownian Man.

The paper concludes with Section 5, which discusses possible extensions of these results, further questions, and conjectures.

We now state the main results of this paper, using terms defined in Section 2. Here and elsewhere in the paper, we consider only domains in Euclidean space of dimensions 2 or higher.

Theorem 1. *Suppose that D is a bounded domain with boundary satisfying uniform exterior sphere and interior cone conditions, and which is CAT(0) in its intrinsic metric. There can be no shy co-adapted coupling for reflected Brownian motion in D .*

Examples of CAT(0) domains include convex domains and domains that are the unions of a pair of convex domains. See, for instance, Bridson and Haefliger (1999) and Alexander et al. (2006), where more general examples are also provided; in particular, a large range of examples follows from iterated application of the result that if two CAT(0) domains have a geodesically convex intersection then their union is CAT(0). The exterior sphere and interior cone conditions in the theorem are required in order to apply the results of Saisho (1987) to generate reflected diffusions using the Skorokhod transformation.

The three dimensional domain in Figure 1 is CAT(0). There are two different ways to see this. First, it is easy to see that for every point on the boundary of the domain, at most one of the principal curvatures is negative. An alternative way to see that the domain is CAT(0) is to observe that a single dumbbell (the union of two spheres and the connecting tube) is a CAT(0) domain. The whole set is the union of five dumbbells. The non-empty intersections of the dumbbells are balls.




FIGURE 1. A CAT(0) example which is the union of five dumbbells.

Remarkably, all bounded simply-connected planar domains are CAT(0) in their intrinsic metrics. Thus, in the planar case, there is an immediate consequence of Theorem 1 which is a strikingly powerful result depending principally on topological conditions:

Theorem 2. *Suppose that D is a simply-connected bounded planar domain with boundary satisfying uniform exterior sphere and interior cone conditions. There can be no shy co-adapted coupling for reflected Brownian motion in D .*

1.2. Some basic tools for probabilistic coupling. All probabilistic couplings considered here are *co-adapted couplings*, which are defined for general Markov processes in Kendall (2009). In essence, a co-adapted coupling of two Markov processes is a construction of the two Markov processes on the same probability space, which are adapted to the same filtration such that each process possesses the prescribed transition functions *with respect to the common filtration*.

In this paper, it suffices to work with co-adapted couplings of d -dimensional Brownian motions: B and \tilde{B} are said to be *co-adaptively coupled Brownian motions* if they are defined on the same probability space and adapted to the same filtration $\{\mathcal{F}_t : t \geq 0\}$ and if, in addition, both satisfy an independent increments property taken *with respect to the common filtration*:

$$\begin{aligned} B_{t+s} - B_t &\text{ is independent of } \mathcal{F}_t \text{ for all } t, s \geq 0, \\ \tilde{B}_{t+s} - \tilde{B}_t &\text{ is independent of } \mathcal{F}_t \text{ for all } t, s \geq 0. \end{aligned}$$

Note that $B_{t+s} - B_t$ and $\tilde{B}_{t+s} - \tilde{B}_t$ need not be independent of each other. Kendall (2009, Lemma 6) shows that one may represent such a coupling using stochastic calculus, possibly at the cost of augmenting the filtration by adding a further independent Brownian motion C : there exist $(d \times d)$ -matrix-valued predictable random processes \mathbb{J} and \mathbb{K} such that

$$\tilde{B} = \int \mathbb{J}^\top d B + \int \mathbb{K}^\top d C;$$

moreover, one may choose $\mathbb{J}^\top \mathbb{J} + \mathbb{K}^\top \mathbb{K}$ to be equal to the $(d \times d)$ identity matrix at all times.

A pair of processes X and \tilde{X} is said to form a co-adapted coupling if they can be defined by strong solutions of stochastic differential equations driven by B, \tilde{B} respectively. In the paper, we will employ the stochastic differential equation obtained from the Skorokhod transformation for reflected Brownian motion in a domain D of suitable boundary regularity, such as under uniform exterior sphere and uniform interior cone conditions, as discussed in Section 2. For $r > 0$, set $\mathcal{N}_{x,r} = \{\nu \in \mathbb{R}^d : |\nu| = 1, \mathcal{B}(x + r\nu, r) \cap D = \emptyset\}$. The vectors ν can be viewed as “exterior normal unit vectors at $x \in \partial D$ ”; note that there may be more than one such vector at a particular point $x \in \partial D$. The set $\mathcal{N}_{x,r}$ is decreasing in r , and the uniform exterior sphere condition asserts that r can be chosen so that, for all $x \in \partial D$, $\mathcal{N}_{x,r} \neq \emptyset$, with $\mathcal{N}_{x,r} = \mathcal{N}_{x,s}$ for $0 < s \leq r$. Under uniform exterior sphere and uniform interior cone conditions, Saisho (1987) has shown that, given a driving Brownian motion B , there exists a unique solution pair (X, L^X) satisfying

$$\begin{aligned} d X &= d B - \nu_X d L^X, \\ L^X &\text{ is non-decreasing and increases only when } X \in \partial D, \\ \nu_X &\in \mathcal{N}_{X,r}. \end{aligned}$$

Thus L^X may be viewed as the local time of the reflected Brownian motion X on the boundary ∂D .

In this paper, all vectors are assumed to be column vectors unless specified otherwise.

2. CAT(0) GEOMETRY AND THE DETERMINISTIC PURSUIT-EVASION PROBLEM

Recall that the *intrinsic metric* for a domain D is generated by the infimum of lengths $\text{len}(\gamma)$ of smooth connecting paths γ lying wholly within the domain. (The definition is typically formulated in the context of general metric spaces and regularizable paths.)

Definition 3. *The intrinsic distance between two points x and y in a domain D is given by*

$$(1) \quad \text{dist}_{\text{intr}}(x, y) = \inf \{ \text{len}(\gamma) : \gamma \text{ is a smooth path connecting } x \text{ and } y \text{ in } D \}.$$

For a domain D , a standard compactness argument shows that paths attaining the infimum of (1) will always exist in the closure of the domain: these are called *intrinsic geodesics*.

As described in Bridson and Haefliger (1999, §II.1, Definition 1.1, case of $\kappa = 0$), see also Burago, Burago, and Ivanov, 2001, one can define simple curvature conditions for metric spaces such as $(D, \text{dist}_{\text{intr}})$, based on the behaviour of geodesic triangles.

Definition 4. We say that $(D, \text{dist}_{\text{intr}})$ is CAT(0) if the following triangle comparison holds: Suppose that $\Gamma_{a,b}$, $\Gamma_{a,c}$, and $\Gamma_{b,c}$ are unit-speed intrinsic geodesics for D , connecting points a to b , a to c , and b to c respectively. Then, for all such geodesic triangles,

$$\text{dist}_{\text{intr}}(\Gamma_{a,b}(s), \Gamma_{a,c}(t)) \leq r(s, t),$$

where $r(s, t)$ is the distance between points at distance s , respectively, t , from \tilde{a} along the side $\tilde{a}\tilde{b}$, respectively $\tilde{a}\tilde{c}$, of an ordinary Euclidean triangle $\tilde{a}\tilde{b}\tilde{c}$ that has the same side-lengths.




FIGURE 2. Illustration of the CAT(0) condition.

Consequently, chords of triangles in $(D, \text{dist}_{\text{intr}})$ are shorter than comparable chords of the comparable Euclidean triangles, as illustrated in Figure 2.

Remark. CAT is an acronym introduced by Gromov. It stands for **C**artan, **A**leksandrov, **T**oponogov.

Remark. As noted in *Bridson and Haefliger (1999, Proposition II.3.1)* (see also *Burago et al., 2001, §4.3*), in CAT(0) spaces (more generally, Alexandrov spaces of non-positive curvatures), the notion of angle is well-defined for (locally) minimal geodesics.

Consequently, geodesics in a CAT(0) space diverge at least as fast as corresponding geodesics in Euclidean space. Note that CAT(0) is a global condition, applying to all possible geodesic triangles. In particular it can be shown that CAT(0) spaces are always simply-connected and indeed contractible (*Bridson and Haefliger, 1999, Proposition II.1.4*, or *Alexander, Bishop, and Ghrist, 2010, Appendix A*).

Remarkably, bounded *planar* domains are CAT(0) if they are simply-connected; see *Bishop (2008)* for a careful proof. Readers may convince themselves of this at an intuitive level by drawing pictures (as exemplified in Figure 3); as is the case with other foundational results in metric spaces, the rigorous proof requires delicate reasoning.

We now introduce two complementary notions of boundary regularity following *Saisho (1987)*. An *exterior sphere condition* (also called *weak convexity*) requires that every boundary point is touched by at least one external sphere. Here and in the following, let $\mathcal{B}(y, s)$ denote the open Euclidean ball of radius s centered on y .




FIGURE 3. Illustration of the CAT(0) property for a bounded simply-connected planar domain. The effect of the boundary is to make the triangle “skinnier” than its Euclidean counterpart, thus establishing the CAT(0) comparison property.

Definition 5 (Uniform exterior sphere condition, from [Saisho, 1987](#), §1, Condition (A)). *A domain D is said to satisfy a uniform exterior sphere condition, based on radius r if, for every $x \in \partial D$, the set of “exterior normals” $\mathcal{N}_{x,r} = \{\nu \in \mathbb{R}^d : |\nu| = 1, \mathcal{B}(x + r\nu, r) \cap D = \emptyset\}$ is non-empty, with $\mathcal{N}_{x,r} = \mathcal{N}_{x,s}$ for $0 < s \leq r$.*

Thus a uniform exterior sphere condition allows one to move a fixed ball all the way around the outside of the domain boundary. In particular, D can have no “inward-pointing corners”. Here is a simple observation which will be useful later and corresponds to the intuition about being able to move a fixed ball about D ; such D may be represented as intersections of complements of balls, in a manner entirely analogous to the representation of a convex set as the intersection of half-planes (so justifying the alternate term “weak convexity”).

Lemma 6. *Suppose that the domain D satisfies a uniform exterior sphere condition based on radius r . Then*

$$\overline{D} = \bigcap \{ \mathcal{B}(z, r)^c : \mathcal{B}(z, r) \cap D = \emptyset \} .$$

Proof. Let the Minkowski sum $A \oplus B$ of two Euclidean sets A and B be $A \oplus B = \{x + y : x \in A, y \in B\}$. Certainly $F = \bigcap \{ \mathcal{B}(z, r)^c : \mathcal{B}(z, r) \cap D = \emptyset \}$ is closed, since $\mathcal{B}(z, r)$ is an open ball. Moreover $D \subseteq F$; hence $\overline{D} \subseteq F$. Furthermore $F \subseteq \overline{D} \oplus \mathcal{B}(\mathbf{o}, r)$, where \mathbf{o} is the origin of the ambient Euclidean space.

Following [Saisho \(1987, Remark 1.3\)](#), because of the uniform exterior sphere condition, we can define a projection $x \mapsto \bar{x}$ from $\overline{D} \oplus \mathcal{B}(\mathbf{o}, r)$ onto \overline{D} using the Euclidean metric. Consider $x \in \overline{D} \oplus \mathcal{B}(\mathbf{o}, r)$. Then the projection $\bar{x} \in \overline{D}$ is defined; moreover, if $\bar{x} \in \partial D$ and $x \notin \overline{D}$, then

$$\frac{\bar{x} - x}{|\bar{x} - x|} \in \mathcal{N}_{\bar{x}, r}$$

is a unit vector whose offset produces a tangent sphere of radius r at \bar{x} (using the argument of [Saisho, 1987](#)). But this implies that if $x \in (\bar{D} \oplus \mathcal{B}(\mathbf{o}, r)) \setminus D$ then

$$x \in \mathcal{B}\left(\bar{x} - r \frac{\bar{x} - x}{|\bar{x} - x|}, r\right)$$

and so $x \notin F$. Accordingly $\bar{D} = F$ as required. \square

On the other hand, a *uniform interior cone condition* requires that any boundary point supports a bounded cone truncated to the boundary of a ball, and moreover that the cone may be translated locally within the domain.

Definition 7 (Uniform interior cone condition, from [Saisho, 1987](#), §1, Condition (B')). *A domain D is said to satisfy a uniform interior cone condition, based on radius $\delta > 0$ and angle $\alpha \in (0, \pi/2]$, if, for every $x \in \partial D$, there is at least one unit vector \mathbf{m} such that the cone $C(\mathbf{m}) = \{z : \langle z, \mathbf{m} \rangle > |z| \cos \alpha\}$ satisfies*

$$(y + C(\mathbf{m})) \cap \mathcal{B}(x, \delta) \subseteq D \quad \text{for all } y \in D \cap \mathcal{B}(x, \delta).$$

We say that the cone $y + C(m)$ is based on y and angle $\alpha \in (0, \pi/2]$.

Thus a uniform interior cone condition implies that the “outward-pointing corners” must not be too sharp. Note that [Saisho](#) actually uses a slightly weaker condition with less intuitive content ([Saisho, 1987](#), Condition (B)); we do not consider this weaker notion further in what follows.

In fact, the property of a domain satisfying a uniform interior cone condition is equivalent to it being a Lipschitz domain.

Definition 8 (Lipschitz domain). *Recall that a function $f : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ is Lipschitz, with constant $\lambda < \infty$, if $|f(x) - f(y)| \leq \lambda|x - y|$ for all $x, y \in \mathbb{R}^{n-1}$. A domain D is said to be Lipschitz, with constant λ , if there exists $\delta > 0$ such that, for every $x \in \partial D$, there exists an orthonormal basis e_1, e_2, \dots, e_n and a Lipschitz function $f : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$, with constant λ , such that*

$$\{y \in \mathcal{B}(x, \delta) \cap D\} = \{y \in \mathcal{B}(x, \delta) : f(y_1, \dots, y_{n-1}) < y_n\},$$

where we write $y_1 = \langle y, e_1 \rangle, \dots, y_n = \langle y, e_n \rangle$.

The equivalence of Definitions 7 and 8 depends on the fact that the cone axis vector in Definition 7 is chosen to be the same for all $y \in \mathcal{B}(x, \delta)$, and so can be used as e_n in the orthonormal basis for $\mathcal{B}(x, \delta)$ required in Definition 8. The constants λ and α in Definitions 7 and 8 are related by $\lambda = \cot \alpha$, while the two δ 's of Definitions 7 and 8 may be taken to be equal. Note too that if the uniform interior cone / Lipschitz domain property holds for a given $\delta > 0$, then evidently it also holds for all smaller δ .

If a domain satisfies a uniform interior cone condition, then the intrinsic metric and Euclidean metric properties are closely related.

Lemma 9. *A domain D that is bounded in Euclidean metric and satisfies a uniform interior cone condition must have finite intrinsic diameter.*

Proof. Certainly $\text{dist}_{\text{intr}}(x, y)$ is a continuous function of (x, y) in the open set $D \times D$ and takes only finite values there. The domain D is path-connected, being an open connected subset of Euclidean space.

Suppose that D satisfies a uniform interior cone condition based on radius $\delta > 0$ and angle $\alpha \in (0, \pi/2]$. Choosing δ' with $0 < \delta' < \delta \tan \alpha$,

$$D \ominus \mathcal{B}(\mathbf{o}, \delta') \stackrel{\text{def}}{=} (D^c \oplus \mathcal{B}(\mathbf{o}, \delta'))^c = \{x \in D : \mathcal{B}(x, \delta') \subset D\}$$

is closed. Inheriting boundedness from D , it is therefore compact in the Euclidean topology, and hence also in the topology derived from the intrinsic metric, since the two metrics are locally equal away from the boundary of D . Hence $\{\text{dist}_{\text{intr}}(x, y) : x, y \in D \ominus \mathcal{B}(\mathbf{o}, \delta')\}$ attains a maximum value, which is therefore finite. However, for *any* $x', y' \in D$, we have

$$(2) \quad \text{dist}_{\text{intr}}(x', y') \leq 2\delta + \sup\{\text{dist}_{\text{intr}}(x, y) : x, y \in D \ominus \mathcal{B}(\mathbf{o}, \delta')\},$$

because the uniform interior cone condition assures us that, from all points on the boundary, there is a straight-line segment of length δ to $\{\text{dist}_{\text{intr}}(x, y) : x, y \in D \ominus \mathcal{B}(\mathbf{o}, \delta')\}$. Hence the intrinsic diameter must be bounded by the right-hand side of (2). \square

The full force of the uniform interior cone condition is not required for the above result; the proof does not require coordination of the directions of interior cones at different base-points. The full force of the uniform interior cone condition assures us that any path of finite length leading in D to a point x on the boundary of D can be deformed continuously in D into one which in its final phase leads down the segment on which the interior cone at x is based. Moreover, the lengths of the curves throughout this deformation can be constrained to be arbitrarily close to the length of the original path. This allows us to view D as a topological manifold with boundary, which is continuously embedded in the ambient Euclidean space. More than this, it shows that the completion \hat{D} of D under the intrinsic metric can be identified with the Euclidean closure \overline{D} and moreover that the intrinsic metric and the Euclidean metric actually endow \overline{D} with the same topology. Finally, [Bridson and Haefliger \(1999, Corollary II.3.11\)](#) show that the closure \overline{D} , viewed as the completion \hat{D} of D in intrinsic metric, inherits CAT(0) structure from D .

2.1. Regularity for geodesics. We wish to consider pursuit-evasion in a bounded CAT(0) domain. Lion and Man both move with unit speed, with the Lion seeking to draw closer to the Man by using the “greedy” pursuit strategy (which is not necessarily optimal). This Lion strategy can be phrased in terms of a unit-length \mathbb{R}^d -valued vectorfield χ defined on the configuration space $(\overline{D} \times \overline{D}) \setminus \{(x, x) : x \in \overline{D}\}$, such that $\chi(x, y)$ is the initial velocity of the unit-speed geodesic moving from x to y . (This is the vectorfield described pedantically in Section 1 as a section of the pullback of the tangent bundle of D to the configuration space of the pursuer and evader before capture.)

We first show that the combination of uniform exterior sphere and uniform interior cone / Lipschitz conditions implies that every boundary point of the intersection of the domain D with a suitable 2-plane will support an exterior sphere, albeit with smaller radius.




FIGURE 4. Illustration of interior cone C_z and exterior ball $\mathcal{B}(y, r)$ at $z \in D$.

Lemma 10. *Suppose that D is a domain satisfying a uniform exterior sphere condition based on radius $r > 0$, and a uniform interior cone condition based on radius $\delta > 0$ and angle $\alpha \in (0, \pi/2]$. Suppose that $z \in \partial D$ and e_n is the n^{th} vector in the orthonormal basis corresponding to z as in Definition 8. Let P be a 2-plane intersecting D and containing z and $z + e_n$. Then there exists $w \in P$, with $|w - z| = \text{dist}(w, D \cap P) = r \sin \alpha$ and $\mathcal{B}(w, r) \cap (D \cap P) = \emptyset$.*

Note that $z \in \partial(D \cap P)$ implies $z \in \partial D$. Hence the lemma shows that every point in the boundary of $D \cap P$ supports an exterior sphere of radius $r \sin \alpha$.

Proof. Suppose that $z \in \partial(D \cap P)$. Since $z \in \partial D$, there is an exterior sphere touching z , defined by a ball $\mathcal{B}(y, r) \subseteq D^c$ with $z \in \overline{\mathcal{B}(y, r)}$. By Definition 8, the cone

$$C_z = \{w : \langle w - z, e_n \rangle > |w - z| \cos \alpha\}$$

lies locally in D , in the sense that $C_z \cap \mathcal{B}(z, \delta) \subseteq D$ (see Figure 4). If $\frac{\pi}{2} + \beta$ is the angle between e_n and $y - z$, then two-dimensional geometry (Figure 5) shows that

$$\min \{|y - (\gamma e_n + z)| : \gamma \in \mathbb{R}\} = r \cos \beta.$$

But $\beta \geq \alpha$ if $C_z \cap \mathcal{B}(z, \delta) \subseteq D$ and $\mathcal{B}(y, r) \subseteq D^c$; moreover, the line $\{\gamma e_n + z : \gamma \in \mathbb{R}\}$ must lie in P . Hence the distance from y to P is at most $r \cos \beta \leq r \cos \alpha$. Consequently the radius of the disk $\mathcal{B}(y, r) \cap P$ is at least $r \sin \alpha$; since $z \in \partial(D \cap P)$ and $\mathcal{B}(y, r) \cap P$ is an exterior sphere to z in P , the lemma follows. \square

We can now establish some important technical consequences of the uniform exterior sphere and interior cone conditions, employed together with CAT(0); namely, that the Euclidean and intrinsic distances are locally comparable, and that the vectorfield χ is continuous with reference to the common topology of the Euclidean metric and the intrinsic metric, and hence is uniformly continuous over regions for which the two arguments are well-separated. This is spelled out in the following proposition.




FIGURE 5. Two-dimensional section of Figure 4 illustrating the underlying two-dimensional geometry.

Proposition 11. *Suppose that D is a CAT(0) domain, bounded in the Euclidean metric and satisfying a uniform exterior sphere condition based on radius $r > 0$, and a uniform interior cone condition based on radius $\delta > 0$ and angle $\alpha \in (0, \pi/2]$.*

- (1) *Suppose $a, b \in \overline{D}$ are close in the Euclidean metric, in the sense that $|a-b| < \min\{\delta, 2r \sin \alpha\}$. Then*

$$(3) \quad 2r \sin \alpha \sin \left(\frac{\text{dist}_{\text{intr}}(a, b)}{2r \sin \alpha} \right) \leq |a - b| \leq \text{dist}_{\text{intr}}(a, b).$$

- (2) *Intrinsic geodesics for D (necessarily minimal, by the CAT(0) condition) are continuously differentiable and their direction fields satisfy a Lipschitz property with constant $\frac{4}{\sqrt{3}} \frac{1}{2r \sin \alpha}$ that therefore holds uniformly for all minimal intrinsic geodesics in D and hence in \overline{D} (since CAT(0) geodesics depend continuously on their end-points);*
- (3) *For x, y in \overline{D} , let $\chi(x, y)$ be the unit vector at x pointing along the unique intrinsic geodesic $\gamma^{(x,y)}$ from x to y . Then $\chi(x, y)$ depends continuously on $(x, y) \in (\overline{D} \times \overline{D}) \setminus \{(x, x) : x \in \overline{D}\}$ and hence is uniformly continuous over regions for which the two arguments are well-separated.*

Proof of Proposition 11, part (1). As remarked in the discussion after Definitions 7 and 8, the domain D is Lipschitz with constant $\lambda = \cot \alpha$. For each ball of radius δ , we may therefore construct a coordinate system e_1, \dots, e_n and a Lipschitz function f to implement the Lipschitz property of D .

Consider $a, b \in \overline{D}$ with $|a - b| < \min\{\delta, 2r \sin \alpha\}$. If the line segment S between a and b does not intersect ∂D , then it must form the (unique, minimal) intrinsic geodesic between a and b , and (3) follows immediately. If S does not intersect $\text{int}(D^c)$, then we can cover the intersection $S \cap \partial D$

with finitely many balls $\mathcal{B}(z, \delta)$, $z \in \partial D$, and use the unit vector e_n corresponding to each ball (equivalently, the unit vectors defining the cones for each ball) to perturb S by arbitrarily small perturbations with length arbitrarily close to that of S . Since S must be the intrinsic geodesic between a and b , (3) follows immediately. So we can confine our attention to the case when $a \neq b$ and S intersects $\text{int}(D^c)$.

Let x be the point in $S \cap \partial D$ that is closest to a . By the triangle inequality, both $|x - a|$ and $|x - b|$ are smaller than δ and therefore $a, b \in \mathcal{B}(x, \delta)$. Accordingly, by Definition 8 (which is equivalent to the uniform interior cone condition as noted above), there is a Lipschitz function $f : \mathbb{R}^{d-1} \rightarrow \mathbb{R}$, with Lipschitz constant $\lambda = \cot \alpha$, and an orthonormal basis e_1, \dots, e_d , such that

$$\mathcal{B}(x, \delta) \cap D = \{y \in \mathcal{B}(x, \delta) : f(y_1, \dots, y_{d-1}) < y_d\}$$

where $y_1 = \langle y, e_1 \rangle, \dots, y_d = \langle y, e_d \rangle$. Let P be the 2-plane

$$P = a + \text{linear span} \{b - a, e_d\}.$$

Note that $x \in P$, and $D \cap P \neq \emptyset$.


Lemma 10 applies, so every boundary point of $D \cap P$ supports an exterior sphere of radius $r \sin \alpha$. The Lipschitz / uniform interior cone property of D , applied within $\mathcal{B}(x, \delta)$ and using the unit vector e_d , shows that rays from a and b in the e_d direction must lie in D until they leave $\mathcal{B}(x, \delta)$:

$$\begin{aligned} \mathcal{B}(x, \delta) \cap \{a + \gamma e_d : \gamma > 0\} &\subseteq D, \\ \mathcal{B}(x, \delta) \cap \{b + \gamma e_d : \gamma > 0\} &\subseteq D. \end{aligned}$$

Moreover, since $a \neq b$ and $|b - a| < 2r \sin \alpha$, there are exactly two circles in the plane P of radius $r \sin \alpha$ that touch both a and b . Furthermore, since $b - a$ cannot be parallel to e_d (as this would force one of $a - x, b - x$ to lie outside D), it follows that exactly one of these circles fails to intersect either of these rays. Let F be the disk corresponding to this circle and let A be the minor arc from a to b along the boundary circle. (This construction is illustrated in Figure 6.) Then any other disk of radius $r \sin \alpha$ that intersects the minor arc A from a to b along F must contain an initial segment of one or the other of the rays and therefore cannot be an exterior disk. It follows that A must be contained in \overline{D} , and moreover that a small perturbation of A , using the vector e_d , will provide a path in D from a to b of length arbitrarily close to the length of A .

Calculation of the length of the minor arc A now leads to the desired bounds on $\text{dist}_{\text{intr}}(a, b)$ as given in (3).

Proof of Proposition 11, part (2). Consider points a, b and c in \overline{D} , lying in this order along an intrinsic geodesic Γ in \overline{D} . For some positive $d < \min\{\delta, 2r \sin \alpha\}$, suppose that the intrinsic distances between a and b and between b and c are both equal to d . Note that Γ is a minimal geodesic, and therefore the intrinsic distance between a and c must be $2d$. Let $\rho_1 = |a - b|$, $\rho_2 = |b - c|$ and $\rho_3 = |a - c|$ be the Euclidean distances between these three pairs of points and let

FIGURE 6. Illustration of construction of disk F and minor arc A .

$\pi - \theta$ be the interior angle at b in the triangle abc . By the cosine formula,

$$\cos \theta = -\cos(\pi - \theta) = -\frac{\rho_1^2 + \rho_2^2 - \rho_3^2}{2\rho_1\rho_2} = \frac{\rho_3^2 - \rho_1^2 - \rho_2^2}{2\rho_1\rho_2}.$$

The upper bound on d means we can apply (3) to the intrinsic and Euclidean distances between a , b and c . Hence $\rho_1 \leq d$, $\rho_2 \leq d$ and

$$\rho_3 \geq 2r \sin \alpha \sin \left(\frac{2d}{2r \sin \alpha} \right) \geq 2d \left(1 - \frac{1}{6} \left(\frac{d}{r \sin \alpha} \right)^2 \right),$$

where the last step uses $\sin \alpha \geq \alpha - \alpha^3/6$ if $\alpha \geq 0$. Together with the cosine formula, these bounds for ρ_1, ρ_2 and ρ_3 yield

$$\begin{aligned} \cos \theta &\geq \frac{\left(2d \left(1 - \frac{1}{6} \left(\frac{d}{r \sin \alpha} \right)^2 \right) \right)^2 - \rho_1^2 - \rho_2^2}{2\rho_1\rho_2} \\ &\geq \frac{\left(2d \left(1 - \frac{1}{6} \left(\frac{d}{r \sin \alpha} \right)^2 \right) \right)^2 - 2d^2}{2d^2} = 2 \left(1 - \frac{1}{6} \left(\frac{d}{r \sin \alpha} \right)^2 \right)^2 - 1, \end{aligned}$$

hence

$$\cos \frac{\theta}{2} \geq 1 - \frac{1}{6} \left(\frac{d}{r \sin \alpha} \right)^2.$$

Considering $d < \min\{\delta, 2r \sin \alpha\}$, it follows by calculus that there exists a $c(d)$ tending to zero with d such that

$$(4) \quad \theta \leq \frac{4}{\sqrt{3}} \frac{d}{2r \sin \alpha} \left(1 + c(d) \frac{d}{2r \sin \alpha} \right).$$

Suppose now that the intrinsic geodesic Γ has total length K . For any positive integer $m > K/\min\{\delta, 2r \sin \alpha\}$, let $a_0 = a, a_1, \dots, a_{m-1}, a_m = b$ be $m + 1$ points equally spaced along the geodesic, so that $\text{dist}_{\text{intr}}(a_{j-1}, a_j) = d$ for $j = 1, \dots, m$. Define $g_m : [0, K] \rightarrow \mathbb{R}^m$ to be the piecewise-linear curve interpolating $g_m(jK/m) = a_j$ for $j = 0, \dots, m$. By (4), all the angles between successive line-segments of the trajectory of g_m are bounded above by

$$\frac{4}{\sqrt{3}} \frac{d}{2r \sin \alpha} \left(1 + c(d) \frac{d}{2r \sin \alpha} \right).$$

Define the *directional unit vector field* of the curve g_m by $\omega_m(s) = g'_m(s)/|g'_m(s)|$ for s where $g_m(s)$ is linear, and extend to all s using left-limits for $s > 0$ and the right-limit for $s = 0$. Then, by the triangle inequality,

$$|\omega_m(t) - \omega_m(s)| \leq \frac{4}{\sqrt{3}} \frac{d}{2r \sin \alpha} \left(1 + c(d) \frac{d}{2r \sin \alpha} \right) \left(\frac{|t-s|}{d} + 1 \right).$$

From (3),

$$\frac{2r \sin \alpha}{d} \sin \left(\frac{d}{2r \sin \alpha} \right) \leq |g'_m(s)| \leq 1;$$

hence we obtain the inequality

$$(5) \quad |g'_m(t) - g'_m(s)| \leq |\omega_m(t) - \omega_m(s)| + |\omega_m(t) - g'_m(t)| + |g'_m(s) - \omega_m(s)| \\ \leq \frac{4}{\sqrt{3}} \frac{d}{2r \sin \alpha} \left(1 + c(d) \frac{d}{2r \sin \alpha} \right) \left(\frac{|t-s|}{d} + 1 \right) + 2 \left(1 - \frac{2r \sin \alpha}{d} \sin \left(\frac{d}{2r \sin \alpha} \right) \right),$$

from which there follows a uniform bound on the absolute variation of the g'_m functions. Thus we can apply Helly's selection theorem to deduce that g'_m will converge along a subsequence, both pointwise and locally in L^1 , to a continuous limit h . It is immediate that g_m converges uniformly to Γ , and Γ must be almost everywhere differentiable with limit $h = \Gamma'$. Moreover, from (5) (and bearing in mind that $c(d) \rightarrow 0$ with d), we may deduce that the derivative Γ' is Lipschitz with constant

$$\frac{4}{\sqrt{3}} \frac{1}{2r \sin \alpha},$$

and indeed that Γ is continuously differentiable.

Proof of Proposition 11, part (3). As noted above, the CAT(0) property of \overline{D} implies that all geodesics in \overline{D} are unique and minimal. Consider $(x, y), (x_n, y_n) \in (\overline{D} \times \overline{D}) \setminus \{(x, x) : x \in \overline{D}\}$ with $x_n \rightarrow x$ and $y_n \rightarrow y$ in the Euclidean metric; taking subsequences we may suppose that $\chi(x_n, y_n)$ converges to a limit. Part (2) of the lemma establishes the uniform Lipschitz property of the direction fields of all minimal geodesics in \overline{D} so, by the Arzela-Ascoli theorem, we can find a subsequence (x_{n_k}, y_{n_k}) such that the geodesics from x_{n_k} to y_{n_k} must converge to a curve from x to y whose direction field is the limit of the direction fields of these minimal geodesics; hence its direction at x must be $\lim_k \chi(x_{n_k}, y_{n_k})$. By minimality of the geodesics from x_{n_k} to y_{n_k} and taking limits,

the length of the limiting curve can be no greater than that of the unique minimal geodesic from x to y ; therefore the limiting curve must also be a minimal geodesic from x to y . By the CAT(0) property, the two minimal geodesics from x to y must therefore be equal, and therefore it follows that $\lim_k \chi(x_{n_k}, y_{n_k}) = \chi(x, y)$. It follows that any subsequence of $(x_n, y_n) \rightarrow (x, y)$ (convergence in Euclidean metric) must possess a further subsequence for which $\lim_k \chi(x_{n_k}, y_{n_k}) = \chi(x, y)$, and therefore $\lim_n \chi(x_n, y_n) = \chi(x, y)$ must hold. This establishes continuity of χ with reference to the Euclidean metric. \square

Remark. Part (1) of Proposition 11 may be used to show that $\chi(a, b)$ is Hölder($\frac{1}{2}$) in its second argument b when a and b are well-separated. We omit this argument, as the result is not used in this paper.

Remark. Setting $\rho = |a - b|$ and $d = \text{dist}_{\text{intr}}(a, b)$, Inequality (3) can be rewritten as

$$\frac{2r \sin \alpha}{d} \sin \left(\frac{d}{2r \sin \alpha} \right) \leq \frac{\rho}{d} \leq 1.$$

The following is a trivial but useful consequence of the above estimates: for some $c_1 > 0$, depending on D , and all $a, b \in \overline{D}$, with $|x - y| \leq c_1$,

$$(6) \quad |a - b| \leq \text{dist}_{\text{intr}}(a, b) \leq 2|a - b|.$$

Since $\sin \phi \geq \phi - \frac{1}{6}\phi^3$ if $\phi \geq 0$,

$$1 - \frac{1}{6} \frac{d^2}{4r^2 \sin^2 \alpha} \leq \frac{\rho}{d} \leq 1.$$

The last inequality and (6) imply that for some $c_2, c_3 < \infty$, depending on δ, r and α , and for $\rho < \min\{\delta, 2r \sin \alpha\}$,

$$(7) \quad 1 \leq \frac{d}{\rho} \leq 1 + c_2 d^2 \leq 1 + c_3 \rho^2.$$

Proposition 11 makes it possible to quantify the extent to which short intrinsic geodesics may be approximated by Euclidean segments.

Corollary 12. Suppose the assumptions on Proposition 11 hold, and that Γ is a unit-speed intrinsic geodesic with intrinsic length $d < \min\{\delta, 2r \sin \alpha\}$. Then

$$(8) \quad |\Gamma(d) - \Gamma(0) - \Gamma'(0)d| \leq \frac{4}{3} \frac{d^2}{2r \sin \alpha}.$$

Proof. Set $\rho = |\Gamma(d) - \Gamma(0)|$ equal to the Euclidean distance between the two end-points of Γ ; then ρ is bounded above by the intrinsic length d . Let θ be the angle between $\Gamma'(0)$ and $\Gamma(d) - \Gamma(0)$.

Proposition 11, (2) tells us that Γ' is Lipschitz with constant $\frac{4}{\sqrt{3}} \frac{1}{2r \sin \alpha}$. Hence the angular distance between the unit vectors $\Gamma'(0)$ and $\Gamma'(t)$ is bounded above by $\frac{4}{\sqrt{3}} \frac{t}{2r \sin \alpha}$ when $t \geq 0$. Using

$\cos u \geq 1 - \frac{1}{2}u^2$, it follows that

$$(9) \quad \langle \Gamma'(t), \Gamma'(0) \rangle \geq 1 - \frac{8}{3} \frac{t^2}{4r^2 \sin^2 \alpha}$$

and this integrates to

$$\langle \Gamma(d) - \Gamma(0), \Gamma'(0) \rangle \geq \left(1 - \frac{8}{9} \frac{d^2}{4r^2 \sin^2 \alpha}\right) d.$$

Consequently

$$\begin{aligned} |\Gamma(d) - \Gamma(0) - \Gamma'(0)d|^2 &= |\Gamma(d) - \Gamma(0)|^2 + |\Gamma'(0)d|^2 - 2 \langle \Gamma(d) - \Gamma(0), \Gamma'(0)d \rangle \\ &\leq \rho^2 + d^2 - 2 \left(1 - \frac{8}{9} \frac{d^2}{4r^2 \sin^2 \alpha}\right) d^2 \leq \frac{16}{9} \frac{d^4}{4r^2 \sin^2 \alpha}. \end{aligned}$$

The result follows by taking square roots. \square

We recall Gauss' lemma from Riemannian geometry. Cheeger and Ebin (2008, Chapter 1, Section 2) observe that, for smooth Riemannian manifolds, it is equivalent to the assertion that the Riemannian distance $\text{dist}_{\text{intr}}(x, y)$ is continuously differentiable in x when $x \neq y$ and y does not lie in the cut-locus of x , with the gradient being given by the tangent of the geodesic running from y to x . Proposition 11 and Corollary 12 can be used to prove the following Gauss lemma for CAT(0) domains with sufficient boundary regularity. Here, $\text{grad}_x \text{dist}_{\text{intr}}(x, y)$ refers to the Euclidean gradient with respect to x , with $\text{grad} \text{dist}_{\text{intr}}(x, y)$ being the gradient with respect to both variables.

Proposition 13. *Suppose (as in Proposition 11) that D is a CAT(0) domain, bounded in the Euclidean metric, satisfying a uniform exterior sphere condition based on radius $r > 0$, and a uniform interior cone condition based on radius $\delta > 0$ and angle $\alpha \in (0, \pi/2]$. For every $c_1 > 0$, there exist $c_2, c_3 < \infty$ such that, if $x, y \in \overline{D}$ with $\text{dist}_{\text{intr}}(x, y) \geq c_1$ and $\sqrt{|u - x|} \vee |u - x| \leq c_2$, then*

$$(10) \quad |\text{dist}_{\text{intr}}(u, y) - (\text{dist}_{\text{intr}}(x, y) + |u - x| \cos \theta)| \leq c_3 |u - x|^{3/2},$$

where θ is the angle between the geodesic from y to x and the Euclidean segment from x to u that is exterior to the direction from y to x (see Figure 7). Consequently, if $x, y \in \overline{D}$ with $x \neq y$, then

$$(11) \quad \text{grad}_x \text{dist}_{\text{intr}}(x, y) = -\chi(x, y).$$

Moreover,

$$(12) \quad \text{grad} \text{dist}_{\text{intr}}(x, y) = (-\chi(x, y), -\chi(y, x)).$$

Note that Bieske (2010) establishes a similar result for Carnot-Carathéodory spaces. In both cases, the relevant distance function satisfies an eikonal equation.




FIGURE 7. Illustration of the configuration of the triangle referred to in the statement of Proposition 13. The sides running from y to u and from y to x (and of intrinsic lengths d_{yu} and d_{yx} respectively) are intrinsic geodesics. The side running from x to u is a Euclidean segment.

Proof. In order to demonstrate (10), we establish upper and lower bounds on the difference (13)

$$\text{dist}_{\text{intr}}(y, u) - \text{dist}_{\text{intr}}(y, x)$$

when u is close to x . We abbreviate, setting $d_{yx} = \text{dist}_{\text{intr}}(y, x)$.

Let θ' be the exterior angle between the geodesic from x to y and the geodesic from x to u . By the CAT(0) property, the Euclidean triangle with the same side lengths as a triangle in the intrinsic metric has larger interior angles and therefore smaller exterior angles. (The elementary argument is given in Bridson and Haefliger, 1999, Chapter II.1, Proposition 1.7(4).) Thus if θ'' is the exterior angle of the comparison triangle for x, y and u corresponding to the exterior angle θ' , then $\theta'' \leq \theta'$, and so

$$\begin{aligned} d_{yu} &= \sqrt{d_{yx}^2 + d_{xu}^2 + 2d_{yx}d_{xu} \cos \theta''} \geq \sqrt{d_{yx}^2 + d_{xu}^2 + 2d_{yx}d_{xu} \cos \theta'} \\ &\geq \sqrt{d_{yx}^2 + d_{xu}^2 \cos^2 \theta' + 2d_{yx}d_{xu} \cos \theta'} = d_{yx} + d_{xu} \cos \theta' \\ &\geq d_{yx} + |u - x| \cos \theta'. \end{aligned}$$

Corollary 12 implies that $|\theta - \theta'| \leq c_4 d_{xu}$ for small d_{xu} . Hence, $|\theta - \theta'| \leq c_5 |u - x|$ and $|\cos \theta - \cos \theta'| \leq c_5 |u - x|$. We obtain for $|u - x| \leq c_2$, for some $c_2 > 0$,

$$(14) \quad \begin{aligned} d_{yu} &\geq d_{yx} + |u - x| \cos \theta' \geq d_{yx} + |u - x| \cos \theta - |u - x| |\cos \theta - \cos \theta'| \\ &\geq d_{yx} + |u - x| \cos \theta - c_5 |u - x|^2. \end{aligned}$$

This provides a lower bound on (13) and a bound for one direction of (10).

We now establish an upper bound on (13). Fix a point w on the intrinsic geodesic from y to x . Then $d_{yx} = d_{yw} + d_{wx}$ and $d_{yu} \leq d_{yw} + d_{wu}$. We shall require w to be close to x , but not as close as u , with $|w - x| = \sqrt{|u - x|}$ being assumed.

Because w is close to x and thus also close to u , we may replace the intrinsic geodesics from w to u and from w to x by Euclidean segments, without greatly altering lengths and segments. Let θ^* be the exterior angle at x for the Euclidean triangle with sides $d_{x,w}$, $d_{x,u}$ and $d_{u,w}$. From (7),

$$(15) \quad |u - w| \leq d_{wu} = \frac{d_{wu}}{|u - w|} |u - w| \leq (1 + c_6 |u - w|^2) |u - w|,$$

$$(16) \quad |x - w| \leq d_{wx} = \frac{d_{wx}}{|x - w|} |x - w| \leq (1 + c_6 |x - w|^2) |x - w|,$$

when $|u - w|, |x - w| < 2r \sin \alpha$.

As before, by Corollary 12, $|\theta - \theta^*| \leq c_7 d_{xw}$ for small d_{xw} . Hence, $|\theta - \theta^*| \leq c_8 |w - x|$ and

$$(17) \quad |\cos \theta - \cos \theta^*| \leq c_8 |w - x|.$$

These computations allow us to establish an upper bound for $\text{dist}_{\text{intr}}(y, u) - \text{dist}_{\text{intr}}(y, x)$. First note that

$$\begin{aligned} d_{yu} &\leq d_{yw} + d_{wu} \leq d_{yw} + (1 + c_6 |u - w|^2) |u - w| \\ &\leq d_{yw} + |u - w| + c_6 |u - w|^3 \\ &\leq d_{yw} + |u - w| + c_6 (|u - x| + |w - x|)^3. \end{aligned}$$

Now apply the cosine formula to control $|u - w|$, using (17):

$$\begin{aligned} |u - w| &= \sqrt{|w - x|^2 + |u - x|^2 + 2|w - x||u - x| \cos \theta^*} \\ &= \sqrt{(|w - x| + |u - x| \cos \theta)^2 + |u - x|^2 \sin^2 \theta + 2|w - x||u - x|(\cos \theta^* - \cos \theta)} \\ &\leq |w - x| + |u - x| \cos \theta \\ &\quad + \frac{1}{2} \frac{|u - x|^2 \sin^2 \theta}{|w - x| + |u - x| \cos \theta} + \frac{|w - x||u - x|(\cos \theta^* - \cos \theta)}{|w - x| + |u - x| \cos \theta} \\ &\leq |w - x| + |u - x| \cos \theta + \frac{1}{2} \frac{|u - x|^2 \sin^2 \theta}{|w - x| - |u - x|} + \frac{|w - x||u - x|}{|w - x| - |u - x|} c_8 |w - x|. \end{aligned}$$

If we take $|w - x| = \sqrt{|u - x|}$, with $|u - x| < c_9$ for a suitably small $c_9 > 0$, then

$$|u - w| \leq |w - x| + |u - x| \cos \theta + c_{10} |u - x|^{3/2}.$$

Combining these bounds implies

$$\begin{aligned}
d_{yu} &\leq d_{yw} + |u - w| + c_6(|u - x| + |w - x|)^3 \\
&\leq d_{yw} + |w - x| + |u - x| \cos \theta + c_{10}|u - x|^{3/2} + c_6(|u - x| + |w - x|)^3 \\
&\leq d_{yw} + d_{w,x} + |u - x| \cos \theta + c_{11}|u - x|^{3/2} \\
&= d_{yx} + |u - x| \cos \theta + c_{11}|u - x|^{3/2},
\end{aligned}$$

which provides an upper bound on (13). It follows from the above inequality and (14) that

$$|d_{yu} - (d_{yx} + |u - x| \cos \theta)| \leq c_3|u - x|^{3/2},$$

which yields the bound in (10). The formula in (11), for the gradient of the intrinsic distance $\text{dist}_{\text{intr}}(x, y)$ with respect to x , follows immediately.

We still need to demonstrate the formula in (12). Let c_1, c_2 and c_3 be as in the statement of (10). Fix $x, y \in \overline{D}$ and suppose that $\text{dist}_{\text{intr}}(x, y) \geq 2c_1$. Suppose that $u, v \in \overline{D}$, $\sqrt{|u - x|} \vee |u - x| \leq c_2 \wedge c_1/4$ and $\sqrt{|v - y|} \vee |v - y| \leq c_2 \wedge c_1/4$. Let θ_x be the exterior angle between the geodesic from x to y and the Euclidean segment from x to u . Similarly, let θ_y be the exterior angle between the geodesic from y to x and the Euclidean segment from y to v . Also, let θ' be the exterior angle between the geodesic from y to u and the Euclidean segment from y to v . Then by the above reasoning

$$(18) \quad |\text{dist}_{\text{intr}}(u, y) - (\text{dist}_{\text{intr}}(x, y) + |u - x| \cos \theta_x)| \leq c_3|u - x|^{3/2}$$

and

$$(19) \quad |\text{dist}_{\text{intr}}(v, u) - (\text{dist}_{\text{intr}}(y, u) + |v - y| \cos \theta')| \leq c_3|v - y|^{3/2}.$$

Recall that the Euclidean triangle with the same side lengths as a triangle in intrinsic metric has larger interior angles. Hence, $|\theta_y - \theta'|$ is less than the angle at the vertex corresponding to y in the Euclidean triangle with sides d_{xy}, d_{xu} and d_{yu} . It follows that $|\theta_y - \theta'| \leq c_{12}d_{xu}/d_{xy} \leq c_{13}|u - x|$ and therefore $|\cos \theta_y - \cos \theta'| \leq c_{13}|u - x|$. This and (19) yield

$$(20) \quad |\text{dist}_{\text{intr}}(v, u) - (\text{dist}_{\text{intr}}(y, u) + |v - y| \cos \theta_y)| \leq c_3|v - y|^{3/2} + c_{13}|u - x||v - y|.$$

The triangle inequality applied to the left hand sides of (18) and (20) implies that

$$\begin{aligned}
&|\text{dist}_{\text{intr}}(v, u) - (\text{dist}_{\text{intr}}(x, y) + |v - y| \cos \theta_y + |u - x| \cos \theta_x)| \\
&\leq c_3|u - x|^{3/2} + c_3|v - y|^{3/2} + c_{13}|u - x||v - y|.
\end{aligned}$$

Consequently, $\text{grad dist}_{\text{intr}}(x, y)$ exists when $\text{dist}_{\text{intr}}(x, y)$ is viewed as a function of $(x, y) \in (\overline{D} \times \overline{D}) \setminus \{(u, u) : u \in \overline{D}\}$ and is given by

$$\text{grad dist}_{\text{intr}}(x, y) = (-\chi(x, y), -\chi(y, x)).$$

□

In Proposition 14, we consider solutions of the differential equation $dx = \chi(x, y) dt$ for pursuit and evasion. Proposition 11 established partial regularity, for $\chi(x, y)$, which does not automatically guarantee well-posedness of solutions (as defined below). However, the CAT(0) property, together with boundary regularity, will imply well-posedness, even for some discontinuous driving paths y .

Suppose that $y(t)$, $t \in [0, T_1]$, is cadlag, of bounded variation on finite intervals, and takes values in \overline{D} . We will say that $x(t)$, $t \in [0, T_1]$, is a weak solution to $dx = \chi(x, y) dt$ if $x(t) = x(0) + \int_0^t \chi(x(s), y(s)) ds$ for all $t \in [0, T_1]$.

Proposition 14. *Let D be a CAT(0) domain satisfying uniform exterior sphere and interior cone conditions. For distinct $x, y \in \overline{D}$, let $\chi(x, y)$ be the unit tangent vector at x of the geodesic from x to y . We consider the differential equation*

$$(21) \quad dx = \chi(x, y) dt$$

defined in the weak sense for absolutely continuous paths $\{x(t) : t \geq 0\}$ in \overline{D} , driven by paths $\{y(t) : t \geq 0\}$, up until the first time that x and y are equal. The problem is well-posed, in the sense that solutions x exist, are uniquely determined by initial values $x(0)$, and depend continuously on the initial value $x(0)$ and the driving process $\{y(t) : t \geq 0\}$ (using the uniform distance metric in both cases).

Proof. The argument is based on the simpler case when the path y is constant in time, which we for the moment assume. In this case, existence follows directly from the existence of intrinsic geodesics in CAT(0) domains. To show uniqueness, note that, for two solutions x and \tilde{x} of (21), since x and \tilde{x} are absolutely continuous and satisfy the differential equation weakly, for almost all s , the time-derivatives of $x(s)$ and $\tilde{x}(s)$ must exist and be given by $\chi(x(s), y)$ and $\chi(\tilde{x}(s), y)$. Exploiting the differentiability of the intrinsic distance given by Proposition 13, for $x(s) \neq y$ and $\tilde{x}(s) \neq y$, one has

$$(22) \quad \left[\frac{d}{dt} \text{dist}_{\text{intr}}(x(s+t), \tilde{x}(s+t)) \right]_{t=0} = \left[\frac{d}{dt} \text{dist}_{\text{intr}}(\Gamma^{(s)}(t), \tilde{\Gamma}^{(s)}(t)) \right]_{t=0},$$

where $\Gamma^{(s)}, \tilde{\Gamma}^{(s)}$ are unit-speed geodesics running from $x(s), \tilde{x}(s)$ to y . We will show that

$$(23) \quad \left[\frac{d}{dt} \text{dist}_{\text{intr}}(\Gamma^{(s)}(t), \tilde{\Gamma}^{(s)}(t)) \right]_{t=0} \leq 0.$$

Consider a Euclidean triangle abc with side lengths satisfying $|ab| = \text{dist}_{\text{intr}}(\Gamma^{(s)}(0), y)$, $|cb| = \text{dist}_{\text{intr}}(\tilde{\Gamma}^{(s)}(0), y)$ and $|bc| = \text{dist}_{\text{intr}}(\Gamma^{(s)}(0), \tilde{\Gamma}^{(s)}(0))$. Let $z(t) \in ab$ be a point such that $|z(t) - a| = t$, and let $\tilde{z}(t) \in cb$ be a point such that $|\tilde{z}(t) - c| = t$. Then Definition 4 implies that

$$\text{dist}_{\text{intr}}(\Gamma^{(s)}(t), \tilde{\Gamma}^{(s)}(t)) \leq |z(t) - \tilde{z}(t)| \leq |z(0) - \tilde{z}(0)| = \text{dist}_{\text{intr}}(\Gamma^{(s)}(0), \tilde{\Gamma}^{(s)}(0)).$$

This implies (23). It follows that the derivative on the left hand side of (22) is non-positive; therefore $x = \tilde{x}$ if $x(0) = \tilde{x}(0)$, and so uniqueness holds.




FIGURE 8. Illustration of the CAT(0) comparison argument applied to the triangles defined by vertices (a) $x(s)$, $\tilde{x}(s)$, y and (b) $x(s)$, $\tilde{y}(s)$, $y(s)$. A chord crossing the Euclidean quadrilateral as illustrated will be longer than a path making comparable chords with the two CAT(0) triangles, which itself will be longer than a comparable geodesic chord of the CAT(0) quadrilateral.

By considering the behaviour over disjoint time intervals, existence and uniqueness follow for the case when y is piecewise-constant, in which case the solution curve x is piecewise-geodesic.

We will establish continuous dependence on the initial position $x(0)$ and the driving process y , when y is piecewise constant. Suppose that y, \tilde{y} are two piecewise-constant paths in \bar{D} , and x, \tilde{x} solve

$$dx = \chi(x, y) dt, \quad d\tilde{x} = \chi(\tilde{x}, \tilde{y}) dt$$

for prescribed initial positions $x(0) \neq y(0)$ and $\tilde{x}(0) \neq \tilde{y}(0)$. The solutions x, \tilde{x} satisfy the differential equations weakly, and therefore, for almost all s , the time-derivatives of $x(s)$ and $\tilde{x}(s)$ must exist and are given by $\chi(x(s), y(s))$ and $\chi(\tilde{x}(s), \tilde{y}(s))$. Arguing as before, for $x(s) \neq y(s)$ and $\tilde{x}(s) \neq \tilde{y}(s)$, we may construct a CAT(0) comparison for the two triangles defined by (a) vertices $x(s), \tilde{x}(s), y(s)$ and (b) vertices $x(s), \tilde{y}(s), y(s)$. Using this comparison, and continuing until either $x(t) = y(t)$ or $\tilde{x}(t) = \tilde{y}(t)$, the function $\text{dist}_{\text{intr}}(x, \tilde{x})$ is dominated by its Euclidean counterpart for a two-dimensional quadrilateral which is based on a pair of opposing sides of lengths $\text{dist}_{\text{intr}}(x(s), \tilde{x}(s))$ and $\text{dist}_{\text{intr}}(y(s), \tilde{y}(s))$. Consequently, two-dimensional Euclidean geometry (see Figure 8) shows that

$$\text{dist}_{\text{intr}}(x(t), \tilde{x}(t)) \leq \max \{ \text{dist}_{\text{intr}}(x(s), \tilde{x}(s)), \text{dist}_{\text{intr}}(y(s), \tilde{y}(s)) \},$$

for $t \in [s, T]$ and any T such that y and \tilde{y} are constant on $[s, T]$, and $x(t) \neq y(t)$ and $\tilde{x}(t) \neq \tilde{y}(t)$ for $t \in [s, T)$. It follows that the solution x depends continuously on $x(0)$ and y until $x(t) = y(t)$, as specified in the statement of this lemma.

We now consider general y in (21). There exist piecewise-constant functions y_n converging to y uniformly on compact intervals; let x_n be the corresponding solutions to (21), with $x_n(0) = x(0) \in \bar{D}$. If $|y_n(t) - y_m(t)| \leq c_1$ for $t \in [0, T]$, then $|x_n(t) - x_m(t)| \leq c_1$ by the argument given above. Since the sequence y_n is Cauchy in the uniform norm on $[0, T]$, so is the sequence x_n , which therefore converges to a function x .

Recall that we are assuming $x(0) \neq y(0)$. Choose fixed $\varepsilon_1, \varepsilon_2 > 0$ and let $T = \inf\{t > 0 : |x(t-) - y(t-)| \leq 2\varepsilon_1\}$. By part (3) of Proposition 11, there exists $\delta_1 > 0$ such that, if $|u_1 - u_2| \geq \varepsilon_1$, $|v_1 - v_2| \geq \varepsilon_1$, $|u_1 - v_1| \leq \delta_1$ and $|u_2 - v_2| \leq \delta_1$, then $|\chi(u_1, u_2) - \chi(v_1, v_2)| \leq \varepsilon_2/T$. Suppose that

n is large enough so that $|y_n(t) - y(t)| \leq \delta_1 \wedge \varepsilon_1$ for $t \in [0, T]$. Then $|x_n(t) - x(t)| \leq \delta_1 \wedge \varepsilon_1$ for $t \in [0, T]$ and $|\chi(x(t), y(t)) - \chi(x_n(t), y_n(t))| \leq \varepsilon_2/T$. We obtain, for $t \leq T$,

$$\begin{aligned} & \left| x(t) - x(0) - \int_0^t \chi(x(s), y(s)) \, ds \right| \\ & \leq |x(t) - x_n(t)| + \left| x_n(t) - x(0) - \int_0^t \chi(x_n(s), y_n(s)) \, ds \right| \\ & \quad + \left| \int_0^t \chi(x_n(s), y_n(s)) \, ds - \int_0^t \chi(x(s), y(s)) \, ds \right| \\ & \leq \delta_1 + 0 + \int_0^t |\chi(x_n(s), y_n(s)) - \chi(x(s), y(s))| \, ds \\ & \leq \delta_1 + \varepsilon_2. \end{aligned}$$

Since $\varepsilon_1, \varepsilon_2$ and δ_1 can be chosen arbitrarily small, we see that $x(t) = x(0) + \int_0^t \chi(x(s), y(s)) \, ds$ for all $t < \inf\{t > 0 : |x(t-) - y(t-)| = 0\}$. Hence, x is a solution to (21).

Uniqueness of the solution of (21), for given $x(0)$ and general y , follows by reasoning as in (22) and (23). The continuous dependence of solutions on $x(0)$ and y , for general y , follows from the above estimates by approximating y by piecewise constant driving processes. \square

3. CAT(0) AND PURSUIT-EVASION

We consider the Lion and Man problem in a bounded CAT(0) domain D satisfying the uniform exterior sphere and interior cone conditions. [Alexander et al. \(2006\)](#) showed that $\varepsilon/2$ -capture, for given $\varepsilon > 0$, must occur for the discrete-time variant of this problem. As we will see in Section 4, the Lion and Man trajectories x and y will be weak limits of couplings of reflected Brownian motions, with drift and small noise, that arise from our capture problem.

We therefore modify the [Alexander et al. \(2006\)](#) argument to apply to continuous time; the modified argument also supplies an explicit upper bound on the capture time. We will only need to consider trajectories x and y that are Lipschitz with constant 1. Note that Lipschitz trajectories are absolutely continuous, so that the directions dx/dt and dy/dt are defined for almost all times t .

One can express the trajectories of Lion x and Man y as functions of time t in the following differential form:

$$(24) \quad \begin{aligned} dx &= \chi(x, y) \, dt - \nu_x \, dL^x, \\ dy &= H \, dt - \nu_y \, dL^y. \end{aligned}$$

Here, H is assumed to be a pre-assigned, time-varying unit length vector generating the motion of the Man, $\chi(x, y)$ generates the motion of the Lion and is defined as in Proposition 11, for $x \neq y$, as the unit tangent at x for the corresponding intrinsic geodesic, while $\nu_x \in \mathcal{N}_{x,r}$ and $\nu_y \in \mathcal{N}_{y,r}$ (for $r > 0$ satisfying the exterior sphere condition of D as given in Definition 5) determine the reflection

off of the boundary ∂D . The vector H is assumed to be measurable in t ; on account of Proposition 11, χ is continuous on $x \neq y$. The terms $\nu_x \, dL^x$, respectively $\nu_y \, dL^y$, are differentials arising from Skorokhod transformations and are differentials of functions of bounded variation that increase only when x , respectively y , belong to ∂D , and are then directed along an outward-pointing unit normal so as to cancel exactly with the outward-pointing component of the drifts $\chi \, dt$, respectively $H \, dt$.

We note that Skorokhod transformations are uniquely defined for a domain satisfying uniform exterior sphere and interior cone conditions (Saisho, 1987) (also compare earlier results of Lions and Sznitman, 1984), and they then depend continuously on the driving processes (using the uniform path metric). In fact, by the definition of χ , $\nu_x \, dL^x$ vanishes identically, while $\nu_y \, dL^y$ vanishes identically if $\langle H, \nu \rangle < 0$ whenever $y \in \partial D$. In particular, Proposition 14 applies and guarantees the existence of x and its approximation by piecewise-geodesic paths for y determined by H . (We include both the Skorokhod transformation differentials in (24) as they will both appear in the stochastic version in Section 4.)

We base our argument on Alexander et al. (2006, Theorem 12). The proof analyzes the greedy pursuit strategy arising from the definition of the vector field χ , with the Lion always directing its motion along the intrinsic geodesic from Lion to Man. The CAT(0) property forces the distance between Lion and Man to be non-increasing, and the Man must run directly away from the Lion in order to prolong successful evasion. Since the domain is bounded, this will, however, not be achievable indefinitely.

In order to demonstrate the main result in this section, Proposition 16, we will employ the following lemma.

Lemma 15. *Under the greedy pursuit strategy described above, in a CAT(0) domain satisfying uniform exterior sphere and interior cone conditions, and at a time t at which Lion and Man locations $x(t)$ and $y(t)$ are differentiable in t ,*

$$\frac{d}{dt} \text{dist}_{\text{intr}}(x(t), y(t)) = -(1 - |y'(t)| \cos \alpha(t))$$

where $\alpha(t)$ is the angle between the Man's velocity $y'(t)$ and the geodesic running from Lion to Man.

Proof. This follows immediately from the generalization of Gauss' lemma to such domains, as was established in Proposition 13. \square

Proposition 16. *Suppose that D is a bounded CAT(0) domain that satisfies a uniform exterior sphere condition based on a radius $r > 0$ and a uniform interior cone condition based on a radius $\delta > 0$ and angle $\alpha \in (0, \pi/2]$. Under the greedy pursuit strategy described above, there is a positive constant t_c depending only on the diameter of D and $\varepsilon > 0$ (and not on H in (24)) such that the Lion will come within distance $\varepsilon/2$ of the Man before time t_c , regardless of their starting positions within D .*

Remark. We use $\varepsilon/2$ here rather than ε , since a further distance $\varepsilon/2$ will be required by the stochastic part of the argument.

Proof. This proof follows Alexander et al. (2006), but is modified (a) to account for the continuous time context and (b) because we need to derive a specific upper bound t_c on the time of $\varepsilon/2$ -capture. Below, we abbreviate by setting $\mathcal{L}(t) = \text{dist}_{\text{intr}}(x(t), y(t))$.

Let α be the angle defined in Lemma 15. Evidently, the Lion will have come within $\varepsilon/2$ of the Man by time t unless

$$(25) \quad \int_0^t (1 - \cos \alpha) \, ds < \mathcal{L}(0) - \varepsilon/2 \leq \text{diam}_{\text{intr}}(D) - \varepsilon/2.$$

Now consider the total curvature of the Lion's path. By Proposition 14, the Lion's path is uniformly approximated by pursuit paths driven by discretized approximations to the Man's path. If $x^{(n)}$ is the Lion's path driven by a discretized Man's path $y^{(n)}$, then the Lion's path is piecewise-geodesic, with total absolute curvature given by the sum of the exterior angles formed at the points that connect the geodesics that occur when $x^{(n)}$ changes direction. CAT(0) comparison bounds then show the total curvature of $x^{(n)}$ is bounded above by

$$(26) \quad \sum \frac{\sin \alpha^{(n)}}{\text{dist}_{\text{intr}}(x^{(n)}, y^{(n)})} \Delta y^{(n)},$$

where summation is over the jumps of the discretized path $y^{(n)}$, and $\alpha^{(n)}$ is the exterior angle that the jump $\Delta y^{(n)}$ contributes to the geodesic running from $x^{(n)}$ to $y^{(n)}$.

The total curvature of a path is a lower-semicontinuous function of the path (using the uniform topology) for CAT(0) spaces. (This is a special case of a CAT(κ) result of Karuwanapatana and Maneesawang, 2007, referred to in Alexander et al., 2010, Theorem 18.) For the sake of completeness, we indicate the short proof for the CAT(0) case. Consider a curve q of finite length in a CAT(0) space. Its total curvature $\text{TC}(q)$ is the supremum of sums of exterior angles of piecewise-geodesic curves interpolating q ; a CAT(0) comparison argument shows that these sums of exterior angles increase as the interpolating mesh is refined. Let q^n be a sequence of curves converging uniformly to q . Furthermore, let $q^{n,m}$ be the piecewise-geodesic curve interpolating q^n at the points $k2^{-m}$ for $k = 0, 1, \dots$. Then, by definition of total curvature,

$$\text{TC}(q^{n,m}) \nearrow \text{TC}(q^n) \quad \text{as } m \rightarrow \infty.$$

Bridson and Haefliger (1999, Chapter II.3 Proposition 3.3) observe that the CAT(0) property implies that *interior* angles are continuous functions of their end vertices and upper-semicontinuous functions of their centre vertices. This upper-semicontinuity translates into lower-semicontinuity for exterior angles, and hence

$$\limsup_{n \rightarrow \infty} \text{TC}(q^{n,m}) \geq \text{TC}(q^{\infty,m}),$$

where $q^{\infty,m}$ is the uniform limit of $q^{n,m}$ as $n \rightarrow \infty$ (here we use the CAT(0) property again) and is a piecewise-geodesic interpolation of q at the points $k2^{-m}$ for $k = 0, 1, \dots$. Since $\text{TC}(q) =$

$\lim TC(q^{\infty,m})$, lower-semicontinuity now follows from

$$\limsup_{n \rightarrow \infty} TC(q^n) \geq \limsup_{n \rightarrow \infty} TC(q^{n,m}) \geq TC(q^{\infty,m}) \rightarrow TC(q) \quad \text{as } m \rightarrow \infty.$$

Consequently, the upper bound (26) provides an upper bound on the total absolute curvature of the Lion's path in the limit. Bearing in mind the Lipschitz(1) property of y , the total absolute curvature $\tau(t)$ incurred by x between times 0 and t therefore satisfies

$$(27) \quad \tau(t) \leq \int_0^t \frac{|\sin \alpha(s)|}{\mathcal{L}(s)} ds.$$

Assume that $\mathcal{L}(s) \geq \varepsilon/2$ for $s \leq t$. By the Cauchy-Schwartz inequality and Inequality (25),

$$(28) \quad \begin{aligned} \tau(t) &\leq \frac{2}{\varepsilon} \int_0^t |\sin \alpha| ds \leq \frac{2}{\varepsilon} \sqrt{t \int_0^t \sin^2 \alpha ds} \\ &\leq \frac{2\sqrt{2}}{\varepsilon} \sqrt{t \int_0^t (1 - \cos \alpha) ds} \leq \frac{2\sqrt{2}}{\varepsilon} \sqrt{\text{diam}_{\text{intr}}(D) - \varepsilon/2} \cdot \sqrt{t}. \end{aligned}$$

Next, we follow Alexander et al. (2006) in applying Reshetnyak majorization (Rešetnjak, 1968; see also the telegraphic description in Berestovskij and Nikolaev, 1993, Section 7.4) to generate a lower bound on the total absolute curvature of $\{x(s) : 0 \leq s \leq t\}$. We provide details for the sake of completeness.

We argue as follows. Reshetnyak majorization asserts that for every closed curve ζ in \bar{D} (more generally, in any CAT(0) space), one can construct a convex planar set C , bounded by a closed unit-speed curve $\bar{\zeta}$, and a distance-non-increasing continuous map $\phi : C \rightarrow \bar{D}$ such that $\phi \circ \bar{\zeta} = \zeta$; moreover, ϕ preserves the arc-length distances along $\phi \circ \bar{\zeta}$ and ζ . Consequently, ϕ restricted to ∂C will not increase angles and the pre-images under ϕ of geodesic segments in ζ must themselves be Euclidean geodesics (i.e., line segments).

By our assumptions about t , the total absolute curvature of $\{x(s) : 0 \leq s \leq t\}$ is finite (see (28)). Fix an arbitrarily small $\delta_1 \in (0, \pi/2)$. It follows from the definitions of length and curvature of a path that, for each n , we can approximate the unit-speed curve $\{x(s) : 0 \leq s \leq t\}$ by a piecewise-geodesic curve $\{z(s) : 0 \leq s \leq t\}$ with the following properties:

- The curve z is parametrized using arc-length.
- Note that x and y are continuous, so is $\chi(x, y)$, by Proposition 11 (3). Hence, we can choose $0 = t_0 < t_1 < \dots < t_n = t$ such that the total absolute curvature of $\{x(s) : t_{i-1} \leq s \leq t_i\}$ is equal to $\pi/2 - \delta_1$ for all i , with the possible exception of $i = n$.
- For every i , there exist $t_i = t_i^0 < t_i^1 < \dots < t_i^{m_i} = t_{i+1}$ and $s_i = s_i^0 < s_i^1 < \dots < s_i^{m_i} = s_{i+1}$ such that $z(s_i^j) = x(t_i^j)$ and z is geodesic on $[s_i^j, s_i^{j+1}]$, for all i and j . (Notice that the curve z is inscribed in the curve x .)
- The total absolute curvature of $\{z(s) : s_{i-1} \leq s \leq s_i\}$ is less than $\pi/2$. In other words, the sum (over j) of exterior angles between $\{z(s) : s_i^{j-1} \leq s \leq s_i^j\}$ and $\{z(s) : s_i^j \leq s \leq s_i^{j+1}\}$

- at s_i^j is less than $\pi/2$. (This is a consequence of z being inscribed in x and the CAT(0) property.)
- The difference between the lengths of $\{z(s) : 0 \leq s \leq t'\}$ and $\{x(s) : 0 \leq s \leq t\}$ is less than δ_1 .

Then we have

$$(29) \quad \text{total absolute curvature}(\{x(s) : 0 \leq s \leq t\}) \geq \left(\frac{\pi}{2} - \delta_1\right)(n-1).$$

We apply Reshetnyak majorization to the closed curve formed by $\{z(s) : s_{i-1} \leq s \leq s_i\}$ and its chord (the geodesic running from $z(t_i)$ back to $z(t_{i-1})$). Reshetnyak majorization guarantees that the total absolute curvature of $\{z(s) : s_{i-1} \leq s \leq s_i\}$ dominates the curvature of its pre-image in the boundary of a convex planar set C_i . Moreover, the perimeter of its pre-image in the boundary C_i has length $\text{len}(\{z(s) : s_{i-1} \leq s \leq s_i\})$, while the remainder of the boundary of C_i must be a line segment of length $\text{dist}_{\text{intr}}(z(s_i), z(s_{i-1}))$.

The two-dimensional pre-image of $\{z(s) : s_{i-1} \leq s \leq s_i\}$ therefore has total curvature bound of $\frac{\pi}{2}$. By two-dimensional Euclidean geometry, we can maximize the ratio of the length of the pre-image of $\{z(s) : s_{i-1} \leq s \leq s_i\}$ to the length $\text{dist}_{\text{intr}}(z(s_i), z(s_{i-1}))$ of its chord by considering the case of an isocles right-angled triangle, in which case the ratio is $\sqrt{2}$. Accordingly, we obtain the upper bound

$$\text{len}(\{z(s) : s_{i-1} \leq s \leq s_i\}) \leq \sqrt{2} \text{dist}_{\text{intr}}(z(s_i), z(s_{i-1})) \leq \sqrt{2} \text{diam}_{\text{intr}}(D).$$

It follows that a portion of the piece-wise geodesic curve z which turns no more than $\frac{\pi}{2}$ cannot have length exceeding $\sqrt{2}$ times the intrinsic diameter of the region. (Note this is related to the Euclidean diameter by Lemma 9.) This implies that we can control the total length of z and thus the total length of x , with

$$(30) \quad t - \delta_1 = \text{len}(\{x(s) : 0 \leq s \leq t\}) - \delta_1 \leq \text{len}(\{z(s) : 0 \leq s \leq t\}) \leq \sqrt{2} \text{diam}_{\text{intr}}(D) \times n.$$

Combining Inequalities (29) and (30), we deduce that

$$(31) \quad \begin{aligned} \text{total absolute curvature}(\{x(s) : 0 \leq s \leq t\}) &\geq \left(\frac{\pi}{2} - \delta_1\right)(n-1) \geq \\ &\geq \left(\frac{\pi}{2} - \delta_1\right) \left(\frac{t - \delta_1}{\sqrt{2} \text{diam}_{\text{intr}}(D)} - 1\right). \end{aligned}$$

Recall that $\tau(t) = \text{total absolute curvature}(\{x(s) : 0 \leq s \leq t\})$ and $\text{len}(\{x(s) : 0 \leq s \leq t\}) = t$. Letting $\delta_1 \rightarrow 0$ in (31), it follows that

$$\frac{\tau(t)}{t} \geq \frac{\pi}{2} \left(\frac{1}{\sqrt{2} \text{diam}_{\text{intr}}(D)} - \frac{1}{t} \right).$$

In combination with (28), this yields

$$\frac{\pi}{2} \left(\frac{1}{\sqrt{2} \operatorname{diam}_{\text{intr}}(D)} - \frac{1}{t} \right) \leq \frac{2\sqrt{2}}{\varepsilon} \sqrt{\operatorname{diam}_{\text{intr}}(D) - \varepsilon/2} \cdot \frac{1}{\sqrt{t}}$$

and hence the quadratic inequality for $q = \sqrt{t}$,

$$\left(\frac{\pi}{2} \frac{1}{\sqrt{2} \operatorname{diam}_{\text{intr}}(D)} \right) q^2 - \left(\frac{2\sqrt{2}}{\varepsilon} \sqrt{\operatorname{diam}_{\text{intr}}(D) - \varepsilon/2} \right) q - \frac{\pi}{2} \leq 0.$$

The left hand side is negative for $q = 0$ and the coefficient of q^2 is positive, so there is exactly one positive root q_c (which can be written out explicitly in terms of $\operatorname{diam}_{\text{intr}}(D)$ and ε). Combining this with our earlier arguments, it follows that the Lion will come within $\varepsilon/2$ of the Man by time $t_c := q_c^2$. \square

4. FROM BROWNIAN SHY COUPLINGS TO DETERMINISTIC PURSUIT PROBLEMS

This section is devoted to the proof of Theorem 1. Consider a co-adapted coupling of reflecting Brownian motions X and Y in the bounded domain $D \subseteq \mathbb{R}^d$ satisfying uniform exterior sphere and interior cone conditions. Saisho (1987) showed that the reflected Brownian motions can be realized by means of a Skorokhod transformation as strong solutions of stochastic differential equations driven by free Brownian motions. As discussed in Section 1.2, we can use arguments embedded in the folklore of stochastic calculus, and stated explicitly in Émery (2005) and in Kendall (2009, Lemma 6), to represent this coupling as

$$\begin{aligned} (32) \quad dX &= dB - \nu_X dL^X, \\ dY &= (\mathbb{J}^\top dB + \mathbb{K}^\top dA) - \nu_Y dL^Y, \end{aligned}$$

where A and B are independent d -dimensional Brownian motions, and \mathbb{J}, \mathbb{K} are predictable $(d \times d)$ -matrix processes such that

$$(33) \quad \mathbb{J}^\top \mathbb{J} + \mathbb{K}^\top \mathbb{K} = (d \times d) \text{ identity matrix.}$$

Here L^X and L^Y are the local times of X and Y on the boundary.

The advantage of this explicit representation of the coupling is that we can track what happens to X and Y when we modify the Brownian motion B by adding a drift. We will see that the effect of adding a very heavy drift based on the vectorfield $\chi(X, Y)$ will be to convert (32) into a stochastic approximation of the deterministic Lion and Man pursuit–evasion equations (24) over a short time-scale.

Arbitrarily denoting the Brownian particle X as the Brownian Lion or pursuer, and the other Brownian particle Y as the Brownian Man or evader, we define a new process by adding a drift $n\chi(X, Y) dt$ to the Brownian differential dB over a time period $[0, t_c/n]$, where t_c is the maximum time required for the deterministic strategy based on χ to bring the Lion to within $\varepsilon/2$ of the Man, as described in Proposition 16. This has the effect of adding drifts $n\chi(X, Y) dt$ to dX

and $n\mathbb{J}^\top \chi(X, Y) dt$ to dY over the time interval $[0, t_c/n]$. Thus one obtains the new stochastic differential system

$$(34) \quad \begin{aligned} dX &= dB + n\chi(X, Y) dt - \nu_X dL^X, \\ dY &= (\mathbb{J}^\top dB + \mathbb{K}^\top dA) + n\mathbb{J}^\top \chi(X, Y) dt - \nu_Y dL^Y. \end{aligned}$$

By the Cameron-Martin-Girsanov theorem, the distributions of the solutions of (32) and (34) are mutually absolutely continuous.

We will demonstrate that, for appropriate n , there is a positive probability of the Brownian Lion coming within distance ε of the Brownian Man by time t_c/n , for any pair of initial positions $X(0)$ and $Y(0)$ and any coupling strategy (corresponding to the pair of matrices \mathbb{J}, \mathbb{K} in (32)) that might be employed by the Brownian Man.

For comparison with the deterministic system (24), we employ the rescaling $X(t) = \tilde{X}^n(nt)$, $Y(t) = \tilde{Y}^n(nt)$, $B(t) = \tilde{B}^n(nt)/\sqrt{n}$, $A(t) = \tilde{A}^n(nt)/\sqrt{n}$, $\mathbb{J}(t) = \tilde{\mathbb{J}}^n(nt)$, $\mathbb{K}(t) = \tilde{\mathbb{K}}^n(nt)$. Setting $dt = d\tau/n$, (34) becomes

$$(35) \quad \begin{aligned} d\tilde{X}^n &= \frac{1}{\sqrt{n}} d\tilde{B}^n + \chi(\tilde{X}^n, \tilde{Y}^n) d\tau - \nu_{\tilde{X}^n} dL^{\tilde{X}^n}, \\ d\tilde{Y}^n &= \frac{1}{\sqrt{n}} \left((\tilde{\mathbb{J}}^n)^\top d\tilde{B}^n + (\tilde{\mathbb{K}}^n)^\top d\tilde{A}^n \right) + (\tilde{\mathbb{J}}^n)^\top \chi(\tilde{X}^n, \tilde{Y}^n) d\tau - \nu_{\tilde{Y}^n} dL^{\tilde{Y}^n}. \end{aligned}$$

Saisho (1987, Thm. 4.1) shows that the Skorokhod transformation is continuous under uniform exterior sphere and interior cone conditions on D , in the uniform topology. (This is not stated explicitly in Saisho's formulation but is immediate from the proof.) Weak convergence of a suitable subsequence of (X^n, Y^n) to a limiting process (X^∞, Y^∞) therefore follows as a consequence of the weak convergence of the corresponding subsequence of unreflected ("free") semimartingales U^n and V^n , where

$$\begin{aligned} dU^n &= \frac{1}{\sqrt{n}} d\tilde{B}^n + \chi(\tilde{X}^n, \tilde{Y}^n) d\tau, \\ dV^n &= \frac{1}{\sqrt{n}} \left((\tilde{\mathbb{J}}^n)^\top d\tilde{B}^n + (\tilde{\mathbb{K}}^n)^\top d\tilde{A}^n \right) + (\tilde{\mathbb{J}}^n)^\top \chi(\tilde{X}^n, \tilde{Y}^n) d\tau. \end{aligned}$$

The tightness of these families of semimartingales follows immediately from the tightness criterion given by Stroock and Varadhan (1979, §1.4) since the diffusion coefficients and the drifts are bounded by 1. So, on an appropriate subsequence, (U^n, V^n) converges weakly to a limiting process (U^∞, V^∞) and therefore, by Saisho (1987, Thm. 4.1), (X^n, Y^n) converges weakly to a limiting process (X^∞, Y^∞) along the same subsequence.

Consider now the Brownian martingale parts of U^n and V^n . Both converge weakly to the zero process as $n \rightarrow \infty$ since they are Brownian motions run at rate $\frac{1}{n}$. The locally-bounded variation parts must also converge weakly (taking a further subsequence if necessary) since they are both Lip(1); their limits are therefore also Lip(1). We deduce that

- (1) V^∞ is Lip(1);
- (2) U^∞ is Lip(1) and solves the *deterministic* differential equation

$$(36) \quad dU^\infty = \chi(X^\infty, Y^\infty) d\tau.$$

The coefficient $\chi(X^\infty, Y^\infty)$ in the above equation arises as a consequence of the continuous dependence on X^n and Y^n of the bounded vectorfield $\chi(X^n, Y^n)$, which depends continuously on X^n and Y^n (Proposition 11). We may therefore apply the dominated convergence theorem to deduce the integral representation for U^∞ corresponding to (36).

In the following, we will use the Skorokhod representation to replace the weak convergence of the previous sequences by an almost-sure convergence for U^∞ and V^∞ that is uniform over bounded time intervals. We also show that the limit paths for X^n, Y^n (which must be X^∞, Y^∞ by Saisho, 1987, Thm. 4.1) are both still Lip(1), *with respect to the intrinsic metric of D* . (This is required in order to apply the results on CAT(0) Lion and Man problems at the end of Section 2.) The following lemma establishes this Lipschitz property by using the discrete approximation technique of Saisho (1987):

Lemma 17. *Let D be a domain satisfying uniform exterior sphere and interior cone conditions. Suppose that Z is a continuous process on \bar{D} derived by the Skorokhod transformation from a free process S that has Lip(1) sample paths. Then Z itself has Lip(1) sample paths with respect to the intrinsic metric.*

Proof. Following Saisho (1987, §3), consider the step function S_m obtained from S by sampling at instants $k2^{-m}$, for $k = 0, 1, \dots$. Suppose that $2^{-m} < r$, where r is the radius on which the uniform exterior sphere condition is based. Let \bar{z} be the projection onto \bar{D} described in Lemma 6. The Skorokhod transformation of S_m is Z_m , given by projecting increments back onto \bar{D} :

$$(37) \quad Z_m(t) = \begin{cases} \overline{Z_m((k-1)2^{-m}) + \Delta S_m(k2^{-m})} & \text{for } k2^{-m} \leq t < (k+1)2^{-m}, \\ Z(0) & \text{for } 0 \leq t < 2^{-m}. \end{cases}$$

On account of the Lip(1) property of S , this projection is defined when $2^{-m} < r$.

From Saisho (1987, Theorem 4.1), we know that $Z_m \rightarrow Z$ uniformly on bounded time intervals. We compute the maximum possible Euclidean distance between $Z_m(s)$ and $Z_m(t)$, if $0 \leq t - s < 2^{-m}$, when one or both of $2^m s, 2^m t$ are non-negative integers. Since Z_m is constant on intervals $[k2^{-m}, (k+1)2^{-m})$, it suffices to produce an argument for the case when $2^m s = k - 1$ and $2^m t = k$. We therefore proceed to bound the Euclidean distance $|Z_m(k2^{-m}) - Z_m((k-1)2^{-m})|$. We will show that this can only exceed 2^{-m} by an amount which, for large m , will make a negligible contribution to path length when summed over the whole path.

If $Z_m(k2^{-m}) \notin \partial D$, then there is nothing to prove, since the jump is $\Delta S_m(k2^{-m})$, which is bounded in length by 2^{-m} since S is Lip(1). So we instead suppose that $Z_m(k2^{-m}) \in \partial D$. For convenience, set $y = Z_m(k2^{-m}) - (Z_m((k-1)2^{-m}) + \Delta S_m(k2^{-m}))$ to be the Skorokhod correction to be applied at this step, and set $a = |\Delta S_m(k2^{-m})|$ to be the length of the uncorrected jump. Finally, let θ be the angle between the vector y and the negative jump $-\Delta S_m(k2^{-m})$. These definitions

are illustrated in Figure 9, together with the supporting ball B at $Z_m(k2^{-m}) \in \partial D$ whose centre is located at $Z_m(k2^{-m}) - \lambda y$ for some $\lambda = r/|y| > 0$ and whose existence is guaranteed by the construction of the $x \mapsto \bar{x}$ projection map as described in Lemma 6.




FIGURE 9. Illustration of the geometry underlying the argument of Lemma 17.

First note that $|Z_m(k2^{-m}) - Z_m((k-1)2^{-m})| = \sqrt{a^2 + y^2 - 2ay \cos \theta}$ (where we abuse notation by letting y also stand for the length of the vector y). This increases as θ increases to π , so long as $a, y, Z_m(k2^{-m})$ are held fixed. Thus we can assume that θ has increased to the point where $Z_m((k-1)2^{-m})$, as well as $Z_m(k2^{-m})$, belong to ∂B . (This will happen if, as required above, $2^{-m} < r$.) Now observe that the distance $|Z_m(k2^{-m}) - Z_m((k-1)2^{-m})|$ will be bounded above by the smaller of the two distances from $Z_m(k2^{-m})$ to the intercepts of ∂B by a line parallel to y , and at distance a from $Z_m(k2^{-m})$. But two applications of Pythagoras' theorem show that this distance is given by

$$\begin{aligned} \sqrt{a^2 + (r - \sqrt{r^2 - a^2})^2} &= \sqrt{2r^2 - 2r\sqrt{r^2 - a^2}} \\ &= \sqrt{2r} \sqrt{1 - \sqrt{1 - \frac{a^2}{r^2}}} = \sqrt{2r} \sqrt{\frac{a^2}{2r^2} \left(1 + \frac{1}{4}(z^*)^2\right)} \end{aligned}$$

for some z^* in the range $[0, \frac{r^2}{a^2}]$. (The last step arises from a second-order Taylor series expansion.) Therefore

$$\sqrt{a^2 + (r - \sqrt{r^2 - a^2})^2} \leq \sqrt{2r} \sqrt{\frac{a^2}{2r^2} \left(1 + \frac{a^2}{4r^2}\right)} \leq a \left(1 + \frac{a^2}{8r^2}\right)$$

(using $\sqrt{1+z} \leq 1 + \frac{1}{2}z$ for $z \geq -1$).

Thus the total path length over the time interval (s, t) is bounded above by

$$((t-s)2^m + 2) \times 2^{-m} \left(1 + \frac{2^{-2m}}{8r^2}\right),$$

which converges to $t-s$ as $m \rightarrow \infty$. Hence we obtain

$$\text{dist}_{\text{intr}}(Z(s), Z(t)) \leq t-s,$$

thus establishing the Lip(1) property in intrinsic metric for Z . \square

Proposition 16 states that X^∞ comes within distance $\varepsilon/2$ of Y^∞ in the intrinsic metric by time $\tau = t_c$. Since $(\tilde{X}^n, \tilde{Y}^n)$ converges weakly to (X^∞, Y^∞) , it follows that, for sufficiently large n , there is a positive probability of \tilde{X}^n coming within distance ε of \tilde{Y}^n by time $\tau = t_c$ in this metric. On the other hand, $\text{dist}_{\text{intr}}(x, y) \geq \|x - y\|$. Therefore, by the Cameron-Martin-Girsanov argument given at the beginning of the section, for X and Y satisfying (32) and given $X(0), Y(0) \in \overline{D}$, there is a positive probability of X coming within distance ε of Y by time $t = t_c/n$. We will show that this bound is uniform over all $X(0)$ and $Y(0)$.

Proposition 18. *Suppose that there exist $t_1 > 0$ and $\varepsilon_1 > 0$ such that, for any X and Y satisfying (32) with $X(0), Y(0) \in \overline{D}$,*

$$(38) \quad \mathbb{P} \left[\inf_{0 \leq t \leq t_1} |X(t) - Y(t)| \leq \varepsilon_1 \right] > 0.$$

Then

$$(39) \quad \mathbb{P} \left[\inf_{0 \leq t \leq t_1} |X(t) - Y(t)| \leq \varepsilon_1 \right] > p_1$$

for some $p_1 > 0$ not depending on $X(0)$ and $Y(0)$.

Proof. Suppose (39) does not hold. Then there exist $t_1 > 0$, $\varepsilon_1 > 0$, sequences $\{x_n\}_{n \geq 1}$, $\{y_n\}_{n \geq 1}$ of points in \overline{D} , random processes $\{A_t, t \geq 0\}$, $\{B_t, t \geq 0\}$, $\{\mathbb{J}_t^n, t \geq 0\}$ and $\{\mathbb{K}_t^n, t \geq 0\}$, and solutions X^n and Y^n of (32) satisfying the following properties. The processes A and B are d -dimensional Brownian motions starting from 0, and independent of each other. The $(d \times d)$ -matrix-valued processes \mathbb{J}^n and \mathbb{K}^n are predictable with respect to the natural filtration of A and B , such that $(\mathbb{J}_t^n)^\top \mathbb{J}_t^n + (\mathbb{K}_t^n)^\top \mathbb{K}_t^n$ is the $(d \times d)$ identity matrix at all times t . Let X^n and Y^n denote solutions to (32) based on the Brownian motions A and B , using the predictable integrators \mathbb{J}^n and \mathbb{K}^n , and starting from $X^n(0) = x_n \in \overline{D}$ and $Y^n(0) = y_n \in \overline{D}$. Then

$$(40) \quad \mathbb{P} \left[\inf_{0 \leq t \leq t_1} |X^n(t) - Y^n(t)| > \varepsilon_1 \right] > 1 - 2^{-n}.$$

Let $(M_t^{n,1}, M_t^{n,2}) = (\int_0^t d B_s, \int_0^t (\mathbb{J}_s^n)^\top d B_s + \int_0^t (\mathbb{K}_s^n)^\top d A_s)$. The processes $M^{n,1}$ and $M^{n,2}$ are Brownian motions and so the sequence of pairs is tight, which therefore possesses a subsequence converging in distribution. By abuse of notation, we assume that the whole sequence $(M^{n,1}, M^{n,2})$

converges in distribution to, say, $(M^{\infty,1}, M^{\infty,2})$. It is clear that $M^{\infty,1}$ and $M^{\infty,2}$ are Brownian motions.

Let $\mathcal{F}_t = \sigma((M_s^{\infty,1}, M_s^{\infty,2}), s \leq t)$ be the natural filtration for $(M^{\infty,1}, M^{\infty,2})$. We will show that $(M^{\infty,1}, M^{\infty,2})$ are co-adapted Brownian motions relative to $\{\mathcal{F}_t\}$. Since $(M^{n,1}, M^{n,2})$ are co-adapted Brownian motions, for all $0 \leq t_1 \leq t_2 \leq \dots \leq t_n \leq t \leq s_1 \leq s_2$, the random variable $M_{s_2}^{n,1} - M_{s_1}^{n,1}$ is independent of

$$((M_{t_1}^{n,1}, M_{t_1}^{n,2}), (M_{t_2}^{n,1}, M_{t_2}^{n,2}), \dots, (M_{t_n}^{n,1}, M_{t_n}^{n,2})).$$

Independence is preserved by weak limits, so $M_{s_2}^{\infty,1} - M_{s_1}^{\infty,1}$ is independent of

$$((M_{t_1}^{\infty,1}, M_{t_1}^{\infty,2}), (M_{t_2}^{\infty,1}, M_{t_2}^{\infty,2}), \dots, (M_{t_n}^{\infty,1}, M_{t_n}^{\infty,2})).$$

This implies that $M_{s_2}^{\infty,1} - M_{s_1}^{\infty,1}$ is independent of \mathcal{F}_t . Since the same argument applies to $M_{s_2}^{\infty,2} - M_{s_1}^{\infty,2}$, we see that $(M^{\infty,1}, M^{\infty,2})$ are co-adapted relative to $\{\mathcal{F}_t\}$. Recall from Section 1.2 that this implies that there exist Brownian motions $\{A_t^\infty, t \geq 0\}$ and $\{B_t^\infty, t \geq 0\}$ and processes $\{\mathbb{J}_t^\infty, t \geq 0\}$ and $\{\mathbb{K}_t^\infty, t \geq 0\}$ such that $(M_t^{\infty,1}, M_t^{\infty,2}) = (\int_0^t d B_s^\infty, \int_0^t (\mathbb{J}_s^\infty)^\top d B_s^\infty + \int_0^t (\mathbb{K}_s^\infty)^\top d A_s^\infty)$.

Recall that $(M^{n,1}, M^{n,2}) \rightarrow (M^{\infty,1}, M^{\infty,2})$ weakly in the uniform topology on all compact intervals. Going back to the original notation, we see that

$$\left(\int_0^t d B_s, \int_0^t (\mathbb{J}_s^n)^\top d B_s + \int_0^t (\mathbb{K}_s^n)^\top d A_s \right) \rightarrow \left(\int_0^t d B_s^\infty, \int_0^t (\mathbb{J}_s^\infty)^\top d B_s^\infty + \int_0^t (\mathbb{K}_s^\infty)^\top d A_s^\infty \right)$$

weakly in the uniform topology on all compact intervals. By the Skorokhod Lemma, we can assume that the processes converge a.s. in the supremum topology on compact intervals.

Since \overline{D} is compact, we can assume, passing to a subsequence if necessary, that the initial points satisfy $x_n \rightarrow x_\infty \in \overline{D}$ and $y_n \rightarrow y_\infty \in \overline{D}$ as $n \rightarrow \infty$. In view of the representation of coupled reflected Brownian motions using stochastic differential equations (32), employing Saisho (1987, Theorem 4.1), we see that $(X^n, Y^n) \rightarrow (X^\infty, Y^\infty)$ weakly in the uniform topology on all compact intervals, where (X^∞, Y^∞) represents the solution to (32) with $X^\infty(0) = x_\infty$, $Y^\infty(0) = y_\infty$, corresponding to A^∞ , B^∞ , \mathbb{J}^∞ and \mathbb{K}^∞ . We obtain from (40) and weak convergence of (X^n, Y^n) to (X^∞, Y^∞) that, for every n ,

$$\mathbb{P} \left[\inf_{0 \leq t \leq t_1} |X^\infty(t) - Y^\infty(t)| \geq \varepsilon_1 \right] \geq 1 - 2^{-n}.$$

Taking the limit as $n \rightarrow \infty$, this contradicts (38) in the statement of the Proposition. Consequently (39) must hold for some p_1 . \square

We now complete the proof of Theorem 1, applying Proposition 18 together with standard reasoning. Consider processes X and Y starting from any pair of points in \overline{D} and corresponding to any “strategy” \mathbb{J} and \mathbb{K} . Because of the uniform bound in Proposition 18, the probability of X

and Y not coming within distance ε_1 of each other on the interval $[kt_1, (k+1)t_1]$, conditional on not coming within this distance before kt_1 , is bounded above by $1 - p_1$ for any k , by the Markov property. Hence, the probability of X and Y not coming within distance ε_1 of each other on the interval $[0, kt_1]$ is bounded above by $(1 - p_1)^k$. Letting $k \rightarrow \infty$, it follows that X and Y are not ε_1 -shy. Since ε_1 can be taken arbitrarily small, the proof of Theorem 1 is complete.

We remark that the matrices \mathbb{J} and \mathbb{K} employed in (32) are predictable and, consequently, the choice of the pursuer's velocity is made without looking into the future. This is in contrast to the pursuit-evasion problems and associated paradoxes discussed by Bollobas, Leader, and Walters (2009).

5. COMPLEMENTS AND CONCLUSIONS

We conclude this paper by remarking on some supplementary results and concepts, and by considering possibilities for future work.

5.1. Comparison with previous methods. The fundamental idea in this paper turns out in the end to resemble that of Benjamini et al. (2007), but uses simple notions of weak convergence and tightness, rather than detailed large deviation estimates. Moreover, the use of metric geometry notions enables us to finesse many analytical technicalities. (Perhaps this is the first application of modern metric geometry to Euclidean stochastic calculus?) On the other hand, the stochastic control methods of Kendall (2009) are quite different. The stochastic control approach uses potential theory to estimate the value function of an associated stochastic game; consequently the methods of Kendall (2009) may be expected to give sharper information (bounds on expectation of stopping times), but in more limited cases (convexity of domain). However, one can observe that, at least in principle, the stochastic game formulation still applies in the general case. For example, there is a value function to be discovered for a stochastic control reformulation of Theorem 1, and in principle it might be possible to estimate this value function and so gain more information than is supplied by the weak geometric bounds established above.

We note that many promising ideas based on stochastic calculus fail to show non-shyness because they cannot be applied to “perverse” couplings with the property that, on some time intervals, $|X - Y|$ grows at a deterministic rate. (See Example 4.2 of Benjamini et al., 2007.)

Also note that the proof in Kendall (2009), which works in convex domains, does not appear to be (directly) extendable to calculations involving the intrinsic metric – simple manipulation using symbolic Itô calculus (Kendall, 2001) shows that the drift of $\text{dist}_{\text{intr}}(X, Y)$ is unbounded at distances bounded away from zero. In particular, Bessel-like divergences for $\text{dist}_{\text{intr}}(X, Y)$ of magnitude a occur when the geodesic from X to Y touches a concave part of ∂D at x and $|x - Y| = 1/a$. The first-order differential geometry given in Proposition 13 (the generalization of Gauss' lemma) is the best we can do for CAT(0) domains satisfying uniform exterior sphere and interior cone conditions.

5.2. Higher dimensions and the failure of CAT(0). For planar domains, CAT(0) and simple-connectedness are equivalent, in which case, by Theorem 2, there are no shy co-adapted couplings. In higher dimensions, it is natural to ask whether the CAT(0) condition is essential for there to be no shy coupling. We do not at all believe this to be the case. It is possible to give an argument suggesting that star-shaped domains with smooth boundary conditions cannot support shy coupling, by establishing the analogous result for a corresponding deterministic pursuit-evasion problem. To apply this argument to the probabilistic case would require more careful arguments. We therefore leave this as a project for another day.

As a spur to future work, we formulate a bold and possibly rash conjecture:

Conjecture 1. *There can be no shy co-adapted coupling for reflecting Brownian motions in bounded contractible domains in any dimension.*

While resolution of the star-shaped case appears to be largely a technical matter, we believe that new ideas will be required to make substantial progress towards resolving the conjecture.

5.3. When can shyness exist? Many examples of shy coupling can be generated using suitable symmetries. However, we do not know of any examples in which symmetries play no rôle. Accordingly we formulate a further conjecture:

Conjecture 2. *If a bounded domain D supports a shy co-adapted coupling for reflecting Brownian motions, then the coupling can be realized using a rigid-motion symmetry of the domain D .*

Note that Benjamini et al. (2007, Example 3.9) supplies an example based on Brownian motion on graphs, for which there is no fixed-point-free isometry and yet a shy coupling exists. However we do not see how to use the idea of this construction to construct a counterexample to the above conjecture.

5.4. Further questions. We enumerate a short list of additional questions.

- (1) Shyness is interesting for foundational reasons: coupling is an important tool in probability, and shyness informs us about coupling. We do not know of any honest applications of shyness. However, one can contrive a kind of cryptographic context. Suppose one wishes to mimic a target Y , which is a randomly evolving high-dimensional structure, in such a way that the mimic X never comes within a certain distance of the target Y . Shyness concerns the question, whether it is possible to do this in a way that is perfectly concealed from an observer watching the mimic X alone.
- (2) In this formulation, it is not clear why one should restrict consideration to co-adapted couplings. Our methods do not lend themselves to the non-co-adapted case, and the question is open whether or not results change substantially if one is allowed to use such couplings. In particular, it seems possible that Conjecture 2 might have a quite different answer in this context.

- (3) Finally, the Lion and Man problem has been generalized to the case of multiple Lions. (An early instance is given in [Croft, 1964](#).) Can one formulate and prove useful results for a corresponding notion of multiple shyness?

Acknowledgments. We are grateful to Soumik Pal for giving most helpful advice.

REFERENCES

- Alexander, S. B., R. L. Bishop, and R. Ghrist (2006). Pursuit and evasion in non-convex domains of arbitrary dimension. In *Proceedings of Robotics: Science and Systems*, Philadelphia, USA.
- Alexander, S. B., R. L. Bishop, and R. Ghrist (2010). Total curvature and simple pursuit on domains of curvature bounded above.
- Benjamini, I., K. Burdzy, and Z.-Q. Chen (2007). Shy couplings. *Probability Theory and Related Fields* 137(3-4), 345–377.
- Berestovskij, V. N. and I. G. Nikolaev (1993). Multidimensional generalized Riemannian spaces. In *Geometry, {IV}*, Volume 70 of *Encyclopaedia Math. Sci.*, pp. 165–243, 245–250. Berlin: Springer.
- Bieske, T. (2010). The Carnot-Carathéodory distance vis-a-vis the eikonal equation and the infinite Laplacian. *Bulletin of the London Mathematical Society* 42(3), 395–404.
- Bishop, R. L. (2008). The intrinsic geometry of a Jordan domain. *Int. Electron. J. Geom.* 1(2), 33–39.
- Bollobas, B., I. Leader, and M. Walters (2009). Lion and Man – Can Both Win? arXiv 0909.2524.
- Bridson, M. R. and A. Haefliger (1999). *Metric Spaces of Non-Positive Curvature*, Volume 319 of *Grundlehren der Mathematischen Wissenschaften*. Berlin: Springer-Verlag.
- Burago, D., Y. Burago, and S. Ivanov (2001). *A Course in Metric Geometry*, Volume 33 of *Graduate Studies in Mathematics*. Providence, RI: American Mathematical Society.
- Cheeger, J. and D. G. Ebin (2008). *Comparison Theorems in Riemannian Geometry*. AMS Chelsea Publishing, Providence, RI.
- Croft, H. T. (1964). “Lion and man”: A postscript. *The Journal of the London Mathematical Society (Second Series)* 39, 385–390.
- Émery, M. (2005). On certain almost Brownian filtrations. *Ann. Inst. H. Poincaré Probab. Statist.* 41(3), 285–305.
- Isaacs, R. (1965). *Differential Games. A Mathematical Theory with Applications to Warfare and Pursuit, Control and Optimization*. New York: John Wiley & Sons.
- Karuwannapatana, W. and C. Maneesawarng (2007). The lower semi-continuity of total curvature in spaces of curvature bounded above. *East-West J. Math.* 9(1), 1–9.
- Kendall, W. S. (2001). Symbolic Itô calculus: an ongoing story. *Statistics and Computing* 11, 25–35.
- Kendall, W. S. (2009). Brownian couplings, convexity, and shy-ness. *Electronic Communications in Probability* 14 (Paper 7), 66–80.
- Lions, P.-L. and A.-S. Sznitman (1984). Stochastic differential equations with reflecting boundary conditions. *Comm. Pure Appl. Math.* 37(4), 511–537.

- Littlewood, J. E. (1986). *Littlewood's Miscellany*. Cambridge: Cambridge University Press.
- Nahin, P. J. (2007). *Chases and Escapes*. Princeton, NJ: Princeton University Press.
- Rešetnjak, J. G. (1968). Non-expansive maps in a space of curvature no greater than K . *Sibirsk. Mat. Ž.* 9, 918–927.
- Saisho, Y. (1987). Stochastic differential equations for multidimensional domain with reflecting boundary. *Probability Theory and Related Fields* 74(3), 455–477.
- Stroock, D. W. and S. R. S. Varadhan (1979). *Multidimensional Diffusion Processes*, Volume 233 of *Grundlehren der Mathematischen Wissenschaften*. Berlin: Springer-Verlag.

(M.B.) SCHOOL OF MATHEMATICS, VINCENT HALL, 206 CHURCH ST. SE., UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MN 55455, USA

E-mail address: bramson@math.umn.edu

(K.B.) DEPARTMENT OF MATHEMATICS, BOX 354350, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195, USA

E-mail address: burdzy@math.washington.edu

(W.S.K.) DEPARTMENT OF STATISTICS, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UK

E-mail address: w.s.kendall@warwick.ac.uk