

University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): J.D. Reader, M.J. Green, J. Kaler, S.A. Mason, L.E. Green

Article Title: Effect of mobility score on milk yield and activity in dairy cattle

Year of publication: 2011

Link to published article:

<http://dx.doi.org/10.3168/jds.2011-4415>

Publisher statement: NOTICE: this is the author's version of a work that was accepted for publication in Journal of Dairy Science. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Journal of Dairy Science, Vol. 94, Issue 10, October 2011, DOI: 10.3168/jds.2011-4415

19 **ABSTRACT**

20 Previous studies have indicated that lame cows have a reduced milk yield both before
21 and after they are treated. One explanation for the reduction in yield before treatment is that
22 there is a delay to treatment, that is, cows have impaired mobility for some time before they
23 are treated. The aim of this study was to test this hypothesis by investigating temporal
24 associations between change in milk yield and change in mobility score. Mobility score (MS,
25 on a scale 0 to 3), milk yield, treatments for lameness and cow activity were recorded on 312
26 cows in a dairy herd in Somerset, UK for 1 yr. The MS was scored every 2 wk and
27 compared with the daily yield and activity (steps/h) averaged over the previous 16 d.
28 Approximately 44 % of MS changed within 14 d, usually by 1 score. Overall, milk yields of
29 cows with MS 1 were higher than those of cows with other scores. Cows with MS 2 and 3
30 produced 0.7 (0.35 - 0.97) kg and 1.6 (0.98 – 2.23) kg less milk / d, respectively, compared
31 with cows with MS 1. In addition, cows with MS 1 were slightly but significantly more
32 active than cows with MS 0, 2 or 3. Cows with MS 2 and 3 were 0.002 (0.01 – 0.03) and
33 0.03 (0.01 – 0.05) mean log steps less active than cows with MS 1.

34 There was a reduction in yield from 6 - 8 wk before becoming MS 2 0.5 (0.12 – 0.47)
35 or 3 0.9 (0.16 – 1.65) to 4 wk after recovering from MS 2 0.42 (0.09 – 0.75) and non-
36 significantly, score 3. The activity of cows was significantly less but quantitatively small
37 (mean log steps 0.01) with increasing MS; the associations between activity and parity
38 (mean 0.03 – 0.11) and month of lactation (mean 0.03 – 0.36) were quantitatively larger.
39 Results from a multistate model indicated that once cows were lame they remained lame or
40 become lame again despite treatment. We conclude that cows started to reduce milk
41 production before their mobility is visibly impaired. One explanation for this is that MS is not
42 100% sensitive. An alternative hypothesis, using evidence from other studies, is that reduction
43 in milk yield and development of lameness are on a common causal pathway most likely

44 linked to loss in body condition and reduced digital cushion thickness as a result of the
45 demands from producing high milk yields.

46

47 **Key words** Dairy cow, Milk yield, Lameness, Treatment, Multistate model

48

INTRODUCTION

49 The prevalence and incidence of lameness in dairy cows in intensive systems is
50 unacceptably high with estimates of prevalence in the UK ranging from 21 % (Clarkson et al.,
51 1996) to 36 % (Leach et al., 2010). Lamé cows are in pain and their welfare is compromised
52 (Whay et al., 1997).

53 Lameness is associated with a reduction in milk yield (Juarez et al., 2003; Archer et
54 al., 2010). This reduced milk yield is present before and after a treatment event, but varies by
55 the type of lesion (Green et al., 2002; Amory et al., 2008; Bicalho et al., 2008). The reduction
56 in yield detected before a treatment event with non infectious horn lesions (Amory et al., 2008;
57 Green et al., 2010) might occur because of a long pathogenesis in disease before cows become
58 lame or because of delayed treatment. There is less evidence that infectious claw conditions
59 are associated with reduced milk yield before cows are observed lame, although Warnick et al.
60 (2001) reported that interdigital phlegmon was associated with reduced yield before treatment,
61 possibly because the time to lameness from infection is rapid. For both types of disorders,
62 delay in treatment would probably lead to reduced milk yield because of the increased
63 metabolic demands from pain and reduced feed intake. The treatment of lame cows depends
64 on the ability of farmers to recognize a lame cow and to treat affected cows promptly and
65 appropriately. Most dairy cow farmers underestimate the prevalence of lameness on their
66 farms (Whay et al., 2003) and do so inconsistently compared with a trained researcher
67 (Leach et al., 2010), suggesting that most dairy cow herdsmen do not have a logical way
68 of detecting lameness, in contrast to sheep farmers (King and Green, in press).

69 Mobility scoring has been developed to help farmers improve detection of mild
70 lameness and stimulate treatment and prevention as part of a herd health program. The
71 currently accepted system used in the UK is a 4 point mobility scoring (**MS**, on a scale 0 to
72 3) system (Whay et al., 2003). This system is used by many researchers and veterinary
73 practitioners, but has not been evaluated for repeatability. Some authors have reported that
74 daily activity levels are lower in cows with reduced mobility (O'Callaghan et al., 2003;
75 Mazrier et al., 2006; Walker et al., 2008).

76 The current study was designed to test the hypothesis that the reduction in milk yield
77 that occurs before lame cows are treated is as a result of delayed treatment. This was tested
78 by investigating the temporal association between change in milk yield and change in
79 locomotion and time to treatment. The MS, milk yield, and activity in cattle from 1 farm
80 was observed every 2 wk for 1 yr to estimate precise relationships between MS and changes in
81 MS, milk yield, and cow activity.

82 **MATERIALS AND METHODS**

83 A dairy herd that calved all year round, located in Somerset UK, with a milking herd of
84 200 Holstein cows, producing approximately 9,000 kg milk/cow per year was used for the
85 study. The study started on October 24, 2007 and finished on November 5, 2008. Calving was
86 all yr around; The numbers of cows in milk ranged from 168 (November 5, 2008) to 217
87 (April 23, 2008) with a mean of 197 and median of 200. The herd was divided into 2 groups of
88 about equal size based on milk yield, both housed in 1 building with a floor of concrete and 230
89 free stalls fitted with mattresses and bedded with sawdust. Milking cows had access to pasture
90 in summer with high yielding cows only on pasture for a limited period each day. Non-
91 lactating cows were kept in a separate building and their locomotion was not scored. The
92 herd was milked twice daily through an 18/36 Westfalia herringbone parlor. Milking cows
93 walked through a 5% formalin footbath as they exited the parlor once each week.

94 Cows were selected for foot trimming by the herdsman. Approximately 35 cows
95 were trimmed per month; foot trimming was carried out by a paraprofessional foot trimmer
96 from Kingfisher Veterinary Practice (Synergy Farm Health, West Hill Barns, Evershot,
97 Dorset, England. DT2 0LD). The selection criteria for foot trimming were cows that were
98 clinically lame (MS 2 or 3) or cows that were due to be dried off. The farmer intended to
99 trim feet of each cow at least once each year, but this was not cross checked. Lesions were
100 defined using the definitions in the EU Lamecow Project (Barker et al., 2007) and all foot
101 trimming and lameness were recorded on lameness scoring sheets designed by the EU
102 Lamecow project. Cases of lameness treated by the herdsman or veterinarian (who treated
103 severe cases) were recorded in the same way.

104 All cows were individually identified and fitted with pedometers (Westfalia Dairy
105 Plan C21 (GEA Farm Technologies Australia Pty. Ltd. PO Box 39816 Trade Park Drive
106 Tullamarine VIC 3043). Activity readings for each cow were automatically downloaded to
107 the farm computer in the parlor twice daily and onto a lap top once weekly. The mobility of
108 lactating cows was scored (Table 1) every 2 wk after evening milking by JDR using the
109 system described by Whay et al., (2003). The identity of each cow was recorded as she
110 entered the parlor and mobility was scored and recorded on standardized sheets as the cow
111 exited the parlor. The MS was transferred to an Excel 2003 spreadsheet (Microsoft Corp.,
112 Redmond, WA). Milk yield, activity (mean steps/hr), health records, lameness records, and
113 group were downloaded from the farm computer into the spreadsheet.

114 *Data analysis*

115 The mean proportion of cows with each MS by stage of lactation (1 to 90 d, 91 to 180 d,
116 >180 d), mean milk yield, and mean activity over 16 d previously were calculated. The
117 probability of transition between MS from time t to time $t + 1$, 14 d later, was estimated.
118 Two multilevel statistical models were constructed, using conventional methods (Goldstein,

119 1995). In the first model the outcome variable was mean milk yield in the 16 d before a MS and
120 the impact of MS before and after this outcome was investigated. In the second model \log^{10}
121 mean activity score for the previous 16 d was the outcome and the impact of MS on activity
122 was investigated.

123 The models took the form:

$$124 \quad Y_{ij} = \alpha + \beta_1 X_{ij} + \beta_2 X_j + v_j + e_{ij} \quad v_j \sim N(0, \sigma_v^2)$$

$$125 \quad e_{ij} \sim N(0, \sigma_e^2)$$

126 where the subscripts i , and j denote the i^{th} observation of the j^{th} cow, respectively; α is the
127 regression intercept; X_{ij} is the vector of covariates associated with each observation; β_1 the
128 coefficients for covariates X_{ij} ; X_j the vector of covariates associated with each cow; β_2 the
129 coefficients for covariates X_j ; v_j a random effect to reflect residual variation between cows
130 which is normally distributed with mean = 0 and variance = σ^2 ; and e_{ij} a random effect to
131 reflect residual variation between MS which is normally distributed with mean = 0 and
132 variance = σ^2 . The analysis was carried out using MLwiN 2.02 with penalized quasi-
133 likelihood for parameter estimation (Rasbash et al., 2005). Covariates were left in the model
134 when the significance probability was $P < 0.05$ based on the Wald Test. When mean milk
135 yield was the outcome, DIM, the exponential DIM^{0.05} (Wilmink, 1987) and parity 1, 2, 3, and >
136 3, and first or second lactation in the study were forced into the model. Then the discrete
137 variable MS (0, 1, 2, and 3) at time t was added. The impact of MS at time $t - 1$, $t - 2$, ..., $t - 5$
138 and $t + 1$, $t + 2$, ..., $t + 5$, where each time interval i was 14 d, was tested in the model. When
139 log mean activity was the outcome, parity 1, 2, 3, and > 3, second lactation in the study and
140 month in milk were forced into the model and then the mobility score at times t , $t - 1$, ..., $t - 5$
141 and $t + 1$, ..., $t + 5$, where each time interval t was 14 d, were tested in the model. Missing
142 observations were random and so were fitted in the model as discrete variables to minimize
143 loss of data. The model fit was checked.

144 Finally, a multistate model was set up to test the factors associated with cows
145 becoming lame, remaining lame, becoming sound, and remaining sound. Mobility score was
146 categorized into 2 states: not lame (scores 0 and 1) and lame (scores > 1). A cow was in 1 of
147 2 states, not lame or lame. An episode was defined as the continuous period of time a cow
148 spent in either state until a transition to the other state occurred. For each episode j for cow k
149 there was an original state i (0 (not lame), 1 (lame)) the duration spent in that state was
150 categorized into discrete time intervals of 14 d, t_i (measured as $t = 1, 2, \dots, n$ with n being the
151 maximum duration of an episode) and an outcome event at the end of the discrete time
152 interval, y , with 0 = no change in state, and 1 = occurrence of a change in state. A logit link
153 function was used to express the ratio of probability of a change in state to probability of no
154 change in the state and took the form:

$$155 \text{logit} [\pi_{ik(t)}] = \beta_{0_i} + \alpha_i(t) + \beta x_{ik(t)} + u_k^{(i)}$$

156 where β_{0_i} is a state specific intercept, $\alpha_i(t)$ a set of dummy variables for the discrete time
157 interval t depicting duration of state, $\beta x_{ik(t)}$ covariates include a vector of explanatory
158 variables varying by time or cow with a dummy variable for original state. The model was
159 run in MlwiN 2.02 (Rasbash et al., 2005) using Markov chain Monte Carlo estimation. The
160 first 5,000 iterations were discarded and then 500,000 iterations until the chains were visually
161 stable.

162 RESULTS

163 Mobility was scored on 28 occasions, 312 cows (allowing for additions and removals)
164 were scored with 168 to 217 at each observation, the number of scores arranged from 5 to 28 /
165 cow. The percent of scores 0, 1, 2, and 3 were 23, 45, 27, and 5, respectively, with 1, 20 ,
166 48, and 31% of cows with maximum scores of 0, 1, 2, and 3, respectively. The mean number
167 of observations with MS 2 or 3 was 32%, ranging from 24% in October 2008 to 40% in July

168 2008. The mean duration of lameness was 5.5 [s.e. 3] wk (median 4 wk, interquartile range 2
169 to 7 wk). Only 48% of scores remained unchanged from 1 score to the next, but cows were
170 unlikely to move more than 1 score in a 2-wk period. Once cows were a certain MS for 2
171 observations they were more likely to remain at that MS than change score. Patterns of scores
172 are in Table 2.

173 The milk yield was highest in cows with MS 1 (Table 3). Cows produced 0.7 kg/d and
174 1.6 kg/d less milk when MS 2 or 3, respectively, compared with cows with MS 1 ($P <$
175 0.05). There was a reduction in yield from $t - 3$ before becoming MS 2 (0.47 CI (0.11 – 0.82)
176 or MS 3 (0.9 (0.15 – 1.65) and $t + 2$ after recovering from MS 2 (0.85 CI 0.5 – 1.2).

177 First, second and third lactation cows were 58, 48, and 19%, respectively, less active
178 (took fewer steps) than cows parity >3 ($P < 0.05$; Table 3). Cows were less active in early
179 lactation (mean log 1.38 steps/hr in month 1) and became more active as lactation progressed
180 (mean log 1.74 steps/hr in month 10), e.g., cows that were 9 months into lactation were 42%
181 more active than those in the first month of lactation ($P < 0.05$). Cows with MS 0 were 1%
182 less active than a cow with MS 1 ($P < 0.05$). Cows with MS 2 and 3 were 3 and 5 % less
183 active than a cow with MS 1 ($P < 0.05$). Cows had a decreased activity for 42 d before being
184 MS 2 (mean 0.02 (CI 0.01 – 0.03)): they were 3% less active 2 wk before and 2% less active
185 4 wk before they became MS 2 compared with a cow with MS 1 ($P < 0.05$). Cows with MS
186 3 were less active from 28 d before they developed MS 3 (-0.02 CI (0.00 – 0.04)). Similarly,
187 cows that were MS 2 were less active by 3 to 4 % for the following 5 recordings and cows
188 that had MS 3 were less active by 3 to 6 % for the following recordings ($P < 0.05$).

189 A total 444 lesions (185/100 cows per yr) with 385 primary lesions on 258 feet were
190 recorded by the herdsman, veterinarian, and foot trimmer. Over the 12 mo study period 178
191 cows (74%) were treated for at least 1 lesion; 72 (30%) cows had more than 1 foot with a
192 lesion and 81 (31%) feet were treated more than once. The lesions recorded were digital

193 dermatitis (39%) sole ulcer (25%), white line disease (**WLD**) (12%), interdigital growth (9%),
194 and other (15%).

195 From the multistate model (Table 4) the longer the period a cow was not lame (i.e.,
196 not MS 2 or 3) the less likely she was to make a transition to being lame and the longer a cow
197 was lame the less likely she was to recover from being lame. Cows < 90 DIM were less likely
198 to become lame than cows ≥ 90 DIM (Odds Ratio (**OR**) = 0.66) and cows with milk yield >
199 15 to ≤ 35 kg in the previous 16 d were less likely to recover from lameness (OR = 0.73) than
200 cows with milk yield > 35 kg.

201 Cows in parity 1 (OR = 0.49) or 2 (OR = 0.79) were less likely to become lame and
202 they were more likely to recover (OR = 1.26 and 1.32, respectively) once they had become
203 lame compared with cows of parity >2. Lame cows with 'other' lesions that were treated
204 were less likely to recover from being lame (OR = 0.58) than untreated lame cows. Cows
205 treated with a sole ulcer (OR = 1.35), digital dermatitis (OR = 1.51) or 'other' lesions (OR =
206 1.39) were more likely to become lame again in comparison with non lame cows that had not
207 been treated (Table 4).

208 **DISCUSSION**

209 In the current study, milk yield was reduced in cows with MS 2 or 3 for up to 4 to 8 wk
210 before their locomotion moved from MS 1. This period of time was considerably less than the
211 reduction in yield seen 3 to 4 mo before treatments reported by Green et al. (2002) and Amory
212 et al. (2008) and suggests that there was a delay in treatment in these 2 studies. If MS was used
213 to identify lame cattle and they were treated promptly the duration of both lameness and milk
214 loss might be reduced (Green et al., 2010). From the multistate model and patterns of MS
215 (Tables 4 and 2), treatment in the current study herd was not successful, with treated cattle
216 either not recovering (digital dermatitis) or being more likely to become lame again (sole ulcer
217 and other diseases). Note that WLD was not associated with lameness (Table 4) as in other

218 studies (Tadich et al., 2010). Repeated occurrences of lameness might indicate meager
219 treatment strategy or efficacy, but might also indicate that treatment cannot address intrinsic
220 factors such as a thin digital cushion. Treatment was added to the milk yield model; however,
221 it did not alter the associations between yield and MS and so was excluded.

222 That cows with MS 1 had a lower milk yield for 4 to 8 wk before there was a change in
223 mobility score from MS 1 to MS 2 or 3 suggests that the reduction in yield occurred before
224 lameness was detectable. One possible explanation for the reduction in yield before MS
225 changed is that MS was not sufficiently sensitive to detect the initial stages of disease. In
226 other studies of dairy cow lameness authors have reported lesions on sound cows (Manske et
227 al., 2002; Tadich et al., 2010; Bicalho et al., 2008). One hypothesis, drawing evidence from
228 Bicalho et al. (2009), is that lameness and foot lesions are positively associated with a thin
229 digital cushion which is associated with low body condition, this might cause sub clinical
230 disease that is not detectable externally or by MS, but is sufficiently painful to reduce food
231 intake, increase metabolic rate and so reduce milk yield. Low body condition *per se* could
232 also lead to reduced milk yield. It is unfortunate that we did not score the body condition of
233 the cattle in the current study but one could speculate that the cattle that moved from MS 1 to
234 MS 2 or 3 lost body condition before the transition whilst those that remained at MS 1 did not.

235 The fact that high yielding cattle at greater risk of lameness (Green et al., 2002;
236 Amory et al., 2008; Green et al., 2010) might help explain why cows with MS 1 produced
237 more milk than cows with scores 0, 2 or 3. These cows are producing high yields and their
238 locomotion is impaired (they are marginally lame). Over time, a proportion remain at MS 1
239 (Tables 2 and 5) and continue to produce high yields (Table 3) but some move to MS 2 or 3
240 and the pattern of lower yield and higher mobility score ensues. Once a cow is lame, she
241 might continue to have a further reduction in yield because extra energy is required to cope
242 with the pain of the foot lesion and energy is directed to this rather than milk production.

243 Depending on farm layout, lame cows might also feed less frequently and so reduce feed
244 intake, exacerbating the disease process. If this was so, then successful treatment might
245 increase mobility and stabilize milk yield, as seen in Green et al. (2010).

246 A large numbers of transitions in MS were seen between fortnightly scores for
247 individual cows in our study. In the UK farmers often MS their cattle annually or biannually
248 to comply with assurance scheme standards e.g. Tesco scheme, the current results suggest that
249 infrequent MS would give a snap shot of prevalence, but have little value in management of
250 lameness. Cows that had a MS of 2 or 3 had a high probability of remaining a 2 or a 3 (Table
251 2) and becoming lame again (Tables 2 and 4). The effects of this may be seen in terms of milk
252 production, but the effects on cow welfare are not so easy to quantify, although these cows did
253 have lower activity.. This suggests that prevalence, incidence, and repeat cases should be standard
254 recordings.

255 The results demonstrate that it is not only the MS on the day of recording that is
256 important, but that the length of time that a cow has been at a particular MS is highly
257 relevant. Our examples demonstrate that a cow that had been MS 2 for 6 wk lost 4.5 kg of
258 milk per d while at MS 3 lost 6 kg/d of milk . These results support Juarez et al. (2003) who
259 demonstrated a drop in milk yield of 4 kg/d for a lame cow. Extrapolating these results to a
260 cow that is lame for 12 wk equates to 610 kg milk lost, supporting Amory et al. (2008).

261 Results from this herd suggest that activity data may not play a useful role in early
262 identification of lameness because the absolute changes were so small: parity and stage
263 of lactation had a much greater effect on activity than MS (Table 3). Cows became steadily
264 more active as lactation progressed and with increasing parity, contrary to the findings of
265 O'Callaghan et al. (2003) who reported a decreased level of activity as lactation progressed.
266 The average change in activity associated with mobility score was less than 1%/d in our study,
267 while they reported that cows that were lame were 24% less active than non lame cows.

Comment [FCG1]: Over what period of time?

268 There might be large variations in activity between herds, this might depend on the farm
269 layout, and this might be very important when considering the necessary and unnecessary
270 activity of cows.

271 The results suggest that a decrease in milk yield could have a role as an early indicator of
272 lameness, while change in activity is a less sensitive measure. In order to be practically
273 applied on farms, algorithms for milk yield, correcting for parity and stage of lactation, would
274 need to be incorporated into on-farm software alongside daily milk recording. In conjunction
275 with fortnightly MS this could alert the farmer that cows need early intervention. Before this
276 could be achieved, research needs to be repeated across many farms and systems to validate
277 the findings. In addition, unexpected reduction in milk yield might indicate that a cow is not
278 metabolically stable (Bicalho et al., 2009) and lameness is only one of the risks for such
279 cattle.

280 The advantage of this study was the large amount of detailed data that were collected.
281 This farm was chosen because it was similar to many farms in the UK with Holsteins
282 producing large quantities of milk under intensive conditions; the patterns within cow are
283 useful additions to our understanding of the associations between milk yield, MS, activity, and
284 lameness. A disadvantage of this study was that the data were from only 1 farm. It is not
285 possible to generalize prevalence, incidence, and transitions between MS. Whatever the
286 factors initiating lameness it appears that changing external management (Barker et al.,
287 2007, 2009) is likely to be only part of the story to prevent lameness in dairy cows, possibly
288 explaining part of the limited success of intervention studies (Bell et al., 2007; Barker 2007).
289 Further work is required to elucidate when biochemical and pathological changes occur in
290 the development of lameness. If these changes can be identified, then we can move forward
291 in preventing lameness in dairy cows.

292 **CONCLUSIONS**

293 A reduction in mobility occurred 4 to 8 wk after cows had started to reduce milk
294 yield and an increase in milk yield occurred approximately 6 wk after a cow returned to
295 MS 0 or 1, suggesting that either mobility scoring is insufficiently sensitive to detect
296 lameness, that cattle mask lameness despite being diseased, or that a lameness and
297 reduction in yield are linked by a common intrinsic event. Once lame, cows were likely
298 to remain lame or become lame again, suggesting that either treatment was unsuccessful
299 or that the internal origin of lameness overrode treatment. Further work investigating
300 body condition, biochemical profiles, mobility, and lameness longitudinally could have a
301 huge impact on our understanding of the etiology of lameness.

302 **ACKNOWLEDGMENTS**

303 We thank the RCVS Trust for supporting this research and John Hembrow and
304 Chris Kiddle for their assistance with data collection.

305 **REFERENCES**

- 306 Amory, J. R., Barker, Z. E., Wright, J. L., Mason, S. A., Blowey, R. W., and Green, L. E.
307 2008. Associations between sole ulcer, white line disease and digital dermatitis and
308 the milk of 1824 dairy cows on 30 dairy cow farms in England and Wales. *Prev.*
309 *Vet. Med.* 83:381- 391.
- 310 Archer, S. C., Green, M. J., and Huxley, J. N. 2010. Association between milk yield and
311 serial locomotion score assessments in UK dairy cows. *J. Dairy Sci.* 93:4045-4053
- 312 Barker, Z. E. (2007) *Epidemiology of Lameness in Dairy cows*. PhD thesis. University of
313 Warwick.
- 314 Barker, Z. E., Amory, J. R., Wright, J. L., Mason, S. A., Blowey, R. W., and Green, L. E. 2007.
315 Management factors associated with impaired locomotion in dairy cows in England
316 and Wales. *J. Dairy Sci.* 90:3270–3277.
- 317 Barker Z. E., Amory J. R., Wright J. L., Mason S. A., Blowey R. W., and Green L. E. 2009.

318 Risk factors for increased rates of sole ulcers white line disease and digital dermatitis
319 in dairy cattle from twenty-seven farms in England and Wales. *J. Dairy Sci.* 92:1971-
320 1978.

321 Bell N. J., Bell M. J., Knowles T. G., Whay H. R., Main D. C. J., and Webster A. J. F. 2009.
322 The development implementation and testing of a lameness control programme based
323 on HACCP principles and designed for heifers on dairy farms. *Vet. J.* 180:178-188.

324 Bicalho, R. C., Warnick, L. D., and Guard, C. L. 2008. Strategies to analyze milk losses
325 caused by diseases with potential incidence throughout the lactation: a lameness example.
326 *J. Dairy Sci.* 91:2653-61

327 Bicalho, R. C., Machado, V. S., and Caixeta, L. S. 2009. Lameness in dairy cattle: A
328 debilitating disease or a disease of debilitated cattle? A cross-sectional study of
329 lameness prevalence and thickness of the digital cushion *J. Dairy Sci.* 92:3175–3184

330 Clarkson, M. J., Downham, D. Y., Faull, W. B., Hughes, J. W., Manson, F. J., Merritt, J.B.,
331 Murray, R. D., Russell, W. B., Sutherst, J. E., and Ward, W. R. 199 . Incidence and
332 prevalence of lameness in dairy cattle. *Vet. Rec.* 138:563-567.

333 Goldstein, H., 1995. *Multilevel Statistical Models* Ed ., London, Edward Arnold.

334 Green, L. E., Hedges, V. J. Schukken, Y. H., Blowey, R. W., and Packington, A. J. 2002.
335 The impact of clinical lameness on the milk yield of dairy cows. *J. Dairy Sci.*
336 85:2250-2256.

337 Green, L. E., Borkert, J., Monti, G., and Tadich, N. 2010. Associations between lesion-
338 specific lameness and the milk yield of 1635 dairy cows from seven herds in the Xth
339 region of Chile and implications for the management of dairy cows worldwide. *Anim.*
340 *Welfare* 19:419 – 427

341 Juarez, S. T., Robinson, P. H., DePeters, E. J., and Price, E. O. 2003. Impact of lameness on
342 behavior and productivity of lactating Holstein cows. *Appl. Anim. Behav. Sci.* 83:1-14.

343 King E. M., and Green, L. E. Assessment of farmer recognition and reporting of lameness in
344 lowland sheep flocks in England. *Anim. Welfare in press*

345 Leach, K. L., Whay, H. R., Maggs, C. M., Barker, Z. E., Paul, E. S., Bell, A. K., and Main D.
346 C. J. 2010. Working towards a reduction in cattle lameness: 1. Understanding barriers
347 to lameness control on dairy farms. *Res. Vet. Sci.* 89:311-317.

348 Manske, T., Hultgren, J., and Bergsten, C. 2002. Prevalence and interrelationships of hoof
349 lesions and lameness in Swedish dairy cows. *Prev. Vet. Med.* 54:247-263.

350 Mazrier, H., Tal, S., Aizinbud, E., and Bargai, U. 2006. A field investigation of the use of the
351 pedometer for the early detection of lameness in cattle. *Can Vet. J.* 47:883-886

352 O'Callaghan, K. A., Cripps, P. J., Downham, D. Y., and Murray, R. D. 2003. Subjective and
353 objective assessment of pain and discomfort due to lameness in dairy cattle. *Anim.*
354 *Welfare.* 12:605-610.

355 Rasbash J., Browne W. J., Healy M., Cameron B., and Charlton C. 2005. MLwiN Version 2.02.

356 Tadich, N., Flor, E., and Green, L. E. 2010 Associations between hoof lesions and
357 locomotion score in 1098 unsound dairy cows *Vet.J.*, 184:60 – 65

358 Walker, S. L., Smith, R.F., Routly, J. E., Jones, D. N., Morris, M. J., and Dobson, H. 2008.
359 Lameness activity time budgets and estrus expression in dairy cattle. *J. Dairy Sci.*
360 91:4552- 4559

361 Warnick, L. D., Janssen, D., Guard, C. L., and Grohn, Y. T. 2001. The effect of lameness on
362 milk production in dairy cows. *J. Dairy Sci.* 84:1988-1997

363 Whay, H. R., Main, D. C., Green, L. E., and Webster, A. J. 2003. Assessment of the welfare
364 of dairy cattle using animal-based measurements: Direct observations and
365 investigation of farm records. *Vet. Rec.* 153:197-202.

366 Whay, H. R., Waterman, A. E., and Webster, A. J. 1997. Associations between locomotion,
367 claw lesions and nociceptive threshold in dairy heifers during the peri-partum period.

368 Vet. J. 154:155-161.

369 Wilmink, J. B. M. 1987. Adjustment of test-day milk, fat and protein yields. *Livest. Prod. Sci.*

370 16:335-348.

371

372

373 Table 1. Definitions of mobility scores (Whay et al., 2003)

Mobility score	Definition	Description of cow mobility
0	Good mobility / sound	Walks with even weight bearing and rhythm on all 4 feet with a flat back. Long fluid strides possible.
1	Imperfect mobility	Steps unevenly or shortened strides. Affected limbs not immediately identifiable.
2	Impaired mobility	Uneven weight bearing on limb immediately identifiable and/or obviously shortened stride. Usually arched back.
3	Severely impaired mobility	Unable to walk as fast as brisk human pace plus signs of score 2.

374

375

376 Table 2. Transitions in mobility score from time $t - 3$ to time t where $t = 14$ d intervals
 377 illustrating that 50 – 60% of cows remain at a score for 8 weeks but that 40 – 50% cows move
 378 mobility score
 379

$t^1 - 3$	$t - 2$	$t - 1$	t	Probability of score at t	N^2 sequence observed
0	0	0	0	0.57	244
0	0	0	1	0.41	244
1	0	0	1	0.51	182
1	0	0	0	0.44	182
1	1	1	1	0.65	665
1	1	1	0	0.19	665
2	2	2	2	0.64	390
3	3	3	3	0.67	54
3	3	3	1	0.02	54
3	3	3	2	0.31	54
3	3	3	3	0.67	54
3	3	2	1	0.16	31
2	3	2	1	0.09	54
1	3	2	1	0.20	10

380 t^1 = time, $t +/- i$ = time from / to t in 2 wk intervals N^2 = number of occasions,
 381

382

383 Table 3. Random effects model of mean 16 d yield and 16 d mean log activity in 312 cows
 384 from 1 dairy herd in Somerset, UK

385

	Mean yield	lower 95% CI ³	upper 95% CI	mean Log activity	lower 95% CI	upper 95% CI
intercept	41.9	40.685	43.115	1.384	0.972	1.796
parity >3	reference	reference	reference	reference	reference	reference
ce						
parity 1	-5.78	-7.113	-4.447	0.113	0.072	0.154
parity 2	-2	-3.078	-0.922	0.039	0.006	0.072
parity 3	-2.4	-3.282	-1.518	0.072	0.047	0.097
2nd lactation	-0.7	-1.366	-0.034	0.237	0.217	0.257
DIM	-0.05	-0.052	-0.048			
Wilmink	-15.7	-17.013	-14.387			
month in milk 1				reference	reference	reference
month in milk 2				0.033	0.011	0.055
month in milk 3				0.065	0.043	0.087
month in milk 4				0.078	0.056	0.100
month in milk 5				0.113	0.089	0.137
month in milk 6				0.125	0.101	0.149
month in milk 7				0.157	0.132	0.182
month in milk 8				0.205	0.180	0.230
month in milk 9				0.244	0.217	0.271
month in milk 10				0.304	0.277	0.331
month in milk 11				0.361	0.330	0.392
at t ¹						
MS ² 1	reference	reference	reference	reference	reference	reference
ce						
MS 0	-0.45	-0.764	-0.136	-0.004	-0.014	0.006
MS 2	-0.66	-0.974	-0.346	-0.016	-0.026	-0.006
MS 3	-1.61	-2.237	-0.983	-0.025	-0.045	-0.005
at t+1						
MS 1	reference	reference	reference	reference	reference	reference
ce						
MS 0	-0.76	-1.093	-0.427	-0.007	-0.017	0.003
MS 2	-0.43	-0.763	-0.097	-0.012	-0.022	-0.002
MS 3	-0.5	-1.147	0.147	-0.011	-0.031	0.009
at t+2						
MS 1	reference	reference	reference	reference	reference	reference
ce						
MS 0	-0.85	-1.203	-0.497	-0.005	-0.015	0.005

386

387

388 Table 3. Two level random effects model of mean 16 d milk yield and log activity in 312

389 cows from one herd in Somerset, UK *continued*

390

	Mean yield	lower 95% CI	upper 95% CI	mean Log activity	lower 95% CI	upper 95% CI
MS 2	-0.42	-0.753	-0.087	-0.002	-0.012	0.008
MS 3	0.26	-0.387	0.907	0.002	-0.018	0.022
at t+3						
MS 1	reference	reference	reference	reference	reference	reference
MS 0	-0.84	-1.212	-0.468	0.001	-0.011	0.013
MS 2	-0.26	-0.613	0.093	0.007	-0.003	0.017
MS 3	0.47	-0.196	1.136	0.009	-0.011	0.029
at t+4						
MS 1	reference	reference	reference			
MS 0	-0.65	-1.022	-0.278			
MS 2	-0.1	-0.453	0.253			
MS 3	0.28	-0.406	0.966			
at t-1		0.000	0.000			
MS 1	reference	reference	reference	reference	reference	reference
MS 0	-0.4	-0.733	-0.067	-0.005	-0.015	0.005
MS 2	-0.95	-1.283	-0.617	-0.015	-0.025	-0.005
MS 3	-2.67	-3.336	-2.004	-0.031	-0.051	-0.011
at t-2						
MS 1	reference	reference	reference	reference	reference	reference
MS 0	-0.44	-0.773	-0.107	-0.010	-0.020	0.000
MS 2	-0.69	-1.043	-0.337	-0.170	-0.180	-0.160
MS 3	-1.39	-2.096	-0.684	-0.019	-0.041	0.003
at t-3						
MS 1	reference	reference	reference	reference	reference	reference
MS 0	-0.25	-0.603	0.103	-0.013	-0.023	-0.003
MS 2	-0.47	-0.823	-0.117	-0.015	-0.025	-0.005
MS 3	-0.9	-1.645	-0.155	0.010	-0.225	0.245
at t-4						
MS 1	reference	reference	reference			
MS 0	0.09	-0.282	0.462			
MS 2	-0.41	-0.782	-0.038			
MS 3	0.31	-0.474	1.094			

391 ¹t = time, t +/- i = time from / to t in 2-wk intervals392 ²MS = mobility score

393 ³CI = confidence interval

394 Table 4: Multivariable multistate model of transitions between lame (mobility score 2 or 3)
 395 and non lame (mobility score 0 or 1) states in 312 cows from 1 dairy herd observed for 1 yr in
 396 Somerset, UK

397

variables	Transition			
	Non lame to lame		Lame to non lame	
intercept	-5.15	0.21	-4.58	0.37
	OR	CI	OR	CI
Duration spent in state				
≤ 2 wk	4.06	2.96-5.55	3.63	1.90-6.94
> 2-4wk	3.16	2.22-4.49	2.51	1.29-4.89
> 4-18 wk	1.80	1.32-2.47	1.93	1.01-3.69
> 18 wk	reference		reference	
DIM				
0-90	0.66	0.57-0.78	1.25	0.93-1.67
91-180	1.00	0.79-1.26	1.15	0.91-1.46
>180	reference		reference	
Past treatments				
Sole ulcer				
yes	1.35	1.11-1.64	0.84	0.69-1.02
no	reference		reference	
Digital dermatitis				
yes	1.51	1.29-1.76	0.86	0.72-1.03
no	reference		reference	
White line disease				
yes	1.15	0.91-1.46	0.83	0.65-1.05
no	reference		reference	
Other				
yes	1.39	1.10-1.76	0.58	0.45-0.74
no	reference		reference	
Pregnant				
yes	0.87	0.70-1.08	1.67	1.34-2.07
no	reference		reference	
Mean milk yield in previous 16 d				
missing	0.90	0.46-1.80	1.16	0.50-2.70
≤15	1.22	0.84-1.77	0.81	0.54-1.22
>15-35	1.15	0.89-1.48	0.73	0.55-0.98
>35	reference		reference	

Parity				
1	0.49	0.39-0.62	1.26	1.00-1.59
2	0.79	0.63-0.98	1.32	1.05-1.67
3	0.94	0.74-1.19	1.15	0.89-1.48
>3	reference		reference	

398