

Original citation:

Ronconi, Luca, Pizarro, Ana M., McQuitty, Ruth J. and Sadler, P. J.. (2011) Insights into the acid-base properties of PtIV-diazidodiam(m)inedihydroxido complexes from multinuclear NMR spectroscopy. Chemistry - A European Journal, Volume 17 (Number 43). pp. 12051-12058. ISSN 0947-6539

Permanent WRAP url:

<http://wrap.warwick.ac.uk/40374/>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 3.0 Unported (CC BY 3.0) license and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/3.0/>

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

warwick**publications**wrap

highlight your research

<http://go.warwick.ac.uk/lib-publications>

Insights into the Acid–Base Properties of Pt^{IV}–Diaزيدodiam(m)inedihydroxido Complexes from Multinuclear NMR Spectroscopy

Luca Ronconi,^[a] Ana M. Pizarro,^[b] Ruth J. McQuitty,^[b] and Peter J. Sadler*^[b]

Abstract: Platinum(IV) am(m)ine complexes are of interest as potential anticancer pro-drugs, but there are few reports of their acid–base properties. We have studied the acid–base properties of three photoactivatable anticancer platinum(IV)–diaزيدodiam(m)ine complexes (*cis,trans,cis*-[Pt^{IV}(N₃)₂(OH)₂(NH₃)₂], *trans,trans,trans*-[Pt^{IV}(N₃)₂(OH)₂(NH₃)₂], and *cis,trans*-[Pt^{IV}(N₃)₂(OH)₂(en)]) using multinuclear NMR methods and potentiometry. In particular, the combination of

both direct and indirect techniques for the detection of ¹⁵N signals has allowed changes of the chemical shifts to be followed over the pH range 1–11; complementary ¹⁴N NMR studies have been also carried out. A distinct pK_a value of approximately 3.4 was determined

for all the investigated complexes, involving protonation/deprotonation reactions of one of the axial hydroxido groups, whereas a second pH-dependent change for the three complexes at approximately pH 7.5 appears not to be associated with a loss of an am(m)ine or hydroxido proton from the complex. Our findings are discussed in comparison with the limited data available in the literature on related complexes.

Keywords: acid–base properties · anticancer complexes · bioinorganic chemistry · NMR spectroscopy · pK_a values · platinum

Introduction

The platinum(II) complexes cisplatin, carboplatin, and oxaliplatin are currently approved drugs for the treatment of cancer. However, in spite of therapeutic success in the treatment of several types of tumors, their effectiveness is severely hindered by adverse side effects such as nausea, ototoxicity, neurotoxicity, myelosuppression, and nephrotoxicity. A second major drawback is tumor resistance, either acquired during cycles of therapy (as occurs in patients with, for instance, ovarian cancer) or intrinsic resistance (such as in patients with prostate, lung, or breast cancer).^[1] Thus, effort in designing new platinum drugs is aimed at making platinum-based therapy safer to patients, in particular by reducing or removing severe side effects, providing oral administration, and overcoming both intrinsic and acquired resistance.

In this regard, there has long been interest in cytotoxic six-coordinate platinum(IV) analogues, especially since they are often effective against tumors resistant to cisplatin. Plat-

inum(IV) is a classically-inert low-spin d⁶ metal ion. Platinum(IV)-based anticancer drugs might be less toxic and more readily tolerated by normal cells. Increasing evidence suggests that for activity they require in vivo reduction to the corresponding platinum(II) species by biological reductants (e.g. ascorbic acid, intracellular glutathione). Such chemical reduction is highly dependent on the levels of reducing agents present in body fluids and cells, which can be variable. Platinum(IV) complexes may be regarded as inactive pro-drugs, and the coordinative binding of the resulting active platinum(II) species to DNA is widely considered to be responsible for their therapeutic activity.^[2] Several platinum(IV) drugs, including tetraplatin ([Pt^{IV}Cl₄(1,2-diaminohexane)]), satraplatin (*cis,trans,cis*-[Pt^{IV}Cl₂(acetate)₂(NH₃)(cyclohexylamine)]), and iproplatin (*cis,trans,cis*-[Pt^{IV}Cl₂(OH)₂(isopropylamine)₂]), have reached clinical trials and some positive responses have been reported in the early phases.^[3]

Our current interest is in platinum(IV)–diaزيدodiam(m)inedihydroxido complexes, pro-drugs which are relatively inert toward reducing agents under physiological conditions. These non-toxic precursors might be selectively activated by light at the tumor site leading to the corresponding platinum(II) species, thereby potentially improving targeting and minimizing side effects.^[4,5] Their behavior upon irradiation has been recently investigated by both experimental^[6–8] and theoretical^[9] studies, providing insights into the photoreaction pathways that are involved. The photoreaction pathways and types of photoproducts depend on the pH of the solution and other reaction conditions.

It is often assumed that neutral platinum(IV) ammine derivatives remain as neutral complexes in aqueous solution although am(m)ines coordinated to highly charged metals

[a] Dr. L. Ronconi*

School of Chemistry, University of Edinburgh
West Mains Road, Edinburgh EH9 3JJ (UK)

[b] Dr. A. M. Pizarro, R. J. McQuitty, Prof. Dr. P. J. Sadler

Department of Chemistry, University of Warwick
Gibbet Hill Road, Coventry CV4 7AL (UK)
Fax: (+44) 24-76523819
E-mail: P.J.Sadler@warwick.ac.uk

[*] Current address: Department of Chemical Sciences, University of Padova, Via Marzolo 1, Padova 35131 (Italy)

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/chem.201002792>.


Re-use of this article is permitted in accordance with the Terms and Conditions set out at [http://onlinelibrary.wiley.com/journal/10.1002/\(ISSN\)1521-3765/homepage/2111_onlineopen.html](http://onlinelibrary.wiley.com/journal/10.1002/(ISSN)1521-3765/homepage/2111_onlineopen.html).

ions such as platinum(IV), or the isoelectronic gold(III), can be acidic with pK_a values close to 7, that is, within the physiological pH range.^[10] The presence of an anionic $-NH_2^-$ ligand as opposed to a neutral $-NH_3$ ligand, for example, might be expected to have a major effect on cell uptake and distribution of platinum(IV) drugs, including binding to negatively charged DNA, and a detailed knowledge of the acid–base properties of platinum(IV) drugs is therefore important. Despite this, there appear to have been only a few reported studies.

To understand better the solution behavior of these platinum(IV)-diazidodiam(m)inedihydroxido pro-drugs and, consequently, the implications for their possible mechanism of action, we have made a detailed study of the acid–base properties of *cis,trans,cis*-[Pt^{IV}(N₃)₂(OH)₂(NH₃)₂] (**1**), *trans,trans,trans*-[Pt^{IV}(N₃)₂(OH)₂(NH₃)₂] (**2**), and *cis,trans*-[Pt^{IV}(N₃)₂(OH)₂(en)] (en = ethylenediamine) (**3**) using multinuclear NMR methods (combined with ¹⁵N-labeling of the complexes) and potentiometry. We discuss our findings in comparison with the limited literature data on related platinum(IV) complexes.

Results

NMR studies: The acid–base properties of complexes **1–3** were investigated by 1D ¹H, ¹⁵N, and ¹⁴N NMR spectroscopy. For the ¹⁵N NMR spectroscopy, ¹⁵N-labeling of the am(m)ine ligands was essential. The corresponding ¹⁵N-labeled am(m)ine analogues are referred to as ¹⁵N-**1**, ¹⁵N-**2**, and ¹⁵N-**3**.



The pH dependence of the ¹H NMR chemical shifts of the ammine protons of complex ¹⁵N-**1** over the pH range 1.0–6.5 is shown in Figure 1A. No proton signal was detected above pH 6.5 for coordinated ¹⁵NH₃ due to the rapid exchange of ammine protons with the solvent. By fitting the experimental data to Equation (1) (see the Experimental Section), a value of $pK_{a1} = 3.44 \pm 0.01$ was determined.

1D ¹⁵N{¹H} INEPT (Insensitive Nuclei Enhanced by Polarization Transfer) NMR experiments were also carried out over the same pH range but, again, no ¹⁵N peak was observable for ammine ligands above pH 6.5. On the other hand,




Figure 1. pH dependence of NMR chemical shifts for Pt^{IV}-¹⁵NH₃ of complex ¹⁵N-**1**. A) ¹H ($\chi^2 = 0.00002$; $R^2 = 0.99908$; $pK_{a1} = 3.44 \pm 0.01$), and B) ¹⁵N ($\chi^2 = 0.00082$; $R^2 = 0.99920$; $pK_{a2} = 3.34 \pm 0.01$; $pK_{a2} = 7.44 \pm 0.23$).

acquisition of a succession of 1D ¹⁵N{¹H} NMR spectra enabled the direct detection of the ¹⁵N-ammine signals over the entire range, from pH 1 to 11. Combined data for the pH dependence of the ¹⁵N chemical shifts of ammine ligands for complex ¹⁵N-**1** (Figure 1B) were fitted to Equation (2) (see the Experimental Section), giving now two pK_a values $pK_{a1} = 3.34 \pm 0.01$ (in good agreement with that derived above from ¹H NMR data) and $pK_{a2} = 7.44 \pm 0.23$, respectively. The latter is associated with only a small ¹⁵N shift change (ca. 0.1 ppm) and was not apparent from the ¹H NMR data.

Similar NMR experiments were carried out for complex ¹⁵N-**2** (Figure 2). The ¹H data gave rise to $pK_{a1} = 3.46 \pm 0.01$. As a consequence of the fast exchange of ammine protons, no 1D ¹⁵N{¹H} INEPT NMR signals were detected above pH 7.5. Again, ¹⁵N signals of the ammine ligands were recorded over the pH range 1–11 by direct detection 1D ¹⁵N{¹H} NMR experiments. The overall ¹⁵N NMR data for complex ¹⁵N-**2** (Figure 2B) gave rise to $pK_{a1} = 3.42 \pm 0.01$, and an apparent $pK_{a2} = 8.06 \pm 0.04$, although again this pH-induced change in the chemical shift was observable only by a small change in ¹⁵N chemical shift and not for ¹H shifts.

To confirm the observations by direct detection 1D ¹⁵N{¹H} NMR experiments, 1D ¹⁴N{¹H} NMR spectra were recorded at pH 1, 5 and 10 for complex **1** (Figure 3). ¹⁴N is high in natural abundance (99.6%) but quadrupolar ($I = 1$), and therefore resonances tend to be relatively broad, especially for the coordinated nitrogen of azide. Four ¹⁴N resonances were detected at approximately $\delta = 230, 160, 35,$ and -40 ppm (relative to NH₄Cl through N₂ at $\delta = 287.5$ ppm as internal reference). In line with ¹⁵N experiments, the chemical shifts at pH 1 and 5 differ by several ppm (up to 6.32 for Pt^{IV}-NNN, the terminal azido nitrogen, peak *x*), whereas little change was observed between pH 5 and 10. The Pt^{IV}-NH₃ ¹⁴N resonance shifts by 2.15 ppm from pH 1 to pH 5, but by only 0.06 ppm from pH 5 to 10 (upfield), similar to the shift changes for ¹⁵N resonances (2.23 and 0.10 ppm, respectively; see Table S1 in the Supporting Information). Analogous shift changes were observed in the 1D ¹⁴N{¹H} NMR spectra of complex **2** recorded under similar experimental conditions (see Figure S1 in the Supporting Information).




Figure 2. pH dependence of NMR chemical shifts for Pt^{IV}-¹⁵NH₃ of complex ¹⁵N-2. A) ¹H ($\chi^2=0.00001$; $R^2=0.99963$; $pK_{a1}=3.46\pm 0.01$), and B) ¹⁵N ($\chi^2=0.00003$; $R^2=0.99988$; $pK_{a1}=3.42\pm 0.01$; $pK_{a2}=8.06\pm 0.04$).




Figure 3. 1D ¹⁴N{¹H} NMR spectra of complex **1** in D₂O in the dark at 298 K at A) pH 1, B) pH 5, and C) pH 10. Assignments (internal reference N₂ at 287.5 ppm): coordinated azido ligand as labeled in structure ($w = \text{Pt}^{\text{IV}}\text{-NAN}$, $x = \text{Pt}^{\text{IV}}\text{-NNN}$, $y = \text{Pt}^{\text{IV}}\text{-N(NN)}$); ammine ligand ($z = \text{Pt}^{\text{IV}}\text{-NH}_3$). Inserts of signals w and z show their dependence on the pH.

To evaluate the effect of different amine ligands, the acid–base properties of the ethylenediamine (en) derivative *cis,trans*-[Pt^{IV}(N₃)₂(OH)₂(¹⁵N₂-en)] (¹⁵N-**3**) were also investigated. The ¹H NMR chemical shift changes for the ethylenediamine -CH₂- groups over the pH range 1.5–10 (Figure 4A) were fitted to Equation (2), giving $pK_{a1}=3.38\pm 0.03$ and $pK_{a2}=6.99\pm 0.61$. The large error in pK_{a2} is due to the extremely small upfield shift of the -CH₂- peak for this step. The dependence of ¹H (Figure 4B) and ¹⁵N (Figure 4C) NMR chemical shifts of the Pt^{IV}-¹⁵NH₂- group on pH were also determined. Direct 1D ¹⁵N{¹H} NMR detection was successfully used over the entire pH range, whereas 1D ¹H and ¹⁵N{¹H} INEPT techniques were used only below pH 6.5,

due to the fast exchange of the amine protons. From combined ¹⁵N data (Figure 4C) a pK_{a1} value of 3.45 ± 0.31 was obtained, in good agreement with the value of 3.32 ± 0.01 obtained by fitting ¹H NMR amine proton chemical shifts (Figure 4B). From ¹⁵N data, a pK_{a2} of 7.00 ± 0.50 (Figure 4C), was also determined.

HPLC studies: RP-HPLC was used to determine the purity of compounds **1–3** using a near neutral (pH 6) aqueous mobile phase. Compounds **1** and **3** each gave rise to a single major peak under the experimental conditions (see Figure S2 in the Supporting Information). For complex **2**, the major peak with a retention time of approximately 1.6 min was accompanied by a minor peak at approximately 1.7 min, which accounted for about 16% of the total peak area. The minor peak (see Figure S2A in the Supporting Information) seemed to increase in concentration over time and it was not observed when the chromatographic separation started less than 5 min after the compound was dissolved (see Figure S2B in the Supporting Information). Despite working under conditions of minimal light, this species can be attributed to a photoproduct, possibly the *cis*-isomer,^[11] due to the high photosensitivity of this compound.

Additional peaks were observed for complexes **1** and **3** at retention times 5.2 and 4.7 min, respectively, when a more acidic mobile phase was used (0.1% TFA water/0.1% TFA acetonitrile), using the same mobile phase gradient (data not shown). These impurities account for 0.4% for compound **1**, and 0.3% for compound **3**, and were therefore not characterized. However, they can be tentatively assigned to photoreduction products.^[12]

ESI-MS studies: The purity of the investigated platinum(IV)-diazidodiam(m)inedihydroxido complexes was also studied by ESI-MS. As an example, the mass spectra for compound **1** under acidic conditions (pH 4.9) in positive-




Figure 4. pH dependence of NMR chemical shifts of complex ^{15}N -**3**. A) ^1H chemical shift of $-\text{CH}_2-$ ($\chi^2 = 0.00002$; $R^2 = 0.99509$; $\text{p}K_{\text{a}1} = 3.38 \pm 0.03$; $\text{p}K_{\text{a}2} = 6.99 \pm 0.61$), B) ^1H chemical shift of $\text{Pt}^{\text{IV}}\text{-}^{15}\text{NH}_2-$ ($\chi^2 = 0.00002$; $R^2 = 0.99899$; $\text{p}K_{\text{a}1} = 3.32 \pm 0.01$), and C) ^{15}N chemical shift of $\text{Pt}^{\text{IV}}\text{-}^{15}\text{NH}_2-$ ($\chi^2 = 0.00522$; $R^2 = 0.99951$; $\text{p}K_{\text{a}1} = 3.45 \pm 0.31$; $\text{p}K_{\text{a}2} = 7.00 \pm 0.50$).

ion mode and under basic aqueous solution (pH 9.3) in negative-ion mode are shown in Figure 5 A and Figure 5 B, respectively.

The major ESI-MS peak for the observed positive ions at low cone voltage (+15 V) appears at m/z 386.5, consistent with the species $[\mathbf{1} + \text{K}]^+$ (calcd m/z 386.3). An additional peak of lower intensity was detected at m/z 370.2 assignable to the species $[\mathbf{1} + \text{Na}]^+$ (calcd m/z 370.2). ESI-MS spectra obtained in negative-ion mode (cone voltage: -15 V) in basic solution (pH 9.3) show a peak at m/z 346.4, consistent with the species $[\mathbf{1} - \text{H}]^-$ (calcd m/z 346.2). Similar experiments were performed for the *trans*-isomer **2** under the same experimental conditions. Again, the major ESI-MS peaks for the observed positive ions in acidic aqueous solu-




Figure 5. ESI mass spectra of aqueous solutions containing $0.8\text{--}1\text{ mg mL}^{-1}$ of complex **1** in A) positive-ion mode at pH 4.9 and B) negative-ion mode at pH 9.3.

tion (pH 4.7) are consistent with $[\mathbf{2} + \text{Na}]^+$ and $[\mathbf{2} + \text{K}]^+$, whereas negative ion peaks can be assigned to $[\mathbf{2} - \text{H}]^-$ (data not shown), similar to the results obtained for the *cis*-isomer **1**.

Potentiometric titrations: The $\text{p}K_{\text{a}}$ values for complex **1** were determined by potentiometric pH titrations in aqueous solution, and only one $\text{p}K_{\text{a}}$ value at 3.56 ± 0.03 could be determined. No dissociation of protons from the complex at pH values close to 7 was observed.

Discussion

NMR titration methods are widely used to determine proton dissociation constants (K_{a}). Proton dissociation from an ionizable group usually results in shifts of the NMR peaks from nearby nuclei in a molecule. Such shifts occur when the rate of proton ex-

change is ‘fast’ on the NMR time scale and represents the weighted average of the chemical shifts of the protonated and deprotonated forms. pH titration curves are then analyzed by curve-fitting approaches (e.g. Henderson–Hasselbalch equation-based algorithms), allowing $\text{p}K_{\text{a}}$ determinations. A major advantage of this method over other techniques is that the results are not greatly affected by the presence of minor impurities and the shift changes often allow the protonation site to be identified unambiguously.^[13]

NMR titrations are based on the evaluation of the changes of chemical shifts of $I = 1/2$ nuclei such as ^1H , ^{31}P , and, usually upon isotopic enrichment, for example with ^{13}C or ^{15}N . In this regard, ^{15}N NMR spectroscopy has been commonly employed in the study of platinum-based anticancer drugs as ^{15}N -labeled derivatives can often be prepared from readily available starting materials. ^{15}N NMR chemical shifts are sensitive to the nature of the *trans* ligand in platinum-am(m)ine derivatives and provide a powerful method for identifying the ligands in the coordination spheres of both platinum(II) and platinum(IV) complexes.^[17] In particular, 2D [^1H , ^{15}N] HSQC (Heteronuclear Single Quantum Correlation) NMR spectroscopy has been exploited to study hydrolysis profiles and $\text{p}K_{\text{a}}$ values of bound aqua ligands in aqua adducts of platinum am(m)ine compounds.^[14–17] However, there are only a few reports in the literature concerning the acid–base properties of non-aqua ligands bound to platinum and other metal complexes (e.g. am(m)ine protons).^[10,18,19] Coordination of the nitrogen atom of

am(m)ines to a highly charged metal ion can result in a dramatic increase in the acidity of am(m)ine protons so that their pK_a values can be determined in aqueous solution.^[20]

Direct detection is the most common technique for the observation of 1D ¹⁵N NMR spectra of all types of nitrogen atoms. Unfortunately, ¹⁵N suffers from low natural abundance (0.365%), although this can be overcome by ¹⁵N-enrichment, a low and negative magnetogyric ratio ($\gamma = -2.712 \times 10^7 \text{ rad T}^{-1} \text{ s}^{-1}$), low relative sensitivity (0.022 relative to ¹³C), and often also from long relaxation times.^[21] The indirect INEPT method improves the sensitivity of NMR experiments with low-abundance and low magnetogyric ratio nuclei. Specifically, the net effect is the non-selective polarization transfer from protons to ¹⁵N nuclei with an appropriate ¹H–¹⁵N coupling. This polarization transfer method enhances signal intensity by transferring the greater population differences of high- γ spins (¹H) onto their scalar spin-coupled low- γ partners (¹⁵N), leading to the amplification of the heteronucleus signal, increased sensitivity and shorter accumulation times.^[22]

On the basis of these considerations, we assumed that when a ¹⁵N NMR signal is detected by both 1D ¹⁵N{¹H} INEPT and direct detection 1D ¹⁵N{¹H} NMR techniques the exchange of the ammine protons (interpreted in terms of base-catalyzed hydrogen exchange of coordinated am(m)ines in aqueous solution) is ‘slow’ on the NMR time scale. On the contrary, when it is recordable only through direct-detection 1D ¹⁵N{¹H} NMR experiments, proton exchange is likely to be more rapid.

The use of both direct and indirect detection of ¹⁵N NMR parameters allowed detection of two distinct pH-dependent structural changes for all three platinum(IV) diazido complexes studied here. The first, with an associated pK_{a1} of approximately 3.4, can be assigned to the deprotonation of one of the axial hydroxide groups. By fitting the ¹⁵N chemical shifts of Pt^{IV}–¹⁵N–am(m)ine at different pH to Equation (2) (see the Experimental Section) pK_{a1} values of 3.34 ± 0.01 (Figure 1B), 3.42 ± 0.01 (Figure 2B), and 3.45 ± 0.31 (Figure 4C) have been determined for ¹⁵N-**1**, ¹⁵N-**2**, and ¹⁵N-**3**, respectively (in good agreement with the values 3.44 ± 0.01 (Figure 1A), 3.46 ± 0.01 (Figure 2A) and $3.38 \pm 0.03/3.32 \pm 0.01$ (Figure 4A and Figure 4B) obtained by fitting the related ¹H NMR data to Equation (1) (see the Experimental Section).

For all the complexes, pK_{a1} values associated with the protonation/deprotonation reactions of one of the axial hydroxide groups are very similar. Remarkably, the replacement of the two *cis*-ammine ligands with a chelating ethylenediamine has little effect on the acidity of the axial aqua/hydroxide groups. The pK_{a1} values calculated for complexes **1–3** are consistent with the few data available in the literature.^[23–26] Reported Pt–OH₂/Pt–OH pK_a values, obtained from potentiometric measurements, for platinum(IV) diam(m)ine aqua derivatives of the type *cis,trans,cis*-[Pt^{IV}(X)₂(OH)(OH₂)(NH₃)₂]⁺ (X = Cl, Br, I, CN), *trans,trans,trans*-[Pt^{IV}(X)₂(OH)(OH₂)(NH₃)₂]⁺ (X = Cl, Br, I), and *cis,trans*-[Pt^{IV}(X)₂(OH)(OH₂)(en)]⁺ (X = Cl, Br, I) are shown




Figure 6. pK_a values for an axial aqua ligand in the platinum(IV) diam(m)ine complexes: (●) *cis,trans,cis*-[Pt^{IV}(X)₂(OH)(OH₂)(NH₃)₂]⁺ (X = Cl, Br, I, CN, N₃), (▲) *trans,trans,trans*-[Pt^{IV}(X)₂(OH)(OH₂)(NH₃)₂]⁺ (X = Cl, Br, I, N₃), and (◆) *cis,trans*-[Pt^{IV}(X)₂(OH)(OH₂)(en)]⁺ (X = Cl, Br, I, N₃).^[23–26]

in Figure 6, and compared with the values obtained here for complexes **1–3**. It is notable that these values show a trend in accordance with the electron-withdrawing power of the X ligands: CN > N₃ > Cl > Br > I. Thus, the greater the electron-withdrawing power of the equatorial ligands, the lower the Pt^{IV}–OH₂/Pt^{IV}–OH pK_a value.

The second pH-dependent event, occurring in the pH range 7–8, requires a separate discussion and interpretation. For the three complexes ¹⁵N-**1**, ¹⁵N-**2**, and ¹⁵N-**3**, ‘ pK_{a2} ’ values of 7.44 ± 0.23 (Figure 1B), 8.06 ± 0.04 (Figure 2B) and 7.00 ± 0.50 (Figure 4C) have been determined. At first glance, these values might be associated with acid–base reactions involving the equatorial am(m)ine ligands. However, if a pH-induced deprotonation of a coordinated am(m)ine were to occur, a larger shift of the ¹⁵N NMR resonance of the am(m)ine ligand would be expected, as well as the ¹H NMR signals of the –CH₂– ethylenediamine in complex **3**. Again, contrary to the hypothesis of am(m)ine deprotonation is the fact that the ‘ pK_{a2} ’ determined for compound **2** is about 0.6 units greater than that of its *cis*-isomer **1**, whereas the higher *trans* influence of the azido ligand compared to ammonia,^[27] would be expected to give rise to a higher pK_{a2} for an ammine *trans* to an azido ligand.^[18]

¹⁴N NMR has proved useful for the study of platinum(IV) diazido complexes,^[6–8] especially if pulse sequences are used which minimize acoustic ringing so that very broad peaks can be observed. In this regard, results obtained by direct detection 1D ¹⁵N{¹H} NMR studies have been confirmed by 1D ¹⁴N{¹H} NMR experiments carried out for compounds **1** and **2** at pH 1, 5 and 9/10 (Figure 3 and Figure S1 in the Supporting Information); changes of the ¹⁴N chemical shifts for the ammine ligands are consistent with those observed in ¹⁵N spectra recorded at the same pH values (see Tables S1 and S2 in the Supporting Information). In addition to the small chemical shift changes observed for the ammine nitrogens between pH 5 and 10, the greater shifting of the ¹⁴N signals of the azido ligands supports the idea of a pH-dependent change over that pH range.

Only a few pK_a values for coordinated am(m)ines have been reported in the literature (Table 1), and none of them

Table 1. pK_a values reported for coordinated am(m)ines in mixed-ligand platinum(IV) am(m)ine complexes in aqueous solution at 298 K.

Platinum(IV) complexes	pK_a
<i>trans</i> -[Pt ^{IV} (CN) ₄ (NH ₃) ₂]	12.1 ^[10]
<i>trans</i> -[Pt ^{IV} Cl(CN) ₃ (NH ₃) ₂]	12.7 ^[10]
<i>trans</i> -[Pt ^{IV} Br(CN) ₃ (NH ₃) ₂]	12.8 ^[10]
<i>trans</i> -[Pt ^{IV} I(CN) ₃ (NH ₃) ₂]	13.0 ^[10]
<i>mer</i> -[Pt ^{IV} Cl ₃ (NH ₃) ₃] ⁺	11.1 ^[10,18]
<i>trans</i> -[Pt ^{IV} Cl ₂ (NH ₃) ₄] ²⁺	11.3 ^[10,18,28]
<i>cis</i> -[Pt ^{IV} Cl ₂ (NH ₃) ₄] ²⁺	9.5 ^[10,18,28]
<i>trans</i> -[Pt ^{IV} Cl ₂ (en) ₂] ²⁺	10.3 ^[18]
<i>trans</i> -[Pt ^{IV} Br ₂ (en) ₂] ²⁺	9.3 ^[18]
<i>trans</i> -[Pt ^{IV} I ₂ (en) ₂] ²⁺	8.0 ^[18]
<i>trans,cis</i> -[Pt ^{IV} Cl ₂ (NH ₃) ₂ (en)] ²⁺	10.2 ^{[18][a]}
<i>trans,cis</i> -[Pt ^{IV} Br ₂ (NH ₃) ₂ (en)] ²⁺	9.9 ^{[18][a]}
<i>trans,cis</i> -[Pt ^{IV} I ₂ (NH ₃) ₂ (en)] ²⁺	6.8 ^{[18][a]}
[Pt ^{IV} (OH)(NH ₃) ₅] ³⁺	9.5 ^{[18,24,29][b]}
[Pt ^{IV} Cl(NH ₃) ₅] ³⁺	8.4 ^{[10,18,25,28][b]}
[Pt ^{IV} Br(NH ₃) ₅] ³⁺	8.3 ^{[10,18,25][b]}
[Pt ^{IV} Cl(NH ₃) ₃ (en)] ³⁺	7.7/8.9 ^{[30][c]}
[Pt ^{IV} (NH ₃) ₆] ⁴⁺	7.7 ^[10,18,28]
[Pt ^{IV} (en) ₃] ⁴⁺	5.5 ^[18,28]

[a] Not specified whether ammine or ethylenediamine protons are involved. [b] Not specified whether *cis* or *trans* (to either hydroxide or halide) ammine ligands are involved. [c] Values refer to two unspecified isomers.

refers to complexes similar to those investigated here. Comparing the available data with those for compounds **1–3**, only complexes of the type [Pt^{IV}(Am)_{*n*}]⁴⁺ (Am = NH₃ (*n* = 6) or en (*n* = 3)) show lower or, at least, comparable pK_a values to those associated with the second inflexion point. However, this is attributed to the fact that these complexes carry a high positive charge. Compared to the neutral platinum(IV) am(m)ine derivatives in Table 1, the ' pK_{a2} ' value for the am(m)ine protons in the diazido complexes should be dramatically more basic. For example, if we compare complex **2** with complex *trans*-[Pt^{IV}(CN)₄(NH₃)₂], due to the lower electron-withdrawing properties of the azido and hydroxido ligands in comparison with the cyanido ligand, it would be predicted that the pK_a of the ammine ligands in **2** is greater than 12.1, and not consistent with a value close to 7.5.

Taken together, these observations suggest that the ' pK_{a2} ' values of approximately 7.5 calculated for complexes **1–3** at the second inflexion points are unlikely to be associated with the deprotonation of an equatorial am(m)ine ligand. This is consistent with the potentiometric titration for compound **1**. From this, only one pK_a value of 3.56 ± 0.03 could be determined, assignable to the protonation/deprotonation of an axial hydroxido group, in good agreement with the pK_{a1} values determined by NMR spectroscopy (3.34/3.44). No other acid–base equilibrium was detected potentiometrically, in particular within the range 7–8.

The lack of detection of proton release for this step in combination with NMR data, suggests that the observed NMR effects are due to a pH-dependent proton movement but not release from the complex, for example, an am(m)ine proton adopting an H-bonding position with an axial hydroxido perhaps mediated by a water molecule or hydroxide ion. Hambley and co-workers^[31] have recently determined

pK_a values for axial hydroxido ligands in platinum(IV) complexes using cyclic voltammetry and obtained values that lie in the same range as those reported here. Moreover, they have suggested that intermolecular bridging mechanisms are likely to be involved in both reduction and ligand exchange, which seem to be influenced by protonation of the hydroxido ligands. In this regard, the possible formation of bridged complexes upon deprotonation of the am(m)ine ligands cannot be ruled out, including amido-bridged species, although the limited relevant data available in the literature do not aid interpretation of our experimental results.^[32,33]

The possibility can be considered that such pH-dependent behavior near pH 7 is due to deprotonation of a minor species in solution which interacts rapidly on the NMR time-scale with the investigated complexes. In this respect, it has been reported that association of trace platinum(II) impurities (which might arise for example, from the starting platinum(II) precursors, or from photoreduction of the platinum(IV) starting materials when briefly exposed to light during sample preparation) with the corresponding platinum(IV) complexes can increase the substitution rates of ligands coordinated to the platinum(IV) center.^[34] Since HPLC studies have shown that the complexes used in this NMR study have a high state of purity (>99%, see Figure S2 in the Supporting Information), it seems reasonable to suppose therefore that the NMR data can be interpreted as arising from one predominant species in solution.

Conclusion

In this work, we have investigated the acid–base properties of three platinum(IV) diazidodiam(m)inedihydroxido complexes, *cis,trans,cis*-[Pt^{IV}(N₃)₂(OH)₂(NH₃)₂] (**1**), *trans,trans,trans*-[Pt^{IV}(N₃)₂(OH)₂(NH₃)₂] (**2**), *cis,trans*-[Pt^{IV}(N₃)₂(OH)₂(en)] (**3**), by means of multinuclear NMR methods and potentiometry. These complexes belong to a class of photoactivatable platinum(IV) anticancer prodrugs.^[4–9] In this regard, it is of particular interest to investigate whether the acid–base behavior of these platinum(IV) derivatives can influence their chemical and biological properties. Platinum(IV) complexes are relatively inert, so they are not expected to undergo facile ligand substitution reactions, but are likely to reach cellular target sites intact. To exert their antitumor activity, it is normally thought that platinum(IV) complexes must be first reduced by biological reductants to the corresponding more reactive platinum(II) species, so any pH influence is generally taken into account only with respect to their reduction pathways.^[35]

pH titrations followed by NMR spectroscopy allowed us to determine two distinct pH-dependent events with mid-points at approximately pH 3.4 and pH 7.5; this was made possible by the combination of both direct and indirect techniques for the detection of ¹⁵N and ¹⁴N chemical shifts. By comparison with the few data reported in the literature, it is reasonable to conclude that the lower pK_a value involves protonation/deprotonation of one of the axial hydroxido

groups. In contrast, the second pH-dependent phenomenon cannot be reasonably be attributed to the deprotonation of one of the equatorial am(m)ine ligands, since a larger shift of the ¹⁵N and ¹⁴N NMR resonances upon deprotonation of the coordinated am(m)ine would be expected. In addition, no proton release was detected potentiometrically at around pH 7.5. Also there are no reported pK_a values lower than 10 for neutral platinum(IV)-am(m)ine complexes. Only complexes with overall charges of 2+, 3+, and 4+ have pK_a values below 8 (Table 1).

It seems likely therefore that the second inflexion point in the titration curve at approximately pH 7–8 is associated with an intramolecular structural change in which a proton changes its environment in the complex but does not dissociate. The nature of the structural change will require further investigation. Perhaps it could involve for example a movement of the axial OH proton (or an NH proton) towards a terminal azido nitrogen—effectively pH-induced deprotonation resulting in an increase in basicity of bound azide. Such a structural change could influence the photoreduction pathways for these complexes and their biological activity.

Experimental Section

Materials: K₂[Pt^{IV}Cl₄] was purchased from Precious Metals Online; KI, NaN₃, KOH, NaCl, NaOH, KClO₄, and HCl from Fisher Scientific; AgNO₃, NH₄Cl, ¹⁵NH₄Cl, ethylenediamine dihydrochloride, ¹⁵N₂-ethylenediamine dihydrochloride, 1,4-dioxane, and D₂O from Sigma–Aldrich; H₂O₂ (30%) from Prolabo; CH₃CN and HClO₄ from Fisons.

Syntheses: Complexes *cis,trans,cis*-[Pt^{IV}(N₃)₂(OH)₂(NH₃)₂] (1), *trans,trans,trans*-[Pt^{IV}(N₃)₂(OH)₂(NH₃)₂] (2) and *cis,trans*-[Pt^{IV}(N₃)₂(OH)₂(en)] (3) were synthesized according to literature procedures.^[36,37] ¹⁵NH₄Cl and ¹⁵N₂-ethylenediamine dihydrochloride were used as the source of ¹⁵N for the synthesis of the corresponding ¹⁵N-labeled analogues (¹⁵N-1, ¹⁵N-2, ¹⁵N-3).

1. Yield: 78–80%; 1D ¹H NMR (500.1 MHz, 90% H₂O/10% D₂O, pH 5.4, TSP): δ = 5.12 ppm (Pt^{IV}-¹⁵NH₃, 6H, ¹J(¹⁵N–H) = 73 Hz, ²J(¹⁹⁵Pt–H) = 46 Hz); 2D [¹H, ¹⁵N] HSQC NMR (500.1/50.7 MHz, 90% H₂O/10% D₂O, pH 5.4, ¹⁵NH₄Cl): δ = –40.8 ppm (Pt^{IV}-¹⁵NH₃, ¹J(¹⁹⁵Pt–¹⁵N) = 261 Hz).

2. Yield: 72–73%; 1D ¹H NMR (500.1 MHz, 90% H₂O/10% D₂O, pH 5.4, TSP): δ = 5.31 ppm (Pt^{IV}-¹⁵NH₃, 6H, ¹J(¹⁵N–H) = 73 Hz, ²J(¹⁹⁵Pt–H) = 48 Hz); 2D [¹H, ¹⁵N] HSQC NMR (500.1/50.7 MHz, 90% H₂O/10% D₂O, pH 5.4, ¹⁵NH₄Cl): δ = –42.1 ppm (Pt^{IV}-¹⁵NH₃, ¹J(¹⁹⁵Pt–¹⁵N) = 282 Hz).

3. Yield: 88–90%; 1D ¹H NMR (500.1 MHz, 90% H₂O/10% D₂O, pH 5.5, TSP): δ = 2.95 ppm (Pt^{IV}-¹⁵NH₂–CH₂, 4H, ²J(¹⁵N–H) = 1.4 Hz, ³J(¹⁹⁵Pt–H) = 24 Hz); δ = 6.39 ppm (Pt^{IV}-¹⁵NH₂, 4H, ¹J(¹⁵N–H) = 76 Hz, ²J(¹⁹⁵Pt–H) = 44 Hz); 2D [¹H, ¹⁵N] HSQC NMR (500.1/50.7 MHz, 90% H₂O/10% D₂O, pH 5.5, ¹⁵NH₄Cl): δ = –7.7 ppm (Pt^{IV}-¹⁵NH₂, ¹J(¹⁹⁵Pt–¹⁵N) = 277 Hz).

Warning! Heavy metal-azido complexes are known to be shock-sensitive detonators. We encountered no problems in this study, but these materials should be handled with extreme caution, especially not to put pressure on them in the crystalline form.

All experiments were carried out with minimal exposure to light due to the high photosensitivity of the complexes.

NMR spectroscopy: All NMR spectra were acquired in 90% H₂O/10% D₂O at 298 K on a Bruker DMX500 spectrometer using a TBI [¹H, ¹³C,X] probe-head equipped with z-field gradients or a Bruker AV111-600 spectrometer. Data processing was carried out using MestReNova version

5.2.5 (Mestrelab Research S.L.) or with TOPSPIN version 2.1 (Bruker U.K. Ltd.).

Typical acquisition parameters for 1D ¹H NMR spectra (¹H: 500.13 MHz): 16 transients, spectral width 7.5 kHz, using 32k data points and a delay time of 2.0 s. Water suppression was achieved using a 55.0 dB power level presaturation. Data were processed by using exponential weighting with a resolution of 0.5 Hz and a line-broadening threshold of 0.1 Hz. ¹H chemical shifts were referenced to TSP by using internal 1,4-dioxane at 3.764 ppm.

Typical acquisition parameters for 1D ¹⁵N[¹H] INEPT and 1D ¹⁵N[¹H] NMR spectra (¹⁵N: 50.70 MHz): 128 transients, spectral width 15 kHz, using 64k data points and a delay time of 4.0 s. Sequences were optimized for ¹J(¹⁵N–H) = 76 Hz, and ¹H decoupling was achieved by using WALTZ16 pulse sequence. Data were processed by using exponential weighting with a resolution of 0.3 Hz and a line-broadening threshold of 6.0 Hz. ¹⁵N chemical shifts were referenced to external ¹⁵NH₄Cl 1.5 M in 1 M HCl at 0.00 ppm.


Typical acquisition parameters for 2D [¹H, ¹⁵N] HSQC NMR spectra (¹H: 500.13/¹⁵N: 50.70 MHz): 512 transients of 16 scans/block, spectral width 7.5/4.6 kHz, using 2k/2k data points and a delay time of 1.5 s. Sequences were optimized for ¹J(¹⁵N–H) = 76 Hz, and ¹H decoupling was achieved by using WALTZ16 pulse sequence. Data were processed using cosine-square weighting with a resolution of 2.2/3.7 Hz and a line-broadening threshold of 0.7/1.0 Hz. ¹H chemical shifts were referenced to TSP by using internal 1,4-dioxane at 3.764 ppm. ¹⁵N chemical shifts were referenced to external 1.5 M ¹⁵NH₄Cl in 1 M HCl at 0 ppm.

Typical acquisition parameters for 1D ¹⁴N[¹H] NMR spectra (¹⁴N: 43.35 MHz): 32k transients, spectral width 24 kHz, 16k data points and a delay time of 0.20 s. Due to acoustic resonance, the anti-ringing proton-decoupled ARINDEC pulse sequence was used.^[38] Data were processed using an exponential line-broadening of 5 Hz and a resolution of 1.5 Hz. ¹⁴N chemical shifts were referenced to internal N₂ at 287 ppm (relative to external 1.5 M NH₄Cl in 1 M HCl).

pH measurement: pH values were measured at room temperature directly in the NMR tube with a Corning 145 pH-meter equipped with an Aldrich microcombination electrode, calibrated with standard Aldrich buffer solutions at pH 4, 7 and 10.

pH titrations followed by NMR spectroscopy and pK_a determination: The NMR samples for pH titrations contained 15–20 mm platinum(IV) complex in 0.6 mL 90% H₂O/10% D₂O. The pH values of the solutions were adjusted by using dilute KOH or HClO₄, and NMR spectra were recorded.

The experimental pH titration data were fitted to the formulae in Equations (1) and (2) derived from Henderson–Hasselbalch equations (for one or two pK_a values, respectively) according to the general acid–base equilibria in Scheme 1.



Scheme 1. General scheme for a two-step acid–base equilibrium.

$$\delta = \frac{\delta_A \times 10^{-pH} + \delta_B \times 10^{-pK_{a1}}}{10^{-pH} + 10^{-pK_{a1}}} \quad (1)$$

$$\delta = \left(\delta_B + \frac{\delta_A \times 10^{-pH}}{10^{-pK_{a1}} + 10^{-pH}} + \frac{\delta_C \times 10^{-pK_{a2}}}{10^{-pH}} \right) / \left(1 + \frac{10^{-pH}}{10^{-pK_{a1}}} + \frac{10^{-pK_{a2}}}{10^{-pH}} \right) \quad (2)$$

K_{a1} and K_{a2} are the dissociation constants for species LH₂⁺ and LH, respectively, and δ_A, δ_B, and δ_C are the limiting chemical shifts of species LH₂⁺, LH and L[–], respectively. The pH titration curves (chemical shift vs. pH) were fitted using the program ORIGIN version 6.0 (Microcal Software Ltd.).

Mass spectrometry (ESI-MS): Positive- and negative-ion electrospray mass spectra were obtained on a Micromass Platform II mass spectrometer. The samples were prepared in water to a final concentration of 0.8–1.0 mg mL⁻¹, and pH was adjusted using dilute KOH or HClO₄ solutions. Samples (100 μL) were injected directly into the source by a syringe pump at a flow rate of 0.5 mL h⁻¹, and the ions were produced in an atmospheric pressure ionization (API) ESI ion source. The source temperature was 383 K, and the drying gas (N₂) flow rate was 300 L h⁻¹. A potential of 3.5 kV (capillary voltage) was applied to the probe tip (HV lens voltage: 0.5 kV; multiplier: 650 V), and the cone voltage was varied between (±) 5 and 25 V, depending on sensitivity. Mass spectra were recorded over the scan range 200–800 Da at a scan time rate of 2 s and a resolution of 2 *m/z*. Data acquisition and processing were carried out using MassLynx Software version 3.5 (Waters Corp.).

Potentiometric titrations: Potentiometric studies were carried out with a 721 NET Titrino autotitrator (Metrohm) equipped with a Metrohm combined pH glass electrode (Ag/AgCl) with 3 M KCl internal filling solution. The NaOH aqueous solution was standardized against potassium hydrogen phthalate with phenolphthalein as indicator. The electrode was calibrated with dilute standard acid and alkali solutions, thus defining pH = -log[H⁺] (pK_w = 14.28).^[39] The linearity of the electrode response and carbonate contamination of the standardized NaOH solution (0.0933 M) was determined by Gran's method^[40] and was found to be less than 1%. In a typical pH-metric determination, an aqueous solution (57 mL) of complex **1** (1.021 mM) treated with 4.45 equivalents of HClO₄ was titrated with the standardized NaOH solution. The temperature of the solutions (298 K) in the covered water-jacketed cell was kept constant by a Julabo circulating bath. The ionic strength (*I*) was adjusted to 0.1 with KClO₄. The protonation/deprotonation constants were calculated from the titration data by using the program HyperQuad 2000 version 2.1 (HyperQuad Solutions).^[41]

HPLC studies: HPLC investigations of complexes **1–3**, were carried out on an Agilent 1100 system using a reverse-phase C18 column (150 × 4.6 mm, 5 μm particle size, Agilent Zorbax Eclipse Plus). Solvents were either H₂O (solvent A) and CH₃CN (solvent B), at a flow rate of 1.0 mL min⁻¹ eluting with a gradient 5–80% B over 60 min. All injections were 50 μL of aqueous solutions of approximately 20–30 μM of each platinum(IV) complex. All solutions were made up and handled with minimal exposure to light. The detection wavelength was 254 nm (close to the λ_{max} of the azide-to-Pt^{IV} charge-transfer bands of these complexes).

Acknowledgements

We thank the EC (Marie Curie Fellowship for L.R.), EPSRC (EP/G006792/1), ERC (grant no. 247450), and ERDF/AWM (Science City) for support, Juraj Bella (Edinburgh) and Dr Nicola Farrer (Warwick) for assistance with NMR experiments, and Professor Trevor Hambley (University of Sydney) and members of EC COST Action D39 for stimulating discussions.

- [1] L. Kelland, *Nat. Rev. Cancer* **2007**, *7*, 573–584.
 [2] M. D. Hall, T. W. Hambley, *Coord. Chem. Rev.* **2002**, *232*, 49–67.
 [3] M. D. Hall, H. R. Mellor, R. Callaghan, T. W. Hambley, *J. Med. Chem.* **2007**, *50*, 3403–3411.
 [4] L. Ronconi, P. J. Sadler, *Coord. Chem. Rev.* **2007**, *251*, 1633–1648.
 [5] N. J. Farrer, L. Salassa, P. J. Sadler, *Dalton Trans.* **2009**, 10690–10701.
 [6] L. Ronconi, P. J. Sadler, *Chem. Commun.* **2008**, 235–237.
 [7] H. I. A. Phillips, L. Ronconi, P. J. Sadler, *Chem. Eur. J.* **2009**, *15*, 1588–1596.
 [8] L. Ronconi, P. J. Sadler, *Dalton Trans.* **2011**, *40*, 262–268.
 [9] L. Salassa, H. I. A. Phillips, P. J. Sadler, *Phys. Chem. Chem. Phys.* **2009**, *11*, 10311–10316.
 [10] J. Burgess, *Metal Ions in Solution*, Wiley, New York, **1978**.

- [11] F. S. Mackay, J. A. Woods, P. Hernigová, J. Kaspárková, A. M. Pizarro, S. A. Moggach, S. Parsons, V. Brabec, P. J. Sadler, *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 20743–20748.
 [12] L. Cubo, A. M. Pizarro, A. Gómez Quiroga, L. Salassa, C. Navarro-Ranninger, P. J. Sadler, *J. Inorg. Biochem.* **2010**, *104*, 909–918.
 [13] L. Fielding, *Tetrahedron* **2000**, *56*, 6151–6170.
 [14] S. J. Barton, K. J. Barnham, A. Habtemariam, R. E. Sue, P. J. Sadler, *Inorg. Chim. Acta* **1998**, *273*, 8–13.
 [15] Y. Chen, Z. Guo, P. J. Sadler in *Cisplatin. Chemistry and Biochemistry of a Leading Anticancer Drug* (Ed.: B. Lippert), Wiley-VCH, Weinheim, **1999**; pp. 293–318.
 [16] M. S. Davies, J. W. Cox, S. J. Berners-Price, W. Barklage, Y. Qu, N. Farrell, *Inorg. Chem.* **2000**, *39*, 1710–1715.
 [17] S. J. Berners-Price, L. Ronconi, P. J. Sadler, *Prog. Nucl. Magn. Reson. Spectrosc.* **2006**, *49*, 65–98.
 [18] L. E. Erickson, *J. Am. Chem. Soc.* **1969**, *91*, 6284–6290.
 [19] L. G. Sillén, E. A. Martell, J. Bjerrum, *Stability Constants of Metal-Ion Complexes. (Special Publication No. 17)*, 2nd ed., RCS, London, **1964**.
 [20] J. W. Palmer, F. Basolo, *J. Inorg. Nucl. Chem.* **1960**, *15*, 279–286.
 [21] L. Ronconi, P. J. Sadler, *Coord. Chem. Rev.* **2008**, *252*, 2239–2277.
 [22] N. A. Kratochwil, J. A. Parkinson, P. J. Bednarski, P. J. Sadler, *Angew. Chem.* **1999**, *111*, 1566–1569; *Angew. Chem. Int. Ed.* **1999**, *38*, 1460–1463.
 [23] N. N. Zheligovskaya, M. G. Felin, V. I. Spitsyn, *Dokl. Akad. Nauk SSSR* **1970**, *195*, 113–114.
 [24] N. N. Zheligovskaya, M. G. Felin, *Vestn. Mosk. Univ. Ser. 2: Khim.* **1971**, *12*, 679–682.
 [25] M. G. Felin, N. N. Zheligovskaya, V. I. Spitsyn, *Dokl. Akad. Nauk SSSR* **1971**, *199*, 665–666.
 [26] M. G. Felin, D. A. Murashov, N. N. Zheligovskaya, V. I. Spitsyn, *Izv. Akad. Nauk SSSR Ser. Khim.* **1972**, 790–792.
 [27] W. Beck, K. Z. Feldl, *Naturforsch. B* **1966**, *21*, 588–588.
 [28] D. Humphreys, P. J. Staples, *J. Chem. Soc. Dalton Trans.* **1973**, 897–900.
 [29] N. N. Zheligovskaya, M. G. Felin, *Vestn. Mosk. Univ. Ser. 2: Khim.* **1971**, *12*, 618–619.
 [30] K. I. Gil'dengershel, S. I. Pechenyut, A. I. Stetsenko, V. F. Budanova, *Zhurnal Neorganicheskoi Khimii*, **1971**, *16*, 2029–2030.
 [31] M. S. Davies, M. D. Hall, S. J. Berners-Price, T. W. Hambley, *Inorg. Chem.* **2008**, *47*, 7673–7680.
 [32] N. W. Alcock, P. Bergamini, T. J. Kemp, P. G. Pringle, S. Sostero, O. Traverso, *Inorg. Chem.* **1991**, *30*, 1594–1598.
 [33] S. A. Delp, C. Munro-Leighton, C. Khosla, J. L. Templeton, N. M. Alsop, T. Brent Gunnoe, T. R. Cundari, *J. Organomet. Chem.* **2009**, *694*, 1549–1556.
 [34] S. Choi, L. Vastag, Y. C. Larrabee, M. L. Personick, K. B. Schalberg, B. J. Fowler, R. K. Sandwick, G. Rawji, *Inorg. Chem.* **2008**, *47*, 1352–1360.
 [35] L. T. Ellis, H. M. Er, T. W. Hambley, *Aust. J. Chem.* **1995**, *48*, 793–806.
 [36] P. Müller, B. Schröder, J. A. Parkinson, N. A. Kratochwil, R. A. Coxall, A. Parkin, S. Parsons, P. J. Sadler, *Angew. Chem.* **2003**, *115*, 349–353; *Angew. Chem. Int. Ed.* **2003**, *42*, 335–339.
 [37] F. S. Mackay, J. A. Woods, H. Moseley, J. Ferguson, A. Dawson, S. Parsons, P. J. Sadler, *Chem. Eur. J.* **2006**, *12*, 3155–3161.
 [38] N. J. Farrer, P. Gierth, P. J. Sadler, *Chem. Eur. J.* DOI: 10.1002/chem.201101409.
 [39] E. M. Woolley, D. G. Hurkot, L. G. Hepler, *J. Phys. Chem.* **1970**, *74*, 3908–3913.
 [40] A. E. Martell, R. J. Motekaitis, *Determination and Use of Stability Constants*, Wiley-VCH, Weinheim, **1992**.
 [41] P. Gans, A. Sabatini, A. Vacca, *Talanta* **1996**, *43*, 1739–1753.

Received: September 29, 2010

Revised: May 22, 2011

Published online: September 16, 2011