

Original citation:

Wand, B. M. et al. (2012). Should we abandon cervical spine manipulation for mechanical neck pain? Yes. BMJ, 344(jun07 3), pp. e3679

Permanent WRAP url:

<http://wrap.warwick.ac.uk/47935>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes the work of researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

3 X E O L V K H U ¶ V V W D W H P H Q W
© 2012 BMJ Publishing Group Ltd
DOI: 10.1136/bmj.e3679

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to FR Q V X O W W K H S X E R Please see U ¶ V Y H U V L R W K H µ S H U P D Q H Q W : 5 \$ 3 X U O ¶ D E R Y H I R U G H W D and Note R Q D F F H V V L that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

Should we abandon cervical spine manipulation for mechanical neck pain? Yes

Benedict M Wand, Peter J Heine, Neil E O'Connell

BMJ 2012; 344 doi: 10.1136/bmj.e3679

Cervical spine manipulation (a high velocity, low amplitude, end range thrust manoeuvre) is a common treatment option for mechanical neck pain yet may carry the potential for serious neurovascular complications, specifically vertebral artery dissection and subsequent vertebrobasilar stroke. The non-superiority of manipulation to alternative treatments, coupled with concerns regarding safety, renders cervical spine manipulation unnecessary and inadvisable.

The controversy surrounding the association between manipulation and neurovascular complications is longstanding and not fully resolved, partly because it is difficult to obtain conclusive evidence on rare adverse events. What can be accepted is that the incidence of vertebral artery dissection is low, with estimates between 1 (95% confidence interval 0.5 to 1.4) and 1.7 (1.1 to 2.3) per 100 000 person years in the United States.¹ The estimates for stroke resulting from vertebral artery dissection are lower still, ranging from 0.75 to 1.12 per 100 000 person years,² and many are unlikely to be the result of cervical manipulation.

Nevertheless, numerous case studies report neurovascular complications immediately after cervical manipulation,³ and more robust case-control studies provide consistent evidence of an association between neurovascular injury and recent exposure to cervical manual therapy, particularly manipulation.^{4, 5, 6} Although absolute risk cannot be accurately estimated, these studies have reported large effects in general populations (adjusted odds ratios 6.62, 95% confidence interval 1.4 to 304; 12.67, 1.43 to 112.05) and in patients under 45 (5.03, 1.58 to 16.076). However, the causal nature of this association has recently been called into question by the findings of one case-crossover study.⁷ Although the study found an association between vertebrobasilar stroke and chiropractic care in patients under 45 (3.60, 1.46 to 10.84), a comparable association was found between vertebrobasilar stroke and primary care practitioner visits (2.99, 1.81 to 4.96). The authors suggest that the increased risk after chiropractic treatment may be an artefact of patients seeking care for neck pain resulting from existing vertebral artery dissection rather than the result of treatment itself. Although the results suggest that some cases of vertebrobasilar stroke may be misattributed to manipulation, this does not rule out that some patients have dissection induced by manipulation or that the clinical sequelae are worsened by manipulation in some patients with spontaneous dissection.

To conclude that all adverse neurovascular events seen after manipulation are the manifestation of a pre-existing spontaneous dissection is at odds with several findings. A previous case-control study found that manipulation remained an independent risk factor for dissection after controlling for the previous presence of neck pain (adjusted odds ratio 6.62, 95% confidence interval 1.4 to 30),⁴ and another study reported that patients with vertebral artery dissection and previous exposure to manipulation are more likely to present with damage to the more mechanically vulnerable upper cervical portion of the artery than those without exposure (increase in prevalence ratio attributable to manipulation 4.14).⁸ Furthermore, patients presenting with conditions that do not share symptoms with vertebral artery dissection (such as low back pain) have reported neurovascular complications after neck manipulation,⁹ and it seems most reported cases of vertebral artery dissection and stroke after manual therapy have followed chiropractic care rather than osteopathy or physiotherapy, where manipulation is used less often.⁹

Though causality is not proved, legitimate concerns remain regarding the risk of such serious events. Whether there are factors that leave some patients more susceptible to dissection remains a matter of conjecture,^{1, 5} and there are no satisfactory screening procedures that acceptably mitigate this risk.⁵ It

follows that neck manipulation should be used only if there is substantial and unique benefit associated with this technique.

On this point the literature is clearer. A recent Cochrane review of randomised controlled trials of neck manipulation or mobilisation concluded that as a stand alone treatment, manipulation provides only moderate short term pain relief versus waiting list control, sham manipulation, or muscle relaxants (standardised mean difference -0.90 , 95% confidence interval -1.78 to -0.02), is unlikely to offer meaningful long term benefit for people with neck pain, and does not seem to be better than other manual therapy techniques such as cervical mobilisation (-0.07 , -0.47 to 0.32).¹⁰ A recent clinical trial suggests this equivalence remains even in patients whom the clinician deemed particularly suitable for manipulation.¹¹ Other recent large, high quality randomised trials reinforce the message that manipulation is not superior when directly compared with other physical interventions such as exercise and confers no additional benefit when added to them.^{12, 13}

Given the equivalence in outcome with other forms of therapy, manipulation seems to be clinically unnecessary. The potential for catastrophic events and the clear absence of unique benefit lead to the inevitable conclusion that manipulation of the cervical spine should be abandoned as part of conservative care for neck pain. In the interests of patient safety, the regulatory and professional bodies associated with professions that use manual therapy should consider adopting this as a formal policy.

References

1. DeBette S, Leys D. Cervical-artery dissections: predisposing factors, diagnosis and outcome. *Lancet Neurol*2009;8:668-78.
2. Boyle E, Côte P, Grier AR, Cassidy JD. Examining vertebrobasilar artery stroke in two Canadian provinces. *J Manipulative Physiol Ther*2009;32:S194-200.
3. Ernst E. Adverse effects of spinal manipulation: a systematic review. *J R Soc Med*2007;100:330-8.
4. Smith WS, Johnston SC, Skalabrin EJ, Weaver M, Azari P, Albers GW, et al. Spinal manipulative therapy is an independent risk factor for vertebral artery dissection. *Neurology*2003;60:1424-8.
5. Thomas LC, Rivett DA, Attia JR, Parsons M, Levi C. Risk factors and clinical features of craniocervical arterial dissection. *Manual Therapy*2011;16:351-6.
6. Rothwell DM, Bondy SJ, Williams JI, Bousser MG. Chiropractic manipulation and stroke: a population-based case-control study. *Stroke*2001;32:1054-60.
7. Cassidy JD, Boyle E, Côté P, He Y, Hogg-Johnson S, Silver FL, Bondy SJ. Risk of vertebrobasilar stroke and chiropractic care. Results of a population-based case-control and case-crossover study. *Spine*2008;33 (suppl 4):S176-83.
8. Kawchuk GN, Jhangri GS, Hurwitz EL, Wynd S, Haldeman S, Hill MD. The relation between the spatial distribution of vertebral artery compromise and exposure to cervical manipulation. *J Neurol*2008;255:371-7.
9. Ernst E. Vascular accidents after neck manipulation: cause or coincidence? *Int J Clin Pract*2010;64:673-7.
10. Gross A, Miller J, D'Sylva J, Burnie SJ, Goldsmith CH, Graham N, et al. Manipulation or mobilisation for neck pain. *Cochrane Database Syst Rev*2010;1:CD004249.

11. Leaver AM, Maher CG, Herbert RD, Latimer J, McAuley JH, Jull G, et al. A randomized control trial comparing manipulation with mobilization for recent onset neck pain. Arch Phys Med Rehabil 2010;91:1313-8.
12. Evans R, Bronfort G, Schulz C, Maiers M, Bracha Y, Svendsen K, et al. Supervised exercise with and without spinal manipulation perform similarly and better than home exercise for chronic neck pain: a randomized controlled trial. Spine 2012;37:903-14.
13. Bronfort G, Evans R, Anderson AV, Svendsen KH, Bracha Y, Grimm RH. Spinal manipulation, medication, or home exercise with advice for acute and subacute neck pain: a randomized trial. Ann Intern Med 2012;156:1-10.