

University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): Cheng Qin, Nongnong Shi, Mei Gu, Hang Zhang, Bin Li, Jiajia Shen, Atef Mohammed, Eugene Ryabov, Chunyang Li, Huizhong Wang, Yule Liu, Toba Osman, Manu Vatish & Yiguo Hong

Article Title: Involvement of RDR6 in short-range intercellular RNA silencing in *Nicotiana benthamiana*

Year of publication: 2012

Link to published article:

<http://dx.doi.org/10.1038/srep00467>

Publisher statement: License: This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported

License. To view a copy of this

license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>

.



SUBJECT AREAS:

PLANT SCIENCES

RNAI

VIROLOGY

SMALL RNAS

Involvement of *RDR6* in short-range intercellular RNA silencing in *Nicotiana benthamiana*

Cheng Qin^{1*}, Nongnong Shi^{1,2*}, Mei Gu^{3*}, Hang Zhang^{2,4}, Bin Li¹, Jiajia Shen¹, Atef Mohammed^{2,5}, Eugene Ryabov², Chunyang Li^{2,4}, Huizhong Wang¹, Yule Liu⁶, Toba Osman^{2,5}, Manu Vatish³ & Yiguo Hong^{1,2}

Received
16 May 2012

Accepted
11 June 2012

Published
22 June 2012

Correspondence and requests for materials should be addressed to Y.H. (yghongabc@googlemail.com or yiguo.hong@hznu.edu.cn or yiguo.hong@warwick.ac.uk)

* These authors contributed equally to this work.

¹Research Centre for Plant RNA Signalling, College of Life and Environmental Sciences, Hangzhou Normal University, Hangzhou 310036, China, ²Warwick HRI, University of Warwick, Warwick CV35 9EF, UK, ³Clinical Sciences Research Institute, University of Warwick, Coventry CV2 2DX, UK, ⁴Chengdu Rongsheng Pharmaceuticals, Chengdu 610041, China, ⁵Department of Botany, Faculty of Agriculture, Fayoum University, Fayoum 63514, Egypt, ⁶School of Life Sciences, Tsinghua University, Beijing 100084, China.

In plants, non-cell autonomous RNA silencing spreads between cells and over long distances. Recent work has revealed insight on the genetic and molecular components essential for cell-to-cell movement of RNA silencing in *Arabidopsis*. Using a local RNA silencing assay, we report on a distinct mechanism that may govern the short-range (6–10 cell) trafficking of virus-induced RNA silencing from epidermal to neighbouring palisade and spongy parenchyma cells in *Nicotiana benthamiana*. This process involves a previously unrecognised function of the RNA-dependent RNA polymerase 6 (*RDR6*) gene. Our data suggest that plants may have evolved distinct genetic controls in intercellular RNA silencing among different types of cells.

RNA silencing is a potent surveillance system to protect against virus infections in plants, fungi and animals. Viruses have evolved both passive and active strategies to escape and effectively suppress RNA silencing. In particular, viruses encode suppressors of RNA silencing (VSRs) that can directly or indirectly target various steps in intra-, intercellular and systemic RNA silencing¹. Indeed, VSRs can subvert antiviral silencing by inhibiting the biosynthesis of small interfering (si)RNAs, by preventing the assembly of RNA-induced silencing complexes (RISC), by repressing the activity of the Argonaute (AGO) proteins, by impeding silencing amplification, and/or by blocking the spread of silencing signals. For example, the P38 coat protein (CP) of *Turnip crinkle virus* (TCV) is an effective VSR^{2,3}. It binds to dsRNA and siRNA and specifically inhibits DCL4 activity to prevent siRNA production. TCV P38 also interacts with (and subsequently inactivates) AGO1 to impede RISC-mediated targeting and the degradation of viral RNAs^{4–6}. The silencing suppression function of TCV P38 also requires the ethylene-inducible host transcription factor RAV2a⁷.

In plants, RNA silencing is non-cell autonomous⁸. The 21-nt siRNA produced by DCL4 from inverted-repeat transgenes, acts as the primary silencing signal and spreads short-range over 10–15 cells. This type of intercellular trafficking of RNA silencing does not require RNA-dependent RNA polymerase 6 (*RDR6*). However, DCL4 also generates an *RDR6*-dependent 21-nt secondary siRNA for long-range (i.e., beyond 10–15 cells) signalling in systemic RNA silencing^{9,10}. Genetic analysis in *Arabidopsis* has revealed three recessive *silencing-movement-deficient* (*smd1-3*) mutations which are compromised in short-range cell-to-cell movement of RNA silencing⁹. *SMD1* and *SMD2* are allelic to *RDR2* and to *NRPD1a*, respectively. Both genes are required for intercellular but not intracellular RNA silencing¹¹. More recently, through cell-specific rescue of the DCL4 function and cell-specific suppression of the movement of RNA silencing, it has been shown that 21-nt siRNA duplexes probably represent a component of the mobile silencing signal that spreads between plant cells¹². Interestingly, mobile silencing signals also include a range of small RNA (sRNA) species. Indeed, transgene-derived and endogenous 22-, 23- and 24-nt siRNAs are able to move between plant cells through plasmodesmata and traffic long distances via phloem transportation¹³. These findings have led to an elegant model that explains the genetic and molecular basis for limited (10–15 cell) spread of RNA silencing from companion cells to neighbouring parenchyma and mesophyll cells outside the vasculature in *Arabidopsis*^{8–12}. It should be noted that in this model RNA silencing is triggered by



a long dsRNA-producing transgene or a synthetic 21-nt siRNA. Interestingly, plants have also been shown to be capable of hijacking viral RNA movement proteins in order to facilitate trafficking of virus-induced and transgene-mediated RNA silencing^{14,15}.

On the other hand, RNA silencing can be initiated by transgenes, high molecular weight RNAs, siRNA or plant RNA and DNA viruses in various cell types. In these scenarios, the initiation of silencing and the subsequent movement of silencing signals between cells may have different requirements. Using a local RNA silencing assay coupled with molecular and cellular analyses, we have uncovered a mechanism associated with short-range (6–10 cell) spread of virus-induced RNA silencing from individual epidermal cells to adjacent cells. This process requires a functional *RDR6* gene in *Nicotiana benthamiana*.

Results

***RDR6* is essential for cell-to-cell trafficking of local virus induced RNA silencing.** We exploited the *Turnip crinkle virus* (TCV)-based

local RNA silencing assay devised to dissect the cell-to-cell spread of antiviral RNA silencing in plants^{3,14}. The movement-deficient TCV-GFPΔCP was constructed by replacing the P38 CP gene with the green fluorescent protein (GFP) coding sequence³. Thus, it lacks the potent silencing suppressor CP and is restricted to single epidermal cells of *N. benthamiana*^{2–5,7,12}. When mechanically applied to the upper epidermis of young leaves of GFP-expressing *N. benthamiana* GFP16c transgenic plants¹⁶, TCV-GFPΔCP was able to initiate *gfp* RNA silencing in single epidermal cells. Then, silencing spread from cell-to-cell to form *gfp*-silenced foci, that were visible under long-wavelength UV light (Fig. 1a, b, e, f, i, j, m, n, q, r). Efficient and multi-dimensional spread of *gfp* silencing from a single epidermal cell to other cell types was evident (Table 1; Fig. 2a). Large numbers of silencing foci per inoculated leaf were counted on both the upper (161 ± 6) and lower (146 ± 12) epidermises of four leaves. Silencing foci, selected at random, appeared to have similar sizes (0.98 ± 0.39 mm, 0.97 ± 0.27 mm; n = 12) on upper and lower epidermis, respectively. Interestingly, we




Figure 1 | Local induction and cell-to-cell spread of virus-induced RNA silencing. The first and second young leaves of *N. benthamiana* GFP16c and GFP16c/RDR6i at the six-leaf stage were mechanically inoculated with an equal amount of TCV-GFPΔCP RNA transcripts. Control plants were mock-inoculated with 10 mM Tris-HCl (pH 8.0) containing 10 mM EDTA. Both the upper and lower epidermises of inoculated leaves were photographed 8 days post-inoculation (dpi) (e–l) or 12 dpi (a–d, m–t) using a Nikon Coolpix995 digital camera under long-wavelength UV illumination through a yellow Kodak No. 58 filter. *Gfp* RNA-silenced tissue (foci) showed red chlorophyll fluorescence and *gfp*-expressing tissue showed green fluorescence. Colour-coded arrows in Panels (g), (h), (o) and (p) show the same *gfp* silencing foci observed from the upper or lower epidermis. A diagram showing the genome of TCV-GFPΔCP is included. A cartoon of a plant shows the positions of the first and second inoculated leaves (I. L.).

Table 1 | Impact of *RDR6* on cell-to-cell spread of virus-induced RNA silencing

		GFP16c	GFP16c/ <i>RDR6i</i>	Student's <i>t</i> test ³
Numbers of silencing foci per leaf ¹	Upper	161 ± 6 (n = 4)	25 ± 25 (n = 4)	p = 0.00001**
	Lower	146 ± 12 (n = 4)	4 ± 4 (n = 4)	p = 0.00001**
		p = 0.124*	p = 0.209*	
Diameters of silencing foci (mm) ²	Upper	0.98 ± 0.39 (n = 12)	0.50 ± 0.35 (n = 12)	p = 0.002**
	Lower	0.97 ± 0.27 (n = 12)	0.51 ± 0.39 (n = 12)	p = 0.003**
		p = 0.952*	p = 0.953*	

¹Number of silencing foci were counted on the upper or lower epidermis of four leaves, one leaf from each of four inoculated plants.

²Diameters of 12 randomly selected silencing foci were measured using a micro-ruler under long-wavelength UV light.

³Student's *t* tests were carried out between the upper and lower epidermis*. The *t* tests were also performed between GFP16c and GFP16c/*RDR6i*** . There are significant differences in the average numbers and sizes of silencing foci between GFP16c and GFP16c/*RDR6i*, but not between the upper and lower epidermis of plants with a same genetic background.

noticed that the number of silencing foci was always higher in the 2nd inoculated leaf than in the 1st inoculated leaf (Fig. 1), suggesting that the age of leaf tissues might have an effect of local RNA silencing. These data were consistent with previous results^{3,14,17}. However, in GFP16c/*RDR6i* plants in which *RDR6* was silenced¹⁶, we found a marked reduction in the numbers of visible *gfp* silencing foci, on the upper (25 ± 25) and, particularly the lower (4 ± 4) epidermis of four treated leaves (Table 1; Fig. 1c, d, g, h, k, l, o, p, s, t; Fig. 2a). There was also a significant decrease in the average size (diameter) of the *gfp* silencing foci on the upper (0.50 ± 0.35 mm) and lower (0.51 ± 0.39 mm) epidermises (Table 1; Fig. 2b). This phenomenon was observed in all repeated (more than three separate) experiments. Thus, in *RDR6*-“knockdown” plants, the silencing initiated by TCV-GFPΔCP in individual upper epidermal cell was incompetent to spread between epidermal cells and to penetrate to palisade and spongy parenchyma cells in order to reach lower epidermis. These findings suggest that *RDR6* was required for the cell-to-cell communication of RNA silencing in *N. benthamiana*. It should be noted that sizes of individual silencing foci could vary because the initiation of intracellular silencing may not be synchronous and the development of each silencing focus is probably a dynamic process.

***RDR6* is required for the short-range (6 – 10 cell) intercellular spread of virus-induced RNA silencing.** We examined 12 randomly selected silencing foci by fluorescent microscopy and estimated the numbers of epidermal cells in which *gfp* RNA silencing had occurred (Fig. 2c–l). A typical silencing focus that formed in GFP16c leaves inoculated with TCV-GFPΔCP consisted of 100–300 epidermal cells, equivalent to a circular zone with a radius of 6 – 10 epidermal cells (Fig. 2c, d, g, h, k). These data demonstrate that virus-induced RNA silencing can move from a single upper epidermal cell over a short-range of 6 – 10 upper epidermal cells. However, intracellular RNA silencing originating from a single upper epidermal cell could move, simultaneously, in three-dimensions to and through many more cell types, including palisade, mesophyll and lower epidermal cells, to form a visible focus of silencing (Fig. 1). Such RNA silencing was unlikely to move into the vasculatures as systemic RNA silencing was not observed in distal leaf tissues^{3,14,17}. Consistent with their reduced size (Fig. 2b), the *gfp* RNA silencing foci that formed on GFP16c/*RDR6i* leaves encompassed only 30 – 100 epidermal cells, equivalent to a circular zone with a radius of 3–6 epidermal cells (Fig. 2e, f, i, j, l). These data provide additional evidence that a functional *RDR6* gene plays an important role in short-range (6 – 10 cell) trafficking of virus-induced RNA silencing.

***RDR6* does not influence the replication and defective movement of TCV-GFPΔCP in *N. benthamiana*.** To address whether *RDR6* gene expression could affect replication and cell-to-cell spread of TCV-GFPΔCP, we first performed quantitative RT-PCR (qRT-PCR) assays to detect and compare the levels of *RDR6* mRNA in

wild-type and *RDR6*-silenced lines N*bRDR6i* and GFP16c/*RDR6i*¹⁶ using two housekeeping gene *GAPDH* and *EF1α* transcripts as internal controls. Our data showed that *RDR6* gene expression was statistically significantly reduced in N*bRDR6i* and GFP16c/*RDR6i* plants when compared to that in the wild-type *RDR6* (Nb and GFP16c) plants (Fig. 3a, b; Table S1). Moreover, qRT-PCR analyses showed that TCV-GFPΔCP RNA accumulation reached equivalent levels with no statistical significance in wild type *N. benthamiana* (Nb) and *RDR6*-silenced N*bRDR6i* (Fig. 3a, b; Table S1). We also showed that the levels of TCV-GFPΔCP RNA in the inoculated leaves of GFP16c and GFP16c/*RDR6i* were similar, but decreased substantially when compared to the viral RNA levels in wild-type Nb or *RDR6*-silenced N*bRDR6i* plants, respectively (Fig. 3a, b; Table S1). This is suggesting a partial resistance to TCV-GFPΔCP in the GFP16c lines. Such resistance may be derived from homologous sequence-dependent specific targeting to the GFP marker gene, which then resulted in degradation of the recombinant virus by RNA silencing. It is worthwhile noting that the transgenic GFP16c line and the wild-type *N. benthamiana* plants are susceptible to infection with TCV, *Tobacco mosaic virus* and *Potato virus X*^{3,17,18}, suggesting the GFP16c line should not have abnormalities in viral immunity at least for these viruses. Furthermore, TCV-GFPΔCP was always found to be restricted to single epidermal cells with a wild-type or *RDR6*-silenced genetic background (Fig. 3c, d). These data indicate that TCV-GFPΔCP was an effective inducer and the target of intracellular RNA silencing, and that *RDR6* was not required for the induction of antiviral RNA silencing in individual epidermal cells.

Taken together, our findings support the hypothesis that *RDR6* is probably involved in promoting the cell-to-cell spread of TCV-GFPΔCP induced RNA silencing, and the significant decrease in the intercellular RNA silencing in GFP16c/*RDR6i* plants is not due to any impact from *RDR6* on viral RNA replication and intercellular virus trafficking.

Discussion

Plants have multiple *RDR* genes that function in different biological processes. In *Arabidopsis*, there are six *RDR* genes (*RDR1*–6) that encode potential RNA dependent RNA polymerases (RDRs), of which *RDR6* is the major *RDR* acting in post-transcriptional gene silencing and virus induced RNA silencing^{11,19,20}, while *RDR2* plays an important role in transcriptional gene silencing²¹. *RDR6*-dependent RNA silencing forms an important antiviral defence. It defends both differentiated and meristematic tissues from viral invasion. *RDR6* silencing causes plants to be hyper-susceptible to virus infection^{16,22,23}. *RDR1* is also involved in antiviral RNA silencing, but its effects on plant responses to viral invasion tend to be virus specific^{24–27}. Recently, small RNA deep sequencing revealed a role for *RDR1* and *RDR6* in the biogenesis of viral siRNA^{24,28} and showed *RDR6* was essential for the maintenance and transitivity of RNA silencing^{19,29–31}. In *Arabidopsis*, *RDR6* is also thought to play a role in long-range




Figure 2 | Influence of *RDR6* on intercellular RNA silencing. (a–b) The *RDR6* gene affects the intercellular spread of TCV-GFPΔCP-induced *gfp* silencing. The average numbers of silencing foci per inoculated leaf for four leaves in a typical experiment using recombinant RNA transcripts produced from 2.5 μg of TCV-GFPΔCP DNA template are shown (a). Numbers of foci were counted 8 days post-inoculation (dpi). The average sizes of 12 randomly selected silencing foci from the upper and lower epidermises were analysed (b). (c–l) Dark *gfp* RNA silencing foci from the upper and lower epidermises of *N. benthamiana* GFP16c and GFP16c/RDR6i were examined under a fluorescence microscope using a green filter (c–f) or through bright field illumination (g–l). The boxed areas were used to estimate the numbers of epidermal cells in which *gfp*-silencing occurred. Individual epidermal cells are outlined and marked with asterisks. Scale bars are indicated.

(> 10 – 15 cell) intercellular and systemic RNA silencing, but not in short-range (10 – 15 cell) cell-to-cell signalling of RNA silencing that was initiated in companion and phloem cells^{10,16,32}.

However, considering the fact that TCV-GFPΔCP is restricted to single epidermal cell of mechanically inoculated leaves, our data suggest that the short-range (6 – 10 cell) intercellular trafficking of virus-induced RNA silencing from a single epidermal cell to adjacent palisade and spongy parenchyma cells in *N. benthamiana* may require a functional *RDR6* gene (Table 1; Fig. 1; Fig. 2). It is possible that spread of this type of RNA silencing may be associated with a

small RNA signal⁸. The RDR6 protein is known to bind ssRNA, dsRNA and even dsDNA in plants³³. Thus, RDR6 and small RNAs could form an RNA-protein complex that may directly promote 6 – 10 cell-to-cell movement of silencing signal. This model is consistent with the localization of the RDR6 protein in the cytoplasm, the nucleus and the cell membrane^{34,35}. Alternatively, RDR6 may be required to produce a signal for limited intercellular silencing. Single cells with a normal RDR6 function in GFP16c plants could generate sufficient quantities of mobile signal molecules to enable the formation of silencing foci. In contrast, reduced RDR6 activity in




Figure 3 | *RDR6*-independent intracellular RNA silencing. (a and b) The *RDR6* gene does not affect replication of TCV-GFP Δ CP. The *RDR6* mRNA were analysed 12 days post-inoculation (dpi) by quantitative (q) RT-PCR, in triplicate, using total RNAs extracted from mock- or TCV-GFP Δ CP-inoculated leaves of wild-type *N. benthamiana* (Nb) and NbRDR6i (*RDR6*i), transgenic GFP 16c (GFP16c) and GFP16c/*RDR6*i plants. Virus-induced intracellular RNA silencing targets TCV-GFP Δ CP. TCV-GFP RNA was analysed by qRT-PCR in triplicate, using total RNAs extracted from mock- or TCV-GFP Δ CP-inoculated GFP16c and GFP16c/*RDR6*i leaves at 12 dpi. *N. benthamiana* housekeeping *GAPDH* (a) and *EF1 α* (b) transcripts were used as internal controls. Relative RNA levels of *RDR6* or TCV were obtained by normalising against the baseline expression levels of *GAPDH* (a) and *EF1 α* (b) mRNA, respectively, and showed similar tendencies between the two internal controls. Student's *t*-tests were carried out to evaluate whether there would be any statistical significance in RNA levels between different biological samples (Table S1). (c and d) Fluorescence microscopic examination of *gfp* expression in a single epidermal cell in TCV-GFP Δ CP-inoculated Nb (c) or NbRDR6i (d) leaves 6 dpi using a Zeiss Axiophot microscope through a green filter. Bar = 100 μ m.

GFP16c/*RDR6*i plants could lead to decreased synthesis of such a signal. Thus, the level of the mobile signal may be an important factor for intercellular RNA silencing. In this scenario, *RDR6* may play an indirect role in short-range (6 – 10 cell) trafficking of RNA silencing. Also supporting a basic role of *RDR6*, not only the size but also the number of foci is lower in the *RDR6*-deficient plants. Moreover, some silencing foci were detected in the lower epidermis of leaves inoculated in their upper faces (Fig. 1). However, this could be caused by that *RDR6* was not completely knocked out and residual *RDR6* activity might persist in the *RDR6*i plants. Indeed, low but quantifiable levels of *RDR6* gene expression were readily detectable in all *RDR6*-silenced plants (Fig. 3a, b). A third possibility could be that indirect impacts of the knockdown of *RDR6* on the plant defensive responses, or even on the physiological transport of macro-molecules might affect the cell-to-cell spread of virus induced RNA silencing. It

should be noted that the cell-to-cell spread of RNA silencing in TCV-GFP Δ CP/GFP16c-based local RNA silencing assay depends on the expression of functional P8 and P9 proteins of TCV¹⁴. This assay is also susceptible to different VSRs¹⁷. These factors may affect how *RDR6* protein is recruited to promote 6 – 10 cell-to-cell spread of virus-induced RNA silencing, a process that may have different genetic and molecular requirements in *Arabidopsis* and other plants^{8–12}. Regardless of the underpinning mechanism, our data have revealed a previously unknown function for the *RDR6* gene in short-range (6 – 10 cell) intercellular RNA silencing in *N. benthamiana*.

Our unexpected finding contradicts previous reports that *RDR6* was not involved in the limited 10 – 15 cell-to-cell spread of RNA silencing in *Arabidopsis*^{8–12}. Such a discrepancy may be due to a combination of factors (Table S2). Compared with *Arabidopsis*, *N. benthamiana* is permissive and can be a non-native host for a wide range of plant viruses, supposedly due to a naturally occurring mutation in the *N. benthamiana RDR1* gene^{26,27}. Indeed, *N. benthamiana* and *Arabidopsis* plants respond differently to TCV-GFP Δ CP. For instance, TCV CP is essential for cell-to-cell virus movement in *N. benthamiana*^{3,36}, but not in *Arabidopsis*⁵. Consequently, TCV-GFP Δ CP moved effectively from cell-to-cell to form multi-cellular lesions in inoculated *Arabidopsis* leaves (unpublished data). However, the same virus was restricted to single cell in wild-type and *RDR6*i *N. benthamiana* plants (Fig. 3c, d). Such distinctive virus-host responses and different genetic backgrounds may affect the biosynthesis and spread of the silencing signal, thus have a significant impact on TCV-GFP Δ CP-mediated local RNA silencing in these plant species.

On the other hand, confinement of virus infection to individual epidermis cells was only demonstrated in plants not expressing GFP (Fig. 3c, d), but not directly in the centre of the silenced foci of the GFP16c and GFP16c/*RDR6*i plants or in single infected cells that are not surrounded of silenced foci in the GFP16c/*RDR6*i plants. However, we had checked hundreds of silencing foci under epifluorescent and confocal microscopes, no individual epidermal cells with viral transient GFP fluorescence in the centre of silencing foci were ever found, although GFP-expressing cells of Nb and NbRDR6i leaves inoculated with TCV-GFP Δ CP were readily observed at 2 – 12 dpi. Virus can be a trigger and a target of virus-induced RNA silencing (Fig. 3a, b). Once a silencing focus became visible, virus-induced RNA silencing would be certainly well-established and it could target viral RNAs for degradation. Therefore, accumulation of viral RNAs and proteins in single epidermal cells could be reduced to an undetectable level. Nevertheless, our findings emphasise that the signalling of intercellular RNA silencing is a more complex process than previously proposed (Table S2). This process may have different molecular components in different cell types in same plants and exhibit distinct genetic requirements in different plant species.

Methods

Plasmid construction, inoculation and plant maintenance. Plasmid TCV-GFP Δ CP has been described³. Wild-type (Nb) and *RDR6*-knocked down (NbRDR6i)¹⁶ *N. benthamiana* and *Arabidopsis* plants were inoculated with TCV-GFP Δ CP as described^{3,37}. Plants were maintained in an insect-free glasshouse at 25°C with supplementary lighting to give a 16-hour photoperiod. To measure the accumulation of viral RNA, total RNAs were extracted from tissues using the RNeasy Plant Minikit (Qiagen) as described¹⁷ and assayed by Quantitative RT-PCR (qRT-PCR).

Local RNA silencing assay. Two-to-four seedlings of *gfp*-expressing transgenic lines of *N. benthamiana* (GFP16c and GFP16c/*RDR6*i)¹⁶ were mechanically inoculated with RNA produced by *in vitro* transcription from linearised TCV-GFP Δ CP plasmid DNA in three separate experiments. The induction of intercellular *gfp* RNA silencing was examined under long-wavelength UV light and recorded photographically through a Kodak No. 58 filter using a Nikon Digital Camera Coolpix995 as described¹⁴. Regions of leaf lamina in which silencing of *gfp* RNA occurred show only red chlorophyll fluorescence, while tissues expressing GFP show green fluorescence.

qRT-PCR assay. Virus or mock-inoculated leaves of 3 plants (Nb, NbRDR6i, GFP16c and GFP16c/*RDR6*i) were separately taken at 12 days post inoculation in repeated



experiments for RNA extraction¹⁷. The first-stranded cDNA was synthesized using total RNAs treated with RNase-free DNase I as templates by the M-MLV Reverse Transcriptase (Promega). The qRT-PCR analyses of viral and *RDR6* mRNA levels were performed using specific primers (Table S3) and the iQTM SYBR Green Supermix. The amplification program for SYBR Green I was performed at 95°C for 10 seconds and 55°C for 30 seconds on the CFX96 machine (Bio-Rad), following the manufacturer's instructions. Triplicate quantitative assays were performed on cDNA of each biological sample. The relative quantification of viral or *RDR6* mRNA was calculated using the formula $2^{-\Delta\Delta Ct}$ and normalized to the amount of *GAPDH* (Genbank accession number TC17509) or *EF1 α* (Genbank accession number AY206006) mRNA detected in the same sample, respectively.

Fluorescence and laser confocal microscopy. Mock- or virus-inoculated leaves were collected, examined under a Zeiss Axiopt microscope through a green filter and photographed with a Nikon Coolpix995 digital camera^{38,39}, or under an LSM 710 Laser Scanning Microscope using settings to visualise GFP green and chlorophyll red fluorescence⁴⁰.

- Shi, Y., Gu, M., Fan, Z. & Hong, Y. RNA silencing suppressors: how viruses fight back. *Future Virol.* **3**, 125–133 (2008).
- Qu, F., Ren, T. & Morris, T. J. The coat protein of turnip crinkle virus suppresses posttranscriptional gene silencing at an early initiation step. *J. Virol.* **77**, 511–522 (2003).
- Ryabov, E. V., van Wezel, R., Walsh, J. & Hong, Y. Cell-to-Cell, but not long-distance spread of RNA silencing that is induced in individual epidermal cells. *J. Virol.* **78**, 3149–3154 (2004).
- Azevedo, J. *et al.* Argonaute quenching and global changes in Dicer homeostasis caused by a pathogen-encoded GW repeat protein. *Genes Dev.* **24**, 904–915 (2010).
- Deleris, A. *et al.* Hierarchical action and inhibition of plant Dicer-like proteins in antiviral defense. *Science* **313**, 68–71 (2006).
- Mérai, Z. *et al.* Double-stranded RNA binding may be a general plant RNA viral strategy to suppress RNA silencing. *J. Virol.* **80**, 5747–5756 (2006).
- Endres, M. W. *et al.* Two plant viral suppressors of silencing require the ethylene-inducible host transcription factor RAV2 to block RNA silencing. *PLoS Pathog.* **6**, e1000729 (2010).
- Melnyk, C. W., Molnar, A. & Baulcombe, D. C. Intercellular and systemic movement of RNA silencing signals. *EMBO J.* **30**, 3553–3563 (2011).
- Dunoyer, P., Himber, C. & Voinnet, O. DICER-LIKE 4 is required for RNA interference and produces the 21-nucleotide small interfering RNA component of the plant cell-to-cell silencing signal. *Nature Genet.* **37**, 1356–1360 (2005).
- Himber, C., Dunoyer, P., Moissiard, G., Ritzenthaler, C. & Voinnet, O. Transitivity-dependent and -independent cell-to-cell movement of RNA silencing. *EMBO J.* **22**, 4523–4533 (2003).
- Dunoyer, P., Himber, C., Ruiz-Ferrer, V., Alioua, A. & Voinnet, O. Intra- and intercellular RNA interference in *Arabidopsis thaliana* requires components of the microRNA and heterochromatic silencing pathways. *Nature Genet.* **39**, 848–856 (2007).
- Dunoyer, P. *et al.* Small RNA duplexes function as mobile silencing signals between plant cells. *Science* **328**, 912–916 (2010).
- Molnar, A. *et al.* Small silencing RNAs in plants are mobile and direct epigenetic modification in recipient cells. *Science* **328**, 872–875 (2010).
- Zhou, Y., Ryabov, E., Zhang, X. & Hong, Y. Influence of viral genes on the cell-to-cell spread of RNA silencing. *J. Exp. Bot.* **59**, 2803–2813 (2008).
- Vogler, H. *et al.* Tobacco mosaic virus movement protein enhances the spread of RNA silencing. *PLoS Pathog.* **4**, e1000038 (2008).
- Schwach, F., Vaistij, F. E., Jones, L. & Baulcombe, D. C. An RNA-dependent RNA polymerase prevents meristem invasion by potato virus X and is required for the activity but not the production of a systemic silencing signal. *Plant Physiol.* **138**, 1842–1852 (2005).
- Shi, Y. *et al.* Suppression of local RNA silencing is not sufficient to promote cell-to-cell movement of Turnip crinkle virus in *Nicotiana benthamiana*. *Plant Signal. Behav.* **4**, 15–22 (2009).
- van Wezel, R. & Hong, Y. Virus survival of RNA silencing without deploying protein-mediated suppression in *Nicotiana benthamiana*. *FEBS Lett.* **562**, 65–70 (2004).
- Dalmay, T., Hamilton, A., Rudd, S., Angell, S. & Baulcombe, D. C. An RNA-dependent RNA polymerase gene in Arabidopsis is required for posttranscriptional gene silencing mediated by a transgene but not by a virus. *Cell* **101**, 543–553 (2000).
- Xie, Z., Fan, B., Chen, C. & Chen, Z. An important role of an inducible RNA-dependent RNA polymerase in plant antiviral defense. *Proc. Natl. Acad. Sci. USA.* **98**, 6516–6521 (2001).
- Xie, Z. *et al.* Genetic and functional diversification of small RNA pathways in plants. *PLoS Biol.* **2**, E104 (2004).
- Muangsan, N., Beclin, C., Vaucheret, H. & Robertson, D. Geminivirus VIGS of endogenous genes requires SGS2/SDE1 and SGS3 and defines a new branch in the genetic pathway for silencing in plants. *Plant J.* **38**, 1004–1014 (2004).
- Qu, F. *et al.* RDR6 has a broad-spectrum but temperature-dependent antiviral defense role in *Nicotiana benthamiana*. *J. Virol.* **79**, 15209–15217 (2005).

- García-Ruiz, H. *et al.* Arabidopsis RNA-dependent RNA polymerases and Dicer-Like proteins in antiviral defense and small interfering RNA biogenesis during Turnip mosaic virus infection. *Plant Cell* **22**, 481–496 (2010).
- Rakshandehroo, F., Takeshita, M., Squires, J. & Palukaitis, P. The Influence of RNA-Dependent RNA Polymerase 1 on *Potato virus Y* Infection and on Other Antiviral Response Genes. *Mol. Plant-Microbe Interact.* **22**, 1312–1318 (2009).
- Yang, S.-J., Carter, S. A., Cole, A. B., Cheng, N.-H. & Nelson, R. S. A natural variant of a host RNA-dependent RNA polymerase is associated with increased susceptibility to viruses by *Nicotiana benthamiana*. *Proc. Natl. Acad. Sci. USA.* **101**, 6297–6302 (2004).
- Ying, X.-B. *et al.* RNA-dependent RNA polymerase 1 from *Nicotiana tabacum* suppresses RNA silencing and enhances viral infection in *Nicotiana benthamiana*. *Plant Cell* **22**, 1358–1372 (2010).
- Qi, X., Bao, F. S. & Xie, Z. Small RNA deep sequencing reveals role for *Arabidopsis thaliana* RNA-dependent RNA polymerases in viral siRNA biogenesis. *PLoS ONE* **4**, e4971 (2009).
- Dalmay, T., Horsefield, R., Braunstein, T. H. & Baulcombe, D. C. SDE3 encodes an RNA helicase required for post-transcriptional gene silencing in *Arabidopsis*. *EMBO J.* **20**, 2069–2078 (2001).
- Mourrain, P. *et al.* Arabidopsis SGS2 and SGS3 genes are required for posttranscriptional gene silencing and natural virus resistance. *Cell* **101**, 533–542 (2000).
- Vaistij, P. E., Jones, L. & Baulcombe, D. C. Spreading of RNA targeting and DNA methylation in RNA silencing requires transcription of the target gene and a putative RNA-dependent RNA polymerase. *Plant Cell* **14**, 857–867 (2002).
- Brosnan, C. A. *et al.* Nuclear gene silencing directs reception of long-distance mRNA silencing in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA.* **104**, 14741–14746 (2007).
- Curaba, J. & Chen, X. Biochemical activities of Arabidopsis RNA-dependent RNA polymerase 6. *J. Biol. Chem.* **283**, 3059–3066 (2008).
- Hoffer, P. *et al.* Posttranscriptional gene silencing in nuclei. *Proc. Natl. Acad. Sci. USA.* **108**, 409–414 (2011).
- Luo, Z. & Chen, Z. Improperly terminated, unpolyadenylated mRNA of sense transgenes is targeted by RDR6-mediated RNA silencing in *Arabidopsis*. *Plant Cell.* **19**, 943–958 (2007).
- Cohen, Y., Gisel, A. & Zambryski, P. C. Cell-to-cell and systemic movement of recombinant green fluorescent protein-tagged turnip crinkle viruses. *Virology* **273**, 258–266 (2000).
- Li, C. *et al.* A cis element within flowering locus T mRNA determines its mobility and facilitates trafficking of heterologous viral RNA. *J. Virol.* **83**, 3540–3548 (2009).
- Hong, Y., Stanley, J. & Van Wezel, R. Novel system for the simultaneous analysis of geminivirus DNA replication and plant interactions in *Nicotiana benthamiana*. *J. Virol.* **77**, 13315–13322 (2003).
- Dong, X., Van Wezel, R., Stanley, J. & Hong, Y. Functional characterization of the nuclear localization signal for a suppressor of posttranscriptional gene silencing. *J. Virol.* **77**, 7026–7033 (2003).
- Li, C. *et al.* Mobile FT mRNA contributes to the systemic florigen signalling in floral induction. *Sci. Rep.* **1**, 73; DOI:10.1038/srep00073 (2011).

Acknowledgements

We thank David Baulcombe for providing seed of transgenic lines GFP16c, GFP16c/RDR6i and NbRDR6i, and critical comments on the original manuscript, Andy Maule and Bryan Harrison for constructive suggestions, and Michael Wilson for critical reading of the manuscript. We are grateful to John Carr and Jack Westwood for providing the qRT-PCR protocol. We thank the Special Foundation for Hangzhou Talents, the Science Foundation of the Key Laboratory of Hangzhou (20090232T05) and the China Natural Science Foundation (NSFC30770185, 30870180) for grants and a scholarship to N.S. and the Ministry of Higher Education of Egypt for a PhD studentship to A.M. H.Z. was supported by an R&D grant from Warwick Ventures to Y.H. The work is also in part supported by an innovative R&D grant from Hangzhou Normal University to Y.H.

Author contributions

C.Q., N.S., M.G., H.Z., B.L., J.S. and A.M. designed and performed experiments, C.L. and E.R. performed researches, H.W., Y.L., T.O. and M.V. contributed through discussions and revised the paper. Y.H. initiated the project, designed the experiments and wrote the paper.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/scientificreports>

Competing financial interests: The authors declare no competing financial interests.

License: This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>

How to cite this article: Qin, C. *et al.* Involvement of RDR6 in short-range intercellular RNA silencing in *Nicotiana benthamiana*. *Sci. Rep.* **2**, 467; DOI:10.1038/srep00467 (2012).