

Original citation:

Stonehewer, Stewart E. (Stewart Edward), 1935-. (2012) Generalising quasinormal subgroups. Bulletin of the Australian Mathematical Society , Volume 86 (Number 1). pp. 1-10. ISSN 0004-9727

Permanent WRAP url:

<http://wrap.warwick.ac.uk/52614/>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Copyright © Australian Mathematical Publishing Association Inc. 2012

A note on versions:

The version presented in WRAP is the published version or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: publications@warwick.ac.uk



<http://wrap.warwick.ac.uk/>

GENERALISING QUASINORMAL SUBGROUPS

STEWART STONEHEWER

(Received 16 September 2011)

Dedicated to John Cossey on the occasion of his seventieth birthday

Abstract

In Cossey and Stonehewer [‘On the rarity of quasinormal subgroups’, *Rend. Semin. Mat. Univ. Padova* **125** (2011), 81–105] it is shown that for any odd prime p and integer $n \geq 3$, there is a finite p -group G of exponent p^n containing a quasinormal subgroup H of exponent p^{n-1} such that the nontrivial quasinormal subgroups of G lying in H can have exponent only p , p^{n-1} or, when $n \geq 4$, p^{n-2} . Thus large sections of these groups are devoid of quasinormal subgroups. The authors ask in that paper if there is a nontrivial subgroup-theoretic property \mathfrak{X} of finite p -groups such that (i) \mathfrak{X} is invariant under subgroup lattice isomorphisms and (ii) every chain of \mathfrak{X} -subgroups of a finite p -group can be refined to a composition series of \mathfrak{X} -subgroups. Failing this, can such a chain always be refined to a series of \mathfrak{X} -subgroups in which the intervals between adjacent terms are restricted in some significant way? The present work embarks upon this quest.

2010 *Mathematics subject classification*: primary 20E07; secondary 20E15.

Keywords and phrases: generalising quasinormality in finite p -groups.

1. Introduction

A subgroup Q of a group G is said to be *quasinormal* (sometimes *permutable*) if $\langle Q, H \rangle = QH$ for all subgroups H of G ; that is, if $QH = HQ$. We write $Q \text{ qn } G$. The concept is due to Ore [3], and in [4] he proved that, in finite groups, quasinormal subgroups are always subnormal. He also proved that they are *modular*. Recall that a subgroup M of a group G is *modular* if

$$\langle X, M \rangle \cap Y = \langle X, M \cap Y \rangle \quad \text{for all } X \leq Y \quad (1.1)$$

and

$$\langle X, M \rangle \cap Y = \langle X \cap Y, M \rangle \quad \text{for all } X \text{ and } Y \text{ with } M \leq Y. \quad (1.2)$$

In fact a subgroup M of a finite group G is quasinormal if and only if M is modular and subnormal in G (see [6, Theorem 5.1.1]). So in finite p -groups,

quasinormal subgroups and modular subgroups coincide.

Also they are clearly invariant under subgroup lattice isomorphisms (that is, *projectivities*).

It was explained in [1] why p -groups are so important in the theory of quasinormal subgroups of finite groups. For, suppose that G is finite with Q qn G and Q core-free. Then Q is nilpotent and each Sylow p -subgroup P of Q is quasinormal in G . Also if S is a Sylow p -subgroup G , then P qn S and the complexities of the embedding of Q in G are reduced to those of P in S . Therefore in what follows we shall be mainly interested in finite p -groups. Our aim is to weaken the definition of quasinormality, while retaining its invariance under projectivities in finite p -groups, so as to produce a class of subgroups that are more involved with the structure of the group. We are particularly interested in relating group and lattice properties. An obvious starting point is to adopt just one of the definitions (1.1) and (1.2) of modularity.

DEFINITION 1.1. We say that a subgroup M of a group G is *semimodular* if (1.1) holds (see [6, Exercise 7, p. 49]). Write M sm G .

We should point out (and indeed the referee drew attention to the fact) that *semimodularity of lattices* is a standard concept, namely $X \cap Y$ maximal in X implies Y maximal in $\langle X, Y \rangle$ for all X and Y (see, for example, [7]). However, since we are concerned with individual elements rather than whole lattices, we have taken the liberty (following Schmidt in [6]) of using this terminology, which, while different from the original, will hopefully seem appropriate in the context. (The same applies in Definitions 1.4, 2.1 and 2.2 below.)

Given subgroups $H \leq K$ of a group, denote the lattice of subgroups between H and K by $[K/H]$. Then for subgroups X, Y of a group G , define a map

$$\Phi_{X,Y} : [X/X \cap Y] \rightarrow [\langle X, Y \rangle / Y]$$

by

$$H \mapsto \langle H, Y \rangle.$$

Similarly, define a map

$$\Psi_{X,Y} : [\langle X, Y \rangle / Y] \rightarrow [X/X \cap Y]$$

by

$$K \mapsto X \cap K.$$

In relation to semimodularity we have the following proposition.

PROPOSITION 1.2. *Let M be a subgroup of G . Then:*

(i) M sm G if and only if, for all $X \leq G$,

$$\Phi_{X,M} \Psi_{X,M} = \text{id}_{[X/X \cap M]};$$

(ii) if M sm G , then, for all $X \leq G$,

$$\Psi_{M,X} \Phi_{M,X} = \text{id}_{[\langle X, M \rangle / X]}.$$

Both of these results are left to the reader in [6, Exercise 7, p. 49], and we do the same. However, when G is a finite p -group there is a striking corollary.

PROPOSITION 1.3. *A semimodular subgroup of a finite p -group is quasinormal and therefore modular.*

PROOF. Let G be a finite p -group with M sm G and let $X \leq G$. Then it follows from Proposition 1.2 that the map

$$\Psi_{M,X} : [\langle X, M \rangle / X] \rightarrow [M / X \cap M]$$

is injective. Therefore $|\langle X, M \rangle : X| \leq |M : X \cap M|$. However,

$$|M : X \cap M| = |M| / |X \cap M| = |XM| / |X|,$$

that is, $|\langle X, M \rangle| \leq |XM|$. Hence $\langle X, M \rangle = XM$ and so M qn G . \square

So (1.2) is a redundant requirement in the definition of modular subgroups in finite p -groups. But of course we cannot remove the hypothesis that G is a finite p -group in Proposition 1.3. For example, the subgroups of order two in the alternating group A_4 are semimodular, but not modular. As we stated above, however, our interest is in p -groups.

We consider next the defining property (1.2).

DEFINITION 1.4. We say that a subgroup M of a group G is *weakly modular* if (1.2) holds. Write M wm G .

Corresponding to Proposition 1.2(i) we have the following proposition.

PROPOSITION 1.5. *Let $M \leq G$. Then M wm G if and only if, for all $X \leq G$,*

$$\Psi_{X,M} \Phi_{X,M} = \text{id}_{[\langle X, M \rangle / M]}. \quad (1.3)$$

PROOF. Suppose that M wm G , $X \leq G$ and $M \leq Y \leq \langle X, M \rangle$. Then $Y = \langle X, M \rangle \cap Y = \langle X \cap Y, M \rangle$, by (1.2). Therefore $\Psi_{X,M} : Y \mapsto X \cap Y$ and $\Phi_{X,M} : X \cap Y \mapsto \langle X \cap Y, M \rangle = Y$. Hence we have the necessary requirement.

Conversely, suppose that (1.3) holds and let $M \leq Y \leq G$. Then $X \cap Y = X \cap (\langle X, M \rangle \cap Y)$. Therefore

$$\Psi_{X,M} : \langle X, M \rangle \cap Y \mapsto X \cap Y \quad \text{and} \quad \Phi_{X,M} : X \cap Y \mapsto \langle X \cap Y, M \rangle.$$

Thus, by (1.3), $\langle X, M \rangle \cap Y = \langle X \cap Y, M \rangle$ and so M wm G . \square

Again we have a striking corollary.

PROPOSITION 1.6. *A weakly modular subgroup of a finite p -group is quasinormal and therefore modular.*

PROOF. Let G be a finite p -group with M w.m. G and let X be a cyclic subgroup of G . It is sufficient to show that $\langle X, M \rangle = XM$. By Proposition 1.5, $\Phi_{X,M}$ is a surjective map. Therefore since $[X/X \cap M]$ is a chain, it follows that $[\langle X, M \rangle/M]$ is a chain and

$$|\langle X, M \rangle : M| \leq |X : X \cap M|.$$

Thus $|XM| = |X||M|/|X \cap M|$ implies that $|\langle X, M \rangle : M| \leq |XM|/|M|$ and hence $|\langle X, M \rangle| \leq |XM|$. Then $\langle X, M \rangle = XM$, as claimed. \square

It follows that *either of the two conditions (1.1) and (1.2) for modularity is sufficient in finite p -groups*. However, as before, we cannot remove the hypothesis that G is a finite p -group in Proposition 1.6. For example, a maximal subgroup of any group is weakly modular.

So two possible lattice-invariant generalisations of the concept of quasinormality in finite p -groups have failed. In the next section we consider a third which at least produces a larger class of subgroups.

2. Upper semimodular subgroups

In [6, pp. 46, 47], Schmidt defines two generalisations of the concept of modular lattice, namely *upper semimodular* and *lower semimodular*. Applied to individual subgroups these are as follows. Let M be a subgroup of a group G .

DEFINITION 2.1. We say that M is *upper semimodular* in G (and we write M usm G) if, for all $X \leq G$,

$$X \cap M \text{ maximal in } M \text{ implies } X \text{ maximal in } \langle X, M \rangle. \quad (2.1)$$

DEFINITION 2.2. We say that M is *lower semimodular* in G if, for all $X \leq G$,

$$X \text{ maximal in } \langle X, M \rangle \text{ implies } X \cap M \text{ maximal in } M.$$

It is easy to see that *in all finite p -groups, every subgroup is lower semimodular*. However, an upper semimodular subgroup of a finite p -group is not always modular, as we see in the following example.

EXAMPLE. Let p be an odd prime and let G be the group generated by elements a_1, a_2, a_3, b of order p such that

$$[a_i, b] = a_{i+1}, i = 1, 2 \quad \text{and} \quad [a_3, b] = [a_i, a_j] = 1, \text{ all } i, j.$$

Let $M = \langle a_1, a_2 \rangle$. We claim that

$$M \text{ usm } G. \quad (2.2)$$

For, let $X \leq G$ with $|X \cap M| = p$ and $X \not\leq M$. We show that $|\langle X, M \rangle : X| = p$. Clearly we may assume that $X \not\leq A = \langle a_1, a_2, a_3 \rangle$. Therefore there is an element $ab \in X$ with $a \in A$. We distinguish two cases.

- (i) Suppose that $X \cap M = \langle a_2 \rangle$. Then $a_3 \in X$ and so $|X| \geq p^3$. Hence $|X| = p^3$ and $G = XM$, that is, X is maximal in G .
- (ii) Suppose that $X \cap M \neq \langle a_2 \rangle$. Then there is an element $a_1 a_2^i \in X$, some i . Thus $a_2 a_3^i \in X$ and $a_3 \in X$. So $a_2 \in X$, a contradiction.

Hence (2.2) follows. However, M is not modular in G , since it is not quasinormal, that is, $M\langle b \rangle$ is not a subgroup.

Of course the property of being an upper semimodular subgroup is invariant under projectivities, that is,

upper semimodularity is a lattice-invariant property.

These subgroups also satisfy another important property, as we now show. We shall refer to them as *usm* subgroups.

PROPOSITION 2.3. *The join of two usm subgroups of a finite p -group is a usm subgroup.*

PROOF. Let G be a finite p -group and M_1, M_2 usm G . Let $M = \langle M_1, M_2 \rangle$ and suppose that $X \leq G$ with $X \cap M$ maximal in M . We write $X \cap M \leq M$. So $|M : X \cap M| = p$. We show that

$$X \triangleleft \langle X, M \rangle. \quad (2.3)$$

Clearly $M \not\leq X$ and so without loss of generality we may assume that $M_1 \not\leq X$. But $\langle X, M_1 \rangle = \langle X, X \cap M, M_1 \rangle = \langle X, M \rangle$ and $X \triangleleft \langle X, M_1 \rangle$ since $X \cap M_1 \leq M_1$. So (2.3) follows. \square

We saw in the above example that usm subgroups of finite p -groups are not quasinormal in general. In that example the subgroup is elementary abelian of rank two. It could not have had order p for the following reason.

PROPOSITION 2.4. *A usm subgroup of order p in a finite p -group is quasinormal.*

PROOF. Let G be a finite p -group with M usm G and $|M| = p$. Let $X \leq G$. Then

$$\langle X, M \rangle = XM. \quad (2.4)$$

For we may assume that $M \not\leq X$. Thus $M \cap X = 1$ and so by hypothesis $|\langle X, M \rangle : X| = p$. Hence $X \triangleleft \langle X, M \rangle$ and (2.4) follows. \square

In fact the above result is true even for cyclic usm subgroups. But in order to prove this we need two elementary properties.

PROPOSITION 2.5. *Let M be a usm subgroup of a group G .*

- (i) *If $M \leq H \leq G$, then M usm H .*
- (ii) *If θ is a homomorphism of G , then $M\theta$ usm $G\theta$.*

PROOF. (i) This is clear.

(ii) Let $N \triangleleft G$. Then it suffices to show that $MN/N \text{ usm } G/N$. Let $X/N \leq G/N$ such that $X \cap MN \triangleleft MN$. We have

$$MN/N \cong M/(N \cap M)$$

and under this isomorphism $(X \cap M)N/N = (X \cap MN)/N$ corresponds to $(X \cap M)/(N \cap M)$. Therefore $X \cap M \triangleleft M$ and so $X \triangleleft \langle X, M \rangle$, by hypothesis, that is, $X/N \triangleleft \langle X/N, MN/N \rangle$. Hence $MN/N \text{ usm } G/N$. \square

Now we can show that Proposition 2.4 is just the simplest case of a more general result. We denote the subgroup generated by the elements of order p in a p -group G by $\Omega_1(G)$.

THEOREM 2.6. *A cyclic usm subgroup of a finite p -group is quasinormal.*

PROOF. Let G be a finite p -group with $M \text{ usm } G$ and M cyclic. Suppose for a contradiction that M is not quasinormal in G and let G be a counterexample of minimal order. Then there is a cyclic subgroup X of G such that $XM \neq MX$. Also $G = \langle X, M \rangle$, by Proposition 2.5(i) and choice of G .

Let N be a minimal normal subgroup of G . Then again by choice of G and Proposition 2.5(ii), $G = NXM$. We claim that

$$N = Z_1(G), \tag{2.5}$$

the centre of G . For otherwise there exists an element $xy \in Z_1(G)$, $xy \neq 1$, $x \in X$, $y \in M$. Then $[x, y] = 1$ and so $\langle xy \rangle \subset XM$. But $G = \langle xy \rangle XM$, since $G/\langle xy \rangle$ is not a counterexample. Therefore $G = X\langle xy \rangle M = XM$, contradiction. Thus (2.5) is true.

By [6, Theorem 5.2.13], G/N is metacyclic, being the product of two cyclic groups. Therefore there are cyclic subgroups H, K of G such that

$$NH \triangleleft G = NHK.$$

Let $H = \langle h \rangle$, $K = \langle k \rangle$. So we may assume that

$$h^k = uh^{1+\lambda p}, \tag{2.6}$$

where $u \in N$, λ an integer. Suppose that $|H| = p$. If also $|K| = p$, then $|G| = p^3$. If p is odd, G must have exponent p . But then $|M| = p$, contradicting Proposition 2.4. On the other hand, if $p = 2$, then $G \cong D_8$, the dihedral group of order eight, with M of order four, again by Proposition 2.4. So $M \triangleleft G$, a contradiction. Thus we must have $|K| \geq p^2$. Then $K^p \leq Z_1(G)$, by (2.6), and so $K^p = N$, by (2.5). Therefore again $|G| = p^3$ and we easily obtain contradictions. Thus

$$|H| \geq p^2.$$

It follows from (2.6) that $\Omega_1(H) \leq Z_1(G)$ and so $\Omega_1(H) = N$ and $G = HK$, with $H \triangleleft G$ and H and G/H cyclic. If p were odd, then all its subgroups would be

quasinormal, by [6, Theorem 2.3.1]. So $p = 2$. We shall see in the next lemma that M , not being quasinormal in G , cannot lie in the subgroup of G of index two which contains H . But then we have $G = HM$. Also G/H^4 has a quotient isomorphic to D_8 in which the image of M is noncentral of order two (again by [6, Theorem 2.3.1]). However, this contradicts Propositions 2.4 and 2.5(ii) and completes the proof of the theorem. \square

LEMMA 2.7. *Let G be a finite 2-group with $H \triangleleft G$, H cyclic of order at least four, and let G/H be cyclic and nontrivial. Let L/H be the subgroup of index two in G/H . Then every subgroup of L is quasinormal in G .*

PROOF. Suppose that the lemma is false and let G be a counterexample of minimal order. Let $M \leq L$ with M not quasinormal in G . Then $H \cap M = 1$, otherwise $M/H \cap M$ is quasinormal in $G/H \cap M$, by choice of G , and thus $M \text{ qn } G$, a contradiction. Therefore M is cyclic. Let X be a cyclic subgroup of G such that

$$\langle M, X \rangle \neq MX. \quad (2.7)$$

Again by [6, Theorem 2.3.1], we see that L is a modular group, that is, all its subgroups are modular, therefore quasinormal (in L). Thus $X \not\leq L$ and so $G = HX$. Also we must have $H \cap X = 1$, otherwise $MX/H \cap X$ is a subgroup by choice of G .

Let $M = \langle hx \rangle$, $h \in H$, $x \in X^2$. Then $h \in MX$. Also $h \neq 1$. Now

$$(hx)^2 = hh^{1+4\lambda}x^2,$$

for some integer λ . Squaring this equation repeatedly, we see that some power h_1x_1 ($h_1 \in H$, $x_1 \in X^2$) of hx has $|h_1| = 2$. Thus $\langle h_1 \rangle \in MX \cap Z_1(G)$ and again, by choice of G , $MX/\langle h_1 \rangle$ is a subgroup of $G/\langle h_1 \rangle$. This contradicts (2.7) and so the lemma follows. \square

The above results suggest that the concept of upper semimodularity might be a useful lattice-invariant generalisation of quasinormality in finite p -groups, with a view to understanding more about the connection between group and subgroup-lattice structures. But it turns out that there are two properties of quasinormal subgroups, one extremely useful and the other quite intriguing, neither of which is inherited by usm subgroups. The first concerns the intersection of a quasinormal subgroup Q with an arbitrary subgroup H in a group. The intersection $Q \cap H$ is always quasinormal in H . However, this property is not possessed by usm subgroups. To see this, let G be the group of the previous example. Then $M = \langle a_1, a_2 \rangle \text{ usm } G$. Let $H = \langle a_2, a_3, b \rangle$, so $H \cap M = \langle a_2 \rangle$. Clearly $\langle a_2 \rangle$ is not quasinormal in H . Therefore $\langle a_2 \rangle$ is not a usm subgroup of H , by Proposition 2.4.

In fact the same example illustrates the second property of quasinormal subgroups referred to above. It was shown in [2] that *for odd primes p , a quasinormal subgroup of order p^2 in a finite p -group G always contains a quasinormal subgroup (of G) of order p* . However, it is easy to see that in the example above, where M has order p^2 ,

there is no usm subgroup of G of order p lying in M . For, such a subgroup would have to be quasinormal in G , by Proposition 2.4. Thus upper semimodularity is not a candidate for the class \mathfrak{X} referred to at the beginning of this paper.

There is yet a third property of quasinormal subgroups that is not inherited by usm subgroups. A subgroup M that permutes with all cyclic subgroups is quasinormal. But again take G in our example, with $p \geq 5$, and let $M = \langle a_1, a_3 \rangle$. Since G has exponent p , each nontrivial cyclic subgroup X of G with $X \cap M \triangleleft M$ lies in M and so (2.1) holds. But if M usm G , then modulo $\langle a_3 \rangle$, M would be quasinormal in G (by Propositions 2.4 and 2.5(ii)). Thus M would be quasinormal in G , which is not the case.

3. Joins as restricted products

We conclude this preliminary investigation by considering another concept generalising quasinormality. By definition, H qn G if and only if $\langle H, K \rangle = HK$ for all $K \leq G$. Of course when G is finite, for any two subgroups H and K of G , we always have

$$\langle H, K \rangle = (HK)^n$$

for some finite n . (Here for sets X, Y , $XY = \{xy \mid x \in X, y \in Y\}$.) In this connection there is an interesting elementary result.

PROPOSITION 3.1. *Let G be a finite p -group with subgroups H, K such that $G = HKH$. Then $G = HK$.*

PROOF. We proceed by induction and assume the result is true for groups of order less than $|G|$. Let h_1kh_2 be a central element of order p , with $h_1, h_2 \in H, k \in K$. Thus h_1kh_2 commutes with kh_2 and so h_1 commutes with kh_2 . Therefore $h_1kh_2 = kh_2h_1$. Similarly, h_2 commutes with h_1k . Put $h = h_2h_1$. Then

$$hk = h_2h_1k = h_1kh_2 = kh_2h_1 = kh.$$

Thus $N = \langle hk \rangle \subset HK$ and $N \leq Z_1(G)$. By induction,

$$G/N = HKN/N = HNK/N = HK/N$$

and hence $G = HK$ as required. \square

For our final generalisation of quasinormality, we shall use the fact that a subgroup is quasinormal if and only if its join with every cyclic subgroup is just its product.

DEFINITION 3.2. We call a subgroup H of a group G 4-*quasinormal*, and write H qn₄ G , if

$$\langle H, K \rangle = HKHK,$$

for all cyclic subgroups K of G .

Clearly this concept can be generalised successively to n -quasinormal by increasing the number of factors H, K . In this way, eventually every subgroup H of every finite group will be included.

We shall restrict ourselves to finite p -groups. Given the complexity of these groups, the difficulties of discovering the properties of even 4-quasinormal subgroups will surely be considerable. However, we have been able to make a start when $|H| = p$ by establishing their lattice-invariance.

THEOREM 3.3. *Let p be a prime and let G, \bar{G} be lattice-isomorphic finite p -groups. Suppose that $H \text{ qn}_4 G$ and $|H| = p$. Denote the image of H in \bar{G} by \bar{H} . Then $\bar{H} \text{ qn}_4 \bar{G}$.*

PROOF. We have to show that for every cyclic subgroup \bar{K} of \bar{G} ,

$$\langle \bar{H}, \bar{K} \rangle = \bar{H}\bar{K}\bar{H}\bar{K}.$$

Of course the pre-image K of \bar{K} is cyclic of the same order as \bar{K} , say p^n , $n \geq 1$. Thus we may assume that

$$G = \langle H, K \rangle = HKHK \quad (3.1)$$

and $\bar{G} = \langle \bar{H}, \bar{K} \rangle$ having the same order as G . Let $N = K_G$, the core of K in G . Then (continuing to use bars to denote images) $\bar{N} \text{ qn } \bar{G}$ and so \bar{H} normalises \bar{N} . Therefore $\bar{N} \triangleleft \bar{G}$ and clearly $\bar{N} = \bar{K}_{\bar{G}}$. Since 4-quasinormality is obviously preserved by homomorphisms, we may assume that

$$K_G = 1 \quad \text{and} \quad \bar{K}_{\bar{G}} = 1. \quad (3.2)$$

From (3.1), we see that $|\bar{G}| = |G| \leq p^{2n+1}$. Let $\bar{H} = \langle h \rangle$, $\bar{K} = \langle k \rangle$. Then

$$\text{the double cosets } \bar{H}k^i h \bar{K}, \quad 0 \leq i \leq p^n - 1, \text{ are distinct.} \quad (3.3)$$

For suppose that $\bar{H}k^i h \bar{K} = \bar{H}k^j h \bar{K}$ with $i \neq j$, $0 \leq i, j \leq p^n - 1$. Then

$$k^i h = h_1 k^j h k_1, \quad (3.4)$$

where $h_1 \in \bar{H}$, $k_1 \in \bar{K}$. Therefore $k^i = h_1 k^j k_1 [k_1, h^{-1}]$. It follows from (3.4) that $k_1 \neq 1$. So $[k_1, h^{-1}] \neq 1$, by (3.2), and hence $h_1^{-1} k^{i-j} k_1^{-1} \in \bar{G}' \setminus \{1\}$. Since $\bar{G} = \langle h, k \rangle$, we must have $h_1 = 1$, otherwise $\bar{G} = \langle h_1^{-1} k^{i-j} k_1^{-1}, k \rangle = \langle k \rangle$, a contradiction. Thus, by (3.4),

$$k_1^{h^{-1}} \in \bar{K}.$$

But this contradicts (3.2). Therefore (3.3) follows.

Finally, for $g \in \bar{G}$,

$$|\bar{H}g\bar{K}| = |\bar{H}||\bar{K}|/|\bar{H}^g \cap \bar{K}| = p^{n+1}$$

(see [5, p. 90, Exercise 247]). Thus from (3.3) we see that $|\bar{H}\bar{K}\bar{H}\bar{K}| \geq p^{2n+1}$. Therefore $\bar{G} = \bar{H}\bar{K}\bar{H}\bar{K}$. \square

An obvious question is whether Theorem 3.3 extends to arbitrary cyclic qn_4 -subgroups or indeed to all qn_4 -subgroups, and we have made some progress in this direction. Also, which other properties of quasinormal subgroups are inherited by, or can be generalised to, qn_4 -subgroups? We make no conjectures, but hopefully more results will appear in due course.

References

- [1] J. Cossey and S. E. Stonehewer, 'On the rarity of quasinormal subgroups', *Rend. Semin. Mat. Univ. Padova* **125** (2011), 81–105.
- [2] J. Cossey, S. E. Stonehewer and G. Zacher, 'Quasinormal subgroups of order p^2 ', *Ric. Mat.* **57** (2008), 127–135.
- [3] O. Ore, 'Structures and group theory I', *Duke Math. J.* **3** (1937), 149–173.
- [4] O. Ore, 'On the application of structure theory to groups', *Bull. Amer. Math. Soc.* **44** (1938), 801–806.
- [5] J. S. Rose, *A Course on Group Theory* (Cambridge University Press, Cambridge, 1978).
- [6] R. Schmidt, *Subgroup Lattices of Groups* (de Gruyter, Berlin, 1994).
- [7] M. Stern, *Semimodular Lattices* (Cambridge University Press, Cambridge, 1999).

STEWART STONEHEWER, Mathematics Institute, University of Warwick,
Coventry CV4 7AL, UK

e-mail: S.E.Stonehewer@warwick.ac.uk