

Original citation:

Novak, L. A. and Gibbons, A. M. (1989) Double independent subsets of a graph. University of Warwick. Department of Computer Science. (Department of Computer Science Research Report). (Unpublished) CS-RR-152

Permanent WRAP url:

<http://wrap.warwick.ac.uk/60846>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

A note on versions:

The version presented in WRAP is the published version or, version of record, and may be cited as it appears here. For more information, please contact the WRAP Team at: publications@warwick.ac.uk



<http://wrap.warwick.ac.uk/>

_____Research report 152_____

DOUBLE INDEPENDENT SUBSETS OF A GRAPH

Ladislav Novak*
Alan Gibbons

(RR152)

Department of Computer Science
University of Warwick
Coventry CV4 7AL
United Kingdom

*Department of Electrical Engineering
University of Novi Sad
Yugoslavia

November 1989

Double Independent Subsets of a Graph

Ladislav Novak

Department of Electrical Engineering, University of Novi Sad, Yugoslavia

and

Alan Gibbons

Department of Computer Science, University of Warwick, England

§1 Introduction

It was pointed out by Amari [1] that the set of edges of a graph can be divided into two distinct subsets such that the sum of the rank of one subset and the corank of the other may give a number which is less than both the rank and the corank of the graph. Not long afterwards, this number was formally defined by Ohtsuki, Ishizahi and Watanabe and called the hybrid rank [2]. In the same paper the notion of a minimal hybrid rank taken over all possible partitions of the edge set of a graph was introduced and called the topological degree of freedom. That paper together with the paper of Kishi and Kajitani [3] in which maximally distant pairs of trees and principal partition were introduced provide the foundation of the so-called hybrid approach in graph theory. Since 1967 many papers were published in this area, mostly by Japanese authors [8]-[22].

The appearance of new concepts can be a spurious process and it may not at first be clear which are of value and whether the most useful definitions have been made. In this paper we introduce a concept, that of **double independent subsets** of a graph. This concept, although it has never been given a specific name, has featured from the beginning of the hybrid approach in graph theory. In matroid theory it is called matroid intersection [6]. The concept of double independent subset inherently has a hybrid flavour. In a previous paper [7] we introduced a concept, that of a **perfect pair of trees** of a graph which is closely related to the concept of double independent subset. We hope that both concepts provide excellent intuitive insight within this area of study. Throughout this paper we shall be concerned only with 2-connected graphs.

§2 Preliminaries

This section is devoted to some definitions and assertions related to material that follows. We presume that the reader is familiar with the following basic notions in graph theory : graph, edge, circuit and cutset. We take these to be primary notions that need not be defined. However we will define all other notions on the basis of these. Throughout we denote a graph by G and its edge set by E . The terms circuit, cutset, tree, cotree, forest and coforest will be used here to mean a subset of edges of a graph. A forest is a maximal circuitless subset of edges while a coforest is a maximal cutsetless subset of edges. If the graph is connected then a forest is a tree and a coforest is a cotree. In what follows, a tree will be denoted by t and a cotree by t^* . Given a tree t , any edge in the corresponding cotree t^* forms exactly one circuit with edges in t . Such a circuit is called a **fundamental circuit** of G with respect to t . Similarly, any edge of the tree t defines exactly one cutset with the edges in the corresponding cotree t^* . Such a cutset is called a **fundamental cutset** of G with respect to t^* . If E' is a subset of E then the **rank** of E' is the cardinality of the largest circuitless subset of E' , the **co-rank** of E' is the cardinality of the largest cutsetless subset of E' and the **complement** of E is the set difference $E \setminus E'$ denoted by E^* . By $|E'|$ we denote the number of elements in (that is, the cardinality of) the subset E' .

The **distance** [4] between two trees t_1 and t_2 of a graph, written $|t_1 \setminus t_2|$, is the number of edges which are in t_1 but not in t_2 . A tree t_2 is said to be **maximally distant from** another tree t_1 [5] if $|t_1 \setminus t_2| \geq |t_1 \setminus t|$ for every tree t of G . A pair of trees (t_1, t_2) is defined to be a **perfect pair** of trees [7] if both t_2 is maximally distant from t_1 and t_1 is maximally distant from t_2 .

Assertion 1 [5]

Given a tree t_0 of a graph G , $(\forall t) |t_0 \setminus t| \leq \text{rank } t_0^*$

Assertion 2 [7]

The following five statements are equivalent:

- i) t_2 is maximally distant from t_1
- ii) the fundamental circuit with respect to t_2 defined by an edge in $t_1^* \cap t_2^*$ contains no edges in $t_1 \cap t_2$.
- iii) the fundamental cutset with respect to t_1^* defined by an edge in $t_1 \cap t_2$ contains no edges in $t_1^* \cap t_2^*$.
- iv) $|t_1 \setminus t_2| = \text{rank } t_1^*$.
- v) the number of edges in $t_1 \cap t_2$ is equal to the maximal number of independent cutsets of the graph that belong entirely to the tree t_1 .

Assertion 3 (theorem 1 of [7])

The following five statements are equivalent:

- i) (t_1, t_2) is a perfect pair
- ii) fundamental circuits with respect to t_1 and t_2 defined by edges in $t_1^* \cap t_2^*$ contains no edges in $t_1 \cap t_2$
- iii) fundamental cutsets with respect to t_1^* and t_2^* defined by edges in $t_1 \cap t_2$ contains no edges in $t_1^* \cap t_2^*$
- iv) $\text{rank } t_1^* = |t_1 \setminus t_2| = |t_2 \setminus t_1| = \text{rank } t_2^*$
- v) the following three numbers, associated with the pair of trees (t_1, t_2) are equal:
 - the maximal number of independent cutsets of the graph that belong to t_1
 - the maximal number of independent cutsets of the graph, that belong to t_2
 - the number of common edges in t_1 and t_2 .

§3 Double independent subsets

A subset of edges of a graph G is said to be a **double independent** subset if it contains no circuits and no cutsets of the graph G

According to the preceding definition, we can consider a double independent subset to be a subset of a tree that does not contain cutsets of the graph or (in dual fashion) as a subset of a cotree that does not contain circuits of the graph.

Remark 1

Because a double independent subset does not contain cutsets, removing all the edges of a double independent subset from a graph the rank of the graph remains the same.

Assertion 4

A subset of edges of a graph is a double independent subset iff it can be represented as a set difference of a pair of trees of G .

Proof

\Rightarrow Let d be a double independent subset of a graph G and let t_1 be a tree that contains a double independent subset d . The subgraph G' , obtained by removing all edges of d from G , has the same rank as G (Remark 1). Hence any tree t_2 of the subgraph G' is a tree of G . Therefore $d = t_1 \setminus t_2$.

\Leftarrow Let (t_1, t_2) be a pair of trees of a graph G . Then, $t_1 \setminus t_2$ is a subset of both t_1 and t_2^* . Therefore, $t_1 \setminus t_2$ is a double independent subset of G . □




Figure 1

A double independent subset d of edges of a graph G is a **maximally double independent** if, for an arbitrary edge e in the complement of d , $d \cup \{e\}$ is not a double independent subset of G .

Figure 1 shows six copies of a graph and for each copy a different subset of edges is indicated by the use of bold edges. The subset of edges d_1 is a double independent subset and so is d_2 . However, d_1 is not a maximally double independent whereas d_2 is. Notice also that $d_1 = t_1 \setminus t_2$ and that $d_2 = t_3 \setminus t_4$, where t_1, t_2, t_3 and t_4 are all trees of the graph G .

Assertion 5

A double independent subset d of edges of a graph is a maximally double independent iff every edge in the complement of d form a circuit or/and a cutset with the edges in d only.

Proof

\Rightarrow Given a maximally double independent subset d of G , suppose that there exists an edge e in the complement of d such that for every circuit C_e that contains e , $C_e \setminus (d \cup \{e\})$ is nonempty, and for every cutset S_e that contains e , $S_e \setminus (d \cup \{e\})$ is nonempty. Consequently, $d \cup \{e\}$ is also a maximally double independent subset which contradict the assumption that d is a maximally double independent subset of G .

\Leftarrow Suppose that, given a double independent subset d of edges of a graph, every edge in the complement of d forms a circuit or/and a cutset with edges in d only. Then, for every edge e in the complement of d , $d \cup \{e\}$ is not double independent due to the fact that it contains a circuit or a cutset. □

According to Assertion 4, for any double independent subset d of a graph, there always exists a pair of trees (t_1, t_2) such that $d = t_1 \setminus t_2$. The next two assertions provide a link between a maximally double independent subset and a perfect pair of trees.

Assertion 6

Let (t_1, t_2) be a pair of trees of a graph G . If $t_1 \setminus t_2$ is a maximal double independent subset of G , then (t_1, t_2) is a perfect pair of trees.

Proof

If $t_1 \setminus t_2$ is a maximal double independent subset of G then, according to Assertion 5, each edge from its complement (including the edges in $t_1^* \cap t_2^*$) makes a circuit or/and a cutset with the elements of $t_1 \setminus t_2$ only. But $t_1 \setminus t_2$ together with $t_1^* \cap t_2^*$ belongs to t_2^* and hence the edges in $t_1^* \cap t_2^*$ cannot make cutsets with the edges $t_1 \setminus t_2$ only. Therefore the edges in $t_1^* \cap t_2^*$ make circuits with edges in $t_1 \setminus t_2$ only and consequently $\text{rank } t_2^* = |t_1 \setminus t_2|$. That means that tree t_1 is maximally distant from the tree t_2 . On the other hand, according to Assertion 2, $\text{rank } t_1^* \geq |t_1 \setminus t_2|$. We shall now prove that for the case under consideration, equality must occur. That is, t_2 is also maximally distant from t_1 . Suppose that this is not true. Then, according to Assertion 1, there exists an edge $e' \in t_1^* \cap t_2^*$ such that a fundamental circuit with respect to t_2 , defined by that edge contains an edge $c \in t_1 \cap t_2$. Consequently, $t'_2 = (t_2 \setminus e) \cup \{e'\}$ is again a tree and such that $t_1 \setminus t_2 \subseteq t_1 \setminus t'_2$. But subset $t_1 \setminus t'_2$ is, according to Assertion 4, also a double independent subset that contains as a proper subset the maximal double independent subset $t_1 \setminus t_2$. According to Assertion 5, this is a contradiction.

Thus we have proved that t_1 is maximally distant from t_2 and vice versa. Hence (t_1, t_2) is a perfect pair. □




Figure 2

Remark 2

The converse of Assertion 6 is not generally true. That is, if (t_1, t_2) is a perfect pair of trees then their set difference is not necessarily a maximally independent subset. To see this consider figures 2 and 2. Figure 2 shows four copies of the same graph and within each a subset of edges is indicated using bold lines. Now (t_1, t_2) is a perfect pair and (by inspection) $t_1 \setminus t_2$ is a maximal double independent subset while $t_2 \setminus t_1$ is not. Figure 3 shows four copies of the same graph and again various subsets of edges are indicated using bold lines. Again (t_1, t_2) is a perfect pair while neither $t_1 \setminus t_2$ nor $t_2 \setminus t_1$ is a maximal independent subset. The marked edges form neither circuits nor cutsets.




Figure 3

It is obvious that any double independent subset can be embedded in a maximal double independent subset. Also, any subset of a maximal double independent subset is double independent.

Remark 3

Suppose that for a given perfect pair of trees (t_1, t_2) , $t_1 \setminus t_2$ is not a maximal double independent subset and that we want to enlarge this subset until we obtain a maximal double independent subset. Let (t'_1, t'_2) be another perfect pair such that $t'_1 \setminus t'_2$ is a maximal double independent subset and let $t_1 \setminus t_2$ be a proper subset of $t'_1 \setminus t'_2$. Then $t_2 \setminus t_1$ does not belong to $t'_2 \setminus t'_1$ as a proper subset. To prove this let us consider a set of edges that have to be added to $t_1 \setminus t_2$ in order to obtain $t'_1 \setminus t'_2$. Due to properties of perfect pairs (Assertion 3, parts ii) and iii)) we cannot enlarge $t_1 \setminus t_2$ with elements of $t_1^* \cap t_2^*$ or $t_1 \cap t_2$. So, we have to take some edges from $t_2 \setminus t_1$. This means that $t_2 \setminus t_1$ partly belongs to $t'_1 \setminus t'_2$. But $t'_1 \setminus t'_2$ and $t'_2 \setminus t'_1$ are disjoint and consequently $t_2 \setminus t_1$ only partly belongs to $t'_2 \setminus t'_1$ which completes the proof.

To describe more closely the situation when the set difference of a perfect pair of trees is not a maximal double independent subset we establish the following assertion

Assertion 7

Given a perfect pair of trees (t_1, t_2) , the following three conditions are equivalent.

- (i) $t_1 \setminus t_2$ is not a maximal double independent subset.
- (ii) There exists an edge in $t_2 \setminus t_1$ that belongs to a fundamental circuit with respect to t_2 defined by an edge in $t_1^* \cap t_2^*$ and at the same time forms a fundamental circuit respect to t_1 in which at least one edge is in $t_1 \cap t_2$.
- (iii) There exists an edge in $t_2 \setminus t_1$ that belongs to a fundamental cutset with respect to t_1^* defined by an edge in $t_1 \cap t_2$ and at the same time forms a fundamental cutset respect to t_2^* in which at least one edge is in $t_1^* \cap t_2^*$.

Proof(i) \Leftarrow (ii)

Suppose that condition (ii) holds. That is, there exists an edge $e \in t_2 \setminus t_1$ that forms a fundamental circuit with respect to t_1 in which at least one edge is in $t_1 \cap t_2$ (call this conclusion 1). On the other hand, this edge belongs to the fundamental circuit with respect to t_2 defined by an edge in $a \in t_1^* \cap t_2^*$. Because the pair (t_1, t_2) is a perfect pair, each edge from $t_1^* \cap t_2^*$ forms fundamental circuits with respect to t_2 only with edges in $t_2 \setminus t_1$. So, the fundamental circuit defined by a contains only edges from $t_2 \setminus t_1$, including the edge e . As is well known from general graph theory, the intersection of a cutset and a circuit always contains an even number of edges. Therefore, any cutset that includes the edge e , includes at least one more edge from $t_2 \setminus t_1$. Thus, we conclude that edge e does not form a cutset with edges in $t_1 \setminus t_2$ only (call this conclusion 2). According to Assertion 5, conclusions 1 and 2 imply that $t_1 \setminus t_2$ is not a maximal double independent subset.

(i) \Rightarrow (ii)

Suppose that condition (ii) is not true. That is, suppose that each edge in $t_2 \setminus t_1$ that belongs to a fundamental circuit with respect to t_2 , defined by an edge in $t_1^* \cap t_2^*$ defines a fundamental circuit with respect to t_1 with edges in $t_1 \setminus t_2$ only (call this conclusion 3). The remaining edges in $t_2 \setminus t_1$ that do not belong to fundamental circuits with respect to t_2 defined by an edge in $t_1^* \cap t_2^*$ necessarily form cutsets with edges in $t_1 \setminus t_2$ only (call this conclusion 4). From conclusions 3 and 4 we see that all edges in $t_2 \setminus t_1$ form circuits or cutsets with respect to t_1 with edges only in $t_1 \setminus t_2$. On the other hand, for each perfect pair we have that all edges in $t_1^* \cap t_2^*$ form circuits with edges in $t_1 \setminus t_2$ only and all edges in $t_1 \cap t_2$ form cutsets with edges in $t_1 \setminus t_2$ only. According to Assertion 5, $t_1 \setminus t_2$ is a maximal double independent subset. Using reductio ad absurdum we conclude that (i) \Rightarrow (ii).

(ii) \Leftrightarrow (iii)

This is evident from the following well known statement: two edges belong to a circuit iff they both belong to a same cutset. ((ii) and (iii) are dual statements) Note also that $t_2 \setminus t_1 = t_1^* \setminus t_2^*$. \square

As an immediate consequence of Assertions 6 and 7, we have the following theorem.

Theorem 1

A subset of edges d (of a graph G) is a maximal double independent subset iff the conjunction of the following two statements hold.

- (i) There exists a perfect pair (t_1, t_2) such that $d = t_1 \setminus t_2$.
- (ii) Each edge in $t_2 \setminus t_1$ that belongs to a fundamental circuit with respect to t_2 , defined by an edge in $t_1^* \cap t_2^*$, defines a fundamental circuit with respect to t_1 with edges in $t_1 \setminus t_2$ only.

Assertion 8

If (t_1, t_2) is a maximally distant pair of trees then both $t_1 \setminus t_2$ and $t_2 \setminus t_1$ are maximal double independent subsets.

Proof

Suppose that one of the subsets $t_1 \setminus t_2$ or $t_2 \setminus t_1$ is not maximal double independent, for example the subset $t_1 \setminus t_2$. Then there exists a maximal double independent subset d that contains $t_1 \setminus t_2$ as a proper subset. According to Assertion 6 there is a perfect pair of trees (t'_1, t'_2) such that $t'_1 \setminus t'_2 = d$. Because $t_1 \setminus t_2 \subset d = t'_1 \setminus t'_2$, we conclude that $|t_1 \setminus t_2| < |d| = |t'_1 \setminus t'_2|$ which contradicts the assumption that (t_1, t_2) is a maximally distant pair of trees. \square

Remark 4

The converse of Assertion 8 is not generally true. That is, if $t_1 \setminus t_2$ and $t_2 \setminus t_1$ are both maximal double independent subsets, then (t_1, t_2) is not necessarily a maximally distant pair of trees. In order to see this consider figure 4. This figure shows four copies of the same graph with different subsets of edges indicated with bold lines. Now $t_1 \setminus t_2$ and $t_2 \setminus t_1$ are both maximal double independent subsets but (t_1, t_2) is not a maximally distant pair of trees.




Figure 4

Conclusion

In this paper the notion called maximally double independent subset is considered and related to the concept of perfect pair of trees. Several assertions were stated in order to closely characterise its properties. Also, several examples were included in order to help the reader gain intuitive insight.

References

- [1] Amari, S, "Topological foundation of Kron's tearing of electrical networks", RAAG Memoirs, vol 3, pp. 322-349, 1962.

- [2] Ohtsuki, T, Ishizaki, Y, and Watanabe, H, "Topological degrees of freedom and mixed analysis of electrical networks", **IEEE Trans. on Circuit Theory**, CT-17, pp.491-499, 1970.
- [3] Kishi, G and Kajitani, Y, "Maximally distant trees and principal partition of a linear graph", **IEEE Trans. on Circuit Theory**, vol.CT-16, pp.323-330, 1969.
- [4] Hakimi, S L, "On trees of a graph and their generation", **J. Franklin Inst.**, vol 272, pp.347-359, November 1961.
- [5] Deo, N, "A central tree", **IEEE Trans.on Circuit Theory**, vol CT-13, pp.439-440, 1960.
- [6] Welsh, D J A, **Matroid Theory**, Academic Press, London, 1976.
- [7] Novak, L and Gibbons, A M, "On perfect pairs of trees in a graph", Research Report 151, Department of Computer Science, University of Warwick, October 1989.
- [8] Bruno, J and Weinberg, L, "A constructive graph-theoretic solution of the shannon switching game", **IEEE Trans. on Circuits and Systems**, CAS-17, pp.74-81, 1970.
- [9] Chen, W K, and Chan, F N T, "On the unique solvability of linear active networks", **IEEE Trans. on Circuits and Systems**, pp.26-35, 1974.
- [10] Ozawa, T, "Topological conditions for the solvability of linear active networks", **Int. J. of Circuit Theory and Appl.**, pp.125-136, 1976.
- [11] Shinoda, S, "A minimization problem in network diagnosis", **Trans. IECEJ**, vol.53A, pp.569-570, 1970.
- [12] Tomizawa, N, "An algorithm for determining the maximally suspended pair of bases in a semi-munoid with applications to the diagnosis of networks", **Paper of Tech. Group on Circuit and System Theory of IECEJ**, CST74-80, pp.19-22, 1974.
- [13] Tomizawa, N, "An algorithm for determining the maximally distant pair of bases with matroidal constraints - generalized versions of Menger's theorem for matroids", **Papers of Tech. Group on Circuit and System Theory of IECEJ**, CST74-33, pp.17-24, 1974.
- [14] Kajitani, Y, and Kawamoto, T, "Graph theoretical degree of freedom in network analysis and all-central-tree graphs", **ibid**, CST76-94, pp.65-70, 1976.

- [15] Kawamoto, T, Kajitani, Y, and Shinoda, S, "New theorems on central trees described in connection with the principal partition of a graph", **ibid**, CST77-109, pp.63-68, 1977.
- [16] Kawamoto, T, and Shinoda, S, "Theorems on central trees in connection with the principal partition of a graph", **ibid**, CAS80-49, pp.41-45, 1980.
- [17] Kawamoto, T, Kajitani, Y, and Shinoda, S, "Eccentricity of the tree set", **Proc. of the IEEE Int. Symp. on Circuits and Systems**, paper R14-1, 1981.
- [18] Gao, S, and Chen, W K, "The hybrid method of network analysis and topological degree of freedom", **ibid**, pp.158-161, 1982.
- [19] Gao, S, and Chen, W K, "On graph classification with application", **Proc. 25th -Midwest Symp. on Circuits and Systems**, pp.244-248, 1982.
- [20] Lin, P M, "Complementary trees in circuit theory", **IEEE Trans. on Circuits and Systems**, CAS-27, pp.921-928, 1980.
- [21] Harary, F, Mokken, R J and Plantholt, M J, "Interpolation theorem for diameters of spanning trees", **IEEE Trans.on Circuits and Systems**, vol CAS-30, pp 429-431, 1983.
- [22] Novak, L A, "A few small theorems concerning pairs of matroid bases", to appear in **Int.J of Circuit Theory and Appl.** in 1990.
- [23] Swamy, M N S and Thulasiraman, K, "Graphs, Networks and Algorithms", John Wiley and Sons, 1981.