

Original citation:

Wischer, Daniela, Kumaresan, Deepak, Johnston, Antonia, El Khawand, Myriam, Stephenson, Jason, Hillebrand-Voiculescu, Alexandra M, Chen, Yin and Colin Murrell, J.. (2014) Bacterial metabolism of methylated amines and identification of novel methyloproths in Movile Cave. The ISME Journal . ISSN 1751-7362

Permanent WRAP url:

<http://wrap.warwick.ac.uk/62579>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Link to the published version: <http://dx.doi.org/10.1038/ismej.2014.102>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP url' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: publications@warwick.ac.uk



<http://wrap.warwick.ac.uk>

**Bacterial metabolism of methylated amines and identification of novel methylotrophs in
Movile Cave**

5 Daniela Wischer¹, Deepak Kumaresan^{1*}, Antonia Johnston¹, Myriam El Khawand¹, Jason
Stephenson², Alexandra M. Hillebrand-Voiculescu³, Yin Chen² and J. Colin Murrell¹

¹School of Environmental Sciences, University of East Anglia, Norwich, UK

²School of Life Sciences, University of Warwick, Coventry, UK

10 ³Emil Racoviță Institute of Speleology, Bucharest, Romania

* Current affiliation: School of Earth and Environment, The University of Western Australia,
Crawley, Perth, Australia

Correspondence: J Colin Murrell, School of Environmental Sciences, University of East
15 Anglia, Norwich, NR4 7TJ, UK. Telephone: +44 (0)1603 592959 Fax: +44 (0)1603 591327
Email: j.c.murrell@uea.ac.uk

Subject Category: Microbial ecology and functional diversity of natural habitats

Running title: Methylamine metabolism in Movile Cave

20 **Keywords:** *Catellibacterium* / DNA-SIP / gamma-glutamylmethylamide synthetase / *gmaS* /
Movile Cave

Abstract

Movile Cave, Romania, is an unusual underground ecosystem that has been sealed off from the outside world for several million years and is sustained by non-phototrophic carbon fixation. Methane and sulfur-oxidising bacteria are the main primary producers, supporting a complex food web that includes bacteria, fungi and cave-adapted invertebrates. A range of methylotrophic bacteria in Movile Cave grow on one carbon compounds including methylated amines, which are produced via decomposition of organic-rich microbial mats. The role of methylated amines as a carbon and nitrogen source for bacteria in Movile Cave was investigated using a combination of cultivation studies and DNA stable isotope probing (DNA-SIP) using ^{13}C -monomethylamine (MMA). Two newly-developed primer sets targeting the gene for gamma-glutamylmethylamide synthetase (*gmaS*), the first enzyme of the recently-discovered indirect MMA oxidation pathway, were applied in functional gene probing. SIP experiments revealed that the obligate methylotroph *Methylotenera mobilis* is one of the dominant MMA utilisers in the cave. DNA-SIP experiments also showed that a new facultative methylotroph isolated in this study, *Catellibacterium* sp. LW-1 is probably one of the most active MMA utilisers in Movile Cave. Methylated amines were also used as a nitrogen source by a wide range of non-methylotrophic bacteria in Movile Cave. PCR-based screening of bacterial isolates suggested that the indirect MMA oxidation pathway involving gamma-glutamylmethylamide and *N*-methylglutamate is widespread amongst both methylotrophic and non-methylotrophic MMA utilisers from the cave.

45 **Introduction**

Most ecosystems rely on phototrophic carbon fixation, or, in the absence of light, an external supply of phototrophically-fixed carbon into the ecosystem. Exceptions are deep sea hydrothermal vents, where carbon is derived from chemosynthesis using energy sources other than light (reviewed by Lutz & Kennish, 1993; Van Dover *et al.*, 2002; Campbell 2006).

50 Movile Cave, located near the coast of the Black Sea in Mangalia, Romania, is an underground cave system that has been completely sealed off from the outside world for several million years (Sarbu, 1996). Unlike other cave systems, where dissolved and particulate organic carbon enters the cave with meteoric waters from above, the food web in Movile Cave is sustained exclusively by non-phototrophic carbon fixation. Since its discovery

55 in 1986, Movile Cave has provided an excellent natural ecosystem to study a highly unusual, light-independent, microbially-driven foodweb (Sarbu *et al.*, 1994; Sarbu & Kane, 1995; Sarbu *et al.*, 1996; Rohwerder *et al.*, 2003; Hutchens *et al.*, 2004; Porter *et al.*, 2009; Chen *et al.*, 2009). Movile Cave harbours rich and diverse populations of cave-adapted invertebrates, all of which are sustained by chemolithoautotrophic microorganisms that thrive along the

60 redox interface created between the oxygenated atmosphere and the high concentrations of reduced compounds such as hydrogen sulfide (H₂S) and methane (CH₄) present in the water (Sarbu & Kane, 1995). Microbial mats composed of bacteria, fungi and protists float on the water surface (kept afloat by CH₄ bubbles) and also grow on the limestone walls of the cave (Sarbu *et al.*, 1994).

65 Methylootrophs are organisms capable of using one-carbon (C1) compounds, i.e. compounds lacking carbon-carbon bonds, as their sole source of carbon and energy (Anthony, 1982; Lidstrom, 2006; Chistoserdova *et al.*, 2009). In addition to CH₄, C1 compounds such as methanol and methylated amines are important carbon and energy sources for a range of

methylotrophic bacteria in Movile Cave (Hutchens *et al.*, 2004; Chen *et al.*, 2009).
70 Methylated amines are typically associated with saline environments (Gibb *et al.*, 1999;
Fitzsimons *et al.*, 2006) where they form by degradation of glycine betaine and
trimethylamine *N*-oxide (TMAO), osmolytes commonly found in marine organisms (Barrett,
1985; Lin & Timasheff, 1994). There are fewer studies on the distribution of methylated
75 amines in terrestrial and freshwater environments, although the dissolved organic nitrogen
(DON) fraction as a whole is increasingly being recognised as an important source of
microbial nitrogen nutrition (Berman & Bronk, 2003, Worsfold *et al.*, 2008). Generally,
environments with high concentrations of organic matter have a high potential for DON
generation (Neff *et al.*, 2003). We hypothesise that in Movile Cave, degradation of the
extensive, organic-rich microbial mats produces large amounts of methylated amines, which
80 are used as growth substrates by certain microorganisms that are the subject of this study.

Methylotrophs utilising methylated amines as a carbon source are phylogenetically
diverse, ubiquitous in the environment and often metabolically versatile (e.g. Bellion &
Hersh, 1972; Colby & Zatman, 1973; Levering *et al.*, 1981; Anthony, 1982; Bellion &
Bolbot, 1983; Brooke & Attwood, 1984; Kalyuzhnaya *et al.*, 2006b; Boden *et al.*, 2008). New
85 methylotrophs are still being identified from a wide range of environments, including genera
not previously associated with methylotrophy, and novel metabolic pathways (see recent
reviews by Chistoserdova *et al.*, 2009; Chistoserdova 2011).

Methylated amines are also a nitrogen source for a wide range of non-methylotrophic
bacteria. While utilisation of MMA as a bacterial nitrogen source was reported over 40 years
90 ago (Budd & Spencer, 1968; Bicknell & Owens, 1980; Anthony, 1982; Murrell & Lidstrom,
1983; Glenn & Dillworth, 1984), details of the metabolic pathways involved have only
recently been identified (Chen *et al.* 2010b).

A key intermediate in methylotrophic metabolism is formaldehyde (or formate), the branching point at which carbon is either oxidised further to CO₂, or assimilated into cell carbon via the serine cycle or the ribulose monophosphate cycle (Anthony, 1982; Chistoserdova *et al.*, 2009; Chistoserdova, 2011). In the metabolism of methylated amines, there are two possible pathways for the oxidation of MMA (Supplementary Figures S1a and b): In the well-characterised, direct MMA oxidation pathway, a single enzyme oxidises MMA to formaldehyde, releasing ammonium. In methylotrophic Gram-positive bacteria the enzyme responsible is MMA oxidase, while in Gram-negative methylotrophs it is MMA dehydrogenase (Anthony, 1982). PCR primers are available for *mauA* (Neufeld *et al.*, 2007a), the gene coding for the small subunit of MMA dehydrogenase. However, these primers do not detect all MMA-utilising bacteria. An alternative, indirect pathway oxidises MMA not to formaldehyde but to 5,10-methylenetetrahydrofolate (CH₂=THF), in a stepwise conversion via the methylated amino acids gamma-glutamylmethylamide (GMA) and / or N-methylglutamate (NMG) (Latypova *et al.*, 2010; Chistoserdova, 2011). Although this pathway has been known since the 1960s (Kung & Wagner, 1969), the enzymes and genes involved have only recently been characterised (Latypova *et al.*, 2010; Chen *et al.*, 2010a): MMA is converted to GMA by GMA synthetase (*gmaS*), GMA then to NMG by NMG synthase (*mgsABC*), and finally to CH₂=THF by NMG dehydrogenase (*mgdABCD*). A variation of this pathway is found in *Methyloversatilis universalis* FAM5, where *gmaS* is not essential for oxidation of MMA to CH₂=THF via NMG (Latypova *et al.*, 2010). Importantly, the GMA / NMG mediated pathway is also found in bacteria that use MMA only as a nitrogen source (Chen *et al.*, 2010b; Chen 2012). In a recent study (Chen, 2012), PCR primers targeting *gmaS* from marine *Roseobacter* clade (MRC) bacteria were developed for the detection of MMA utilisers in marine environments, highlighting the potential of the *gmaS* gene as a biomarker for MMA utilisation.

The objectives of this study were to determine the role of methylated amines as carbon and nitrogen sources for microorganisms in Movile Cave, and to identify active MMA
120 utilisers in this unique ecosystem using DNA stable isotope probing (SIP) (Radajewski *et al.*,
2000; Murrell & Whiteley, 2010). DNA-SIP has been successfully applied in the study of
methanotrophic and autotrophic communities in Movile Cave (Hutchens *et al.*, 2004; Chen *et*
al., 2009). Time-course SIP experiments with ¹³C-labelled MMA were set up in order to
125 monitor changes in the methylotrophic community. Cultivation-based studies were also used
to isolate and characterise methylated amine-utilising bacteria from the cave. The distribution
of genes for the GMA-dependent MMA-oxidation pathway in Movile Cave microbes was
examined using new PCR primer sets developed to target *gmaS* from non-marine bacteria.

Material and Methods

130 *Study site and sampling*

Movile Cave near Mangalia on the coast of the Black Sea is located in an area rich in
hydrothermal activity with numerous sulfurous springs and lakes, as well as creeks bubbling
with methane. The cave consists of a network of passages, including a dry, upper level and a
lower level which is partly flooded by thermal sulfidic waters (for a detailed cross-section of
135 the cave see Supplementary Figure S2). A small lake room (ca 3 m in diameter) is located
between the dry and the flooded sections of the cave, and two air bells are located in the
submerged region. The temperature in the cave is a constant 21°C (Sarbu & Kane, 1995). The
atmosphere in the air bells is depleted in O₂ (7-10% v/v) and rich in CO₂ (2.5% v/v) and CH₄
(1-2% v/v) (Sarbu & Kane, 1995). The water contains H₂S (0.2-0.3 mM), NH₄⁺ (0.2-0.3 mM)
140 and CH₄ (0.02 mM) and is buffered by high amounts of bicarbonate from the lime stone walls
at ~pH 7.4 (Sarbu, 2000). Dissolved O₂ decreases to less than 1 μM after the first few cm

from the water surface, with the deeper water being essentially anoxic (Sarbu, 2000).

Methylamine concentrations in the cave water were measured by our recently developed ion chromatography method with a detection limit of around 1 μM for MMA (Lidbury *et al.*, 2014). Preliminary measurements carried out using this assay suggested that the *in situ* concentration of MMA in the Movile Cave water is below the detection limit of 1 μM , which could indicate rapid turnover of MMA by bacteria in the cave.

Water and floating mat samples for enrichment and isolation experiments were collected from the lake room and the two air bells in October 2009, stored at 4°C in the nearby field station and processed within 48 hours. Biofilm covering the limestone walls of both air bells was scraped off into sterile tubes. Similar samples for further isolation experiments, SIP enrichments and nucleic acid extractions were obtained from Movile Cave in April 2010. Material for DNA work was concentrated by centrifugation within one hour of sampling and frozen at -20°C for storage until processing.

DNA-SIP with ^{13}C -MMA

SIP incubations were set up at the field station in Mangalia, within one hour of sampling, using water from Airbell 2. For each incubation, a 20 ml aliquot of cave water was added to a pre-sterilised 120 ml serum vial containing 50 μmol of labelled (^{13}C) or unlabelled MMA-HCl (dissolved in 0.2 ml sterilised distilled water). Control incubations with no added MMA (referred to as “no-substrate controls” from here on) were also set up. All serum vials were immediately sealed with a butyl rubber cap and an aluminium crimping lid and incubated at 21°C in the dark. Samples for t_0 ($t = 0$ days) were prepared by centrifugation of 20 ml of cave water, discarding the supernatant and freezing the pellet at -20°C. SIP incubations and no-substrate controls were harvested in the same way at time intervals of 48 hours (t_1), 96 hours (t_2) and four weeks (t_3). In future SIP experiments, the recently developed

ion chromatography method for measuring MMA (see above, Lidbury *et al.*, 2014) could be used to monitor consumption of substrate at the time of sampling. From each sample, up to 1 µg of total extracted DNA was added to caesium chloride (CsCl) solutions for isopycnic ultracentrifugation and gradient fractionation following published protocols (Neufeld *et al.*, 2007b).

Enrichment and isolation of methylated amine utilising bacteria from Movile Cave

Methylotrophic bacteria capable of using methylated amines as a carbon and nitrogen source were selectively enriched using MMA, dimethylamine (DMA) and trimethylamine (TMA). Separate enrichments were set up for each of the three substrates by adding a final concentration of 1 mM substrate to 20 ml cave water in sterile 120 ml serum vials. For mats and biofilms, 2 g sample material was placed into 27 ml serum vials and made up to a final volume of 4 ml with nitrogen-free dilute basal salts (DBS) medium. DBS medium was modified after Kelly & Wood (1998) and contained (per litre): 0.1 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.05 g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.11 g K_2HPO_4 , 0.085 g KH_2PO_4 , adjusted to pH 7. The medium was supplemented with a vitamins solution as described by Kanagawa *et al.* (1982) and 1 ml of a trace element solution (modified after Kelly & Wood, 1998) containing (per 1 L): 50 g ethylenediaminetetraacetic acid (EDTA), 11 g NaOH, 5 g $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 7.34 g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 2.5 g $\text{MnCl}_2 \cdot 6\text{H}_2\text{O}$, 0.5 g $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 0.5 g $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$, 5 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, adjusted to pH 6.0. After flushing the headspace of each vial with N_2 , the head space was made up to a final concentration of 7% (v/v) O_2 and 3.5% (v/v) CO_2 to resemble the cave atmosphere. Enrichments were incubated at 21°C in the dark. After four weeks, 10 ml (for water samples) or 4 ml (for mat samples) of fresh DBS medium were added and cultures were spiked with 20 mM MMA, 10 mM DMA, or 10 mM TMA, respectively. After amending the headspace as previously, enrichment cultures were incubated at 21°C in the dark. When enrichments became turbid (after a further two weeks),

dilutions were spread onto agar plates (DBS medium, 1.5% agar) containing 5 mM MMA, DMA, or TMA, respectively, as the only added carbon and nitrogen source. Plates were
195 incubated at 21°C in the dark until colonies became visible (2 - 10 days). In order to achieve isolation of a variety of methylotrophs, individual colonies were examined by microscopy and a selection of morphotypes was transferred onto fresh plates containing the same substrates as before. Cells were observed at 1000 x magnification in phase-contrast under a Zeiss Axioskop 50 microscope. Isolates were submitted to a series of transfers on plates and microscopy was
200 used routinely to check purity before transferring individual isolates into liquid media (containing 5 mM MMA, DMA, or TMA). Once grown in liquid (2 - 7 days), isolates were transferred back onto methylated amine plates.

In a separate enrichment approach, non-methylotrophic bacteria capable of using methylated amines as a nitrogen (but not carbon) source were isolated. These enrichments
205 were set up in the same manner and using the same sample material as described above for the methylotrophs. In addition to 1 mM of MMA, DMA or TMA, a mixture of alternative carbon compounds (comprising glucose, fructose, succinate, glycerol, pyruvate and acetate) was added to a final concentration of 5 mM. Isolates obtained in this way were additionally tested for growth in liquid medium containing no alternative carbon source to detect any
210 co-enriched methylotrophs, as well as in liquid medium containing carbon sources but no methylated amines to eliminate the possibility that they might be fixing N₂ rather than using methylated amines as nitrogen source.

DNA extraction and PCR amplification of bacterial 16S rRNA genes

215 DNA from cave samples, SIP enrichments and bacterial isolates was extracted as previously described (Neufeld *et al.*, 2007a). DNA from soil and lake sediment samples retrieved from the University of East Anglia campus (used for *gmaS* primer validation, see later) was extracted using the FastDNA® SPIN Kit for soil by MP Biomedicals LLC. Bacterial 16S

rRNA genes from SIP enrichments were amplified using primer set 341f-GC / 907r (Muyzer
220 *et al.* 1993; Lane, 1991) for analysis by denaturing gradient gel electrophoresis (DGGE). For
cloning and sequencing, bacterial 16S rRNA genes from isolates were amplified with primer
set 27f / 1492r (DeLong, 1992; Lane, 1985).

Denaturing gradient gel electrophoresis (DGGE)

225 DGGE analysis of bacterial 16S rRNA gene fragments was carried out as described by
Neufeld *et al.* (2007a) using the DCode™ Universal Mutation Detection System (Bio-Rad).
After electrophoresis for 16 h at 60°C and 80 V, gels were stained using SYBR® Gold
Nucleic Acid Gel Stain (Invitrogen) and viewed under the Bio-Rad Gel Doc XR gel
documentation system using Amber Filter 5206 (Bio-Rad). For gene sequence analysis, well-
230 defined DNA bands were physically excised from the gel for re-amplification using the same
PCR conditions and primers described above, followed by sequencing analysis using primer
341f (Muyzer *et al.*, 1993).

DGGE, when compared to amplicon pyrosequencing is a relatively low resolution
technique. However, the DGGE technique enabled us to accurately compare SIP enrichments
235 across different CsCl gradient fractions (heavy to light) and also to compare ¹³C-incubations
to ¹²C-incubated controls. In this first study on MMA degraders in Movile Cave, it thereby
allowed us to identify key players in the microbial food web. Building on data obtained in this
study, more detailed studies involving pyrosequencing of amplicons can be carried out in the
future.

240 *Functional gene PCR and development of gmaS primers*

mauA genes were amplified using PCR primer set mauAf1 / mauAr1 (Neufeld *et al.* 2007a).
Currently there is one *gmaS* PCR primer set available (Chen, 2012) which targets specifically
the marine *Roseobacter* clade. This PCR primer set therefore may not detect *gmaS* from non-

marine bacteria. Three new *gmaS* PCR primers were designed in this study, based on multiple
245 alignment of 34 *gmaS* sequences derived from (i) methylotrophic isolates confirmed to use
the NMG / GMA mediated pathway and (ii) bacterial genomes published on the Integrated
Microbial Genomes (IMG) platform of the Joint Genome Institute (JGI). Genomes were
screened for *gmaS*-related sequences using *gmaS* from *Methylocella silvestris* as a query
sequence (Chen *et al.*, 2010a). Corresponding full length sequences included both *gmaS* and
250 glutamine synthetase type III (*glnA*) sequences, due to the high level of sequence similarity
between the two genes. In order to identify genuine *gmaS* sequences, the gene neighbourhood
of all obtained sequences was manually inspected for predicted neighbouring open reading
frames (ORFs) typically found adjacent to *gmaS* (genes encoding NMG dehydrogenase and
NMG synthase). Confirmed *gmaS* sequences included many sequences apparently mis-
255 annotated as *glnA*. For primer design, multiple sequence alignments of chosen sequences were
established with the Clustal X program (Thompson *et al.*, 1997) and viewed using the
GeneDoc software (Nicholas *et al.*, 1997). Because of their sequence similarity to *gmaS*, a
number of *glnA* sequences were included in the alignment in order to identify suitable primer-
binding regions specific only to *gmaS* (for a complete list of all *gmaS* and *glnA* sequences
260 used for primer design, see Supplementary Table S1). The resulting forward primer
*gmaS*_557f (GARGAYGCSAACGGYCAGTT) was used in all cases, with the reverse
primers α _gmaS_970r (TGGGTSCGRTRTTGCCSG) and β _gmaS_1332r
(GTAMTCSAYCCAYTCCATG) being used to target the *gmaS* gene of non-marine
Alphaproteobacteria and that of *Beta*- and *Gammaproteobacteria*, respectively. Touchdown
265 PCR protocols for *gmaS* amplification were used as follows: For *gmaS*_557f / α _gmaS_970r,
an initial step at 94°C for 5 min was followed by 10 cycles of denaturation at 94°C for 45
seconds, annealing at variable temperatures for 45 seconds, and extension at 72°C for 1 min.
In the first cycle, the annealing temperature was set to 60°C, and for each of the 9 subsequent
cycles the annealing temperature was decreased by 1°C. This was followed by 30 cycles of 45

270 sec at 94°C, 45 sec at 56°C and 1 min at 72°C, and a final extension time of 8 min at 72°C.
For *gmaS*_557f / β _ γ _gmaS_1332r, a modification of the first touchdown protocol was used;
the annealing temperature was set to 55°C in the first cycle and decreased by 1 degree for
each of the 9 subsequent cycles. The first 10 cycles were followed by 35 cycles with an
annealing temperature of 52°C.

275 The primer sets were tested for their specificity by (i) amplification and sequencing of
gmaS sequences from genomic DNA of the following bacterial strains known to use the
indirect MMA oxidation pathway: *Sinorhizobium meliloti* 1021, *Mesorhizobium loti*
MAFF303099, *Rhizobium leguminosarum* bv. *viciae* 3841, *Agrobacterium tumefaciens* C58
and *Pseudomonas fluorescens* SBW25 (Chen *et al.*, 2010b). For further validation of the
280 primers, *gmaS* was amplified from DNA extracted from (ii) MMA enrichments from Movile
Cave, (iii) fresh Movile Cave mat and (iv) soil and freshwater sediment from a small lake (the
“Broad”) on the University of East Anglia campus. *gmaS*-based clone libraries were
constructed for (ii) – (iv) and a total of 30 clones were randomly selected for sequencing.

285 *DNA sequencing and phylogenetic analysis*

DNA sequencing employed the Sanger method on a 3730A automated sequencing system (PE
Applied Biosystems). To determine approximate phylogenetic affiliations, partial 16S rRNA
gene sequences were analysed with the Basic Local Alignment Search Tool (BLAST) on the
NCBI GenBank database (Altschul *et al.*, 1990). Amino acid and nucleotide-based
290 phylogenetic trees were established using the MEGA5 program (Tamura *et al.*, 2011). The
evolutionary history was inferred by neighbour-joining (Saitou & Nei, 1987). For nucleotide-
based trees (Supplementary Figures 1a and b), the evolutionary distances were computed
using the maximum composite likelihood method (Tamura *et al.*, 2004). For amino-acid
based trees, the evolutionary distances were computed using the Poisson correction method
295 (Zuckerkindl & Pauling, 1965). All positions containing gaps and missing data were

eliminated. Bootstrap analysis (1000 replicates) was performed to provide confidence estimates for phylogenetic tree topologies (Felsenstein *et al.* 1985). Phylogenetic analysis of *gmaS* genes was carried out at the amino acid level (135 - 250 amino-acyl residues).

Nucleotide sequence accession numbers

300 Nucleotide gene sequences obtained from this study were deposited in the GenBank nucleotide sequence database under the accession numbers xxx-xxx for 16S rRNA gene sequences and xxx-xxx for *gmaS* sequences.

Results

305 *Active methylotrophic bacteria identified by DNA-SIP with ¹³C-MMA*

DNA-SIP enrichments with ¹³C-labelled MMA were set up from Movile Cave water in order to identify active, methylotrophic bacteria capable of using MMA as a carbon source. DNA was extracted from microcosms enriched with ¹³C-labelled and unlabelled MMA after (i) 48 hours (t₁), (ii) 96 hours (t₂) and (iii) 4 weeks (t₃). The bacterial communities in the
310 microcosms were investigated by DGGE analysis of bacterial 16S rRNA gene fragments. Comparison of DGGE profiles from unfractionated DNA from the different time points revealed a significant shift in the bacterial community over time, which was similar between ¹²C-MMA and ¹³C-MMA incubations (Figure 1).

[Figure 1]

315 For identification of active methylotrophs, DNA extracted from all time points was subjected to density gradient centrifugation and fractionation, allowing separation of ¹³C-labelled DNA (contained in heavy fractions) from unlabelled, ¹²C-DNA (contained in light

fractions). Bacterial 16S rRNA gene fragments were amplified from all DNA fractions and analysed by DGGE and sequencing. Time point t_1 (48 hours) did not show any significant enrichment in ^{13}C -DNA and was therefore not further analysed. DGGE analysis of heavy and light DNA fractions from time points t_2 and t_3 (^{13}C -MMA incubation) revealed major differences in the community profiles of the heavy fractions (Figures 2a and c): A single band dominated the heavy fractions at t_2 (96 h, Figure 2a) but was absent at t_3 (4 weeks, Figure 2c). Sequence analysis of the excised band revealed that the sequence affiliated with *Methylotenera mobilis* (99% identity), an obligate methylotroph (Kalyuzhnaya *et al.*, 2006a) known to be abundant in Movile Cave from previous studies (Chen *et al.*, 2009). At t_3 , several different phylotypes appeared in the heavy fractions of the ^{13}C -MMA incubation (Figure 2c), i.e. a more diverse bacterial community had incorporated the label following extended incubation with MMA.

[Figure 2]

Bacterial 16S rRNA gene sequences from these DGGE bands affiliated with well-characterised methylotrophs such as *Methylobacterium extorquens* (100% identity) and *Methylovorus* (97% identity to *Methylovorus menthalis*), but also included *Catellibacterium* (98% identity to *Catellibacterium caeni*), *Cupriavidus* (99% identity to *Cupriavidus necator*), *Porphyrobacter* (99% identity to *Porphyrobacter neustonensis*) and *Altererythro bacter* (99% identity to *Altererythro bacter epoxidivorans*), none of which have previously been reported to grow methylotrophically. The *Catellibacterium* sequence identified from DGGE shared 98-100% sequence identity with a novel organism subsequently isolated from Movile Cave during this study (see below) and cloned 16S rRNA gene sequences from ^{13}C -labelled DNA from t_3 (data not shown, refer to Supplementary Figure S3a).

The non-methylotrophic bacterial community co-enriched in ^{13}C -MMA incubations was investigated by PCR-DGGE of 16S rRNA bacterial genes present in the light fractions

(¹²C-DNA). Light fractions harboured a diversity of mostly heterotrophic bacterial sequences (Figures 2a and b), namely *Rhodobacter*, *Acinetobacter*, *Azospirillum*, *Oleomonas* and
345 *Hydrogenophaga* and a number of sequences not closely related to cultivated representatives (as little as 84-87% identity).

All bacterial sequences obtained from DGGE bands in lanes loaded with heavy DNA (i.e. ¹³C-labelled organisms) were exclusive to MMA enrichments, and not seen in no-substrate controls (data not shown). Two sequences detected in the light fractions from MMA
350 incubations (*Acinetobacter* and *Azospirillum*) also appeared to be absent in the no-substrate controls, suggesting that these bacteria may have been selectively enriched due to their capability of using MMA as a nitrogen source (but not as a carbon source, and so their DNA was not labelled). One of these sequences (*Acinetobacter lwoffii*) did indeed correspond to a bacterium isolated from Movile Cave in this study with MMA as the only nitrogen source
355 (see below).

Methylotrophic and non-methylotrophic isolates from Movile Cave

To complement data from ¹³C-MMA SIP experiments, methylated-amine utilising bacteria were isolated from different locations (lake room, Airbell 1 and Airbell 2) in Movile Cave. Methylotrophs were isolated with DBS medium containing MMA, DMA or TMA as sole
360 added source of carbon, energy and nitrogen. A selection of isolates differing in colony and cell morphology was transferred into liquid DBS medium containing the respective methylated amine (to distinguish true methylotrophs from organisms growing on agar). Six methylotrophic strains were isolated, identified based on 16S rRNA gene sequencing analysis (Table 1, Supplementary Figure S3a). The highest diversity of methylotrophs was obtained on
365 MMA enrichments (based on morphology and 16S rRNA gene sequencing data), while DMA and TMA enrichments were dominated by *Xanthobacter tagetidis* (Padden *et al.*, 1997). Notably, no *Methylotenera* isolates were obtained (even after using a variety of different

cultivation media which are commonly used for methylotrophic bacteria, changing incubation conditions such as temperature, pH, ionic strength of media and dilution-to-extinction experiments), despite the active role of this methylotroph in MMA metabolism as determined by DNA-SIP results (see above), and its apparent abundance in Movile Cave (Chen *et al.*, 2009). In addition to well-characterised methylotrophs such as *Methylobacterium extorquens*, two novel methylotrophs were also isolated. A member of the relatively new genus *Catellibacterium* (Tanaka *et al.*, 2004; Liu *et al.*, 2010, Zheng *et al.*, 2011; Zhang *et al.*, 2012), provisionally named *Catellibacterium* sp. LW-1 was isolated from lake water enrichments with MMA. 16S rRNA gene sequences relating to this organism were also detected in heavy DNA fractions from ¹³C-MMA enrichments (see above, Figure 2c, Supplementary Figure S1b), indicating that *Catellibacterium* may play a significant role in the cycling of methylated amines in Movile Cave. In addition, a new member of the genus *Mesorhizobium* (a genus not currently known to contain any methylotrophic species), was isolated from an MMA enrichment set up with floating mat from Airbell 1. All methylotrophic isolates were facultative, i.e. able to use sugars or carboxylic acids for growth. Notably, all methylotrophs could use all three methylated amines as sole growth substrates, with the exception of *Catellibacterium* sp. LW-1 which did not grow on DMA (Table 1).

In a separate experiment, heterotrophic bacteria capable of using methylated amines as a nitrogen but not carbon source were enriched and isolated using the same sample material as above. MMA, DMA or TMA were the only added nitrogen sources in these enrichments and a mixture of sugars and carboxylic acids were added as carbon and energy source. A diversity of non-methylotrophic methylated amine-utilising bacteria was obtained; in total eight bacterial species, as determined by 16S rRNA gene sequencing analysis (Table 1, Supplementary Figures S3a and b). All of these isolates used MMA as a nitrogen source, while only some could use DMA and TMA (Table 1), suggesting that many lack the enzymes

for de-methylation of secondary and tertiary methylated amines to MMA. None of the isolates grew methylotrophically with MMA, DMA or TMA. While all methylotrophic isolates
395 obtained in this study belonged to the *Alphaproteobacteria*, non-methylotrophic MMA utilisers also included *Beta-* and *Gammaproteobacteria* (Table 1). *Acinetobacter lwoffii*, isolated from Airbell 2 water with MMA as a nitrogen source, was also detected in ¹²C-DNA fractions from MMA-SIP incubations (see above), while not seen in control incubations without added MMA. These results suggest that *Acinetobacter* (and other non-methylotrophs)
400 may play an active role in the cycling of methylated amines in Movile Cave.

[Table 1]

Development and validation of functional gene primer sets targeting gmaS

The gene *gmaS* codes for gamma-glutamylmethylamide synthetase, the enzyme catalysing the first step in the conversion of MMA to formaldehyde in the recently characterised indirect
405 MMA oxidation pathway (Latypova *et al.*, 2010; Chen *et al.*, 2010a, 2010b). We selected *gmaS* as a functional biomarker to assess the distribution of this pathway amongst MMA-utilising bacteria. Since currently available *gmaS* primers are specific to the marine *Roseobacter* clade (MRC) (Chen, 2012), we designed two new primer sets covering *gmaS* of non-marine bacteria. Suitable primer regions were identified by alignment of *gmaS* sequences
410 obtained from (i) isolates confirmed to use the GMAS / NMG mediated pathway and (ii) published bacterial genomes. Due to sequence similarity between the two genes, a number of *glnA* gene sequences were included in the alignment to enable identification of suitable *gmaS* primer binding regions not found in *glnA*.

Sequence alignment and establishment of nucleotide-based and amino acid-based
415 phylogenetic trees clearly separated *glnA* from *gmaS* genes and revealed two distinct *gmaS* clusters dividing (i) *Alphaproteobacteria* and (ii) *Beta-* and *Gammaproteobacteria* (Figure 3).

The alphaproteobacterial *gmaS* cluster was further split into two major subgroups: “Group 1” contained MRC-associated sequences (in a separate sub-cluster), as well as sequences belonging to soil and freshwater bacteria from the orders *Rhodobacterales* and *Rhizobiales*, while “Group 2” contained only *gmaS* sequences from non-marine bacteria of the orders *Rhodospirillales*, *Rhizobiales* and *Sphingomonadales* (Figure 3). For primer design, sequences associated with the MRC were removed from the alignment as they were too divergent from the other sequences to be targeted by the same primers. A common region shared by all remaining *gmaS* sequences was used to design the forward primer (*gmaS_557f*). Two different reverse primers were designed for *Alphaproteobacteria* (α *_gmaS_970r*) and *Beta- / Gammaproteobacteria* (β *_* γ *_gmaS_1332r*), respectively, because no further region of sufficient similarity shared by both groups could be identified (alignments in Supplementary Figures S4a-c).

Specificity of these PCR primer sets was confirmed by amplification and sequencing of *gmaS* from (i) five bacteria known to use the *gmaS*-dependent pathway (ii) MMA enrichments from Movile Cave (iii) Movile Cave biofilm and (iv) soil and lake sediment from a different environment (UEA campus, as described in Material and Methods). All PCR products obtained were of the expected size, i.e. ~410 bp (alphaproteobacterial *gmaS*) and ~770bp (beta- and gammaproteobacterial *gmaS*). With DNA from MMA enrichments, a slightly larger, second band was obtained in addition to the *gmaS* product when using *557f* / *1332r*. This gene fragment shared high sequence identity with a viral coat protein and could not be eliminated by using more stringent PCR conditions due to extremely high similarity with the target gene in the primer binding regions. This alternative amplification product was restricted to Movile Cave enrichment DNA and was avoided by gel excision of the *gmaS* band. All sequences obtained from genomic DNA (i) and clone libraries (a total of 30 randomly selected clones from (ii), (iii) and (iv)) were identified as *gmaS*, confirming specificity of the primers (refer to trees in Figure 3, Supplementary Figure S5).

The *gmaS* sequences obtained from Movile Cave DNA affiliated with *gmaS* from both methylotrophic and non-methylotrophic bacteria identified by DNA-SIP and isolation work in this study, namely *Methylobacterium*, *Catellibacterium*, *Pseudomonas* and *Acinetobacter* (99-100% similarity, Figure 3). A further sequence loosely affiliated with *Methylotenera*, *Methylovorus* and *Methylophaga* (89-90% similarity with all three genera). A final *gmaS* sequence was related to *gmaS* from the methylotroph *Hyphomicrobium* (99% similarity) which had not been detected by DNA-SIP or isolation.

[Figure 3]

Distribution of gmaS and mauA genes in Movile Cave isolates

To assess the distribution of the direct and indirect MMA-oxidation pathways in Movile Cave, bacterial isolates were screened for the presence of *mauA* and *gmaS* genes. While the *mauA*-dependent, direct MMA oxidation pathway is so far only known to exist in bacteria using MMA as a carbon source (i.e. methylotrophs), the *gmaS*-dependent, indirect pathway has recently been shown to also exist in bacteria using MMA for nitrogen nutrition only (i.e. non-methylotrophs) (Chen *et al.*, 2010b). Using the *gmaS* primer sets developed in this study, PCR and sequence analysis of DNA from isolates revealed the presence of *gmaS* in all eight non-methylotrophic MMA-utilising bacteria and in all seven methylotrophic MMA-utilisers (Table 2). Phylogenetic analysis placed the retrieved *gmaS* sequences within the alphaproteobacterial and the beta-/ gammaproteobacterial clusters as expected. Interestingly however, *gmaS* from *Aminobacter*, *Paracoccus*, *Catellibacterium*, *Mesorhizobium* and *Rhodobacter* formed a distinct subgroup within the *Alphaproteobacteria*, separate from the other freshwater and soil group, and separate from the marine group (Figure 3). *mauA* was detected in addition to *gmaS* in four of the seven methylotrophic isolates. These data indicate that the *gmaS* gene is widespread amongst MMA utilising bacteria in Movile Cave.

[Table 2]

Discussion

470 *Methylated amine-utilising methylotrophs in Movile Cave*

The combination of SIP and cultivation proved very effective for the identification of methylotrophs. DNA-SIP results revealed *Methylotenera mobilis* as one of the major MMA-utilising methylotrophs in Movile Cave, which is in agreement with previous studies which showed high abundance of this organism (Chen *et al.*, 2009). While resisting all isolation
475 attempts, at 96 hours in SIP incubations *M. mobilis* was the first organism that responded to addition of MMA.

The combination of cultivation-based studies and SIP furthermore revealed that a new methylotroph, *Catellibacterium* sp. LW-1, is an active MMA utiliser in Movile Cave. Growth studies were essential in consolidating DNA-SIP results and confirming *Catellibacterium* as a
480 novel methylotroph and active MMA-utilising bacterium in Movile Cave. These results also highlight the benefit of analysing SIP enrichments at different time points.

Data from SIP enrichments also suggested that *Cupriavidus*, *Porphyrobacter* and *Altererythrobacter* might play a major role in methylotrophic MMA utilisation alongside known methylotrophs such as *Methylobacterium* and *Methylovorus*. While these organisms
485 were not isolated from the cave and have hence not been tested for growth with methylated amines, published genomes of some *Cupriavidus* / *Ralstonia* species contain *gmaS* (refer to trees in Figure 3, Supplementary Figure S5)

Use of methylated amines by non-methylotrophic bacteria in Movile Cave

490 The large variety of bacterial isolates in Movile Cave using methylated amines as nitrogen sources but not as carbon sources is intriguing, considering the relatively high standing

concentrations of ammonium present in Movile Cave water. It is possible that ammonium-depleted areas exist within the microbial mats where utilisation of MMA is advantageous. The fact that nitrogen in the mat is isotopically light while ammonium in the cave water is heavy (Sarbu *et al.*, 1996) could be explained by isotopic fractionation during ammonium assimilation and nitrification. It may however also indicate that a nitrogen source other than ammonium is used. When growing methylotrophically, some bacterial species have been shown to use the nitrogen of MMA, even when high ammonium concentrations are present (Bellion *et al.*, 1983). The high concentrations of ammonium may even be partly due to release of excess nitrogen by bacteria using MMA as both a carbon and nitrogen source.

Distribution of the gmaS gene and its use as a biomarker

The newly developed PCR primers targeting *gmaS* were successful in the detection of MMA-utilising bacteria not covered by currently available primers which target *mauA*-containing methylotrophs. Results from functional gene screening of non-methylotrophic Movile Cave isolates support previous findings (Chen *et al.*, 2010a) which showed that the *gmaS*-dependent pathway is used by the non-methylotroph *Agrobacterium tumefaciens*. Taken together, these results suggest that the *gmaS* pathway may be the major mode of MMA utilisation in bacteria using MMA as a nitrogen, but not as a carbon, source. Based on our results, the *gmaS*-dependent pathway also appears to be present in the majority of methylotrophic MMA-utilising bacteria. The direct, MMA dehydrogenase (*mauA*) dependent pathway, which was detected in a number of methylotrophic isolates in addition to *gmaS*, seems to be restricted to certain groups of methylotrophic bacteria. It will be interesting to understand how the two pathways are regulated under different growth conditions in organisms containing both.

515

Conclusions

Combining DNA-SIP and isolation studies, key methylotrophs in Movile Cave were identified and it was shown that methylated amines are important intermediates in Movile Cave, serving as a source of carbon, energy and / or nitrogen for a wide range of bacteria. The
520 GMAS / NMG mediated pathway appears to be widespread among both methylotrophic and non-methylotrophic MMA utilisers and newly developed primer sets targeting *gmaS* have great potential as biomarkers for identification of MMA-utilising bacteria.

Acknowledgements

We thank Vlad Voiculescu and Mihai Baci, Rich Boden and Sharmishta Dattagupta for help
525 in sampling Movile Cave, Rich Boden for help and advice in experimental design and discussions, Serban Sarbu for his advice and encouragement and Andy Johnston and Andrew Crombie for their insightful comments on the work. We acknowledge funding from the Natural Environment Research Council to JCM (NE/G017956) and YC (NE/H016236) and University of Warwick and University of East Anglia Earth and Life Systems Alliance
530 postgraduate research scholarships to DW.

We are also grateful to the custodian of Movile Cave, the Group for Underwater and Speleological Exploration (GESS), for letting us use its field station in Mangalia and for providing the logistic support for sampling trips.

Supplementary information is available at *The ISME Journal's* website.

Conflict of interest

The authors declare no conflict of interest.

References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. (1990). Basic local alignment search
540 tool. *J Mol Biol* **215**: 403–410.
- Anthony C. (1982). *The biochemistry of methylotrophs*. Academic Press: London.
- Barrett EL, Kwan HS. (1985). Bacterial reduction of trimethylamine oxide. *Annu Rev
Microbiol* **39**:131–149.
- Bellion E, Bolbot J. (1983). Nitrogen assimilation in facultative methylotrophic bacteria.
545 *Curr Microbiol* **9**: 37-44.
- Bellion E, Hersh LB. (1972). Methylamine metabolism in a *Pseudomonas* species. *Arch
Biochem Biophys* **153**: 368–374.
- Bellion E, Kent ME, Aud JC, Alikhan MY, Bolbot JA. (1983.) Uptake of methylamine and
methanol by *Pseudomonas* sp. strain AM1. *J Bacteriol* **154**: 1168–1173.
- 550 Berman T, Bronk DA. (2003). Dissolved organic nitrogen: a dynamic participant in aquatic
ecosystems. *Aquat Microb Ecol* **31**: 279-305.
- Bicknell B, Owens JD. (1980). Utilization of methyl amines as nitrogen sources by non-
methylotrophs. *J Gen Microbiol* **117**: 89-96.
- Boden R, Thomas E, Savani P, Kelly DP, Wood AP. (2008). Novel methylotrophic bacteria
555 isolated from the River Thames (London, UK). *Environ Microbiol* **10**: 3225-3236.
- Brooke AG, Attwood MM. (1984). Methylamine uptake by the facultative
methylotroph *Hyphomicrobium* X. *J Gen Microbiol* **130**: 459–463.

- Budd JA, Spencer CP. (1968). The utilisation of alkylated amines by marine bacteria. *Mar Biol* **2**: 92-101.
- 560 Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K *et al.* (2009). BLAST+: architecture and applications. *BMC Bioinformatics* **10**: 421.
- Campbell KA. (2006). Hydrocarbon seep and hydrothermal vent paleoenvironments and paleontology: past developments and future research directions. *Palaeogeogr Palaeocl* **232**: 362–407.
- 565 Chen Y. (2012) Comparative genomics of methylated amine utilization by marine *Roseobacter* clade bacteria and development of functional gene markers (*tmm*, *gmaS*). *Environ Microbiol* **14**: 2308–2322
- Chen Y, McAleer K L, Murrell JC. (2010b). Monomethylamine as a nitrogen source for a non-methylotrophic bacterium, *Agrobacterium tumefaciens*. *Appl Environ Microbiol* **76**:
570 4102-4104.
- Chen Y, Scanlan J, Song L, Crombie A, Rahman MT, Schäfer H *et al.* (2010a). Gamma-glutamylmethylamide is an essential intermediate for monomethylamine metabolism in *Methylocella silvestris*. *Appl Environ Microbiol* **76**: 4530-4537.
- Chen Y, Wu L, Boden R, Hillebrand A, Kumaresan D, Moussard H *et al.* (2009). Life
575 without light: microbial diversity and evidence of sulfur- and ammonium-based chemolithotrophy in Movile Cave. *ISME J* **3**: 1093-1104.
- Chistoserdova L. (2011). Modularity of methylotrophy, revisited. *Environ Microbiol* **13**: 2603-2622.

- Chistoserdova L, Kalyuzhnaya MG, Lidstrom ME. (2009). The expanding world of
580 methylotrophic metabolism. *Annu Rev Microbiol* **63**: 477–499.
- Colby J, Zatman LJ. (1973). Trimethylamine metabolism in obligate and facultative
methylotrophs. *Biochem J* **132**: 101-12.
- Delong EF. (1992). Archaea in coastal marine environments. *Proc Natl Acad Sci USA* **89**:
5685–5689.
- 585 Felsenstein J. (1985). Confidence limits on phylogenies: An approach using the bootstrap.
Evolution (N Y) **39**: 783-791.
- Fitzsimons MF, Millward GE, Revitt DM, Dawit MD. (2006). Desorption kinetics of
ammonium and methylamines from estuarine sediments: consequences for the cycling of
nitrogen. *Mar Chem* **101**: 12-26.
- 590 Gibb SW, Mantoura RFC, Liss PS, Barlow RG. (1999). Distributions and biogeochemistries
of methylamines and ammonium in the Arabian Sea. *Deep-Sea Res. II* **46**: 593-615.
- Glenn AR, Dilworth MJ. (1984). Methylamine and ammonium transport system in *Rhizobium*
leguminosarum MNF 3841. *J Gen Microbiol* **130**: 1961–1968.
- Hutchens E, Radajewski S, Dumont MG, McDonald IR, Murrell JC. (2004). Analysis of
595 methanotrophic bacteria in Movile Cave by stable isotope probing. *Environ Microbiol* **6**: 111-
120.
- Kanagawa T, Dazai M, Fukuoka S. (1982). Degradation of organo-phosphorus wastes. 2.
Degradation of O,O-dimethyl phosphorodithioate by *Thiobacillus thioparus* TK-1
and *Pseudomonas* AK-2. *Agric Biol Chem.* **46**: 2571-2578.

- 600 Kalyuzhnaya MG, Bowerman S, Jimmie CL, Lidstrom ME, Chistoserdova L. (2006a). *Methylothermobacter mobilis* gen. nov., sp. nov., an obligately methylamine-utilizing bacterium within the family *Methylophilaceae*. *Int J Syst Evol Microbiol* **56**: 2819–2823.
- Kalyuzhnaya MG, De Marco P, Bowerman S, Pacheco CC, Lara JC, Lidstrom, ME *et al.* (2006b). *Methyloversatilis universalis* gen. nov., sp. nov., a novel taxon within
605 the *Betaproteobacteria* represented by three methylotrophic isolates. *Int J Syst Evol Microbiol* **56**: 2517–2522.
- Kelly DP, Wood AP. (1998). Microbes of the sulfur cycle In: Burlage RS, Atlas R, Stahl D, Geesey G, Sayler C (eds). *Techniques in microbial ecology*. Oxford university press: New York, USA, pp. 31–57.
- 610 Kung HF, Wagner C (1969). Gamma-glutamylmethylamide. A new intermediate in the metabolism of methylamine. *J Biol Chem* **244**: 4136–40.
- Lane DJ. (1991). 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M (eds). *Nucleic Acid Techniques in Bacterial Systematics*. John Wiley & Sons Inc: Chichester, UK, pp 115–175.
- 615 Lane DJ, Pace B, Olsen GJ, Stahl DA, Sogin ML, Pace NR. (1985). Rapid determination of 16S ribosomal RNA sequences for phylogenetic analyses. *Proc Natl Acad Sci USA* **82**: 6955–6959.
- Latypova, E, Yang S, Wang YS, Wang T, Chavkin TA, Hackett M, Schäfer H, Kalyuzhnaya MG. (2010). Genetics of the glutamate-mediated methylamine utilization pathway in the
620 facultative methylotrophic beta-proteobacterium *Methyloversatilis universalis* FAM5. *Mol Microbiol* **75**: 426–439.

Levering PR, van Dijken JP, Veenhuis M, Harder W. (1981). *Arthrobacter* P1, a fast growing versatile methylotroph with amine oxidase as a key enzyme in the metabolism of methylated amines. *Arch Microbiol* **129**: 72–80.

625 Lidbury I, Murrell JC, Chen Y. (2014). Trimethylamine *N*-oxide metabolism by abundant marine heterotrophic bacteria. *Proc Natl Acad Sci USA* **111**: 2710-2715.

Lin TY, Timasheff SN. (1994). Why do some organisms use a urea-methylamine mixture as osmolyte? Thermodynamic compensation of urea and trimethylamine *N*-oxide interactions with protein. *Biochemistry US* **33**: 12695–12701.

630 Liu Y, Xu CJ, Jiang JT, Liu YH, Song XF, Li H *et al.* (2010). *Catellibacterium aquatile* sp. nov., isolated from fresh water, and emended description of the genus *Catellibacterium* Tanaka *et al.* 2004. *Int J Syst Evol Microbiol* **60**: 2027-2031.

Lutz RA, Kennish MJ. (1993). Ecology of deep-sea hydrothermal vent communities: A review. *Rev Geophys* **31**: 211–242.

635 Macalady JL, Lyon EH, Koffman B, Albertson LK, Meyer K, Galdenzi S *et al.* (2006). Dominant microbial populations in limestone-corroding stream biofilms, Frasassi cave system, Italy. *Appl Environ Microbiol* **72**: 5596-609.

Markowitz VM, Chen IMA, Palaniappan K, Chu K, Szeto E, Grechkin Y *et al.* (2010). The integrated microbial genomes (IMG) system: an expanding comparative analysis resource.

640 *Nucleic Acids Res* **38**: D382-D390.

Meyer F, Paarmann D, D'Souza M, Olson R, Glass EM, Kubal M *et al.* (2008). The metagenomics RAST server - a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* **9**:386.

- Murrell JC, Lidstrom ME. (1983). Nitrogen metabolism in *Xanthobacter* H4-14. *Arch Microbiol* **136**: 219-221.
- Murrell JC, Whiteley AS. (2011). *Stable Isotope Probing and Related Technologies*. American Society for Microbiology Press: Washington.
- Muyzer G, De Waal EC, Uitterlinden AG. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* **59**: 695-700.
- Neff JC, Chapin FS, Vitousek PM. (2003). Breaks in the cycle: dissolved organic nitrogen in terrestrial ecosystems. *Front Ecol Environ* **1**: 205-211.
- Neufeld JD, Schäfer H, Cox MJ, Boden R, McDonald IR, Murrell, JC (2007a). Stable-isotope probing implicates *Methylophaga* spp and novel *Gammaproteobacteria* in marine methanol and methylamine metabolism. *ISME J* **6**: 480-491.
- Neufeld JD, Vohra J, Dumont MG, Lueders T, Manefield M, Friedrich MW *et al.* (2007b). DNA stable-isotope probing. *Nat Protoc* **2**: 860–866.
- Nicholas KB, Nicholas HB Jr, Deerfield DW. (1997). GeneDoc: Analysis and Visualization of Genetic Variation. *EMBnet News* **4**:14.
- Padden AN, Rainey FA, Kelly DP, Wood AP. (1997). *Xanthobacter tagetididis* sp. nov., an organism associated with *Tagetes* species and able to grow on substituted thiophenes. *Int J Syst Bacteriol* **47**: 394-401.
- Porter ML, Engel AS, Kane TC, Kinkle BK. (2009). Productivity-diversity relationships from chemolithoautotrophically based sulfidic karst systems. *Int J Speleol* **28**: 27-40.

- 665 Radajewski S, Ineson P, Parekh N, Murrell JC. (2000). Stable-isotope probing as a tool in microbial ecology. *Nature* **403**: 646-649.
- Rohwerder T, Sand W, Lascu C. (2003). Preliminary evidence for a sulphur cycle in Movile Cave, Romania. *Acta Biotechnol* **23**: 101-107.
- Saitou N, Nei M. (1987). The neighbor-joining method: A new method for reconstructing
670 phylogenetic trees. *Mol Biol Evol* **4**:406-425.
- Sarbu SM, Kane TC. (1995). A subterranean chemoautotrophically based ecosystem. *Natl Speleol Soc Bull* **57**: 91-98.
- Sarbu SM, Kane TC, Krinkle BK. (1996). A chemoautotrophically based cave ecosystem. *Science* **272**: 1953-1955.
- 675 Sarbu SM, Kinkle BK, Vlasceanu L, Kane TC. (1994). Microbial characterization of a sulfide-rich groundwater ecosystem. *Geomicrobiology J* **12**: 175-182.
- Tamura K, Nei M, Kumar S. (2004). Prospects for inferring very large phylogenies by using the neighbor-joining method. *P Natl Acad Sci USA* **101**:11030-11035.
- Sun S, Chen J, Li W, Altintas I, Lin A, Peltier S *et al.* (2011). Community cyberinfrastructure
680 for Advanced Microbial Ecology Research and Analysis: the CAMERA resource. *Nucleic Acids Res* **39**: D546–D551.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**: 2731-2739
- 685 Tanaka Y, Hanada S, Manome A, Tsuchida T, Kurane R, Nakamura K *et al.* (2004). *Catellibacterium nectariphilum* gen. nov., sp. nov., which requires a diffusible compound

from a strain related to the genus *Sphingomonas* for vigorous growth. *Int J Syst Evol Microbiol* **54**:955-959.

690 Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**: 4876-4882.

Van Dover CL, German CR, Speer KG, Parson LM, Vrijenhoek RC. (2002). Evolution and biogeography of deep-sea venting and seep invertebrates. *Science* **295**: 1253–1257.

705 Vlasceanu L, Popa R, Kinkle BK. (1997). Characterization of *Thiobacillus thioparus* LV43 and its distribution in a chemoautotrophically based groundwater ecosystem. *Appl Environ Microbiol* **63**: 3123-3127

Worsfold PJ, Monbet P, Tappin AD, Fitzsimons MF, Stiles DA, McKelvie ID. (2008). Characterisation and quantification of organic phosphorus and organic nitrogen components in aquatic systems: a review. *Anal Chim Acta* **624**: 37–58.

700 Zhang J, Chen SA, Zheng JW, Cai S, Hang BJ, He J *et al.* (2012). *Catellibacterium nanjingense* sp. nov., a propanil-degrading bacterium isolated from activated sludge, and emended description of the genus *Catellibacterium*. *Int J Syst Evol Microbiol* **62**: 495-499.

705 Zheng JW, Chen YG, Zhang J, Ni YY, Li WJ, He J *et al.* (2011). Description of *Catellibacterium caeni* sp. nov., reclassification of *Rhodobacter changlensis* Anil Kumar *et al.* 2007 as *Catellibacterium changlense* comb. nov. and emended description of the genus *Catellibacterium*. *Int J Syst Evol Microbiol* **61**:1921-1926.

Zuckerandl E, Pauling L. (1965). Evolutionary divergence and convergence in proteins. In: Bryson V, Vogel HJ (eds). *Evolving Genes and Proteins*, Academic Press: New York, pp. 97-166.

710 **Titles and legends to figures**

Figure 1 Denaturing gradient gel electrophoresis analysis of bacterial 16S rRNA gene fragments in native (unfractionated) DNA from incubations of Movile Cave water with ¹³C-MMA (left) and unlabelled MMA (right), after 48 hours (t1), 96 hours (t2) and 4 weeks (t3).

715 **Figure 2** Denaturing gradient gel electrophoresis analysis of bacterial 16S rRNA gene fragments in light and heavy DNA fractions from ¹³C-MMA incubations of Movile Cave water after 96 hours (a) and 4 weeks (c). DGGE profiles of unfractionated DNA of both time points (b) are shown for reference.

Figure 3 Phylogenetic relationship of *gmaS* sequences (135 - 250 amino acids) derived from published bacterial genomes, methylotrophic (solid rectangles / orange font) and non-methylotrophic (hollow rectangles / blue font) bacterial isolates and clone library sequences (triangles / bold print) from Movile Cave. *glnA* sequences present the outgroup. The tree was established using the neighbour-joining method (1000 bootstrap replicates) and the Poisson correction method for computing evolutionary distances. ¹*Rubrobacter xylanophilus* is a member of the *Actinobacteria* although its *gmaS* sequence affiliates with the beta- and 720 gammaproteobacterial cluster. ²*gmaS* sequences containing a total of more than two mismatches across the forward / reverse primer set designed for the respective *gmaS* clusters are marked with an asterisk.

Tables

Table 1 Growth of bacterial isolates from Movile Cave on methylated amines with and without added carbon.

Isolates	Phylogeny	Identity (%)	Growth on methylated amines					
			MMA +C	DMA +C	TMA +C	MMA	DMA	TMA
<i>Alphaproteobacteria</i>								
2W-7	<i>Methylobacterium extorquens</i>	100	+	+	+	+	+	+
LW-13	<i>Xanthobacter tagetidis</i>	100	+	+	+	+	+	+
A2-1D	<i>Paracoccus yeei</i>	100	+	+	+	+	+	+
2W-61	<i>Paracoccus yeei</i>	98	+	+	+	+	+	+
2W-12	<i>Aminobacter niigataensis</i>	100	+	+	+	+	+	+
LW-1	<i>Catellibacterium caeni</i>	99	+	-	+	+	+	+
1M-11	<i>Mesorhizobium loti</i>	99	+	+	+	+	+	+
A2-41x	<i>Shinella yambaruensis</i>	98	+	-	-	-	na	na
1W-5	<i>Rhodobacter blasticus</i>	96	+	+	+	-	-	-
O1	<i>Oleomonas sagaranensis</i>	98	+	+	-	-	-	-
O3	<i>Oleomonas sagaranensis</i>	99	+	-	-	-	-	-
<i>Gammaproteobacteria</i>								
1W-58	<i>Acinetobacter johnsonii</i>	100	+	+	+	-	-	-
2W-62	<i>Acinetobacter lwoffii</i>	100	+	+	-	-	-	-
1W-57Y	<i>Pseudomonas oryzihabitans</i>	99	+	-	-	-	-	-
<i>Betaproteobacteria</i>								
A2-14M	<i>Zoogloea caeni</i>	100	+	+	+	-	-	-

Abbreviations: MMA, monomethylamine; DMA, dimethylamine; TMA, trimethylamine; C, carbon mixture (consisting of sucrose, glucose, fructose, glycerol, pyruvate, acetate); na, not analysed. Carbon sources were supplied at 5 mM, nitrogen sources at 1 mM.

Table 2 Methylated amine metabolism and the presence of functional gene markers in Movile Cave isolates.

Isolate	Methylated amines used as		Functional genes		
	N-source	C-source	<i>gmaS</i>	<i>mauA</i>	
<i>Alphaproteobacteria</i>					
1	<i>Methylobacterium extorquens</i> 2W-7	+	+	+	+
2	<i>Xanthobacter tagetidis</i> LW-13	+	+	+	+
3	<i>Paracoccus yeei</i> A2-1D	+	+	+	+
4	<i>Paracoccus</i> sp. 1W-61	+	+	-	+
5	<i>Aminobacter niigataensis</i> 2W-12	+	+	+	-
6	<i>Catellibacterium</i> sp. LW-1	+	+	+	-
7	<i>Mesorhizobium</i> sp. 1M-11	+	+	+	-
8	<i>Shinella</i> sp. A2-41x	+	-	+	-
9	<i>Rhodobacter</i> sp. 1W-5	+	-	+	-
10	<i>Oleomonas</i> sp. O1	+	-	+	-
11	<i>Oleomonas</i> sp. O3	+	-	+	-
<i>Gammaproteobacteria</i>					
12	<i>Acinetobacter johnsonii</i> 1W-6	+	-	+	-
13	<i>Acinetobacter lwoffii</i> 2W-62	+	-	+	-
14	<i>Pseudomonas</i> sp. 1W-57Y	+	-	+	-
<i>Betaproteobacteria</i>					
15	<i>Zoogloea caeni</i> A2-14M	+	-	+	-

Overview of bacterial isolates from Movile Cave, their capability of using methylated amines as a carbon (C) and / or nitrogen (N) source, and presence of functional genes indicating the direct (*mauA*) or indirect (*gmaS*) methylamine oxidation pathway.

Titles and legends to figures

Figure 1 Denaturing gradient gel electrophoresis analysis of bacterial 16S rRNA gene fragments in native (unfractionated) DNA from incubations of Movile Cave water with ¹³C-monomethylamine (left) and unlabelled monomethylamine (right), after 48 hours (t1), 96 hours (t2) and 4 weeks (t3).


Figure 2 Denaturing gradient gel electrophoresis analysis of bacterial 16S rRNA gene fragments in light and heavy DNA fractions from ¹³C-monomethylamine incubations of Movile Cave water after 96 hours (a) and 4 weeks (c). DGGE profiles of unfractionated DNA of both time points (b) is shown for reference.

Figure 3 Phylogenetic relationship of *gmaS* sequences derived from published bacterial genomes, methylotrophic (solid rectangles) and non-methylotrophic (hollow rectangles) bacterial isolates and clone library sequences (triangles) from Movile Cave. *glnA* sequences present the outgroup. The tree was established using the neighbour-joining method (1000 bootstrap replicates) and the Poisson correction method for computing evolutionary distances. Translated *gmaS* sequences had a length of ~135 amino acids (alphaproteobacterial isolates) and ~250 amino acids for (beta- and gammaproteobacterial isolates).

¹ *Rubrobacter xylophilus* is a member of the *Actinobacteria* although its *gmaS* sequence affiliates with the *Beta*- and *Gammaproteobacterial* cluster.

² *gmaS* sequences containing a total of more than two mismatches across the forward / reverse primer set designed for the respective clusters in this study are marked with an asterisk.

Figures (black and white)



25 **Figure 1**




Figure 2

