

Original citation:

Nayak, M., Libby, J., Malde, S., Thomas, C., Wilkinson, G., Briere, R. A., Naik, P., Gershon, T. J. and Bonvicini, G.. (2015) *First determination of the CP content of $D \rightarrow \pi+\pi-\pi^0$ and $D \rightarrow K+K-\pi^0$* . Physics Letters B, Volume 740 . pp. 1-7. ISSN 0370-2693

Permanent WRAP url:

<http://wrap.warwick.ac.uk/64181>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 3.0 (CC BY 3.0) license and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/3.0/>

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: publications@warwick.ac.uk



<http://wrap.warwick.ac.uk>



First determination of the CP content of $D \rightarrow \pi^+\pi^-\pi^0$ and $D \rightarrow K^+K^-\pi^0$



M. Nayak^a, J. Libby^{a,*}, S. Malde^b, C. Thomas^b, G. Wilkinson^{b,c}, R.A. Briere^d, P. Naik^e, T. Gershon^f, G. Bonvicini^g

^a Indian Institute of Technology Madras, Chennai 600036, India

^b University of Oxford, Denys Wilkinson Building, Keble Road, OX1 3RH, United Kingdom

^c European Organisation for Nuclear Research (CERN), CH-1211, Geneva 23, Switzerland

^d Carnegie Mellon University, Pittsburgh, PA 15213, USA

^e University of Bristol, Bristol, BS8 1TL, United Kingdom

^f University of Warwick, Coventry, CV4 7AL, United Kingdom

^g Wayne State University, Detroit, MI 48202, USA

ARTICLE INFO

Article history:

Received 15 October 2014

Received in revised form 12 November 2014

Accepted 13 November 2014

Available online 18 November 2014

Editor: L. Rolandi

Keywords:

Charm decay

Quantum correlations

CP violation

ABSTRACT

Quantum-correlated $\psi(3770) \rightarrow D\bar{D}$ decays collected by the CLEO-c experiment are used to perform first measurements of F_+ , the fractional CP -even content of the self-conjugate decays $D \rightarrow \pi^+\pi^-\pi^0$ and $D \rightarrow K^+K^-\pi^0$. Values of $0.968 \pm 0.017 \pm 0.006$ and $0.731 \pm 0.058 \pm 0.021$ are obtained for $\pi^+\pi^-\pi^0$ and $K^+K^-\pi^0$, respectively. It is demonstrated how modes of this sort can be cleanly included in measurements of the unitarity triangle angle γ using $B^\mp \rightarrow DK^\mp$ decays. The high CP -even content of $D \rightarrow \pi^+\pi^-\pi^0$, in particular, makes this a promising mode for improving the precision on γ .

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP³.

1. Introduction

Improved knowledge of the unitarity triangle angle γ (also denoted $\phi_3 = \arg(-V_{ud}V_{ub}^*/V_{cd}V_{cb}^*)$) is necessary for testing the Standard Model description of CP violation. Sensitivity to γ can be obtained by studying CP -violating observables in $B^\mp \rightarrow DK^\mp$ decays, where D indicates a neutral charm meson reconstructed in a final state common to both D^0 and \bar{D}^0 mesons. Examples include CP -eigenstates [1], quasi-flavour specific states such as $K^\pm\pi^\mp$ [2], self-conjugate modes such as $K_S^0\pi^+\pi^-$ [3,4] and singly Cabibbo-suppressed decays such as $K_S^0K^\pm\pi^\mp$ [5]. The current world average precision on γ is significantly worse than that of the other angles of the unitarity triangle [6]. Therefore, including additional D -meson final states is desirable to reduce the statistical uncertainty on γ at current and future facilities.

In the case that the D does not decay to a pure CP eigenstate, information is required on the strong decay dynamics in order to relate the CP -violating observables to γ . This information can be obtained from studies of quantum-correlated $D\bar{D}$ mesons pro-

duced in e^+e^- collisions at an energy corresponding to the mass of the $\psi(3770)$ [3,7–9], and relevant measurements have been performed for several decay channels exploiting data collected by CLEO-c [10–15] and BESIII [16].

The decay $D \rightarrow \pi^+\pi^-\pi^0$ is a promising candidate to be added to the suite of modes used in the γ measurement. Its Dalitz plot has been studied by the CLEO and BaBar Collaborations using flavour-tagged D^0 decays and exhibits a strikingly symmetric distribution that suggests the decay may be dominated by a single CP eigenstate [17,18].¹ An isospin analysis [19] of the amplitude model for $D^0 \rightarrow \pi^+\pi^-\pi^0$ presented in Ref. [18] concludes that the final state is almost exclusively $I = 0$. Therefore, given that the parity and G -parity of the three-pion final state is odd and $G = (-1)^I C$, the final state is expected to be $C = -1$ and $CP = +1$. As its branching ratio of $1.43 \pm 0.06\%$ [6] is significantly larger than those of the pure two-body CP -even modes, it has the potential to contribute strongly in any analysis making use of such

¹ Furthermore, the model-dependent analysis in Ref. [18] uses an amplitude model derived from the D^0 sample to search for CP violation in $B^\pm \rightarrow DK^\pm$ decays. This is a different analysis strategy to that made possible through the results presented in this Letter.

* Corresponding author.

E-mail address: libby@iitm.ac.in (J. Libby).

decays. The channel $D \rightarrow K^+K^-\pi^0$ is a similar, but less abundant, self-conjugate mode that has also attracted interest [20,21]. This Letter presents the first analysis of these decays using quantum-correlated $D\bar{D}$ decays, and measurements of their CP content, making use of the CLEO-c $\psi(3770)$ data set. These measurements allow the inclusive decays to be included in future $B^\mp \rightarrow DK^\mp$ analyses in a straightforward and model-independent manner, thus allowing for an improved determination of the angle γ . Throughout the effects of CP violation in charm mesons are neglected, which is a good assumption given theoretical expectations and current experimental limits [6,22].

The remainder of the Letter is structured as follows. Section 2 describes how quantum-correlated D decays are used to determine the CP content. In addition, predictions for the CP content of the state from existing amplitude models are presented. The data set and event selection are described in Section 3. The results and the determination of the systematic uncertainties are presented in Section 4. In Section 5 the implications for the measurement of the unitarity triangle angle γ are discussed. Section 6 gives the conclusions.

2. Measuring the CP content

Consider a $\psi(3770) \rightarrow D\bar{D}$ analysis in which the signal decay mode is $D \rightarrow h^+h^-\pi^0$. Let M^+ designate the number of “double-tagged” candidates, after background subtraction, where one D meson is reconstructed in the signal mode of interest, and the other is reconstructed in a CP -odd eigenstate. The quantum-numbers of the $\psi(3770)$ resonance then require that the signal mode is in a CP -even state, hence the $+$ superscript. The observable M^- is defined in an analogous manner. Let S^+ (S^-) designate the number of “single-tagged” CP -odd (CP -even) candidates in the data sample, where a D meson is reconstructed decaying to a CP eigenstate, with no requirement on the final state of the other D meson in the event. The small effects of $D^0\bar{D}^0$ mixing are eliminated from the measurement by correcting the measured single-tagged yields S_{meas}^\pm such that $S^\pm = S_{\text{meas}}^\pm / (1 - \eta_\pm y_D)$, where η_\pm is the CP eigenvalue of the mode, and $y_D \sim 10^{-2}$ is one of the well-known $D^0\bar{D}^0$ mixing parameters [23]. For a time-integrated measurement at the $\psi(3770)$ there are no effects on the double-tagged yields at leading order in the mixing parameters.

On the assumption that for double-tagged candidates the reconstruction efficiencies of each D meson are independent, then the quantity $N^+ \equiv M^+/S^+$ has no dependence on the branching fractions or reconstruction efficiencies of the CP -eigenstate modes, and can be directly compared with the analogous quantity N^- to gain insight into the CP content of the signal mode. The CP fraction is defined

$$F_+ \equiv \frac{N^+}{N^+ + N^-} \quad (1)$$

and is 1 (0) for a signal mode that is fully CP -even (CP -odd). The notation $F_+(\pi^+\pi^-\pi^0)$ and $F_+(K^+K^-\pi^0)$ is used in the discussion when it is necessary to distinguish between the two final states.

It is also instructive to interpret the observable F_+ making use of the formalism developed in Ref. [8] for binned analyses of self-conjugate three-body final states. Consider the situation where the $D^0 \rightarrow h^+h^-\pi^0$ Dalitz plot is divided into two bins by the line $m^2(h^+\pi^0) = m^2(h^-\pi^0)$. The bin for which $m^2(h^+\pi^0) > m^2(h^-\pi^0)$ is labelled -1 and the opposite bin is labelled $+1$. The CP -tagged populations of these bins, N_i^\pm , normalised by the corresponding single CP -tag yields, is given by

$$N_i^\pm = h_D (K_1 \pm 2c_1 \sqrt{K_1 K_{-1}} + K_{-1}),$$

$$N_{-1}^\pm = h_D (K_{-1} \pm 2c_{-1} \sqrt{K_{-1} K_1} + K_1). \quad (2)$$

Here h_D is a normalisation factor independent of bin number and CP tag. The parameter K_i is the flavour-tagged fraction, being the proportion of decays to fall in bin i in the case that the mother particle is known to be a D^0 meson, for example through tagging the other D meson in the event with a semileptonic decay. The parameter c_i is the cosine of the strong-phase difference between D^0 and \bar{D}^0 decays averaged in bin i and weighted by the absolute decay rate (a precise definition may be found in Ref. [8]). By making use of the relations $N^\pm = \sum_i N_i^\pm$, $\sum_i K_i = 1$ and $c_1 = c_{-1}$ it follows that

$$F_+ = \frac{1}{2} (1 + 2c_1 \sqrt{K_1 K_{-1}}). \quad (3)$$

Therefore the inclusive decay tends to a pure CP eigenstate in the limit that the flavour-tagged Dalitz plot is symmetric, with $K_1 = K_{-1} = 1/2$, and c_1 is -1 or 1 .

Amplitude models of $D^0 \rightarrow \pi^+\pi^-\pi^0$ and $D^0 \rightarrow K^+K^-\pi^0$ are available from studies of flavour-tagged D^0 decays performed by the BaBar Collaboration [18,21]. These models, together with Eq. (3), can be used to calculate predictions for the CP content for each decay. Values of $F_+(\pi^+\pi^-\pi^0) = 0.92$ and $F_+(K^+K^-\pi^0) = 0.64$ are obtained.² The amplitude models are fitted to time-integrated data and include the effects of $D^0\bar{D}^0$ mixing. The biases in the predicted values of F_+ arising from mixing effects are < 0.01 . Other possible biases, associated with the uncertainties in the fitted model components, are expected to be larger, but have not been evaluated.

The CP content of the state $D \rightarrow h^+h^-\pi^0$ also has consequences for the number of self tags M^{self} , which are events containing two $D \rightarrow \pi^+\pi^-\pi^0$ or two $D \rightarrow K^+K^-\pi^0$ candidates. Using the formalism of Ref. [8] for self-tagged events, and once more considering a Dalitz plot divided into two, the number of self-tag candidates in bins i and j is given by

$$M_{ij}^{\text{self}} = 0.5R (K_i K_{-j} + K_{-i} K_j - 2\sqrt{K_i K_{-j} K_{-i} K_j} (c_i c_j + s_i s_j)). \quad (4)$$

Here $R = N_{D\bar{D}} (BR_{h^+h^-\pi^0})^2 \epsilon$, where $N_{D\bar{D}}$ is the number of $D\bar{D}$ pairs in the sample, $BR_{h^+h^-\pi^0}$ is the branching fraction of $D^0 \rightarrow h^+h^-\pi^0$ and ϵ is the detection efficiency. The parameter s_i is the sine of the strong-phase difference between D^0 and \bar{D}^0 decays averaged in bin i and weighted by the absolute decay rate. Employing the same relations as previously, together with $s_1 = -s_{-1}$ and $M^{\text{self}} = \sum_{i,j} M_{ij}^{\text{self}}$, it follows that

$$M^{\text{self}} = R (1 - 4c_1^2 K_1 K_{-1}) = 4RF_+(1 - F_+), \quad (5)$$

where Eq. (3) has been used to express M^{self} in terms of F_+ . Hence the number of self tags vanishes in the case that the signal mode is a CP eigenstate.

3. Data set and event selection

The data set analysed consists of e^+e^- collisions produced by the Cornell Electron Storage Ring (CESR) at $\sqrt{s} = 3.77$ GeV and collected with the CLEO-c detector. The integrated luminosity of the data set is 818 pb^{-1} . The CLEO-c detector is described in detail elsewhere [24]. Simulated Monte Carlo (MC) samples of signal

² The value of $x_0 = 0.850$ reported in Ref. [18] corresponds to a value of F_+ that is very close to the model-derived result given in this Letter.

Table 1
D final states reconstructed in this analysis.

Type	Final states
Signal	$\pi^+\pi^-\pi^0, K^+K^-\pi^0$
CP-even	$K^+K^-, \pi^+\pi^-, K_S^0\pi^0\pi^0, K_L^0\pi^0, K_L^0\omega$
CP-odd	$K_S^0\omega, K_S^0\eta, K_S^0\eta', K_S^0\eta'$

decays are used to estimate selection efficiencies. Possible background contributions are determined from a generic MC sample corresponding to approximately ten times the integrated luminosity of the data set. The EVTGEN generator [25] is used to simulate the decays. The detector response is modelled using the GEANT software package [26].

Table 1 lists the reconstructed D^0 and \bar{D}^0 final states. The unstable final state particles are reconstructed in the following decay modes: $\pi^0 \rightarrow \gamma\gamma$, $K_S^0 \rightarrow \pi^+\pi^-$, $\omega \rightarrow \pi^+\pi^-\pi^0$, $\eta \rightarrow \gamma\gamma$, $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta(\gamma\gamma)\pi^+\pi^-$. The π^0 , K_S^0 , ω , η and η' reconstruction procedure is identical to that used in Ref. [14].

Final states that do not contain a K_L^0 are fully reconstructed via two kinematic variables: the beam-constrained candidate mass, $M_{bc} \equiv \sqrt{s/4c^4 - \mathbf{p}_D^2/c^2}$, where \mathbf{p}_D is the D -candidate momentum, and $\Delta E \equiv E_D - \sqrt{s}/2$, where E_D is the D -candidate energy. The M_{bc} and ΔE distributions of correctly reconstructed D -meson candidates will peak at the nominal D^0 mass and zero, respectively. Neither ΔE nor M_{bc} distributions exhibit any peaking structure for combinatoric background. The double-tagged yield is determined from counting events in signal and sideband regions of M_{bc} after placing requirements on ΔE . The yield determination procedure is identical to that presented in Refs. [10,14] for all modes apart from $h^+h^-\pi^0$ vs. $h^+h^-\pi^0$, which is described separately later in this section.

The selection procedures are almost identical to those presented in Refs. [10,14] apart from that for $h^+h^-\pi^0$ vs. h^+h^- . An additional requirement is placed on the $D^0 \rightarrow h^+h^-$ candidates such that events with the h^\pm compatible with the electron or muon particle identification hypothesis are removed; this criterion removes background from the single-tag sample arising from cosmic ray muons and radiative Bhabha events. These backgrounds distort the combinatoric background shape of the single-tag M_{bc} distribution, such that it cannot be fit readily by an analytic function. (The single-tag yield determination is described later in this section.)

The only final state not considered in Refs. [10,14] is $h^+h^-\pi^0$, for which the criterion $-58.3 < \Delta E < 35.0$ MeV is applied. This ΔE criterion corresponds to a range of approximately three times the experimental resolution around zero. In addition, to suppress background from $D^0 \rightarrow K_S^0\pi^0$ in the $D^0 \rightarrow \pi^+\pi^-\pi^0$ selection, requirements are placed on the vertex of the $\pi^+\pi^-$ pair to be consistent with originating from the e^+e^- collision point. Fig. 1 shows the M_{bc} distributions for CP-tagged signal candidates, summed over all CP-even and CP-odd tags, respectively, where the CP-tag final state does not contain a K_L^0 meson.

Many K_L^0 mesons produced do not deposit any reconstructible signal in the detector. However, double-tag candidates can be fully reconstructed using a missing-mass squared (M_{miss}^2) technique [27] for tags containing a single K_L^0 meson. Yields are extracted from the signal and sideband regions of the M_{miss}^2 distribution. Fig. 2 shows the M_{miss}^2 distributions for candidates tagged with either a $K_L^0\pi^0$ or $K_L^0\omega$ tag.

In the selection of $\pi^+\pi^-\pi^0$ vs. $\pi^+\pi^-\pi^0$ candidates an additional K_S^0 veto is applied to remove $D \rightarrow K_S^0(\pi^+\pi^-)\pi^0$ decays that are otherwise found to dominate the sample. Candidates are rejected if they have a $\pi^+\pi^-$ invariant mass within 30 MeV/ c^2

of the nominal K_S^0 mass. The selected sample of $\pi^+\pi^-\pi^0$ vs. $\pi^+\pi^-\pi^0$ candidates contains a significant combinatoric background from continuum $e^+e^- \rightarrow u\bar{u}, d\bar{d}$ events that hadronise to six pions. Furthermore, this combinatoric background does not follow a uniform distribution in M_{bc} as in the other double-tag modes. Therefore, an alternative strategy is used to determine the signal yield. A maximum-likelihood fit to the distribution of the average M_{bc} of the two $D \rightarrow \pi^+\pi^-\pi^0$ candidates is used to determine the signal yield. The probability density functions (PDFs) are parametrised by a Crystal Ball function [28] and a threshold ARGUS function [29] for the signal and combinatoric background components, respectively. Apart from the signal yield all other parameters of the signal PDF are fixed to those obtained from the signal MC sample. All parameters for the background PDF are obtained from the fit to data. The average M_{bc} distribution for $\pi^+\pi^-\pi^0$ vs. $\pi^+\pi^-\pi^0$ candidates is shown in Fig. 3(a), along with the result of the fit. No significant signal is observed. Even though the combinatoric background in the sample of $K^+K^-\pi^0$ vs. $K^+K^-\pi^0$ candidates is smaller, the same method is applied to determine the signal yield as for $\pi^+\pi^-\pi^0$ vs. $\pi^+\pi^-\pi^0$. The average M_{bc} distribution and fit result for $K^+K^-\pi^0$ vs. $K^+K^-\pi^0$ candidates is shown in Fig. 3(b); again no significant signal is observed.

The peaking background estimates are determined from the generic MC sample of $D\bar{D}$ events. For final states without a K_L^0 the peaking backgrounds are found to be extremely small or negligible. The peaking backgrounds are significant in the states tagged by $K_L^0\pi^0$ and $K_L^0\omega$ as shown in Fig. 2. The dominant source of peaking background is $D^0 \rightarrow K_S^0X, K_S^0 \rightarrow \pi^0\pi^0$ ($X = \pi^0, \omega$) events where the π^0 mesons from the K_S^0 decay are not reconstructed.

The measured event yields after background subtraction are given in Table 2. No significant signal is seen in any of the modes tagged by a CP-even eigenstate, whereas significant signals are seen in most modes tagged by CP-odd eigenstates.

It is also necessary to know the single-tag yield for the CP-eigenstates to normalise the double-tagged yields appropriately to obtain a value of F_+ . The selection of single tags is only possible for modes without a K_L^0 in the final state. The selection criteria are identical to those for the double-tag selection. The signal yield is estimated using a maximum likelihood fit to the M_{bc} distribution where the signal is modelled by the sum of a Gaussian and an asymmetric Gaussian and the combinatoric background is modelled by an ARGUS function. Apart from the signal yield all other parameters of the signal PDF are fixed to those obtained from the signal MC sample. All parameters for the background PDF are obtained from the fit to data. The signal yield is estimated by integrating the best-fit PDF within the interval $1.86 < M_{bc} < 1.87$ GeV/ c^2 . Peaking backgrounds are estimated from the generic $D\bar{D}$ simulation. Significant contributions are only found for $D^0 \rightarrow K_S^0\omega$ and $D^0 \rightarrow K_S^0\eta'(\pi^+\pi^-\pi^0)$ candidates, corresponding to 1.9% and 3.8% of the signal, respectively; in both cases the dominant sources of peaking background are states with intermediate K^*, K_1 and K_2 resonances which lead to $K_S^0\pi^+\pi^-\pi^0$ in the final state. The background-subtracted single-tag yields are given in Table 2.

4. Results, systematic uncertainties and consistency checks

The yields of double-tagged and single CP-tag candidates are used to determine the quantities N^+ and N^- , and from these the CP fraction F_+ . The values for N^+ and N^- are calculated from the ensemble of CP-odd and CP-even tags, respectively, accounting for statistical and systematic uncertainties, and allowing for the correlations that exist between certain systematic components.

There is an uncertainty in the single-tag yields S_{meas}^\pm associated with the fit function used to model the M_{bc} distribution




Fig. 1. M_{bc} distributions for $D \rightarrow \pi^+\pi^-\pi^0$ candidates tagged by CP-even (a) and CP-odd (b) eigenstates; corresponding plots for $D \rightarrow K^+K^-\pi^0$ for CP-even (c) and CP-odd (d). Tags involving a K_L^0 are not included. The vertical dotted lines indicate the applied signal window.




Fig. 2. M_{miss}^2 distributions for $D \rightarrow \pi^+\pi^-\pi^0$ (a) and $D \rightarrow K^+K^-\pi^0$ candidates tagged by CP eigenstates that contain a K_L^0 . The shaded histogram indicates the peaking background. The vertical dotted lines indicate the applied signal window.

of the signal. This shape of the distribution varies depending on whether there are no electromagnetic neutral final-state particles present (K^+K^- and $\pi^+\pi^-$), whether the neutrals are relatively hard ($K_S^0\pi^0(\gamma\gamma)$ and $K_S^0\eta(\gamma\gamma)$) or soft (all other modes). Uncertainties are assigned of 2.0%, 2.5% and 5.0%, respectively. These assignments also adequately cover those uncertainties related to the assumption of the double-tag efficiency factorising into the product of the two single-tag efficiencies. S_{meas}^\pm is corrected for the effects of $D^0\bar{D}^0$ mixing using $y_D = 0.62 \pm 0.08$ [23].

Tags involving a K_L^0 require special treatment as it is not possible to measure a single-tag yield for these modes. The expected

value for the tag $K_L^0\pi^0$ without mixing effects, $S^-(K_L^0\pi^0)$, is given by $2N_{D\bar{D}}\epsilon_{K_L^0\pi^0}\mathcal{B}_{h^+h^-\pi^0}$. Here $\epsilon_{K_L^0\pi^0}$ is an effective single tag efficiency, taken to be equal to the ratio of the double-tagged efficiency to the single-tagged signal efficiency, as determined from simulation and $\mathcal{B}_{h^+h^-\pi^0}$ is the $D \rightarrow h^+h^-\pi^0$ branching fraction [6]. The number of $D\bar{D}$ pairs in the sample, $N_{D\bar{D}}$, can be measured from the double-tagged yield of decays into Cabibbo-favoured final states. It is found that $S^-(K_L^0\pi^0) = 24433 \pm 3934$, where the assigned error reflects the uncertainties in the input factors and assumptions of this calculation. A similar procedure for $K_L^0\omega$ yields $S^-(K_L^0\omega) = 8923 \pm 4015$.




Fig. 3. Average M_{bc} distributions (points with error bars) for (a) $\pi^+\pi^-\pi^0$ vs. $\pi^+\pi^-\pi^0$ and (b) $K^+K^-\pi^0$ vs. $K^+K^-\pi^0$. Superimposed are the total (solid line), signal (dashed line) and background (dotted line) fit results. The shaded histogram indicates the peaking background.

Table 2

Background subtracted signal yields and statistical uncertainties for double tags and the CP single-tag yields.

Tag	$\pi^+\pi^-\pi^0$	$K^+K^-\pi^0$	Single
$\pi^+\pi^-\pi^0$	34.3 ± 20.0	–	–
$K^+K^-\pi^0$	–	2.7 ± 3.6	–
K^+K^-	3.9 ± 5.5	11.3 ± 4.1	11970 ± 116
$\pi^+\pi^-$	13.3 ± 6.6	1.7 ± 3.8	5595 ± 109
$K_S^0\pi^0\pi^0$	1.9 ± 3.4	2.8 ± 2.0	7306 ± 125
$K_L^0\pi^0$	14.6 ± 7.9	10.6 ± 4.0	–
$K_L^0\omega$	-4.3 ± 4.0	3.5 ± 4.4	–
$K_S^0\pi^0$	214.0 ± 15.0	26.0 ± 5.2	20069 ± 146
$K_S^0\omega$	95.5 ± 9.9	11.6 ± 3.4	7960 ± 99
$K_S^0\eta(\gamma\gamma)$	33.0 ± 5.8	3.5 ± 2.4	2903 ± 71
$K_S^0\eta(\pi^+\pi^-\pi^0)$	8.8 ± 3.1	1.0 ± 1.0	1161 ± 48
$K_S^0\eta'$	17.6 ± 4.2	3.0 ± 1.7	1405 ± 38

Finally, there is a possible source of bias arising from non-uniformities in the Dalitz acceptance. The efficiency of reconstruction at CLEO-c is rather flat across phase space, but residual variations are parameterised and used to weight the amplitude models for the two signal modes, and the resulting effective values of F_+ are then calculated. The potential bias is assessed to be 0.001 for $\pi^+\pi^-\pi^0$, and 0.010 for $K^+K^-\pi^0$. The significant difference between the two values is attributed to the larger fraction of events in the CP -odd Dalitz plot for $D \rightarrow K^+K^-\pi^0$, which will be distributed differently than those in the CP -even Dalitz plot. Therefore, the measured value of F_+ is affected more significantly by efficiency variations than for $D \rightarrow \pi^+\pi^-\pi^0$.

The measured values for N^+ and N^- for the two signal modes are displayed in Fig. 4. It can be seen that there is consistency between the individual tags for each measurement. From these results it is determined that $F_+(\pi^+\pi^-\pi^0) = 0.968 \pm 0.017 \pm 0.006$ and $F_+(K^+K^-\pi^0) = 0.731 \pm 0.058 \pm 0.021$, where the first uncertainty is statistical and the second is systematic. These values are slightly higher than, but compatible with, the model predictions reported in Section 2.

The results for the number of self tags are not used in the determination of the CP -even content because they provide much weaker constraints than the CP -tags, and because the K_S^0 veto imposed on the invariant mass of $\pi^+\pi^-$ pairs for the $\pi^+\pi^-\pi^0$ vs. $\pi^+\pi^-\pi^0$ selection distorts the Dalitz space for this mode. Instead, the self-tagged yields are employed as a cross-check. Making use of Eq. (5) and taking the above values for the $F_+(\pi^+\pi^-\pi^0)$ and $F_+(K^+K^-\pi^0)$ leads to an expectation of 2.6 ± 1.6 and 0.4 ± 0.1

candidates for the number of $D \rightarrow \pi^+\pi^-\pi^0$ and $D \rightarrow K^+K^-\pi^0$ self tags, respectively. These predictions are compatible with the measured values reported in Table 2.

5. Implications for the measurement of γ

Sensitivity to the unitarity triangle angle γ is obtained by measuring the relative rates of $B^\mp \rightarrow D(h^+h^-\pi^0)K^\mp$ decays and related observables. Once more considering the Dalitz plot to be divided into a pair of symmetric bins, as introduced in Section 2, and making use of the relations of Ref. [8] and Eq. (3), it follows that

$$\begin{aligned} \Gamma(B^\mp \rightarrow D(h^+h^-\pi^0)K^\mp) &= h_B \left([1 + r_B^2] [1 - 2c_1 y_D \sqrt{K_1 K_{-1}}] \right. \\ &\quad \left. + 2x_\mp [2c_1 \sqrt{K_1 K_{-1}} - y_D] \right) \\ &= h_B \left([1 + r_B^2] [1 - (2F_+ - 1)y_D] \right. \\ &\quad \left. + 2x_\mp [(2F_+ - 1) - y_D] \right). \end{aligned} \quad (6)$$

Here h_B is a normalisation factor, r_B is the ratio of the magnitudes of the $B^+ \rightarrow D^0 K^+$ and $B^+ \rightarrow \bar{D}^0 K^+$ amplitudes, δ_B is the strong-phase difference between these amplitudes and $x_\pm = r_B \cos(\delta_B \pm \gamma)$. This expression includes the effects of $D^0 \bar{D}^0$ oscillations at leading order in the mixing parameters [30].

These partial widths and those involving flavour-specific D meson decays can be used to construct the partial-widths ratio R_{F_+} and CP -asymmetry A_{F_+} :

$$R_{F_+} \equiv \frac{\Gamma(B^- \rightarrow D_{F_+} K^-) + \Gamma(B^+ \rightarrow D_{F_+} K^+)}{\Gamma(B^- \rightarrow D^0 K^-) + \Gamma(B^+ \rightarrow \bar{D}^0 K^+)}, \quad (7)$$

$$A_{F_+} \equiv \frac{\Gamma(B^- \rightarrow D_{F_+} K^-) - \Gamma(B^+ \rightarrow D_{F_+} K^+)}{\Gamma(B^- \rightarrow D_{F_+} K^-) + \Gamma(B^+ \rightarrow D_{F_+} K^+)}, \quad (8)$$

where D_{F_+} indicates a D meson of CP -even content F_+ , established through its decay into the final state $h^+h^-\pi^0$. These observables are directly analogous to the usual so-called GLW [1] observables $R_{CP\pm}$ and $A_{CP\pm}$, where the D meson is reconstructed in a pure CP eigenstate.³

In order to make explicit the relationship to the pure CP -eigenstate case, the effects of mixing are now neglected. Then R_{F_+} and

³ Experimentally, and following the usual procedure established in GLW analyses for measuring $R_{CP\pm}$ [31–34], it is more convenient to determine R_{F_+} from the double ratio $R_{F_+} \approx \frac{\Gamma(B^- \rightarrow D_{F_+} K^-) + \Gamma(B^+ \rightarrow D_{F_+} K^+)}{\Gamma(B^- \rightarrow D_{F_+} \pi^-) + \Gamma(B^+ \rightarrow D_{F_+} \pi^+)} / \frac{\Gamma(B^- \rightarrow D_{K^-\pi^+} K^-) + \Gamma(B^+ \rightarrow D_{K^+\pi^-} K^+)}{\Gamma(B^- \rightarrow D_{K^-\pi^+} \pi^-) + \Gamma(B^+ \rightarrow D_{K^+\pi^-} \pi^+)}$, where in the second ratio the D mesons are reconstructed in the Cabibbo-favoured decay.




Fig. 4. $D \rightarrow \pi^+\pi^-\pi^0$ results for N^+ (a) and N^- (b). $D \rightarrow K^+K^-\pi^0$ results for N^+ (c) and N^- (d). In each plot the vertical (yellow in the web version) band indicates the value obtained from the combination of all tags.

A_{F_+} are found to have the following dependence on the underlying physics parameters:

$$R_{F_+} = 1 + r_B^2 + (2F_+ - 1) \cdot 2r_B \cos \delta_B \cos \gamma, \quad (9)$$

$$A_{F_+} = (2F_+ - 1) \cdot 2r_B \sin \delta_B \sin \gamma / R_{F_+}, \quad (10)$$

which reduces to the equivalent expressions for $R_{CP\pm}$ and $A_{CP\pm}$ in the case F_+ is 1 or 0. Therefore inclusive final states such as $h^+h^-\pi^0$ may be cleanly interpreted in terms of γ and the other parameters of interest, provided that F_+ is known. At leading order the only difference that the CP asymmetry A_{F_+} has with respect to the pure CP -eigenstate case is a dilution factor of $(2F_+ - 1)$, which is 0.936 ± 0.036 for $D \rightarrow \pi^+\pi^-\pi^0$ and 0.462 ± 0.124 for $D \rightarrow K^+K^-\pi^0$. The measurement of F_+ presented here assumes a uniform acceptance across the Dalitz plot; any non-uniformity is considered as a potential source of systematic uncertainty. Therefore, any non-uniformity of the acceptance over the Dalitz plot for an experiment determining R_{F_+} and A_{F_+} must be corrected for, if necessary, and a suitable systematic uncertainty assigned.

6. Conclusion

Data corresponding to an integrated luminosity of 818 pb^{-1} collected by the CLEO-c experiment in e^+e^- collisions at the $\psi(3770)$ resonance have been analysed for the decays $D \rightarrow \pi^+\pi^-\pi^0$ and $D \rightarrow K^+K^-\pi^0$. Measurements of F_+ , the fractional CP -even content of each decay have been performed. Values of $F_+(\pi^+\pi^-\pi^0) = 0.968 \pm 0.017 \pm 0.006$ and $F_+(K^+K^-\pi^0) =$

$0.731 \pm 0.058 \pm 0.021$ are obtained, where the first uncertainty is statistical, and the second is systematic. It has been demonstrated that such self-conjugate inclusive channels can be cleanly included in measurements of the unitarity-triangle angle γ , using $B^\pm \rightarrow DK^\mp$ decays. The high value of F_+ obtained for $D \rightarrow \pi^+\pi^-\pi^0$ makes this channel, in particular, a valuable addition to the suite of D -decay modes used in the measurement of γ at LHCb and Belle-II. Furthermore, given the large branching fraction, the $D \rightarrow \pi^+\pi^-\pi^0$ state can provide an additional source of CP -even tags for quantum-correlated measurements at the $\psi(3770)$. The sample of $D \rightarrow \pi^+\pi^-\pi^0$ tags would be approximately twice as large as the $D \rightarrow h^+h^-$ sample. However, the formalism needs to be adjusted to incorporate F_+ to account for the small CP -odd component in the final state. Improved precision on the F_+ parameters can be obtained using the larger $\psi(3770)$ data set available at BESIII, and similar measurements can also be performed for other self-conjugate final states.

Acknowledgements

This analysis was performed using CLEO-c data, and as members of the former CLEO Collaboration we thank it for this privilege. We thank Brian Meadows and Kalanand Mishra for useful advice, and for helping us implement the BaBar amplitude models. We have benefited from valuable discussions with Mateo Rama, Anton Poluektov, Bhuvanajyoti Bhattacharya and Jonathan Rosner. We are grateful for support from the UK Science and Technology Facilities Council and the UK-India Education and Research Initiative.

References

- [1] M. Gronau, D. London, How to determine all the angles of the unitarity triangle from $B_d^0 \rightarrow DK_S$ and $B_s^0 \rightarrow D\phi$, Phys. Lett. B 253 (1991) 483; M. Gronau, D. Wyler, On determining a weak phase from CP asymmetries in charged B decays, Phys. Lett. B 265 (1991) 172.
- [2] D. Atwood, I. Dunietz, A. Soni, Enhanced CP violation with $B \rightarrow KD^0(\bar{D}^0)$ modes and extraction of the Cabibbo–Kobayashi–Maskawa angle γ , Phys. Rev. Lett. 78 (1997) 3257, arXiv:hep-ph/9612433; D. Atwood, I. Dunietz, A. Soni, Improved methods for observing CP violation in $B^+ \rightarrow KD$ and measuring the CKM phase γ , Phys. Rev. D 63 (2001) 036005, arXiv:hep-ph/0008090.
- [3] A. Giri, Y. Grossman, A. Soffer, J. Zupan, Determining γ using $B^\pm \rightarrow DK^\pm$ with multibody D decays, Phys. Rev. D 68 (2003) 054018, arXiv:hep-ph/0303187.
- [4] A. Bondar, in: Proceedings of BINP Special Analysis Meeting on Dalitz Analysis, 24–26 Sep. 2002, unpublished.
- [5] Y. Grossman, Z. Ligeti, A. Soffer, Measuring γ in $B^\pm \rightarrow K^\pm(KK^*)_D$ decays, Phys. Rev. D 67 (2003) 071301, arXiv:hep-ph/0210433.
- [6] K.A. Olive, et al., Particle Data Group, Review of particle physics, Chin. Phys. C 38 (2014) 090001.
- [7] M. Gronau, Y. Grossman, J.L. Rosner, Measuring $D^0\bar{D}^0$ mixing and relative strong phases at a charm factory, Phys. Lett. B 508 (2001) 37, arXiv:hep-ph/0103110.
- [8] A. Bondar, A. Poluektov, The use of quantum correlated D^0 decays for ϕ_3 measurements, Eur. Phys. J. C 55 (2008) 51, arXiv:0801.0840 [hep-ex]; A. Bondar, A. Poluektov, Feasibility study of model-independent approach to ϕ_3 measurement using Dalitz plot analysis, Eur. Phys. J. C 47 (2006) 347, arXiv:hep-ph/0510246.
- [9] D. Atwood, A. Soni, Role of charm factory in extracting CKM phase information via $B \rightarrow DK$, Phys. Rev. D 68 (2003) 033003, arXiv:hep-ph/0304085.
- [10] D.M. Asner, et al., CLEO Collaboration, Determination of the $D^0 \rightarrow K^+\pi^-$ relative strong phase using quantum-correlated measurements in $e^+e^- \rightarrow D^0\bar{D}^0$ at CLEO, Phys. Rev. D 78 (2008) 012001, arXiv:0802.2268 [hep-ex].
- [11] D.M. Asner, et al., CLEO Collaboration, Updated measurement of the strong phase in $D^0 \rightarrow K^+\pi^-$ decay using quantum correlations in $e^+e^- \rightarrow D^0\bar{D}^0$ at CLEO, Phys. Rev. D 86 (2012) 112001, arXiv:1210.0939 [hep-ex].
- [12] J. Libby, et al., CLEO Collaboration, Model-independent determination of the strong-phase difference between the decays D^0 and $\bar{D}^0 \rightarrow K_S^0 h^+ h^-$ ($h = \pi, K$) and its impact on the measurement of the CKM angle γ , Phys. Rev. D 82 (2010) 112006, arXiv:1010.2817 [hep-ex]; R.A. Briere, et al., CLEO Collaboration, First model-independent determination of the relative strong-phase between D^0 and $\bar{D}^0 \rightarrow K_S^0 \pi^+ \pi^-$ and its impact on the CKM angle γ/ϕ_3 measurement, Phys. Rev. D 80 (2009) 032002, arXiv:0903.1681 [hep-ex].
- [13] J. Libby, et al., New determination of the $D^0 \rightarrow K^-\pi^+\pi^0$ and $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$ coherence factors and average strong-phase differences, Phys. Lett. B 731 (2014) 197, arXiv:1401.1904 [hep-ex].
- [14] N. Lowrey, et al., CLEO Collaboration, Determination of the $D^0 \rightarrow K^-\pi^+\pi^0$ and $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$ coherence factors and average strong-phase differences using quantum-correlated measurements, Phys. Rev. D 80 (2009) 031105(R), arXiv:0903.4853 [hep-ex].
- [15] J. Insler, et al., CLEO Collaboration, Studies of the decays of the $D^0 \rightarrow K_S^0 K^-\pi^+$ and $D^0 \rightarrow K_S^0 K^+\pi^-$, Phys. Rev. D 85 (2012) 092016, arXiv:1203.3804 [hep-ex].
- [16] M. Ablikim, et al., BESIII Collaboration, Measurement of the $D \rightarrow K^-\pi^+$ strong phase difference in $\psi(3770) \rightarrow D^0\bar{D}^0$, Phys. Lett. B 734 (2014) 227, arXiv:1404.4691 [hep-ex].
- [17] D. Cronin-Hennessey, et al., CLEO Collaboration, Searches for CP violation and $\pi\pi$ S wave in the Dalitz-plot analysis of $D^0 \rightarrow \pi^+\pi^-\pi^+$, Phys. Rev. D 72 (2005) 031102(R), arXiv:hep-ex/0503052 (Erratum); D. Cronin-Hennessey, et al., CLEO Collaboration, Phys. Rev. D 75 (2007) 119904.
- [18] B. Aubert, et al., BaBar Collaboration, Measurement of CP violation parameters with a Dalitz plot analysis of $B^\pm \rightarrow D_{\pi^+\pi^-} K^\pm$, Phys. Rev. Lett. 99 (2007) 251801, arXiv:hep-ex/0703037.
- [19] M. Gaspero, B. Meadows, K. Mishra, A. Soffer, Isospin analysis of D^0 decay to three pions, Phys. Rev. D 78 (2008) 014015, arXiv:0805.4050 [hep-ex].
- [20] C. Cawlfeld, et al., CLEO Collaboration, Measurement of interfering $K^{*+}K^-$ and $K^{*-}K^+$ amplitudes in the decay $D^0 \rightarrow K^+K^-\pi^0$, Phys. Rev. D 74 (2006) 031108(R), arXiv:hep-ex/0606045.
- [21] B. Aubert, et al., BaBar Collaboration, Amplitude analysis of the decay $D^0 \rightarrow K^-K^+\pi^0$, Phys. Rev. D 76 (2007) 011102, arXiv:0704.3593 [hep-ex].
- [22] R. Aaij, et al., LHCb Collaboration, A model-independent unbinned search for CP violation in $D^0 \rightarrow \pi^+\pi^-\pi^0$ decays, LHCb-PAPER-2014-054, Phys. Lett. B (2014), in press.
- [23] Y. Amhis, et al., HFAG, Averages of b-hadron, c-hadron, and τ -lepton properties as of early 2012, arXiv:1207.1158, online updates at <http://www.slac.stanford.edu/xorg/hfag>, numbers taken from June 2014 charm physics update.
- [24] Y. Kubota, et al., CLEO Collaboration, The CLEO II detector, Nucl. Instrum. Methods Phys. Res., Sect. A, Accel. Spectrom. Detect. Assoc. Equip. 320 (1992) 66; D. Peterson, et al., The CLEO III drift chamber, Nucl. Instrum. Methods Phys. Res., Sect. A, Accel. Spectrom. Detect. Assoc. Equip. 478 (2002) 142; M. Artuso, et al., Construction, pattern recognition and performance of the CLEO III LiF-TEA RICH detector, Nucl. Instrum. Methods Phys. Res., Sect. A, Accel. Spectrom. Detect. Assoc. Equip. 502 (2003) 91; R.A. Briere, et al., CLEO-c/CESR-c Taskforces, CLEO-c Collaboration, CLEO-c and CESR-c: a new frontier of weak and strong interactions, Cornell LEPP Report CLNS Report No. 01/1742, 2001.
- [25] D.J. Lange, The EvtGen particle decay simulation package, Nucl. Instrum. Methods Phys. Res., Sect. A, Accel. Spectrom. Detect. Assoc. Equip. 462 (2001) 152.
- [26] R. Brun, et al., GEANT 3.21, CERN Program Library Long Writeup W5013, unpublished.
- [27] Q. He, et al., CLEO Collaboration, Comparison of $D \rightarrow K_S^0 \pi$ and $D \rightarrow K_L^0 \pi$ decay rates, Phys. Rev. Lett. 100 (2008) 091801, arXiv:0711.1463 [hep-ex].
- [28] T. Skwarnicki, A Study of the radiative cascade transitions between the Υ and Υ' resonances, Ph.D. Thesis, DESY F31-86-02, 1986, Appendix E.
- [29] H. Albrecht, et al., ARGUS Collaboration, Search for hadronic $b \rightarrow u$ decays, Phys. Lett. B 241 (1990) 278.
- [30] M. Rama, Effect of $D-\bar{D}$ mixing in the extraction of γ with $B^- \rightarrow D^0 K^-$ and $B^- \rightarrow D^0 \pi^-$ decays, Phys. Rev. D 89 (2014) 014021, arXiv:1307.4384 [hep-ex].
- [31] P. del Amo Sanchez, et al., BaBar Collaboration, Measurement of CP observables in $B^\pm \rightarrow D_{CP} K^\pm$ decays and constraints on the angle γ , Phys. Rev. D 82 (2010) 072004, arXiv:1007.0504 [hep-ex].
- [32] K. Abe, et al., Belle Collaboration, Study of $B^\pm \rightarrow D_{CP} K^\pm$ and $D_{CP}^* K^\pm$ decays, Phys. Rev. D 73 (2006) 051106, arXiv:hep-ex/0601032.
- [33] T. Aaltonen, et al., CDF Collaboration, Measurements of branching fraction ratios and CP asymmetries in $B^\pm \rightarrow D_{CP} K^\pm$ decays in hadron collisions, Phys. Rev. D 81 (2010) 031105, arXiv:0911.0425 [hep-ex].
- [34] R. Aaij, et al., LHCb Collaboration, Observation of CP violation in $B^+ \rightarrow DK^+$ decays, Phys. Lett. B 712 (2012) 203, arXiv:1203.3662 [hep-ex]; R. Aaij, et al., LHCb Collaboration, Phys. Lett. B 713 (2012) 351 (Erratum).