

Original citation:

Thorne, C. J., Lockett, A. S., Bullock, I., Hampshire, S., Begum-Ali, S. and Perkins, G. D. (2015) E-learning in advanced life support--an evaluation by the Resuscitation Council (UK). Resuscitation, 90. pp. 79-84.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/68448>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

© 2015, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

1 e-learning in Advanced Life Support – What factors influence assessment outcome?

2 Thorne CJ^{1,2}, Lockey AS^{2,3}, Kimani PK⁵, Bullock I^{2,4}, Hampshire S², Begum-Ali S², Perkins GD^{1,2,5} on behalf of the
3 Advanced Life Support Subcommittee of the Resuscitation Council (UK)

4

5 ¹ Department of Critical Care Medicine, Heart of England NHS Foundation Trust, Birmingham, UK, B9 5SS

6 ² Resuscitation Council (UK), Tavistock House North, Tavistock Square, London, UK, WC1H 9HR

7 ³ Calderdale & Huddersfield NHS Foundation Trust, Halifax, UK, HX3 0PW

8 ⁴ Royal College of Physicians, London, UK, NW1 4LE

9 ⁵ University of Warwick, Warwick Medical School, Warwick, UK, CV4 7AL

10

11

12 Corresponding Author:

13 Dr CJ Thorne

14 Department of Critical Care

15 Heart of England NHS Foundation Trust

16 Bordesley Green East

17 Birmingham

18 B9 5SS

19 cj.thorne@doctors.org.uk

20

21

22

23

24 Word Count: 3,063

25

26 Abstract Word Count: 257

27

28 Tables: 3

29

30 Figures: 3

31

32 Supplementary material: 1

33

34

35

36

37

38

39

40

41

42

43

44

45 **ABSTRACT**

46 **Aim**

47 To establish variables which are associated with favourable Advanced Life Support (ALS) course assessment
48 outcomes, maximising learning effect.

49
50 **Method**

51 Between 1 January 2013 and 30 June 2014, 8,218 individuals participated in a Resuscitation Council (UK) e-learning
52 Advanced Life Support (e-ALS) course. Participants completed 5-8 hours of online e-learning prior to attending a one
53 day face-to-face course. e-learning access data were collected through the Learning Management System (LMS). All
54 participants were assessed by a multiple choice questionnaire (MCQ) before and after the face-to-face aspect
55 alongside a practical cardiac arrest simulation (CAS-Test). Participant demographics and assessment outcomes were
56 analysed.

57
58 **Results**

59 The mean post e-learning MCQ score was 83.7 (SD 7.3) and the mean post-course MCQ score was 87.7 (SD 7.9). The
60 first attempt CAS-Test pass rate was 84.6% and overall pass rate 96.6%. Participants with previous ALS experience,
61 ILS experience, or who were a core member of the resuscitation team performed better in the post-course MCQ,
62 CAS-Test and overall assessment. Median time spent on the e-learning was 5.2 hours (IQR 3.7-7.1). There was a large
63 range in the degree of access to e-learning content. Increased time spent accessing e-learning had no effect on the
64 overall result (OR 0.98, P=0.367) on simulated learning outcome.

65
66 **Conclusion**

67 Clinical experience through core membership of cardiac arrest teams and previous ILS or ALS training were
68 independent predictors of performance on the ALS course whilst time spent accessing e-learning materials did not
69 affect course outcomes. This supports the blended approach to e-ALS which allows participants to tailor their e-
70 learning experience to their specific needs.

71

72

73

74 INTRODUCTION

75 The Formula for Survival¹ identifies three factors that influence survival from cardiac arrest: high-quality research,
76 efficient education of patient caregivers and an effective chain of survival from the early recognition of cardiac arrest
77 through to post resuscitation care.² Advanced Life Support (ALS) courses, which address both the second and third
78 aspects of this formula, are used internationally to train healthcare personnel how to manage patients in cardiac
79 arrest. Previous studies have linked participation on ALS courses to improved outcomes from cardiac arrest.³⁻⁵
80 Courses use multimodal delivery methods to equip participants with background scientific knowledge, targeted
81 clinical skills and non-technical skill development. This blended learning approach is from course manuals, online e-
82 learning material, didactic lectures, hands-on skill stations and formative assessment. In the United Kingdom (UK)
83 and many other countries, successful completion of an ALS course (or similar) is required for healthcare
84 professionals who manage acutely unwell patients on a regular basis.

85
86 The Resuscitation Council (UK) has a 25 year history in delivering ALS courses.⁶ A total of 20,268 individuals
87 participated in an ALS course between January 2015 and December 2015.^{6,7} In 2011, a strategic decision was taken
88 to meet increasing demand, and to increase the flexibility of learning for participants. The Resuscitation Council (UK)
89 launched a novel e-learning ALS course (e-ALS), as an alternative to the conventional two day face-to-face (c-ALS)
90 course, valuing this key educational approach of blended learning. This constitutes 5-8 hours of pre-course online e-
91 learning, followed by a condensed, focussed one day face-to-face element. A multi-centre randomised control trial
92 (RCT) in 2012⁸ and a large observational study of 27,170 participants in 2015⁹ demonstrated almost identical
93 assessment outcomes for participants enrolled upon either c-ALS or e-ALS. The findings of these two studies
94 consolidated the emerging role of the Resuscitation Council (UK) e-ALS course. Whilst outcome data were
95 comparable in the observational study,⁹ it did not assess the extent to which those participants enrolled on the e-ALS
96 course actually accessed the e-learning material, or its effect on assessment outcomes.

97
98 Previous studies investigating the utility of e-learning all display a common limitation, whereby participants often do
99 not fully access the e-learning material.^{10,11} Jensen et al. investigated e-learning as a means for retaining ALS
100 competency but found that only 57.5% of candidates accessed all of the stipulated modules.¹⁰ Similarly Perkins et al.
101 found that only 64% of candidates accessed pre-course e-learning via a CD prior to attending an ALS course.¹¹ This
102 limitation was acknowledged by the authors, who postulated that any true difference between the control and

103 intervention groups may not have been detected because the intervention had not been implemented effectively.
104 Secondly, it provides challenges for ALS course organisers to establish exactly what extent of e-learning has been
105 undertaken by the participants prior to attending a face-to-face course. Whilst this allows personalisation of the
106 learning experience, it also reduces the standardisation of content delivered to those on an ALS course.
107 Consequently, it is unknown whether making e-learning non-compulsory adversely affects candidate outcome.

108 This study was designed to access the aforementioned observational study data set,⁹ analysing the extent to which
109 participants access pre-requisite e-learning material, establishing the effect on candidate ALS assessment outcome.
110 In doing this, study authors intend to highlight independent predictors of successful ALS course outcome.
111
112

113 **METHODS**

114 ***Setting and Participants***

115 ALS participants voluntarily enrolled on a one-day e-ALS course at one of 94 national training centres. Each
116 candidate registered on the Resuscitation Council (UK) Learning Management System (LMS) prior to attending the
117 course. Participants were from a wide range of healthcare professions and stages of training.
118

119 ***The e-ALS Course***

120 The e-ALS course consists of 5-8 hours of e-learning content covering essential ALS topics. Each candidate is given
121 access to the LMS 8 weeks prior to their course and is asked to complete the 12 electronic learning modules.
122 Additionally, participants receive a physical copy of the ALS course manual at least four weeks before the course
123 date. e-learning progress is monitored by the course centres. Participants are free to choose to personalise their
124 learning experience – undertaking as little or as much of the e-learning preparation as they feel necessary although
125 there are three compulsory modules: ALS in perspective; advanced life support algorithm; non-technical skills
126 (progress data are not routinely collected on the LMS for this module as it was only introduced in 2013).
127 There are nine non-compulsory modules: causes and prevention of cardiac arrest; acute coronary syndromes;
128 monitoring, rhythm recognition and 12 lead ECG; bradycardia, pacing and drugs; tachycardia, cardioversion and
129 drugs; special circumstances; post resuscitation care; arterial blood gas analysis; and decisions relating to
130 resuscitation.

131

132 On completion of the e-learning, participants undertake a compulsory multiple choice questionnaire (MCQ),
133 although their results in this do not affect the participants' post-course outcome. After completing the one-day face
134 to face aspect, each candidate undertakes a post-course MCQ and a practical cardiac arrest management simulation
135 test (CAS-Test). In order to achieve ALS competency participants need to pass both of these aspects. Participants are
136 permitted two attempts at the MCQ and three attempts at the CAS-Test. The pre and post-course MCQs comprise 30
137 different stem questions, with each having four true/false answers, creating a total of 120 questions. The pass mark
138 is 75%. The CAS-Test simulations are criterion based and are well validated.^{12,13} They assess participants' abilities in
139 patient assessment, formulating a treatment plan and leadership of the cardiac arrest team. Overall scores and
140 pass/fail data are recorded.

141

142 ***Statistical analysis***

143 Demographic data were collected on the LMS. Anonymised data were transferred to Microsoft Excel (*Microsoft*
144 *Corporation, Redmond, USA*) and analysed using SPSS 23 (*IBM, Armonk, USA*) and R statistical program Version
145 3.3.1.¹⁴ Categorical baseline characteristics were summarised using counts and percentages while continuous
146 baseline characteristics were summarised using mean, median (IQR, interquartile range) and ranges. Independent t-
147 tests, one-way ANOVAs and linear regression models were utilised to determine differences between continuous
148 variables. Logistic regression was used for dichotomous outcome variables.

149

150 A multivariable logistic regression model was fitted to assess which variables predict whether a trainee passes the
151 CAS-Test on the first attempt. Trainees attending the same course session tend to have similar outcomes⁸ and so the
152 multivariable logistic regression model included a random effects term for course session. A similar model was fitted
153 to assess which variables predict whether a trainee passes the overall test. Odds ratios (OR), 95% confidence
154 intervals and p-values from the multivariable random effects logistic regression models were reported. To assess
155 which variables predict the MCQ score of a trainee in the first attempt, MCQ scores were analysed by fitting a linear
156 mixed model with a random effects term for course session. Mean difference in MCQ scores, 95% confidence
157 intervals and p-values from the linear mixed model were reported. An analysis of standard residuals was carried out
158 and outliers removed. Co-linearity was assessed by independently entering each independent variable into a logistic
159 regression with the remaining variables entered as dependent variables. Collinearity diagnostics were calculated and

160 the variance inflation factor (VIF) in all instances was <1. In all models, missing data were excluded from the
 161 complete case analysis by a listwise deletion. Statistical significance was set at P-values of <0.05.

162

163 **RESULTS**

164 **Demographics**

165 8,218 participants were enrolled on one of 450 e-ALS courses during the study period. Mean age was 32.0 years (SD
 166 8.2). 15 participants started but failed to complete the course. 1.8% of the total participants had a degree of missing
 167 data and these were excluded from the analysis. Any missing data occurred due to incomplete data entry by
 168 participants or local course facilitators on the LMS. Stratified participant demographics are displayed below in table 1
 169 in addition to time spent accessing the e-learning and corresponding pass rates.

Table 1: Participant demographics on the e-ALS course and time spent on e-learning

Characteristics/outcomes	n, (%)	Hours spent on compulsory modules	Hours spent on non-compulsory modules	Total hours spent on e-Learning	Overall pass rate (%)
Healthcare background					
Doctor	6236 (75.9)	0-13.2	0-21.0	0-24.0	6095 (97.8)
Range					
Mean (SD)		1.1 (0.8)	4.1 (2.5)	5.3 (3.0)	
Median (IQR)		0.9 (0.7-1.4)	3.8 (2.6-5.3)	4.9 (3.4-6.7)	
Nurse	1244 (15.1)	0-8.9	0-17.2	0-24.0	1122 (90.9)
Range					
Mean (SD)		1.3 (0.9)	5.4 (3.4)	6.9 (3.9)	
Median (IQR)		1.1 (0.8-1.6)	4.8 (3.4-6.6)	6.2 (4.5-8.5)	
Medical student	534 (6.5)	0-4.7	0-16.0	0-17.6	525 (98.3)
Range					
Mean (SD)		1.1 (0.7)	4.4 (2.2)	5.6 (2.6)	
Median (IQR)		0.9 (0.7-1.3)	4.1 (2.9-5.6)	5.3 (4.0-6.9)	
Operating Department Practitioner	73 (0.9)	0-6.9	0-11.5	0.2-21.4	67 (93.1)
Range					
Mean (SD)		1.3 (1.1)	5.3 (2.7)	7.0 (3.7)	
Median (IQR)		1.0 (0.8-1.4)	5.2 (3.5-7.2)	6.4 (4.8-8.8)	
Ambulance staff/ Paramedic	40 (0.5)	0-6.4	0-18.7	0-22.7	39 (97.5)
Range					
Mean (SD)		1.3 (1.2)	4.7 (3.1)	6.5 (4.0)	
Median (IQR)		1.1 (0.7-1.9)	4.8 (3.3-5.7)	6.4 (4.4-8.0)	
Resuscitation Officer	15 (0.2)	0.6-3.0	4.3-9.5	5.1-10.4	15 (100.0)
Range					
Mean (SD)		1.3 (0.7)	6.1 (1.5)	7.5 (1.7)	
Median (IQR)		1.0 (0.8-2.1)	6.1 (4.8-7.1)	7.5 (5.7-9.2)	
Other	74 (0.9)	0-5.5	0-18.0	0-20.6	62 (84.9)
Range					
Mean (SD)		1.4 (0.9)	6.0 (3.4)	7.8 (4.1)	

Median (IQR)		1.2 (0.9-1.5)	4.8 (3.7-7.5)	6.7 (5.0-9.7)	
Not available	2				

Stage of training

Medical Student	537 (6.5)	0-4.7 1.1 (0.7) 0.9 (0.7-1.3)	0-16.0 4.4 (2.2) 4.1 (2.9-5.6)	0-17.6 5.6 (2.6) 5.3 (4.0-6.9)	526 (98.0)
Foundation Year 1 Doctor	1650 (20.1)	0-7.0 1.1 (0.7) 0.9 (0.7-1.3)	0-21.0 4.0 (2.2) 3.8 (2.7-5.2)	0-21.7 5.2 (2.6) 4.9 (3.6-6.5)	1624 (98.4)
Foundation Year 2 Doctor	1663 (20.2)	0-10.0 1.1 (0.8) 0.9 (0.7-1.3)	0-18.4 4.1 (2.3) 3.9 (2.7-5.2)	0-20.8 5.3 (2.8) 5.0 (3.6-6.6)	1639 (98.6)
Junior Grade Doctor (ST1/ST2)	794 (9.7)	0-9.4 1.2 (0.8) 1.0 (0.7-1.5)	0-20.6 4.3 (2.7) 3.7 (2.6-5.4)	0-24.0 5.5 (3.3) 4.9 (3.5-7.0)	768 (96.8)
Middle Grade Doctor[#]	1465 (17.8)	0-13.2 1.1 (0.8) 0.9 (0.7-1.4)	0-20.8 3.9 (2.5) 3.5 (2.3-5.0)	0-23.5 5.1 (2.9) 4.7 (3.2-6.5)	1434 (97.9)
Senior Grade Doctor[§]	488 (5.9)	0-5.1 1.2 (0.9) 1.0 (0.8-1.5)	0-17.7 4.1 (2.7) 3.7 (2.5-5.3)	0-21.2 5.4 (3.4) 4.9 (3.3-7.1)	469 (96.1)
Junior Nurse (Band 4-6)	1002 (12.2)	0-8.9 1.3 (0.9) 1.1 (0.8-1.6)	0-17.2 5.0 (3.2) 4.9 (3.5-6.7)	0-23.1 7.1 (3.9) 6.4 (4.7-8.7)	886 (88.4)
Senior Nurse (Band 7-9)	395 (4.8)	0-6.8 1.3 (0.9) 1.1 (0.8-1.6)	0-15.4 5.0 (3.2) 4.5 (3.1-6.5)	0-24.0 6.6 (3.8) 5.9 (4.2-8.1)	378 (95.5)
Other	223 (2.7)	0-8.3 1.6 (1.2) 1.2 (0.9-1.9)	0-18.7 5.9 (3.3) 5.3 (3.5-7.7)	0-22.7 7.6 (4.2) 6.9 (4.9-9.5)	202 (90.2)
Not available	1				

Previous ALS experience

No	4615 (56.2)	0-10.0 1.2 (0.8) 1.0 (0.7-1.4)	0-21.0 4.5 (2.7) 4.1 (3.9-7.2)	0-24.0 5.8 (3.2) 5.3 (3.8-7.2)	4411 (95.6)
Yes	3593 (43.8)	0-13.2 1.2 (0.8) 1.0 (0.7-1.4)	0-21.0 4.1 (2.6) 3.8 (2.5-5.3)	0-24.0 5.4 (3.2) 5.3 (3.9-7.2)	3515 (98.0)
Not available	10				

Previous ILS experience*

No	2704				2624
Range	(32.9)	0-8.3	0-21.0	0-24.0	(95.5)
Mean (SD)		1.2 (0.9)	4.5 (2.8)	5.8 (3.4)	
Median (IQR)		1.0 (0.8-1.5)	4.1 (2.7-5.8)	5.3 (3.7-7.4)	
Yes	5466				5302
Range	(67.1)	0-13.2	0-20.9	0-24.0	(97.2)
Mean (SD)		1.1 (0.8)	4.3 (2.6)	5.5 (3.1)	
Median (IQR)		1.0 (0.7-1.4)	4.2 (2.9-5.7)	5.4 (3.8-7.3)	
Not available	48				

Core member of resuscitation team

No	4373				4173
Range	(53.8)	0-9.4	0-21.0	0-23.5	(95.7)
Mean (SD)		1.2 (0.8)	4.5 (2.7)	5.8 (3.2)	
Median (IQR)		1.0 (0.8-1.5)	4.2 (2.9-5.7)	5.4 (3.9-7.3)	
Yes	3759				3668
Range	(46.2)	0-13.2	0-21.0	0-24.0	(97.7)
Mean (SD)		1.1 (0.8)	4.1 (2.6)	4.9 (3.1)	
Median (IQR)		0.9 (0.7-1.4)	3.8 (2.6-5.3)	4.9 (3.5-6.8)	
Not available	86				
Total	8218				7926
Range		0-13.2	0-21.0	0-24.0	(96.6%)
Mean (SD)		1.2 (2.8)	4.3 (2.7)	5.6 (3.2)	
Median (IQR)		1.0 (0.74-1.4)	4.0 (2.7-5.5)	5.2 (3.7-7.1)	

*Immediate Life Support

ST3+, middle grade equivalent

§ Consultant or associate specialist

170

171

Assessment outcomes

172

Assessment outcome data are displayed in Table 2. 99.1% of participants completed the post e-learning MCQ, with a

173

mean score of 83.7 (SD 7.3). The mean post-course MCQ score was 87.7 (SD 7.9). Resuscitation officers had the

174

highest mean score in the post-course MCQ (90.5, SD 5.5), with operating department practitioners (ODP) the lowest

175

(79.2, SD 17.0). Those participants who had previous ALS experience or were a core member of the resuscitation

176

team performed better in the post-course MCQ ($P < 0.001$, $P < 0.001$ respectively), as did the more senior doctors and

177

nurses. Participants with previous ILS experience performed worse in the post-course MCQ ($P < 0.001$).

Table 2: Univariate predictors of assessment outcomes

Independent variables	Mean post e-learning MCQ score	Mean post-course MCQ score	P-value	CAS-Test pass (%)	Odds ratio (95% CI)	P-value	Overall course pass (%)	Odds ratio (95% CI)	P-value
Healthcare profession									
Doctor (comparison)	84.7	88.7	$< 0.001^{\text{f}}$	5352 (86.0)			6095 (97.8)		
Nurse	79.7	80.0		1005 (81.3)	0.71 (0.60-0.83)	< 0.001	1122 (90.9)	0.22 (0.17-0.29)	< 0.001
Medical student	83.4	86.5		425	0.64	< 0.001	525 (98.3)	1.31	0.435

				(79.6)	(0.51-0.79)			(0.66-2.59)	
Operating Department Practitioner	73.0	79.2		51 (70.8)	0.40 (0.24-0.66)	<0.001	67 (93.1)	0.30 (0.12-0.76)	0.011
Ambulance staff/ Paramedic	81.4	85.4		37 (92.5)	2.00 (0.62-6.62)	0.247	39 (97.5)	0.88 (0.12-6.43)	0.897
Resuscitation Officer	86.6	90.5		13 (86.7)	1.06 (0.24-4.69)	0.941	15 (100.0)	3.6x10 ⁶	<0.001
Other	79.9	83.6		46 (66.7)	0.33 (0.20-0.54)	<0.001	62 (84.9)	0.12 (0.06-0.24)	<0.001

Stage of training

Medical Student	83.3	86.4		426 (79.5)	0.72 (0.56-0.92)	0.010	526 (98.0)	0.70 (0.34-1.44)	0.332
Foundation Year 1 Doctor	83.0	86.6		1394 (84.7)	1.03 (0.85-1.24)	0.754	1624 (98.4)	0.92 (0.52-1.60)	0.754
Foundation Year 2 Doctor (comparison)	83.2	87.7		1401 (84.3)			1639 (98.6)		
Junior Grade Doctor (ST1/ST2)	85.2	89.1		667 (85.6)	1.11 (0.87-1.40)	0.406	768 (96.8)	0.45 (0.26-0.79)	0.006
Middle Grade Doctor[#]	87.0	91.1	<0.001 [£]	1322 (90.4)	1.75 (1.40-2.17)	<0.001	1434 (97.9)	0.70 (0.41-1.20)	0.197
Senior Grade Doctor[§]	87.9	92.0		425 (87.3)	1.28 (0.95-1.72)	0.107	469 (96.1)	0.40 (0.22-0.76)	0.005
Junior Nurse (Band 4-6)	78.8	82.8		777 (78.3)	0.67 (0.55-0.82)	<0.001	886 (88.4)	0.12 (0.08-0.19)	<0.001
Senior Nurse (Band 7-9)	81.4	86.6		346 (87.8)	1.34 (0.97-1.87)	0.080	378 (95.5)	0.31 (0.17-0.57)	<0.001
Other	82.6	86.6		163 (74.1)	0.53 (0.38-0.74)	<0.001	202 (90.2)	0.14 (0.08-0.26)	<0.001

Previous life support course experience

Previous ALS experience	85.5	89.7	<0.001 [#]	3204 (89.3)	1.97 (1.73-2.24)	<0.001	3515 (98.0)	2.27 (1.73-2.98)	<0.001
No previous ALS experience	82.3	86.1		3727 (81.0)			4411 (95.6)		
Previous ILS experience	83.2	87.4	<0.001 [#]	4666 (85.6)	1.24 (1.09-1.40)	0.001	5302 (97.2)	1.64 (1.29-2.09)	<0.001
No previous ILS experience	84.5	88.3		2265 (82.7)			2624 (95.5)		
Core member of	84.4	88.8	<0.001 [#]	3305	1.67	<0.001	3668	1.91	<0.001

resuscitation team				(88.0)	(1.48-1.90)		(97.7)	(1.48-2.47)	
Not a core member of resuscitation team	83.0	86.6		3540 (81.4)			4173 (95.7)		
Age (years)			-0.33 ([-0.52]- [-0.11])*	0.003		0.98 (0.97- 0.98)	<0.001	0.93 (0.93- 0.94)	<0.001
Time spent on e-Learning (hours)			-0.24 ([-0.30]- [-0.19])*	<0.001		0.93 (0.91- 0.94)	<0.001	0.90 (0.87- 0.93)	<0.001

#Independent samples t-test

£ One way ANOVA

*Linear regression to predict post course MCQ score (B value with 95% confidence intervals)

#ST3+, registrar equivalent

§ Consultant or associate specialist

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

The first attempt pass rate for CAS-Test was 84.6%. Univariate analysis found that paramedic and resuscitation officer pass rates were similar to physicians whilst nurses, medical students and those in the 'other' category had lower pass rates. Those participants with previous ALS experience were 1.97 times more likely to pass the CAS-Test assessment on the first attempt (OR 1.97 (95% CI 1.73-2.24), P<0.001) compared to those with no previous ALS experience. Those who were core members of the resuscitation team were 1.67 times more likely to pass the CAS-Test scenario, compared with those who were not core members (95% CI 1.48-1.90), P<0.001). Middle grade doctors were 1.75 times more likely to pass the CAS-Test compared to Foundation Year 2 doctors. (95% CI 1.40-2.17, P<0.001).

The overall course pass rate was 96.6%. Resuscitation officers demonstrated the highest pass rate at 100%. Junior nurses had the lowest pass rate of 88.4%. When compared to doctors in the univariate analysis; nurses (OR 0.22, 95% CI 0.17-0.29, P<0.001), ODPs (OR 0.30, 95% CI 0.12-0.76, P=0.011) and participants from the 'other' category (OR 0.12, 95% CI 0.06-0.24, P<0.001) had significantly lower overall pass rates. Participants were more likely to pass if they had previously undertaken ALS training (OR 2.27, 95% CI 1.73-2.98, P<0.001), ILS training (OR 1.64, 95% CI 1.29-2.09, P<0.001) or were a core member of the resuscitation team (OR 1.91, 95% CI 1.48-2.47, P<0.001).

The significant independent variables from the univariate analyses were assessed for co-linearity. Grade of training was removed due to co-linearity with healthcare background. The remaining independent variables were entered into multivariate analyses. Figures 1-3 present the findings from the multivariate analyses, with full data in supplementary material. Previous ILS and ALS experience and being a core member of a resuscitation team were

199 independent predictors of CAS-Test performance, post course MCQ score and overall success rates. Increasing age
 200 was associated with worse post course MCQ score, CAS-Test outcome and overall result.

201
 202 **Time spent accessing e-learning**

203 Median time spent on the e-learning was 5.2 hours (IQR 3.7-7.1). Resuscitation officers spent the longest time
 204 (median 7.5 hours, IQR 5.7-9.2). Doctors spent the least amount of time (median 4.9 hours, IQR 3.4-6.7). In general,
 205 those doctors with more clinical experience spent less time accessing the e-learning material. This is demonstrated
 206 below in table 3 where middle grade doctors spend the least time on every module. In the univariate analysis,
 207 increased hours spent accessing e-learning was a statistically significant predictor of failing the post-course MCQ (B=
 208 0.24, 95% CI [-0.30]-[-0.19], P<0.001), the CAS-Test assessment (OR 0.93, 95% CI 0.91-0.94, P<0.001) and the overall
 209 course (OR 0.90, 95% CI 0.87-0.93, P<0.001). When all other co-variables were controlled for in the multivariate
 210 regression, time spent accessing e-learning remained a significant predictor of CAS-Test failure (OR 0.96, 95% CI
 211 0.95-0.98, P<0.001) but was not a significant predictor of overall course failure (OR 0.98, 95% CI 0.95-1.02, P=0.367).

Table 3: Duration spent on individual ALS modules stratified by grade, profession and specialty background (minutes)

	ALS in perspective	ALS algorithm	Causes and Prevention of Cardiac Arrest	Acute Coronary Syndromes	Post Resuscitation Care	Monitoring, Rhythm Recognition and 12-lead ECG	Tachycardia, Cardioversion and Drugs	Bradycardia, Pacing and Drugs	Special Circumstances	Decisions Relating to Resuscitation	Arterial Blood Gas Analysis
Grade/healthcare profession											
Foundation year doctor	9.2	44.0	17.0	27.1	22.5	34.3	32.3	15.7	25.1	8.0	14.5
Junior grade doctor (ST1/ST2)	9.8	45.3	17.7	26.6	22.7	32.5	30.4	14.6	24.6	8.9	15.3
Middle grade doctor	9.5	43.8	17.0	26.4	21.8	30.7	27.8	13.6	22.8	8.0	12.4
Senior grade doctor	10.1	48.0	17.8	25.8	21.4	33.5	31.6	14.2	26.1	9.0	15.4
Junior nurse	11.0	51.0	21.4	31.1	24.9	53.5	39.6	19.9	32.7	10.3	25.1
Senior nurse	10.6	50.1	19.7	29.9	24.8	46.9	38.2	17.6	31.0	9.7	22.4
Paramedic	10.5	42.9	19.4	29.7	25.2	42.4	36.4	17.6	28.9	10.2	19.8
Operating department practitioner	10.6	49.5	22.6	29.5	24.8	57.8	43.8	20.3	33.0	12.1	28.6
Resuscitation officer	13.3	41.7	20.0	40.0	25.9	83.8	42.2	25.6	41.4	11.4	29.9
Medical student	9.3	45.0	17.8	28.1	24.1	38.5	35.8	16.5	28.7	9.3	15.6
Specialty background											
Anaesthetics	9.7	45.5	17.9	27.5	23.0	36.2	32.9	16.0	26.1	8.6	16.0
Cardiology	10.0	44.6	17.9	25.7	21.7	33.1	33.9	15.4	31.8	9.0	19.1
Surgery	9.3	45.0	17.9	28.0	23.0	35.9	33.7	15.5	25.5	8.1	15.5
Medicine	9.3	44.2	17.2	26.5	22.4	33.0	30.9	14.8	25.3	8.1	14.3

Emergency	10.0	45.2	18.2	27.6	23.4	38.3	32.6	16.4	25.6	9.1	18.3
Critical Care	11.1	52.1	20.8	30.7	23.8	46.1	38.2	18.9	32.0	9.8	18.5

212

213

214

215

216

217

DISCUSSION

218

219

220

221

222

223

This study has shown that previous experience in life support courses and being a core member of the resuscitation team predicts a favourable outcome on an e-ALS course. It also identifies the extent to which different candidate groups access the e-learning material and highlights particular modules that may be more challenging. Time spent accessing e-learning material was not related to course outcome; this was thought to be because participants who utilise these skills on a daily basis are already familiar with the material and thus require less time to re-familiarise themselves.

224

225

226

227

228

229

There are increasing pressures to minimise time spent on courses for both participants and faculty and to improve outcomes. It has been postulated that pre-course preparation could lead to either better outcomes or a reduced amount of face-to-face time needed on the course. This could in theory lead to equivalent or better participant outcomes with less resources (time off work for faculty/participants, venue hire etc.). There is very little evidence relating specifically to pre-learning for advanced life support courses, so this study goes some way towards filling that void.

230

231

232

233

234

235

Perkins et al.¹¹ looked at one example of pre-course preparation. This open label, multicentre RCT was a study of 572 participants on Resuscitation Council (UK) ALS courses. The control group received the course manual four weeks before the course. The intervention group received the course manual and also a CD with an interactive e-learning simulation programme. Although there were no significant differences in the primary outcome (performance during a standard cardiac arrest simulation), user evaluations were favourable. The results however cannot necessarily be generalised to all other types of pre-course learning or pre-course learning for other populations/course groups.

236

237

A multi-centre RCT demonstrated equivalence in outcome when comparing e-ALS and c-ALS learning methods and was significantly less costly to deliver.⁸ The findings of this were corroborated by a large observational study of

238 27,170 participants which demonstrated almost identical assessment outcomes for participants enrolled on either a
239 c-ALS or e-ALS course.⁹ These studies were a comparison of a standard life support course against specific pre-course
240 e-learning associated with a shorter duration hybrid life support course.

241 The topic of pre-course learning was addressed during the 2015 ILCOR international consensus on science process. It
242 was felt that a specific recommendation for or against pre-course preparation in ALS courses was too speculative
243 due to the lack of evidence in the literature.¹⁵ These findings were balanced with a statement highlighting the
244 considerable ambiguity in the definition of “pre-course learning” and the difficulty in comparing single interventions
245 like a pre-course CD¹¹ with an intervention followed by a hybrid version of the face-to-face element.^{8,9}

246 With regard to the findings from this study, we found some unexpected and interesting results. The most surprising
247 result was that time spent accessing prerequisite e-learning material was actually associated with worse assessment
248 and overall course outcome in the univariate regression. On further analysis however, this is explained by the fact
249 that those with greater clinical experience spent less time accessing the e-learning but paradoxically performed
250 better in the course assessments. This demonstrates the educational notion that when learning can be based on
251 previous experience; it will normally lead to improved outcomes. This is demonstrated in the multivariate regression
252 where time spent on e-learning was no longer a significant predictor of overall course outcome. Increased age was
253 associated with significantly poorer assessment outcomes. Whilst there is a paucity of evidence for the literature
254 regarding the effect of age on ALS outcomes, this pattern has been found in BLS studies and has been attributed to
255 skill decline over time^{16,17} and psychological factors where younger participants are more motivated to learn.¹⁸ It has
256 been found that those working in a high risk area for area for cardiac arrest were more motivated to learn life
257 support skills.¹⁹

258

259 Participants with greater experience in managing critically unwell patients (paramedics, middle grade doctors,
260 previous ALS/ILS experience, core member of the resuscitation team) performed substantially better in the CAS-Test
261 and overall result. This should not come as a surprise, but is a useful insight for course organisers when identifying
262 participants at the start of a course who do not fall into these groups and may benefit from additional support.

263

264 The e-learning package allows participants to dictate their own level of access dependent upon their prior
265 knowledge, experience and specialty background. They can access material at an appropriate time for them and

266 dedicate a greater amount of time to their weaker knowledge areas. The need for this degree of flexibility is
267 demonstrated by the vastly different durations spent accessing the online content. This is exemplified in table 3
268 which highlights that certain candidate groups (junior nurses and operating department practitioners) spent twice as
269 long on the 'Monitoring, rhythm recognition and 12-lead ECG' module compared to middle grade doctors, perhaps
270 because they do not routinely utilise such skills on a daily basis. The flexibility that the e-ALS course creates is just
271 one reason amongst many why participant satisfaction is greater on e-learning courses than compared to traditional
272 didactic courses.^{20,21}

274 ***Limitations and Further Research***

275 The main limitation of this exploratory study is its observational nature. This means that the authors are only able to
276 suggest causality when determining whether independent variables influence assessment outcome. A specifically
277 designed RCT would be needed to establish a cause-effect relationship on assessment outcome.

278
279 Time is not necessarily an accurate marker of whether participants have truly engaged with the material and as this
280 study has shown, it is significantly confounded by clinical experience (ie if participants are already well versed in ECG
281 interpretation they will spend less time on this module). Furthermore, different individuals possess a spectrum of
282 learning abilities with some participants learning faster than others. A proportion of participants may have chosen to
283 preferentially utilise the course manual as opposed to the e-learning package and others may leave the e-learning
284 running whilst not at the computer, providing a falsely elevated time spent accessing the material. There remains a
285 need for more specific markers for determining whether participants have truly engaged with the e-learning
286 material.

287
288 A final limitation is that it does not determine whether accessing e-learning actually affects patient outcome from
289 cardiac arrest. Whilst this should be the overriding aim behind all resuscitation-related research, such studies are
290 very difficult to achieve. The authors believe however, that by critically appraising course outcome data and
291 continuously improving the delivery methods of resuscitation courses this will ultimately improve the care of the
292 critically unwell patient.

294 ***Conclusion***

295 Clinical experience through core membership of cardiac arrest teams and previous ILS or ALS training were
296 independent predictors of performance on the e-ALS course whilst time spent accessing e-learning materials did not
297 affect course outcomes. The large variation in time spent accessing e-learning reflects the diverse nature of
298 participants on our e-ALS courses and the spectra of learning needs that they possess. This supports the blended
299 approach to e-ALS which allows participants to tailor their e-learning experience to their specific needs.

301 **CONFLICTS OF INTEREST**

302 CJT is a Trainee Representative for the ALS Subcommittee for the Resuscitation Council (UK). ASL is Honorary
303 Secretary of the Resuscitation Council (UK) and a member of the European Resuscitation Council ALS Course
304 Committee. IB is an Educator for the Resuscitation Council (UK). SH is Director of Course Development and Training
305 for the Resuscitation Council (UK). SB-A is Project and Development Manager for the Resuscitation Council (UK). GDP
306 is Chair of the ALS Subcommittee for the Resuscitation Council (UK) and member of the European Resuscitation
307 Council ALS Course Committee.

309 **FUNDING**

310 GDP is supported by the National Institute for Health Research (Senior Investigator) and Intensive Care Foundation
311 (Director of Research).

313 **ACKNOWLEDGEMENTS**

314 The authors would like to acknowledge the ALS instructors and candidates who have participated in an e-ALS course
315 over the duration of this study. They also acknowledge the ongoing work of the ALS subcommittee of the
316 Resuscitation Council (UK) to oversee the hundreds of ALS courses that take place each year.

324

325

326

327

REFERENCES

328

1. Søreide E, Morrison L, Hillman K, et al. The formula for survival in resuscitation. *Resuscitation* 2013;84:1487–93.

329

330

2. Nolan J, Soar J, Eikeland H. The chain of survival. *Resuscitation* 2006;71:270–1.

331

332

333

3. Spearpoint KG, Gruber PC, Brett SJ. Impact of the Immediate Life Support course on the incidence and outcome of in-hospital cardiac arrest calls: an observational study over 6 years. *Resuscitation* 2009;80:638–43.

334

335

4. Moretti MA, Cesar LAM, Nusbacher A, et al. Advanced cardiac life support training improves long-term survival from in-hospital cardiac arrest. *Resuscitation* 2007;72:458–65.

336

337

5. Yule SJ, Walls RM. Advanced life support training: does online learning translate to real-world performance? *Ann Intern Med* 2012;157:69–70.

338

6. Nolan J. Advanced life support training. *Resuscitation* 2001;50:9–11.

339

7. Perkins G, Lockey A. The advanced life support provider course. *BMJ* 2002;325:S81.

340

341

8. Perkins GD, Kimani PK, Bullock I, et al. Improving the efficiency of advanced life support training: a randomized, controlled trial. *Ann Intern Med* 2012;157:19–28.

342

343

9. Thorne CJ, Lockey AS, Bullock I, Hampshire S, Begum-ali S. e-Learning in advanced life support – An evaluation by the Resuscitation Council (UK) & *Resuscitation* 2015;90:79–84.

344

345

10. Jensen ML, Mondrup F, Lippert F, Ringsted C. Using e-learning for maintenance of ALS competence. *Resuscitation* 2009;80:903–8.

346

347

11. Perkins GD, Fullerton JN, Davis-gomez N, et al. The effect of pre-course e-learning prior to advanced life support training: a randomised controlled trial. *Resuscitation* 2010;81:877–81.

348

349

12. Napier F, Davies RP, Baldock C, et al. Validation for a scoring system of the ALS cardiac arrest simulation test (CASTest). *Resuscitation* 2009;80:1034–8.

350

351

13. Perkins GD, Davies RP, Stallard N, et al. Advanced life support cardiac arrest scenario test evaluation. *Resuscitation* 2007;75:484–90.

352

353

14. R Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>. 2016.

354

355

356

15. Finn JC, Bhanji F, Lockey A, et al. Part 8: Education, implementation, and teams 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2015;95:e203–24.

357

358

16. Papalexopoulou K, Chalkias A, Dontas I, et al. Education and age affect skill acquisition and retention in lay rescuers after a European Resuscitation Council CPR/AED course. *Hear Lung J Acute Crit Care* 2014;43:66–71.

359

360

17. Smith KK, Gilcreast D, Pierce K. Evaluation of staff’s retention of ACLS and BLS skills. *Resuscitation* 2008;78:59–65.

361

362

18. Eisenburger P, Safar P. Life supporting first aid training of the public--review and recommendations. *Resuscitation* 1999;41:3–18.

363

364

19. Hopstock LA. Motivation and adult learning: a survey among hospital personnel attending a CPR course. *Resuscitation* 2008;76:425–30.

365

20. Cook DA, McDonald FS. E-learning: is there anything special about the “E”? *Perspect Biol Med* 2008;51:5–21.

366

367

368

21. Soar J, Mancini ME, Bhanji F, et al. Part 12: Education, implementation, and teams: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2010;81 Suppl 1:e288–330.

369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410

LEGENDS TO FIGURES

Table 1: Participant demographics on the e-ALS course and time spent on e-learning

Table 2: Univariate predictors of assessment outcomes

Table 3: Duration spent on individual ALS modules stratified by grade, profession and specialty background

Figure 1: Multivariate analysis demonstrating factors that influence CAS-Test outcome

Figure 2: Multivariate analysis demonstrating factors that influence post-course MCQ score

Figure 3: Multivariate analysis demonstrating factors that influence overall course outcome

Supplementary material 1: Multivariate predictors of assessment outcomes

411
412
413
414
415
416

Supplementary material 1: Multivariate predictors of assessment outcomes												
Independent variables	Mean post e-learning MCQ score	Mean post-course MCQ score	Mean difference (95% CI)	P-value	CAS-Test result		Odds ratio of CAS-Test Pass (95% CI)	P-value	Overall course result		Odds ratio of course Pass (95% CI)	P-value
					Pass (%)	Fail (%)			Pass (%)	Fail (%)		

Healthcare profession

Doctor (comparison)	84.7	88.7			5352 (86.0)	871 (14.0)			6095 (97.8)	137 (2.2)		
Nurse	79.7	80.0	-4.35 ([-4.85]-[-3.85])	<0.001	1005 (81.3)	231 (18.7)	0.92 (0.76-1.10)	0.356	1122 (90.9)	113 (9.1)	0.27 (0.20-0.37)	<0.001
Medical student	83.4	86.5	-0.43 ([-1.31]-[-0.45])	0.334	425 (79.6)	109 (20.4)	0.87 (0.63-1.20)	0.390	525 (98.3)	9 (1.7)	2.16 (0.96-4.48)	0.063
Operating Department Practitioner	73.0	79.2	-9.41 ([-11.13]-[-7.69])	<0.001	51 (70.8)	21 (29.2)	0.44 (0.25-0.78)	0.005	67 (93.1)	5 (6.9)	0.36 (0.13-1.01)	0.052
Ambulance staff/ Paramedic	81.4	85.4	-2.42 ([-4.71]-[0.12])	0.039	37 (92.5)	3 (7.5)	3.75 (1.10-12.85)	0.035	39 (97.5)	1 (2.5)	2.34 (0.27-20.54)	0.444
Resuscitation Officer	86.6	90.5	0.98 ([-3.18]-[5.14])	0.644	13 (86.7)	2 (13.3)	0.79 (0.17-3.73)	0.769	15 (100.0)	0 (0)	78518 (0-infinity)	0.986
Other	79.9	83.6	-4.27 ([-6.00]-[-2.53])	<0.001	46 (66.7)	23 (32.4)	0.47 (0.27-0.81)	0.007	59 (84.3)	11 (15.7)	0.19 (0.09 - 0.42)	<0.001

Previous life support experience

Previous ALS experience	85.5	89.7	3.83 (3.44 - 4.21)	<0.001	3204 (89.3)	383 (10.7)	2.61 (2.22-3.07)	<0.001	3515 (98.0)	72 (2.0)	5.13 (3.66-7.19)	<0.001
No previous ALS experience	82.3	86.1			3727 (81.0)	877 (19.0)			4411 (95.6)	205 (4.4)		
Previous ILS experience	83.2	87.4	-0.27 ([-0.66]-[-0.12])	0.172	4666 (85.6)	787 (14.4)	1.19 (1.02-1.39)	0.024	5302 (97.2)	153 (2.8)	2.18 (1.61-2.95)	<0.001
No previous ILS experience	84.5	88.3			2265 (82.7)	473 (17.3)			2624 (95.5)	124 (4.5)		
Core member of resuscitation team	84.4	88.8	1.28 (0.94-1.62)	<0.001	3305 (88.0)	451 (12.0)	1.39 (1.21-1.59)	<0.001	3668 (97.7)	87 (2.3)	1.47 (1.10-1.98)	0.009
Not a core member of resuscitation	83.0	86.6			3540 (81.4)	809 (18.6)			4173 (95.7)	189 (4.3)		

team									
Age (years)		-0.06 ([-0.09]-[-0.04])	<0.001		0.96 (0.95-0.97)	<0.001		0.93 (0.92-0.94)	<0.001
Time spent on e-Learning (hours)		-0.05 ([-0.11]-0.00)	0.047		0.96 (0.95-0.98)	<0.001		0.98 (0.95-1.02)	0.367

ST3+, middle grade equivalent

§ Consultant or associate specialist