

Original citation:

Petrou, Stavros, Rivero-Arias, Oliver, Dakin, Helen, Longworth, Louise, Oppe, Mark, Froud, Robert J. and Gray, Alastair, 1953-. (2015) Preferred reporting items for studies mapping onto preference-based outcome measures : the MAPS statement. Medical Decision Making, 35 (6). NP1-NP8. ISSN 0272-989X

Permanent WRAP url:

<http://wrap.warwick.ac.uk/71037>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

This article is distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>) which permits any use, reproduction and distribution of the work without further permission provided the original work is attributed.

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: publications@warwick.ac.uk

warwick**publications**wrap

highlight your research

<http://wrap.warwick.ac.uk/>

Preferred Reporting Items for Studies Mapping onto Preference-Based Outcome Measures: The MAPS Statement

Stavros Petrou, PhD, Oliver Rivero-Arias, DPhil, Helen Dakin, DPhil,
Louise Longworth, PhD, Mark Oppe, PhD, Robert Froud, PhD, Alastair Gray, PhD

Background. “Mapping” onto generic preference-based outcome measures is increasingly being used as a means of generating health utilities for use within health economic evaluations. Despite the publication of technical guides for the conduct of mapping research, guidance for the reporting of mapping studies is currently lacking. The MAPS (MApping onto Preference-based measures reporting Standards) statement is a new checklist that aims to promote complete and transparent reporting of mapping studies. **Methods.** In the absence of previously published reporting checklists or reporting guidance documents, a *de novo* list of reporting items was created by a working group comprised of 6 health economists and 1 Delphi methodologist. A 2-round, modified Delphi survey with representatives from academia, consultancy, health technology assessment agencies, and the biomedical journal editorial community was used to identify a list of essential reporting

items from this larger list. **Results.** From the initial *de novo* list of 29 candidate items, a set of 23 essential reporting items was developed. The items are presented numerically and categorized within 6 sections, namely: (i) title and abstract; (ii) introduction; (iii) methods; (iv) results; (v) discussion; and (vi) other. The MAPS statement is best applied in conjunction with the accompanying MAPS explanation and elaboration document. **Conclusions.** It is anticipated that the MAPS statement will improve the clarity, transparency, and completeness of reporting of mapping studies. To facilitate dissemination and uptake, the MAPS statement is being co-published by 7 health economics and quality-of-life journals, and broader endorsement is encouraged. The MAPS working group plans to assess the need for an update of the reporting checklist in 5 years. **Key words:** mapping; outcome measures; preferences; reporting. (*Med Decis Making* 2015;35:NP1–NP8)

INTRODUCTION

The process of “mapping” onto generic preference-based outcome measures is increasingly being used as a means of generating health utilities for application within health economic evaluations.¹ Mapping involves the development and use of an algorithm (or algorithms) to predict the primary outputs of generic preference-based outcome measures (i.e., health utility values), using data on other indicators or measures of health. The source predictive measure may be a non-preference-based indicator or measure of health outcome or, more exceptionally, a preference-based outcome measure that is not preferred by the local health technology assessment agency. The

algorithm(s) can subsequently be applied to data from clinical trials, observational studies, or economic models containing the source predictive measure(s) to predict health utility values in contexts in which the target generic preference-based measure is absent. The predicted health utility values can then be analyzed using standard methods for individual-level data (e.g., within a trial-based economic evaluation) or summarized for each health state within a decision-analytic model.

Throughout recent years, there has been a rapid increase in the publication of studies that use mapping techniques to predict health utility values, and databases of published studies in this field are beginning to emerge.² Some authors³ and agencies⁴ concerned with technology appraisals have issued technical guides for the conduct of mapping research. However, guidance for the *reporting* of mapping studies is currently lacking. In keeping with health-related research more broadly,⁵ mapping studies should be reported fully and transparently to allow

© The Author(s) 2015
Reprints and permission:
<http://creativecommons.org/licenses/by/4.0/>
DOI: 10.1177/0272989X15598332

readers to assess the relative merits of the investigation.⁶ Moreover, there may be significant opportunity costs associated with regulatory and reimbursement decisions for new technologies informed by misleading findings from mapping studies. This has led to the development of the MAPS (Mapping onto Preference-based measures reporting Standards) reporting statement, which we summarize in this article.

The aim of the MAPS reporting statement is to provide recommendations, in the form of a checklist of essential items, which authors should consider when reporting a mapping study. It is anticipated that the checklist will promote complete and transparent reporting by researchers. The focus, therefore, is on promoting the quality of reporting of mapping studies, rather than the quality of their conduct, although it is possible that the reporting statement will also indirectly enhance the methodological rigor of the research.⁷ The MAPS reporting statement is primarily targeted at researchers developing mapping algorithms, the funders of the research, and peer reviewers and editors involved in the manuscript review process for mapping studies.^{5,6} In developing the reporting statement, the term *mapping* is used to cover all approaches that predict the outputs of generic preference-based outcome measures using data on other indicators or measures of health, and it encompasses related forms of nomenclature used by some researchers, such as “cross-walking” or “transfer to utility.”^{1,8} Similarly, the term *algorithm* is used in its broadest sense to

encompass statistical associations and more complex series of operations.

THE DEVELOPMENT OF THE MAPS STATEMENT

The development of the MAPS reporting statement was informed by recently published guidance for health research reporting guidelines⁵ and broadly modeled other recent reporting guideline developments.^{9–14} A working group comprised of 6 health economists (SP, ORA, HD, LL, MO, and AG) and 1 Delphi methodologist (RF) was formed following a request from an academic journal to develop a reporting statement for mapping studies. One of the working group members (HD) had previously conducted a systematic review of studies mapping from clinical or health-related quality-of-life measures onto the EQ-5D.² Using the search terms from this systematic review, as well as other relevant articles and reports already in our possession, a broad search for reporting guidelines for mapping studies was conducted. This confirmed that no previous reporting guidance had been published. The working group members therefore developed a preliminary *de novo* list of 29 reporting items and accompanying explanations. Following further review by the working group members, this was subsequently distilled into a list of 25 reporting items and accompanying explanations.

Members of the working group identified 62 possible candidates for a Delphi panel from a pool of active researchers and stakeholders in this field. The candidates included individuals from academic and consultancy settings with considerable experience in mapping research, representatives from health technology assessment agencies that routinely appraise evidence informed by mapping studies, and biomedical journal editors. Health economists from the MAPS working group were included in the Delphi panel. A total of 48 of the 62 (77.4%) individuals agreed to participate in a Delphi survey aimed at developing a minimum set of standard reporting requirements for mapping studies with an accompanying reporting checklist.

The Delphi panelists were sent a personalized link to a Web-based survey, which had been piloted by members of the working group. Nonresponders were sent up to 2 reminders after 14 and 21 days. The panelists were anonymous to each other throughout the study, and their identities were known only to 1 member of the working group. The panelists were invited to rate the importance of each of the 25

Received 7 July 2015 from the Warwick Clinical Trials Unit, Warwick Medical School, University of Warwick, Coventry, UK (SP, RF); National Perinatal Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK (ORA); Health Economics Research Centre, Nuffield Department of Population Health, University of Oxford, Oxford, UK (HD, AG); Health Economics Research Group, Brunel University London, Uxbridge, UK (LL); EuroQol Research Foundation, Rotterdam, The Netherlands (MO); and Norges Helsehøyskole, Campus Kristiania, Oslo, Norway (RF). Funding: No external funding was received for this study. All members of the working group acted on a voluntary basis. Conflicts of interest: RF is a shareholder and a director of a company (Clinvivo Limited) that provides services for Delphi studies. However, these services were not used in this study, and thus notwithstanding the connection RF declares that he has no conflicts of interest. The remaining authors (SP, ORA, HD, LL, MO, AG) have no conflicts of interest to declare. All authors contributed to the study design and conduct, interpretation of results, and report writing. SP acts as the guarantor.

Address correspondence to Professor Stavros Petrou, Warwick Medical School, University of Warwick, Coventry CV4 7AL, UK; phone: 02476 151124; fax: 02476 151586; e-mail: S.Petrou@warwick.ac.uk.

candidate reporting items identified by the working group on a 9-point rating scale (from 1, *not important*, to 9, *extremely important*), describe their confidence in their ratings (*not confident*, *somewhat confident*, or *very confident*), comment on the candidate items and their explanations, suggest additional items for consideration by the panelists in subsequent rounds, and provide any other general comments. The candidate reporting items were ordered within 6 sections: (i) title and abstract, (ii) introduction, (iii) methods, (iv) results, (v) discussion, and (vi) other. The panelists also provided information about their geographical area of work, gender, and primary and additional work environments. The data were imported into Stata (version 13; Stata Corp., College Station, TX) for analysis.

A modified version of the Research and Development (RAND)/University of California, Los Angeles (UCLA) appropriateness method was used to analyze the round 1 responses.¹⁵ This involved calculating the median score, the interpercentile range (IPR; 30th and 70th), and the interpercentile range adjusted for symmetry (IPRAS) for each item (j) being rated. The IPRAS includes a correction factor for asymmetric ratings, and panel disagreement was judged to be present in cases if $IPR_j > IPRAS_j$.¹⁵ We modified the RAND/UCLA approach by asking panelists about “importance” rather than “appropriateness” per se. Assessment of importance followed the classic RAND/UCLA definitions, categorized simply as whether the median rating fell between 1 and 3 (*unimportant*), 4 and 6 (*neither unimportant nor important*), or 7 and 9 (*important*).¹⁵

The results of round 1 of the Delphi survey were reviewed at a face-to-face meeting of the working group. A total of 46 of the 48 (95.8%) individuals who agreed to participate completed round 1 of the survey. Of the 25 items, 24 were rated as important, with one item (source of funding) rated as neither unimportant nor important. There was no evidence of disagreement on ratings of any items according to the RAND/UCLA method. These findings did not change when the responses of the MAPS working group were excluded. Based on the qualitative feedback received in round 1, items describing “Modeling Approaches” and “Repeated Measurements” were merged, as were items describing “Model Diagnostics” and “Model Plausibility.” In addition, amendments to the wording of several recommendations and their explanations were made in the light of qualitative feedback from the panelists.

Panelists participating in round 1 were invited to participate in a second round of the Delphi survey.

A summary of revisions made following round 1 was provided. This included a document in which revisions to each of the recommendations and explanations were displayed in the form of track changes. Panelists participating in round 2 were provided with group outputs (mean scores and their standard deviations, median scores and their IPRs, histograms, and RAND/UCLA labels of importance and agreement level) summarizing the round 1 results (and disaggregated outputs for the merged items). They were also able to view their own round 1 scores for each item (and disaggregated scores for the merged items). Panelists participating in round 2 were offered the opportunity to revise their rating of the importance of each of the items and informed that their rating from round 1 would otherwise hold. For the merged items, new ratings were solicited. Panelists participating in round 2 were also offered the opportunity to provide any further comments on each item or any further information that might be helpful to the group. Nonresponders to the second round of the Delphi survey were sent up to 2 reminders after 14 and 21 days. The analytical methods for the round 2 data mirrored those for the first round.

The results of the second round of the Delphi survey were reviewed at a face-to-face meeting of the working group. A total of 39 of the 46 (84.8%) panelists participating in round 1 completed round 2 of the survey. All 23 items included in the second round were rated as important with no evidence of disagreement on ratings of any items according to the RAND/UCLA method. Qualitative feedback from the panelists participating in round 2 led to minor modifications to the wording of a small number of recommendations and their explanations. This was fed back to the round 2 respondents, who were given a final opportunity to comment on the readability of the final set of recommendations and explanations. Based on these methods, a final consensus list of 23 reporting items was developed.

THE MAPS STATEMENT

The MAPS statement is a 23-item checklist of recommendations (Table 1) that we consider essential for complete and transparent reporting of studies that map onto generic preference-based outcome measures. The 23 reporting items are presented numerically and categorized within 6 sections: (i) title and abstract (2 items), (ii) introduction (2 items), (iii) methods (9 items), (iv) results (6 items), (v)

Table 1 Checklist of Items to Include When Reporting a Mapping Study

Section or Topic	Item Number	Recommendation	Reported on Page Number or Line Number
Title and Abstract			
Title	1	Identify the report as a study mapping between outcome measures. State the source measure(s) and generic, preference-based target measure(s) used in the study.	_____
Abstract	2	Provide a structured abstract including, as applicable: objectives; methods, including data sources and their key characteristics, outcome measures used, and estimation and validation strategies; results, including indicators of model performance; conclusions; and implications of key findings.	_____
Introduction			
Study rationale	3	Describe the rationale for the mapping study in the context of the broader evidence base.	_____
Study objective	4	Specify the research question with reference to the source and target measures used and the disease or population context of the study.	_____
Methods			
Estimation sample	5	Describe how the estimation sample was identified, why it was selected, the methods of recruitment and data collection, and its location(s) or setting(s).	_____
External validation sample	6	If an external validation sample was used, the rationale for selection, the methods of recruitment and data collection, and its location(s) or setting(s) should be described.	_____
Source and target measures	7	Describe the source and target measures and the methods by which they were applied in the mapping study.	_____
Exploratory data analysis	8	Describe the methods used to assess the degree of conceptual overlap between the source and target measures.	_____
Missing data	9	State how much data were missing and how missing data were handled in the sample(s) used for the analyses.	_____
Modeling approaches	10	Describe and justify the statistical model(s) used to develop the mapping algorithm.	_____
Estimation of predicted scores or utilities	11	Describe how predicted scores or utilities are estimated for each model specification.	_____
Validation methods	12	Describe and justify the methods used to validate the mapping algorithm.	_____
Measures of model performance	13	State and justify the measure(s) of model performance that determine the choice of the preferred model(s), and describe how these measures were estimated and applied.	_____
Results			
Final sample size(s)	14	State the size of the estimation sample and any validation sample(s) used in the analyses (including both number of individuals and number of observations).	_____

(continued)

Table 1 (continued)

Section or Topic	Item Number	Recommendation	Reported on Page Number or Line Number
Descriptive information	15	Describe the characteristics of individuals in the sample(s) (or refer back to previous publications giving such information). Provide summary scores for source and target measures, and summarize the results of analyses used to assess overlap between the source and target measures. State which model(s) is (are) preferred, and justify why this (these) model(s) was (were) chosen.	_____
	16	Provide all model coefficients and standard errors for the selected model(s). Provide clear guidance on how a user can calculate utility scores based on the outputs of the selected model(s).	_____
	17	Report information that enables users to estimate standard errors around mean utility predictions and individual-level variability.	_____
	18	Present results of model performance, such as measures of prediction accuracy, and fit statistics for the selected model(s) in a table or in the text. Provide an assessment of face validity of the selected model(s).	_____
	19		_____
Discussion			
Comparisons with previous studies	20	Report details of previously published studies developing mapping algorithms between the same source and target measures, and describe differences between the algorithms, in terms of model performance, predictions, and coefficients, if applicable.	_____
	21	Outline the potential limitations of the mapping algorithm.	_____
Study limitations	22	Outline the clinical and research settings in which the mapping algorithm could be used.	_____

Other			
Additional information	23	Describe the source(s) of funding and nonmonetary support for the study, and the role of the funder(s) in its design, conduct, and report. Report any conflicts of interest surrounding the roles of authors and funders.	_____

discussion (3 items), and (vi) other (1 item). The reporting of each item does not necessarily have to follow the order within the MAPS statement. Rather, what is important is that each recommendation is addressed in either the main body of the report or its appendices. Several biomedical journals have endorsed the MAPS statement, including *Applied Health Economics and Health Policy*, *Health and Quality of Life Outcomes*, *International Journal of Technology Assessment in Health Care*, *Journal of Medical Economics*, *Medical Decision Making*, *Pharmacoeconomics*, and *Quality of Life Research*. We encourage other journals and research interest groups to endorse the MAPS statement and authors to adhere to its principles.

THE MAPS EXPLANATION AND ELABORATION PAPER

In addition to the MAPS reporting statement, we have produced a supporting Explanation and Elaboration paper¹⁶ modeled on those developed for other reporting guidelines.^{9–14} The reporting items contained within the MAPS statement are best understood by referring to the information contained within this accompanying document. The Explanation and Elaboration paper provides exemplars of good reporting practice identified from the published literature for each reporting item. In addition, it provides a detailed explanation to accompany each recommendation, supported by a rationale and relevant evidence when available. The development of the Explanation and Elaboration paper was completed following several iterations produced by members of the working group, after which the examples and explanations were shared with the Delphi panelists for final revisions to improve readability and their approval. The Explanation and Elaboration paper also summarizes the characteristics of the Delphi panelists and provides detailed statistics for item ratings at each Delphi round.

DISCUSSION

Throughout recent years, there has been a rapid increase in the publication of studies that use mapping techniques to predict health utility values. One recent review article identified 90 studies published up to the year 2013 reporting 121 mapping algorithms between clinical or health-related quality-of-life measures and the EQ-5D.² That review article excluded mapping algorithms targeted at other generic

preference-based outcome measures that can generate health utilities, such as the SF-6D¹⁷ and the Health Utilities Index (HUI),¹⁸ which have been the target of numerous other mapping algorithms.^{1,19–24} Moreover, the popularity of the mapping approach for estimating health utilities is unlikely to wane given the numerous contexts within health economic evaluation in which primary data collection is challenging. However, mapping introduces additional uncertainty, and collection of primary data with the preferred utility instrument is preferable.

The MAPS reporting statement was developed to provide recommendations, in the form of a checklist of essential items, which authors should consider when reporting mapping studies. Guidance for the reporting of mapping studies was not previously available in the literature. The overall aim of MAPS is to promote clarity, transparency, and completeness of reporting of mapping studies. It is not intended to act as a methodological guide, nor as a tool for assessing the quality of study methodology. Rather, it aims to avoid misleading conclusions being drawn by readers, and ultimately policy makers, as a result of suboptimal reporting. In keeping with other recent health research reporting guidelines, we have also produced an accompanying Explanation and Elaboration paper¹⁶ to facilitate a deeper understanding of the 23 items contained within the MAPS reporting statement. That paper should hopefully act as a pedagogical framework for researchers reporting mapping studies.

The development of the MAPS reporting statement, and its Explanation and Elaboration document, was framed by recently published guidance for health research reporting guidelines.⁵ The Delphi panel was composed of a multidisciplinary, multinational team of content experts and journal editors. The panel members included people experienced in conducting mapping studies; of the 84 researchers who were first authors on papers included in a recent review of EQ-5D mapping studies,² 31 (36.9%) were included as panelists. We have no evidence to believe that a larger panel would have altered the final set of recommendations. The Delphi methodologies that we applied included analytical approaches only recently adopted by developers of health-reporting guidelines.¹⁵ We are unable to assess whether a strict adherence to the MAPS checklist will increase the word counts of mapping reports. It is our view that the increasing use of online appendices by journals should permit comprehensive reporting even in the context of strict word limits for the main body of reports.

Evidence for other health research reporting guidelines suggests that reporting quality improved

after the introduction of reporting checklists,^{25–27} although there is currently no empirical evidence that adoption of MAPS will improve the quality of reporting of mapping research. Future research planned by the MAPS working group will include a before-and-after evaluation of the benefits (and indeed possible adverse effects) of the introduction of the MAPS reporting statement. It will also be necessary to update the MAPS reporting statement in the future to address conceptual, methodological, and practical advances in the field. Potential methodological advances that might be reflected in an update might include shifts toward more complex model specifications, better methods for dealing with uncertainty, and guidance on the appropriate use of measures of prediction accuracy, such as the mean absolute error (MAE) and mean square error (MSE). The MAPS working group plans to assess the need for an update of the reporting checklist in 5 years.

In conclusion, this article summarizes a new reporting statement developed for studies that map onto generic preference-based outcome measures. We encourage health economic and quality-of-life journals to endorse MAPS, promote its use in peer review, and update their editorial requirements and “Instructions to Authors” accordingly.

ACKNOWLEDGMENTS

We are grateful to the following individuals, who consented to being acknowledged publicly, for their comments and participation in the Delphi panel: Roisin Adams, Roberta Ara, Nick Bansback, Garry Barton, Andy Briggs, Chris Carswell, Gang Chen, Doug Coyle, Ralph Crott, Richard Edlin, Alastair Fischer, Juan Manuel Ramos-Goñi, Ola Ghatnekar, Patrick Gillard, Ning Yan Gu, Annie Hawton, Haomiao Jia, Kamran Khan, Billingsley Kaambwa, Nick Kontodimopoulos, Quang Le, Jason Madan, Katherine MacGilchrist, David Meads, Duncan Mortimer, Bhash Naidoo, Angus Ng, Nalin Payakachat, Simon Pickard, Rafael Pinedo-Villanueva, Eleanor Pullenayegum, Jeff Richardson, Janet Robertson, Jeff Round, Donna Rowen, Sean Rundell, Paulos Teckle, Matthijs Versteegh, Feng Xie, and Tracey Young. This article is a joint publication by Applied Health Economics and Health Policy, Health and Quality of Life Outcomes, International Journal of Technology Assessment in Health Care, Journal of Medical Economics, Medical Decision Making, Pharmacoeconomics, and Quality of Life Research.

REFERENCES

1. Brazier JE, Yang Y, Tsuchiya A, Rowen DL. A review of studies mapping (or cross walking) non-preference based measures of health to generic preference-based measures. *Eur J Health Econ.* 2010;11:215–25.
2. Dakin H. Review of studies mapping from quality of life or clinical measures to EQ-5D: an online database. *Health Qual Life Outcomes.* 2013;11:151.
3. Chuang LH, Whitehead SJ. Mapping for economic evaluation. *Br Med Bull.* 2012;101:1–15.
4. Longworth L, Rowen D. NICE DSU Technical Support Document 10: The Use of Mapping Methods to Estimate Health State Utility Values. Sheffield, UK: Decision Support Unit, ScHARR, University of Sheffield; 2011.
5. Moher D, Schulz KF, Simera I, Altman DG. Guidance for developers of health research reporting guidelines. *PLoS Med.* 2010;7:e1000217.
6. Moher D, Simera I, Schulz KF, Hoey J, Altman DG. Helping editors, peer reviewers and authors improve the clarity, completeness and transparency of reporting health research. *BMC Medicine.* 2008;6:13.
7. Narahari SR, Ryan TJ, Aggithaya MG, Bose KS, Prasanna KS. Evidence-based approaches for the ayurvedic traditional herbal formulations: toward an ayurvedic consort model. *J Altern Complement Med.* 2008;14:769–76.
8. Mortimer D, Segal L, Sturm J. Can we derive an “exchange rate” between descriptive and preference-based outcome measures for stroke? Results from the transfer to utility (TTU) technique. *Health Qual Life Outcomes.* 2009;7:33.
9. Bossuyt PM, Reitsma JB, Bruns DE, et al. Standards for reporting of diagnostic A: the Stard statement for reporting studies of diagnostic accuracy: explanation and elaboration. *Ann Intern Med.* 2003;138:W1–12.
10. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (coreq): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care.* 2007;19:349–57.
11. Vandenbroucke JP, von Elm E, Altman DG, et al. Strengthening the reporting of observational studies in epidemiology (STROBE): explanation and elaboration. *PLoS Med.* 2007;4:e297.
12. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol.* 2009;62:e1–34.
13. Moher D, Hopewell S, Schulz KF, et al. Consort 2010 explanation and elaboration: updated guidelines for reporting parallel group randomised trials. *BMJ.* 2010;340:c869.
14. Husereau D, Drummond M, Petrou S, et al. Consolidated health economic evaluation reporting standards (CHEERS)—explanation and elaboration: a report of the ISPOR health economic evaluation publication guidelines good reporting practices task force. *Value Health.* 2013;16:231–50.
15. Fitch K, Bernstein S, Aguilar M, et al. The RAND/UCLA Appropriateness Method User’s Manual. Santa Monica, CA: RAND; 2001.
16. Petrou S, Rivero-Arias O, Dakin H, et al. The maps reporting statement for studies mapping onto generic preference-based outcome measures: explanation and elaboration. *Pharmacoeconomics.* DOI: <http://dx.doi.org/10.1007/s40273-015-0312-9>
17. Brazier J, Roberts J, Deverill M. The estimation of a preference-based measure of health from the SF-36. *J Health Econ.* 2002;21:271–92.

18. Feeny D, Furlong W, Boyle M, Torrance GW. Multi-attribute health status classification systems. Health utilities index. *Pharmacoeconomics*. 1995;7:490–502.
19. Yang Y, Wong MY, Lam CL, Wong CK. Improving the mapping of condition-specific health-related quality of life onto SF-6D score. *Qual Life Res*. 2014;23:2343–53.
20. Lee L, Kaneva P, Latimer E, Feldman LS. Mapping the gastrointestinal quality of life index to short-form 6D utility scores. *J Surg Res*. 2014;186:135–41.
21. Roset M, Badia X, Forsythe A, Webb SM. Mapping cushingqol scores onto SF-6D utility values in patients with Cushing's syndrome. *Patient*. 2013;6:103–11.
22. Payakachat N, Tilford JM, Kuhlthau KA, et al. Predicting health utilities for children with autism spectrum disorders. *Autism Res*. 2014;7:649–63.
23. Goldfeld KS, Hamel MB, Mitchell SL. Mapping health status measures to a utility measure in a study of nursing home residents with advanced dementia. *Med Care*. 2012;50:446–51.
24. Chen G, McKie J, Khan MA, Richardson JR. Deriving health utilities from the macnew heart disease quality of life questionnaire. *Eur J Cardiovasc Nurs*. 2014. DOI: 10.1177/1474515114 536096
25. Moher D, Jones A, Lepage L, Group C. Use of the consort statement and quality of reports of randomized trials: a comparative before-and-after evaluation. *JAMA*. 2001;285:1992–5.
26. Delaney A, Bagshaw SM, Ferland A, Manns B, Laupland KB, Doig CJ. A systematic evaluation of the quality of meta-analyses in the critical care literature. *Critical care*. 2005;9:R575–82.
27. Plint AC, Moher D, Morrison A, et al. Does the consort checklist improve the quality of reports of randomised controlled trials? A systematic review. *Med J Aust*. 2006;185:263–7.