

**Original citation:**

Taylor, Andrew, Vágány, Viktória, Jackson, Alison C., Harrison, Richard J., Rainoni, Alessandro and Clarkson, John P.. (2016) Identification of pathogenicity-related genes in *Fusarium oxysporum* f. sp. *cepae*. *Molecular Plant Pathology*, 17 (7). pp. 1032-1047.

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/76986>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 4.0 International license (CC BY 4.0) and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/4.0/>

**A note on versions:**

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)

# Identification of pathogenicity-related genes in *Fusarium oxysporum* f. sp. *cepae*

ANDREW TAYLOR<sup>1,\*</sup>, VIKTÓRIA VÁGÁNY<sup>1</sup>, ALISON C. JACKSON<sup>1</sup>, RICHARD J. HARRISON<sup>2</sup>, ALESSANDRO RAINONI<sup>1,†</sup> AND JOHN P. CLARKSON<sup>1</sup>

<sup>1</sup>Warwick Crop Centre, School of Life Sciences, University of Warwick, Wellesbourne, Warwick, CV35 9EF, UK

<sup>2</sup>East Malling Research, New Road, East Malling, Kent, ME19 6BJ, UK

## SUMMARY

Pathogenic isolates of *Fusarium oxysporum*, distinguished as *formae speciales* (f. spp.) on the basis of their host specificity, cause crown rots, root rots and vascular wilts on many important crops worldwide. *Fusarium oxysporum* f. sp. *cepae* (FOC) is particularly problematic to onion growers worldwide and is increasing in prevalence in the UK. We characterized 31 *F. oxysporum* isolates collected from UK onions using pathogenicity tests, sequencing of housekeeping genes and identification of effectors. In onion seedling and bulb tests, 21 isolates were pathogenic and 10 were non-pathogenic. The molecular characterization of these isolates, and 21 additional isolates comprising other f. spp. and different *Fusarium* species, was carried out by sequencing three housekeeping genes. A concatenated tree separated the *F. oxysporum* isolates into six clades, but did not distinguish between pathogenic and non-pathogenic isolates. Ten putative effectors were identified within FOC, including seven *Secreted In Xylem* (*SIX*) genes first reported in *F. oxysporum* f. sp. *lycopersici*. Two highly homologous proteins with signal peptides and RxLR motifs (*CRX1/CRX2*) and a gene with no previously characterized domains (*C5*) were also identified. The presence/absence of nine of these genes was strongly related to pathogenicity against onion and all were shown to be expressed *in planta*. Different *SIX* gene complements were identified in other f. spp., but none were identified in three other *Fusarium* species from onion. Although the FOC *SIX* genes had a high level of homology with other f. spp., there were clear differences in sequences which were unique to FOC, whereas *CRX1* and *C5* genes appear to be largely FOC specific.

**Keywords:** effector genes, *Fusarium* basal rot, *Fusarium oxysporum* f. sp. *cepae*, onion, pathogenicity, *Secreted In Xylem* (*SIX*).

## INTRODUCTION

*Fusarium oxysporum* is a major pathogen of many important crops worldwide, causing crown and root rots as well as vascular wilts (Leslie and Summerell, 2006). The soil-borne fungus affects a very wide range of crop hosts, including onion, leek, lettuce, tomato, brassicas, asparagus, cucurbits, peppers, coriander, spinach, basil, beans, peas, strawberry, watermelon and banana, and also important non-food crops, such as carnation and narcissus (Leslie and Summerell, 2006; Michielse and Rep, 2009). Its wide host range, as well as its economic and scientific impact, means that *F. oxysporum* was recently identified as the fifth most important plant-pathogenic fungus (Dean *et al.*, 2012). *Fusarium oxysporum* is a species complex and includes both non-pathogenic and pathogenic isolates. Non-pathogenic *F. oxysporum* isolates commonly occur in the soil as saprophytes, while some have been identified as biocontrol agents and endophytes (Alabouvette *et al.*, 2009). Pathogenic *F. oxysporum* isolates are distinguished as *formae speciales* (f. spp.) on the basis of their host specificity (Leslie and Summerell, 2006), and more than 120 have been identified (Michielse and Rep, 2009). Recent advances in the understanding of the pathogenicity in *F. oxysporum* have been made following publication of the genome of *F. oxysporum* f. sp. *lycopersici* (FOL), which infects tomato (Ma *et al.*, 2010). This led to the discovery of lineage-specific mobile pathogenicity chromosomes which contain pathogenicity-related genes. These include *Secreted In Xylem* (*SIX*) genes, the products of which are small effector proteins secreted by FOL during the colonization of tomato plants (Ma *et al.*, 2010). So far, 14 *SIX* genes have been identified in FOL, which show no homology with each other or with any other sequenced gene (Houterman *et al.*, 2007; Schmidt *et al.*, 2013), except for *SIX6* which has homologues in *Colletotrichum* (Gawehns *et al.*, 2014). *SIX1* (also known as *Avr3*), *SIX3* (*Avr2*), *SIX4* (*Avr1*) and *SIX5* are recognized by resistance genes which have been introgressed into tomato (Houterman *et al.*, 2008, 2009; Ma *et al.*, 2015; Rep *et al.*, 2004; Takken and Rep, 2010), whereas gene knock-outs have demonstrated that *SIX1*, *SIX3*, *SIX5* and *SIX6* contribute directly to virulence (Gawehns *et al.*, 2014; Houterman *et al.*, 2009; Ma *et al.*, 2015; Rep, 2005; Takken and Rep, 2010). *SIX* genes have also been found in other f. spp. of *F. oxysporum* (Fraser-Smith *et al.*, 2014; Sasaki *et al.*, 2015b),

\*Correspondence: Email: andrew.taylor@warwick.ac.uk

†Present address: Enza Zaden, Via San Leonardo, 13, Tarquinia, (Viterbo) c.a.p. 01016, Italy.

notably *SIX1*, *SIX4*, *SIX8* and *SIX9* in an *F. oxysporum* isolate infecting *Arabidopsis* and *Brassica* (Thatcher *et al.*, 2012), *SIX6* in *F. oxysporum* f. sp. *vasinfectum* (Chakrabarti *et al.*, 2011), *SIX1* and *SIX6* in f. sp. *betae* (Covey *et al.*, 2014), *SIX1*, *SIX7* and *SIX10* in f. sp. *canariensis* and *lini* (Laurence *et al.*, 2015), *SIX1*, *SIX7* and *SIX8* in f. sp. *cubense* (Meldrum *et al.*, 2012) and *SIX3*, *SIX5* and *SIX7* in f. sp. *cepae* (Sasaki *et al.*, 2015b). In addition, *SIX4* has been shown to play a role in the virulence of *F. oxysporum* f. sp. *conglutinans*, the cause of cabbage yellows (Kashiwa *et al.*, 2013).

The genetically heterogeneous nature and lack of reliable morphological characters in the *F. oxysporum* complex have meant that distinguishing between pathogenic and non-pathogenic isolates and between different f. spp. has been challenging, and has previously relied on pathogenicity tests on different host plants. Molecular methods based on standard approaches, such as DNA fingerprinting and multilocus genotyping using housekeeping genes, have failed to reliably distinguish between different f. spp. (O'Donnell *et al.*, 1998), but the presence/absence of certain *SIX* genes or sequence differences in these genes may form the basis for more reliable detection and identification. *SIX* genes have been used to identify and distinguish races in FOL, where race 2 and 3 isolates, which lack the *SIX4* gene found in race 1, are identified on the basis of variation in *SIX3* sequence (Lievens *et al.*, 2009). Sequence differences in *SIX8* have also been used to identify and distinguish races of *F. oxysporum* f. sp. *cubense*, including tropical race 4 (Fraser-Smith *et al.*, 2014).

Bulb onion (*Allium cepa* L.) is an important crop globally with a total production of 83 million tonnes (FAOSTAT, 2012). Production is often affected by *Fusarium* basal rot (FBR) caused by *F. oxysporum* f. sp. *cepae* (FOC), which is increasing in prevalence, particularly in the UK (Taylor *et al.*, 2013). FOC infects the roots and basal plates of onions, causing symptoms at all stages of plant development, ranging from damping off and delayed seedling emergence to bulb rot at pre- and post-harvest stages (Entwistle, 1990). Infection is favoured by warm temperatures (28–32 °C optimum) and disease incidence is predicted to increase as a result of climate change (Abawi and Lorbeer, 1972; Cramer, 2000; Kehr *et al.*, 1962). FOC also produces chlamydozoospores which can survive for many years in the soil, making disease management very challenging (Brayford, 1996; Cramer, 2000). Other *Fusarium* species have also been associated with root rots of onions or other alliums, including *F. proliferatum*, *F. redolens* and *F. avenaceum*, but are generally less common than FOC (Bayraktar and Dolar, 2011; Du Toit *et al.*, 2003; Galván *et al.*, 2008; Ghanbarzadeh *et al.*, 2013; Shinmura, 2002; Stankovic *et al.*, 2007; Yamazaki *et al.*, 2013), particularly in the UK (Vágány, 2012).

Few studies have attempted the molecular characterization of FOC isolates specifically, but phylogenetic analyses based on *translation elongation factor 1 $\alpha$*  (*EF-1 $\alpha$* ) sequencing and amplified fragment length polymorphism (AFLP) markers have suggested

that *F. oxysporum* can be divided into three distinct clades, with FOC isolates appearing in two of these (Galván *et al.*, 2008; O'Donnell *et al.*, 1998; Sasaki *et al.*, 2015b; Taylor *et al.*, 2013). A more recent study using the intergenic spacer region (IGS) identified eight clades among FOC isolates from bulb onion and *Allium fistulosum* (Sasaki *et al.*, 2015b). However, few molecular studies with FOC, or indeed other f. spp., have associated pathogenicity tests which has confused identification, although a partial association was observed between pathogenicity on Welsh onion and IGS sequence (Dissanayake *et al.*, 2009a; Sasaki *et al.*, 2015b). Studies have also shown that there is genetic diversity amongst FOC isolates based on AFLP markers, inter-simple sequence repeat (ISSR) markers, random amplified polymorphic DNA (RAPD) markers, rRNA, *EF-1 $\alpha$*  or IGS sequencing (Bayraktar and Dolar, 2011; Bayraktar *et al.*, 2010; Dissanayake *et al.*, 2009a, b; Galván *et al.*, 2008; Sasaki *et al.*, 2015b; Southwood *et al.*, 2012a, b; Vágány, 2012). More recently, homologues of *SIX3*, *SIX5* and *SIX7* have been identified in FOC and their presence has been associated with pathogenicity on Welsh onion seedlings (Sasaki *et al.*, 2015b). To date, this is the only record of any putative effector genes in FOC.

The aim of this study was to characterize *F. oxysporum* isolates from onion through the sequencing of housekeeping and pathogenicity-related genes with a particular emphasis on *SIX* genes. The presence/absence of pathogenicity genes was then compared with the ability of the isolates to cause disease in both onion seedlings and bulbs. Isolates of *F. oxysporum* f. spp. *pisi* (pea), *dianthi* (carnation), *narcissi* (daffodil), *cubense* (banana), *lycopersici* (tomato) and other *Fusarium* species isolated from onion/leek (*F. avenaceum*, *F. proliferatum* and *F. redolens*) were also included in the molecular characterization for comparison.

## RESULTS

### Pathogenicity testing

In the onion seedling tests, significant differences were observed in the pathogenicity of the 32 *F. oxysporum* isolates (Table 1) for the two experiments using Napoleon and HZS onion cultivars (Fig. 1,  $P < 0.001$ ). The pathogenicity of each *F. oxysporum* isolate was highly correlated between the two cultivars ( $r = 0.97$ ,  $P < 0.001$ ). Across both onion cultivars, 18 of the 32 isolates resulted in a significant reduction in seedling survival compared with the uninoculated control and were classed as pathogenic, whereas the remaining 14 isolates (including Fo47) were non-pathogenic and had little or no effect (Fig. 1). Two isolates (A1\_2 and 55) caused significant seedling mortality on cv. Napoleon, but not on HZS. Over all the isolates, Napoleon was more susceptible than HZS to *F. oxysporum*.

In the onion bulb test (cv. Napoleon), significant differences were observed in disease levels amongst the 32 *F. oxysporum* isolates ( $P < 0.001$ , Fig. 2), with 21 pathogenic isolates resulting in

**Table 1** *Fusarium* isolates used for pathogenicity testing and/or molecular characterization in this study.

<i>Fusarium</i> species	Isolate code	Location	Origin	Source*	Year isolated
<b>Isolates used for pathogenicity testing and/or molecular characterization</b>					
<i>F. oxysporum</i>	A13	Bedfordshire, UK, site 1	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	A23	Bedfordshire, UK, site 2	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	A28	Bedfordshire, UK, site 2	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	A35	Bedfordshire, UK, site 3	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	F1	Bedfordshire, UK, site 4	Onion bulb	V. Vagany, WCC	2010
<i>F. oxysporum</i>	195	Suffolk, UK, site 1	Onion bulb	C. Handy, WCC	2012
<i>F. oxysporum</i>	224	Suffolk, UK, site 1	Onion bulb	C. Handy, WCC	2012
<i>F. oxysporum</i>	244	Suffolk, UK, site 1	Onion bulb	C. Handy, WCC	2012
<i>F. oxysporum</i>	A21	Suffolk, UK, site 2	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	R3	Suffolk, UK, site 3	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	M1	Suffolk, UK, site 4	Onion bulb	V. Vagany, WCC	2010
<i>F. oxysporum</i>	M9	Suffolk, UK, site 4	Onion bulb	V. Vagany, WCC	2010
<i>F. oxysporum</i>	G12	Suffolk, UK, site 5	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	K3b	Suffolk, UK, site 6	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	S1B	Essex, UK, site 1	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	A14	Essex, UK, site 2	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	A19	Essex, UK, site 2	Onion bulb	V. Vagany, WCC	2009
<i>F. oxysporum</i>	NL70/7	Essex, UK, site 3	Onion bulb	V. Vagany, WCC	2010
<i>F. oxysporum</i>	A1_2	Warwickshire, UK	Onion bulb	V. Vagany, WCC	2008
<i>F. oxysporum</i> f. sp. <i>cepae</i>	FUS2	Lincolnshire, UK	Onion bulb	R. Noble, East Malling Research	Unknown
<i>F. oxysporum</i>	55	Lincolnshire, UK, site 1	Onion bulb	C. Handy, WCC	2012
<i>F. oxysporum</i>	84	Lincolnshire, UK, site 1	Onion bulb	C. Handy, WCC	2012
<i>F. oxysporum</i>	125	Lincolnshire, UK, site 1	Onion bulb	C. Handy, WCC	2012
<i>F. oxysporum</i>	RO2	Lincolnshire, UK, site 2	Onion bulb	V. Vagany, WCC	2010
<i>F. oxysporum</i>	FUS1	Nottinghamshire, UK	Onion bulb	R. Noble, East Malling Research	Unknown
<i>F. oxysporum</i>	FUS3	Nottinghamshire, UK	Onion bulb	R. Noble, East Malling Research	Unknown
<i>F. oxysporum</i>	PG	Cambridgeshire, UK	Onion bulb	T. O'Neill, ADAS	Unknown
<i>F. oxysporum</i>	CB3	UK	Onion set	C. Handy, WCC	2012
<i>F. oxysporum</i>	HB17	UK	Onion set	C. Handy, WCC	2012
<i>F. oxysporum</i>	HB6	UK	Onion set	C. Handy, WCC	2012
<i>F. oxysporum</i>	JB4	UK	Onion set	C. Handy, WCC	2012
<i>F. oxysporum</i>	NRRL 54002 (FO47)	France	Soil	ARS collection	Unknown
<i>F. oxysporum</i>	HAZ	USA	Onion bulb	H. van den Biggelaar, Hazera seeds	Unknown
<i>F. oxysporum</i>	L2-1	UK, site 1	Leek	A. Taylor, WCC	2011
<i>F. oxysporum</i>	L9-1	UK, site 2	Leek	A. Taylor, WCC	2011
<i>F. oxysporum</i>	ATCC90245	Colorado, USA	Pinto bean	ATCC collection	1990
<i>F. oxysporum</i> f. sp. <i>pisi</i> race 1	FOP1	UK	Pea	C. Linfield, WCC	Unknown
<i>F. oxysporum</i> f. sp. <i>pisi</i> race 2	FOP2	UK	Pea	C. Linfield, WCC	Unknown
<i>F. oxysporum</i> f. sp. <i>pisi</i> race 5	FOP5	UK	Pea	C. Linfield, WCC	Unknown
<i>F. oxysporum</i> f. sp. <i>pisi</i>	NRRL36311	The Netherlands	Pea	ARS collection	Unknown
<i>F. oxysporum</i> f. sp. <i>lini</i>	FOLIN	UK	Linseed	C. Linfield, WCC	2010
<i>F. oxysporum</i> f. sp. <i>dianthi</i>	R207	UK	Carnation	C. Linfield, WCC	Unknown
<i>F. oxysporum</i> f. sp. <i>narcissi</i>	FOXN7	UK	Daffodil	C. Handy, WCC	2013
<i>F. oxysporum</i> f. sp. <i>narcissi</i>	FOXN139	UK	Daffodil	C. Handy, WCC	2013
<i>F. oxysporum</i> f. sp. <i>freesia</i>	NRRL26990	The Netherlands	Freesia	ARS collection	Unknown
<i>F. oxysporum</i> f. sp. <i>freesia</i>	NRRL26988	The Netherlands	Freesia	ARS collection	Unknown
<i>F. oxysporum</i> f. sp. <i>cubense</i>	E421A3	UK	Banana	C. Nellist, WCC	Unknown
<i>F. avanaceum</i>	L5	UK, site 1	Leek	A. Taylor, WCC	2011
<i>F. proliferatum</i>	A8	Bedfordshire, UK, site 3	Onion bulb	V. Vagany, WCC	2009
<i>F. proliferatum</i>	A40	Bedfordshire, UK, site 3	Onion bulb	V. Vagany, WCC	2009
<i>F. proliferatum</i>	SP1-2	Spain	Onion bulb	V. Vagany, WCC	2010
<i>F. redolens</i>	NL96	Essex, UK, site 3	Onion bulb	V. Vagany, WCC	2010
<i>F. oxysporum</i> f. sp. <i>lycopersici</i> race 3	NRRL54003 (MN25)	USA	Tomato	ARS Collection	Unknown
<b>Genome sequenced isolates used for comparison in molecular characterization</b>					
<i>F. oxysporum</i> f. sp. <i>pisi</i>	NRRL37622 (HDV247)	Unknown	Pea	ARS Collection	Unknown
<i>F. oxysporum</i>	NRRL32931 (FOSC 3-a)	USA	Human	ARS Collection	Unknown
<i>F. oxysporum</i> f. sp. <i>conglutinans</i>	NRRL54008 (PHW808)	USA	Brassica	ARS Collection	Unknown

Table 1 Continued

<i>Fusarium</i> species	Isolate code	Location	Origin	Source*	Year isolated
<i>F. oxysporum</i> f. sp. <i>raphani</i>	NRRL54005 (PHW815)	France	Radish	ARS Collection	Unknown
<i>F. oxysporum</i> f. sp. <i>radicis-lycopersici</i>	NRRL26381 (CL57)	USA	Tomato	ARS Collection	Unknown
<i>F. oxysporum</i> f. sp. <i>cubense</i>	NRRL54006 (II5)	Indonesia	Banana	ARS Collection	Unknown
<i>F. oxysporum</i> f. sp. <i>melonis</i>	NRRL26406	USA	Melon	ARS Collection	Unknown
<i>F. oxysporum</i> f. sp. <i>vasinfectum</i>	NRRL25433	China	Cotton	ARS Collection	Unknown
<i>F. oxysporum</i> f. sp. <i>lycopersici</i> race 2	NRRL34936 (FOL4287)	USA?	Tomato	ARS Collection	Unknown
<i>F. oxysporum</i>	Fo5176	Australia	<i>Brassica oleracea</i>	ARS Collection	Unknown

\*ATCC, American Type Culture Collection, USA; ARS, Agricultural Research Service culture collection, USA; WCC, Warwick Crop Centre, University of Warwick, UK.




Fig. 1 Pathogenicity of 32 *Fusarium oxysporum* isolates on onion seedlings (cv. Napoleon and Hazera Seeds standard susceptible line, HZS). Data shown are the percentage survival values relative to germination after 42 days in a glasshouse. Error bars represent the least significant difference (LSD) (5%) level for each onion cultivar. An 'S' indicates the value below which there is a significant difference from control plants.


8.7%–58.6% bulb area affected compared with the control (0%), and 11 non-pathogenic isolates (including Fo47) having no significant effect (0%–5.0% bulb area affected). Of the 21 pathogenic isolates, A1\_2 (8.7%) and HB6 (19.1%) showed lower levels of pathogenicity, whereas isolate 55 showed an intermediate level of pathogenicity (29.1%). Highly significant correlations were observed between *F. oxysporum* isolate pathogenicity in the bulb test and the seedling tests with cv. Napoleon ( $r = -0.91$ ,  $P = 0.001$ ) and cv. HZS ( $r = -0.88$ ,  $P = 0.001$ ). Pathogenic isolates resulting in significantly greater disease levels compared with the uninoculated controls in onion seedling or bulb tests were considered to be FOC.

#### Molecular characterization: housekeeping genes


The concatenated tree for *EF-1 $\alpha$* , *RNA polymerase II second largest subunit (RPB2)* and  *$\beta$ -tubulin (TUB2)* sequences resulted in the

majority of the 53 *F. oxysporum* isolates being separated into six clades (Fig. 3). Isolates from onion were represented in clades 1, 3 and 5, and clade 1 contained all those that showed some level of pathogenicity in either the onion seedling or bulb tests (or both), with the exception of A1\_2. Clade 1 also included isolate L2-1 from infected leeks, which has also been shown to be pathogenic on onion bulbs (A. Taylor, A. Jackson & J. P. Clarkson, unpublished data). However, clade 1 also included the non-pathogenic isolate HB17 (from onion sets) and isolates of *F. oxysporum* f. sp. *pisi* race 5 (FOP5), f. sp. *freesia* (NRRL26990) and f. sp. *pisi* (NRRL36311). Two non-pathogenic onion isolates, A28 and CB3, showed high similarity to Fo47 in clade 3, whereas four other non-pathogenic onion isolates (A13, M9, R3 and JB4) were placed in clade 5, together with several other f. spp. *Fusarium oxysporum* f. sp. *cubense* isolates were all placed in a separate and distinct clade (clade 6) and the other *Fusarium* species (*F. proliferatum*, *F. redolens* and *F. avenaceum*) formed distinct outgroups.





**Fig. 2** Pathogenicity of 32 *Fusarium oxysporum* isolates on onion bulbs (cv. Napoleon). Data shown are the percentage bulb areas diseased on bisected bulbs after 9 weeks at 20 °C. Error bar represents the least significant difference (LSD) (5%) level. An 'S' indicates the value above which there is a significant difference from the uninoculated control bulbs.



**Fig. 3** Maximum likelihood tree of *Fusarium* isolates from onion and other hosts based on a concatenated alignment of *translation elongation factor 1 $\alpha$*  (*EF-1 $\alpha$* ) (GenBank accession numbers KP964857–KP964909), *RNA polymerase II second largest subunit* (*RPB2*) (GenBank accession numbers KP964804–KP964856) and  *$\beta$ -tubulin* (*TUB2*) (GenBank accession numbers KP964910–KP964962) genes. Numbers represent bootstrap values from 1000 replicates. Scale bar indicates 0.01 substitutions per site. The tree is rooted through L5 (*F. avenaceum*) and this branch has been collapsed because of its distance from *F. oxysporum*. BI refers to a sequence derived from the genomes on the Broad Institute *Fusarium* database (Broad Institute/MIT, 2007).

Additional trees for each of the housekeeping genes were constructed using neighbour-joining, minimum evolution and UPGMA (Unweighted Pair Group Method with Arithmetic Mean) methods, and similar topography was observed (data not shown).

### Molecular characterization: SIX genes

#### *Fusarium oxysporum* isolates from onion

Using primers based on the FOL or FOC genomes, homologues of *SIX3*, *SIX5*, *SIX7*, *SIX10*, *SIX12* and *SIX9*, *SIX14*, respectively were


identified within the 31 *F. oxysporum* isolates from UK onions (Table 2) and USA isolate HAZ. All seven of these *SIX* genes were present in the 18 highly pathogenic isolates identified from the onion seedling and bulb tests (Table 2). In contrast, all *SIX* genes were absent in the 11 non-pathogenic isolates, with the exception of isolate PG which contained *SIX9* (Table 2). Isolate 55, which had an intermediate level of pathogenicity on onion bulbs, and was mildly pathogenic on seedlings (cv. Napoleon), contained only *SIX9* and *SIX14*, whereas isolates HB6 and A1\_2, which were weakly pathogenic on Napoleon bulbs (and on seedlings for

**Table 2** Presence/absence of *Secreted In Xylem (SIX) 1–14* and three putative novel effectors in *Fusarium oxysporum* and other selected species.

<i>Fusarium</i> species	Host	Isolate code	Pathogenicity*	SIX genes†																
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	C5‡	CRX1§	CRX2¶
<i>F. oxysporum</i> (FOC)	Onion	A23	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	A19	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	–
<i>F. oxysporum</i> (FOC)	Onion	RO2	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	A14	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	–
<i>F. oxysporum</i> (FOC)	Onion	K3B	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	195	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	FUS2	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	125	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	FUS3	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	NL70/7	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	–
<i>F. oxysporum</i> (FOC)	Onion	84	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	M1	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	224	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	F1	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	S1B	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	–
<i>F. oxysporum</i> (FOC)	Onion	A35	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	A21	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	–
<i>F. oxysporum</i> (FOC)	Onion	244	B/S1/S2	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	1
<i>F. oxysporum</i> (FOC)	Onion	55	B/S1	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	1
<i>F. oxysporum</i> (FOC)	Onion	HB6	B	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i> (FOC)	Onion	A1_2	B/S1	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i>	Onion	G12	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	2
<i>F. oxysporum</i>	Onion	CB3	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i>	Onion	PG	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	3
<i>F. oxysporum</i>	Onion	R3	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i>	Onion	A13	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i>	Onion	FUS1	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	1
<i>F. oxysporum</i>	Onion	M9	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i>	Onion	JB4	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i>	Onion	HB17	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i>	Onion	A28	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	2
<i>F. oxysporum</i>	–	FO47	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i>	Onion	HAZ	(+)	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	+
<i>F. oxysporum</i>	Leek	L2-1	(+)	–	–	+	–	+	–	+	–	+	+	–	+	–	+	+	+	+
<i>F. oxysporum</i>	Leek	L9-1	(–)	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i> f. sp. <i>lycopersici</i>	Tomato (race 3)	NRRL54003 (MN25)	NT	+	+	+	–	+	+	+	+	+	+	+	+	+	–	–	–	–
<i>F. oxysporum</i> f. sp. <i>lycopersici</i>	Tomato (race 1)	FOL1	NT	NT	NT	+	NT	NT	NT	NT	NT	NT	NT	NT	NT	–	–	–	–	–
<i>F. oxysporum</i> f. sp. <i>phaseoli</i>	Pinto bean (race 4)	ATCC90245	NT	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i> f. sp. <i>pisi</i>	Pea (race 1)	FOP1	NT	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>F. oxysporum</i> f. sp. <i>pisi</i>	Pea (race 2)	FOP2	NT	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	+
<i>F. oxysporum</i> f. sp. <i>pisi</i>	Pea (race 5)	FOP5	NT	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–







**Fig. 4** Maximum likelihood trees of *Fusarium* isolates from onion and other hosts based on (a) *SIX 7*, (b) *SIX 9*, (c) *SIX 10*, and (d) *SIX 12* gene sequences. Numbers represent bootstrap values from 1000 replicates. Scale bars indicate the number of substitutions per site. All FO *lycopersici* refers to the genome sequenced isolates listed in Table 1 as well as additional identical sequences obtained from a BLAST search. All FO *cubense* refers to the genome sequenced isolate I15 as well as identical BLAST hits. Sequences of other NRRL isolates were extracted from genome sequences (Broad Institute/MIT, 2007). All FO *canariensis* and FO *lini* isolates (with the exception of FOLIN, *SIX 12*) are as described by Laurence *et al.* (2015).

*SIX 8* was detected in FOL and *F. oxysporum* f. sp. *cubense* and f. sp. *phaseoli*, whereas *SIX 9* was detected in FOL using the FOL primers, and in FOC, *F. oxysporum* f. sp. *dianthi* and f. sp. *narcissi* using the FOC primers. *SIX 11* was found in *F. oxysporum* f. sp. *phaseoli*, f. sp. *pisi* race 1 and FOL. *SIX 13* was detected in *F. oxysporum* f. sp. *pisi* races 2 and 5, f. sp. *dianthi*, f. sp. *narcissi*, f. sp. *freesia* and FOL. *SIX 14* was detected in FOC, FOL, *F. oxysporum* f. sp. *pisi* and f. sp. *freesia*. None of the other *Fusarium* species tested contained any of the *SIX* genes.

*SIX* gene sequence variation was observed across the different *F. oxysporum* f. spp., and phylogenetic trees showed that FOC isolates were clearly separated from the other f. spp. based on *SIX 7*, *SIX 10* and *SIX 12*, but not *SIX 9* (Fig. 4). The FOL *SIX 9* gene formed part of a separate clade to the *SIX 9a* and *SIX 9b* genes identified in the *Arabidopsis/brassica*-infecting isolate (Fo5176), whereas the *SIX 9* sequence from FOC was in the same clade as *SIX 9a/SIX 9b* and was closer to *SIX 9a* than *SIX 9b*. The separation of FOC isolates from other *F. oxysporum* f. spp. was less clear for *SIX 9* as the sequence was very similar to that from both *F. oxysporum* f. spp. *narcissi* and *dianthi*. *Fusarium oxysporum* isolates from carnation and *Narcissus* could not be distinguished on the basis of any of the *SIX* gene sequences. For *SIX 5*, six FOC isolates (S1b, A14, A19, A21, NL70/7 and HAZ) had a single base change at position 316 (G to A, Fig. S1, see Supporting Information) causing a single amino acid change from R to K. All of these isolates were in clade 1b in the

housekeeping gene tree (Fig. 3). For *SIX 9*, two onion isolates, A21 (pathogenic) and PG (non-pathogenic), had a different sequence type, differing by a single base pair (T instead of A), resulting in an amino acid change from D to V.

### Molecular characterization: putative novel effectors

Two putative novel effectors, *C5* and *CRX1*, were detected in all of the 18 FOC isolates that were highly pathogenic across both seedling and bulb assays (Table 2), although *CRX1* was also found in isolate A28 which was non-pathogenic. The presence of a third putative effector (*CRX2*) was partially associated with pathogenicity, being detected in 13 of the 18 highly pathogenic isolates (Table 2) and in four non-pathogenic isolates (A28, FUS1, G12 and PG). FOC isolate 55, which demonstrated 'intermediate' pathogenicity on onion bulbs, contained *C5*, *CRX1* and *CRX2* (as well as *SIX 9* and *SIX 14*), whereas FOC isolate A1\_2, which was weakly pathogenic, had no putative effectors. *C5* and *CRX1* were not present in any of the other *F. oxysporum* f. spp., whereas *CRX2* was detected in f. sp. *pisi* race 2, *F. redolens* and all the *F. proliferatum* isolates. Genes with homology to *CRX1/CRX2* were also found in the genome sequences of *F. oxysporum* f. sp. *pisi* (HDV247: FOVG\_18080) and f. sp. *radicis-lycopersici* (CL57: FOCG\_17596 and FOCG\_16735) (Broad Institute/MIT, 2007). *C5* was not present in any of the published *Fusarium* genomes (Table 1) and there are

	Signal Peptide	RxLR domain	dEER
CRX2 (isolate FUS2)	MAEVI <del>GLVSGLLT</del> LATFA	<b>HKSVTKVQKAVQSFRTLP</b> <b><i>PRQLRELLS</i></b> ELAE <del>LS</del> TVL Q <del>D</del>	
CRX1 (FOC)	MAE <del>LIGLVSGLLT</del> LATFA	<b>HQSVTKVQEAVQSFQSLP</b> <b><i>PRQLRELLS</i></b> ELTEL <del>GT</del> TVL Q <del>D</del>	
FOCG_17596 CL57	MAEVI <del>GLVSGLLT</del> LATFA	<b>HKSVTKVQKAVQSFRTLP</b> <b><i>PRQLRELLS</i></b> ELAE <del>LS</del> TVL Q <del>D</del>	
FOCG_16735 CL57	MAEAIGLVSGLLALATFA	<b>HQSVTKG- - - T</b> PFRVS- - - RTKYSTPRTLPSTWR-	
FOVG_18080 HDV247	MAEVI <del>GLVSGLLT</del> LATFA	<b>HKSVTKVQEAVQSFRC</b> L <b><i>PRQLRELLS</i></b> ELAE <del>LS</del> TVL Q <del>D</del>	
<i>F. proliferatum</i>	MAEVI <del>GLVSGLLAL</del> LATFA	<b>HKSVTKVQEAVQSFRTLP</b> <b><i>RQVRELLS</i></b> ELAE <del>LS</del> TVL Q <del>D</del>	

**Fig. 5** Amino acid alignment of putative RxLR effectors from *Fusarium oxysporum* and *F. proliferatum*. The signal peptide (as predicted by SignalP) is shaded in light grey, whereas the RxLR domain is shown in bold (bold and italics for an incomplete RxLR domain). Amino acids that often occur after an RxLR domain (dEER) are underlined. Sequences from HDV247 and CL57 were obtained from the Broad Institute *Fusarium* database (Broad Institute/MIT, 2007).

no sequence matches at the National Center for Biotechnology Information (NCBI).


The putative effectors *CRX1* and *CRX2* in *F. oxysporum* contained RxLR domains close to the N-terminus (at amino acid positions 37–40, Fig. 5), whereas *F. proliferatum* isolates had a modified (RQVR) sequence in this position. The RxLR domains were flanked by modified dEER domains, defined by the presence of >10% D or E residues (Jiang *et al.*, 2008). All *C5* and *CRX1* sequences were identical across the *F. oxysporum* isolates and, based on the partial coding DNA sequence, a phylogenetic tree clearly separated *CRX1* and *CRX2* (Fig. 6). The majority of the 13 (pathogenic) FOC isolates that contained *CRX2* had identical sequences, although the non-pathogenic isolate FUS1 also had the same sequence (Fig. 6). Three non-pathogenic isolates from onion (A28, G12 and PG) had slightly different *CRX2* sequence types and, in the case of isolates PG and G12, this resulted in a stop codon in the middle of the coding region (data not shown). *CRX2* sequences from *F. oxysporum* f. sp. *radicis-lycopersici* (FOCG\_17596) and *F. redolens* (NL96) were very similar to the predominant FOC sequence, whereas the three *F. proliferatum* isolates, which also contained *CRX2*, formed a distinct clade.

### Expression of putative effectors

Using real-time reverse transcription-polymerase chain reaction (RT-PCR), the seven *SIX* genes and the three putative effectors *C5*, *CRX1*, *CRX2* were all shown to be expressed *in planta* across the time course following inoculation with FOC isolate FUS2. *SIX3*, *SIX5*, *SIX7*, *SIX9*, *SIX10*, *SIX12* and *CRX1* showed significant increases in expression levels at 36–72 h post-inoculation (hpi) compared with the first time point at 8 h (Fig. 7), whereas *SIX14* only showed a significant increase at 72 hpi. There was no increase in *C5* expression for any of the time points compared with 8 hpi, although a significant increase was detected between 16 and 72 hpi. *CRX2* expression levels did not change significantly across the time course.

### DISCUSSION


In this study, we have demonstrated, for the first time, a clear association between the presence of seven *SIX* genes and two



**Fig. 6** Neighbour-joining tree of *Fusarium oxysporum* *CRX1* and *CRX2* genes and their homologues. Numbers represent bootstrap values from 1000 replicates. The scale bar indicates 0.02 substitutions per site. Sequences from HDV247 and CL57 were obtained from the Broad Institute *Fusarium* database (Broad Institute/MIT, 2007). The sequence from Fa05001 was obtained from an assembled genome (GenBank accession number GCA\_000769215).

other putative effectors (*C5*, *CRX1*) in FOC and pathogenicity on onion. A third putative effector (*CRX2*) showed a partial association with pathogenicity. Pathogenicity tests on onion seedlings and bulbs were consistent across all the *F. oxysporum* isolates, and a strong correlation between pathogenicity results using two different onion cultivars in the seedling tests supports the suggestion that there is no cultivar × isolate interaction (Taylor *et al.*, 2013).

Three of the seven *SIX* genes identified in FOC (*SIX3*, *SIX5* and *SIX7*) had been identified previously, but their presence was only related to pathogenicity on onion seedlings (Sasaki *et al.*, 2015b), and not in a bulb test, which is more appropriate to disease expression in the field. In addition, this study did not provide evidence of expression *in planta*. The remaining four *SIX* genes



**Fig. 7** Quantitative expression of a set of putative effector genes in onion roots following inoculation with *Fusarium oxysporum* f. sp. *cepae* (FOC) isolate FUS2. Expression was calculated relative to *translation elongation factor 1 $\alpha$*  (*EF-1 $\alpha$* ) and  *$\beta$ -tubulin* (*TUB2*). Error bars show the standard error of the mean (SEM) of three replicates; hpi, hours post-inoculation. Asterisks indicate expression levels significantly different from 8 hpi based on analysis of variance (ANOVA) followed by Tukey's test (\* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ ). + indicates that the expression of *C5* was significantly higher at 72 hpi relative to 16 hpi ( $P < 0.05$ ).

(*SIX9*, *SIX10*, *SIX12* and *SIX14*) are reported here for the first time in FOC. The presence of *SIX* genes has similarly been associated with the pathogenicity of *F. oxysporum* isolates on tomato (*SIX1–7*; Lievens *et al.*, 2009), cotton (*SIX6*; Chakrabarti *et al.*, 2011) and banana (*SIX1*, *SIX7* and *SIX8*; Meldrum *et al.*, 2012). The function of the 14 *SIX* genes detected so far in *F. oxysporum* is unclear as they show little or no homology with any known proteins (Fraser-Smith *et al.*, 2014). However, *SIX1*, *SIX3*, *SIX4*, *SIX5* and *SIX6* have all been shown to make a direct contribution to pathogenicity (Gawehns *et al.*, 2014; Houterman *et al.*, 2009; Ma *et al.*, 2015; Rep, 2005; Takken and Rep, 2010; Thatcher *et al.*, 2012).

One *F. oxysporum* isolate from leek (L2-1) also shared the same effector gene profile as the pathogenic onion isolates. Preliminary work (Taylor A, Jackson A.C, Clarkson J.P, unpublished.) has demonstrated that this leek isolate is also pathogenic on onion bulbs, whereas another leek isolate (L9-1), which was non-pathogenic, lacked any of the effector genes. FOC has also been shown to cause basal rot in Welsh onion (*A. fistulosum*; Dissanayake *et al.*, 2009a,b; Sasaki *et al.*, 2015b) and garlic (*A. sativum*; Rout *et al.*, 2014), suggesting that FOC might be more appropriately named *F. oxysporum* f. sp. *allii*. However, it should be noted that, although FOC isolates from *A. fistulosum* can be

pathogenic on *A. cepa* seedlings, they lack *SIX3*, *SIX5* and *SIX7* (Sasaki *et al.*, 2015b), and the pathogenicity of these isolates on onion bulbs has yet to be tested.

It is likely that the FOC *SIX* genes, and possibly the other putative effectors, are located on dispensable/supernumerary chromosomes, as has been reported for FOL (Ma *et al.*, 2010). FOC *SIX3* has been shown to be located on a small (4-Mb) chromosome, which may be equivalent to a FOL supernumerary chromosome (Sasaki *et al.*, 2015b). It has also been shown recently that *SIX3* and *SIX5* share the same promoter and appear to act as a gene pair, which is recognized by the *I-2* resistance gene (Houterman *et al.*, 2009; Ma *et al.*, 2015). It has been suggested that this is unique to FOL, but our data and those of Sasaki *et al.* (2015a) confirm that these genes are also present in FOC. For all of the seven *SIX* genes in FOC, sequences were mostly identical and, similarly, there was also a high level of sequence conservation with those reported in FOL, suggesting a conservation of function. Indeed, five of the seven FOC *SIX* genes (*SIX3*, *SIX7*, *SIX10*, *SIX12* and *SIX14*) showed no intraspecific sequence differences within isolates originating from the UK, USA and Japan. This supports the suggestion that *SIX* genes in FOC have been acquired by horizontal gene transfer of the supernumerary chromosomes, as proposed for other *F. oxysporum* f. spp. (Fraser-Smith *et al.*, 2014; Laurence *et al.*, 2015; Ma *et al.*, 2010). A naturally occurring isolate of *F. oxysporum* from onion (isolate 55) identified in this study had only two of the seven *SIX* genes (*SIX9* and *SIX14*) and a corresponding intermediate level of pathogenicity. This phenomenon has not been observed so far in other f. spp., and may suggest a missing or mutated copy of a supernumerary chromosome.

In this study, we also report, for the first time, three novel putative effectors in FOC (*C5*, *CRX1* and *CRX2*), and the presence of *C5* and *CRX1* had an almost complete correlation with pathogenicity. *C5* has no homology to any sequenced gene, suggesting that it may be FOC specific, but both *CRX1* and *CRX2* contained RxLR domains, a motif found in many diverse oomycete candidate effector genes which, in some reports, has been shown to facilitate entry into host cells (Kale, 2012; Kale *et al.*, 2010; Rehmany *et al.*, 2005). Putative RxLR effectors have also been reported in FOL (Ma *et al.*, 2013), but this is the first report of putative RxLR effectors in FOC. *CRX2* only had a partial association with pathogenicity in FOC, and also has very close homologues in *F. oxysporum* f. sp. *radicis-lycopersici*, *F. proliferatum* and *F. redolens*. We also demonstrated, for the first time, that the *SIX* genes and the putative FOC-specific effector *CRX1* are up-regulated during the infection process, with levels of expression *in planta* increasing over a time course, further supporting the hypothesis that these genes play a role in pathogenicity. In addition, with the exception of *C5*, all other putative effectors and *SIX* genes had an associated signal peptide, supporting the hypothesis that they are secreted. However, *SIX12* in FOL, which is also reported to have no associ-

ated signal peptide, was found in the xylem sap of infected tomato plants, leading to the conclusion that it was secreted by a different mechanism (Schmidt *et al.*, 2013). Additional functional analyses are now required to reveal the potential roles of the new putative effectors in pathogen virulence.

The characterization of all the *F. oxysporum* isolates using the three housekeeping genes *EF-1 $\alpha$* , *RPB2* and *TUB2* showed that there was considerable genetic variation between isolates from onion, as reported previously using AFLP markers, *EF-1 $\alpha$*  and IGS sequencing (Galván *et al.*, 2008; Sasaki *et al.*, 2015b). Although all pathogenic isolates, with the exception of A1\_2 (weakly pathogenic), were placed in clade 1, the same clade also contained a small number of non-pathogenic isolates and some other f. spp., indicating that housekeeping genes are not useful for distinguishing between different *F. oxysporum* f. spp. or between pathogenic and non-pathogenic isolates. However, one study found a partial association between IGS sequence and pathogenicity (Sasaki *et al.*, 2015b), but it was not clear whether other *F. oxysporum* f. spp. share this sequence type. The non-pathogenic *F. oxysporum* isolates from onion were generally scattered throughout the phylogenetic tree, suggesting that they are more diverse than the pathogenic isolates. Although all *F. oxysporum* isolates in this study were originally from diseased onion tissue, the isolation of non-pathogenic isolates is common and they are likely to be either endophytes or saprophytes, which are secondary colonizers of infected roots or bulbs (Alabouvette *et al.*, 2009).

Our main reason for selecting the additional *F. oxysporum* f. spp. for characterization in this study was to expand the knowledge concerning the distribution of *SIX* genes. A study of *SIX1–SIX7* in a wider range of *F. oxysporum* f. spp. suggested that only *SIX6* (in f. sp. *melonis* and *radicis-cucumerinum*) and *SIX7* (in f. sp. *lilii*) were present in f. spp. other than FOL (Lievens *et al.*, 2009). However, more recent work (and the release of whole genome sequences for selected *F. oxysporum* f. spp.) has shown that all of the 14 *SIX* genes are present in various complements in other f. spp. (Broad Institute/MIT, 2007; Fraser-Smith *et al.*, 2014; Laurence *et al.*, 2015; Sasaki *et al.*, 2015b). FOC isolates in this study contained seven of the 14 *SIX* genes, whereas five were identified for *F. oxysporum* f. sp. *freesia* (NRRL26988) and f. sp. *pisi* race 1. Interestingly, a second *freesia* isolate (NRRL26990) examined here contained no *SIX* genes at all, suggesting that it may not be pathogenic or may have been misidentified. We also found a range of *SIX* genes present in isolates of *F. oxysporum* f. sp. *pisi* and there appeared to be distinct variation between races in terms of *SIX* gene complement. The only previous literature on *SIX* genes in *F. oxysporum* f. sp. *pisi* showed an absence of *SIX6* in races 1, 2, 5 and 6 (Chakrabarti *et al.*, 2011), which is in agreement with our findings.

In this study, an *F. oxysporum* f. sp. *dianthi* isolate (R207) contained *SIX7*, *SIX9*, *SIX10* and *SIX12*, which is in contrast with previous findings that reported an absence of *SIX1–7* in f. sp. *dianthi*



isolates from the USA and the Netherlands (Lievens *et al.*, 2009). Overall, our results, in combination with other reports, suggest that *SIX* gene complements and/or sequences can be used to separate *F. oxysporum* f. spp. without the need for pathogenicity testing. Recently, a quantitative PCR assay was developed for the detection of FOC *in planta* based on *SIX3* (Sasaki *et al.*, 2015a). Although this test requires more rigorous validation against a range of *F. oxysporum* and other soil-borne fungi, it could potentially be useful as a soil or plant test for FOC.

Finally, the other *Fusarium* species *F. proliferatum*, *F. avenaceum* and *F. redolens*, which were also isolated from diseased onions and leeks in this study, did not contain any *SIX* genes. This confirms the results for the isolates of *F. proliferatum* and *F. avenaceum* from sugar beet, where *SIX1* and *SIX6* were absent (Covey *et al.*, 2014). This was also the case for *F. graminearum*, *F. solani* and *F. javanicum*, which lacked *SIX1–7* (Lievens *et al.*, 2009). Whole genome sequences have recently become available for *F. avenaceum* isolates from barley and wheat (Lysøe *et al.*, 2014), and these genomes were examined for the presence of all putative effectors. The only hit was for CRX2 (80% identity), which was only present in an isolate from barley (Fa05001, Fig. 6). The *F. proliferatum* and *F. avenaceum* isolates used in our study were also shown to be pathogenic on onion (A. Taylor, A. C. Jackson, and J. P. Clarkson, unpublished data) and must therefore possess a different mechanism of infection compared with FOC. Overall, our findings provide a greater understanding of pathogenicity in FOC, and could potentially improve the diagnosis and control of *Fusarium* basal rot of onion in the future.

## EXPERIMENTAL PROCEDURES

### *Fusarium* isolates

*Fusarium* isolates were obtained from diseased onion roots and bulbs from different locations in the UK collected between 2008 and 2012 (Table 1), as described by Taylor *et al.* (2013). Thirty-one *F. oxysporum* isolates were selected for subsequent molecular characterization and pathogenicity tests based on sample location, colony morphology, preliminary pathogenicity data (where available) and *EF-1α* sequences of a few isolates (Taylor *et al.*, 2013; Vágány, 2012), as well as the genome sequenced non-pathogenic biocontrol agent Fo47 (NRRL54002). In addition, a genome-sequenced isolate of FOL race 3 (MN25, NRRL54003), a FOC isolate from the USA (HAZ), two *F. oxysporum* isolates from diseased UK leeks (L2-1, L9-1) and 17 isolates of *F. oxysporum* f. spp. *pisi*, *dianthi*, *narcissi*, *cubense* and *lycopersici*, as well as three other *Fusarium* species from diseased onions/leeks (*F. avenaceum*, *F. proliferatum* and *F. redolens*), were also obtained from various researchers and culture collections for comparison in the molecular characterization studies (Table 1).

### Pathogenicity testing: seed inoculation

The 31 *F. oxysporum* isolates from UK onions and the non-pathogenic isolate Fo47 were assessed for their pathogenicity on onion seedlings as

described by Taylor *et al.* (2013). Two experiments were carried out: one using cv. Napoleon (Syngenta, Cambridge, Cambridgeshire UK) and the second using cv. HZS (a standard FOC susceptible line from Hazera Seeds, Made, The Netherlands). For each experiment, there were four independent replicates over time, each consisting of a tray of 28 onion seeds per isolate, which were positioned in a glasshouse using an alpha design (Genstat v.12, VSN International, Hemel Hempstead, Hertfordshire, UK). Three trays of uninoculated control treatments (seeds soaked in sterile distilled water (SDW)) were included in each replicate experiment. The number of surviving seedlings was recorded after 6 weeks and the percentage survival was calculated relative to germination to allow for any variation in germination between replicates. Significant differences between treatments (isolates) for these data were assessed using residual (or restricted) maximum likelihood (REML) analysis (Welham & Thompson, 1997) in GenStat. The Pearson product moment correlation coefficient was also calculated to determine the correlation between the two cultivars.

### Pathogenicity testing: bulb inoculation

The same 32 *F. oxysporum* isolates used in the seedling tests were assessed for pathogenicity on healthy, stored onion bulbs (cv. Napoleon). The outer scales of the bulbs were removed to leave a single brown layer of skin, after which the basal plate of each onion was cut off and the bulb surface was sterilized with 70% ethanol. A potato dextrose agar (PDA) plug (8 mm) taken from the edge of an actively growing colony of each *F. oxysporum* isolate (grown for 7 days at 20 °C) was then positioned on the basal plate of each bulb. Control bulbs were inoculated with a sterile plug of PDA. Bulbs were placed on moist tissue in a plastic box (four per box) inside a sealed plastic bag to maintain high humidity and incubated at 20 °C in the dark. Boxes were randomized in trays following an alpha design. After 48 h, each bulb was wrapped with cling-film to ensure that the agar plug did not dry out. After 9 weeks, each bulb was bisected longitudinally and a digital image was taken (including a 10-cm scale bar). Images were then analysed using ImageJ software (Schneider *et al.*, 2012) to quantify the area of infection as a percentage of the total bulb area. Three independent replicates (four onion bulbs per replicate) were set up for each *F. oxysporum* isolate and significant differences between isolate disease area data were analysed using REML in GenStat. Pearson product moment correlation coefficients were calculated to determine correlations between disease data from seedling and bulb tests.

### Molecular characterization of *Fusarium* isolates: housekeeping genes

Molecular characterization through sequencing of housekeeping genes was carried out for the 32 *F. oxysporum* isolates used in the pathogenicity tests, as well as 21 other f. spp. of *F. oxysporum* and *Fusarium* species (Table 1). Each isolate was grown on PDA for 4–7 days at 25 °C, and three agar plugs (5 mm) taken from the leading edge were placed in a 50-mL tube containing 25 mL of sterile 50% potato dextrose broth (PDB). After incubation at 25 °C for 5 days and centrifugation at 3000 × g for 5 min, excess liquid was removed and the mycelium was rinsed twice with SDW. Finally, excess water was removed and the mycelium was flash frozen in liquid nitrogen before lyophilizing for 24 h. DNA was then extracted from approximately 20 mg of mycelium using a DNeasy plant mini kit (Qiagen, Hilden,



**Table 4** Primer pairs used for molecular characterization of *Fusarium* isolates with product size, annealing temperature and relevant publications.

Gene*	Primers	Sequence 5'–3' (forward primer/reverse primer)	Product size	Annealing temperature (°C)	Publication
<i>TUB2</i>	T1/T22	AACATGCGTGAGATTGTAAGT/TCTGGATGTTGTTGGGAATCC	~1500	60	O'Donnell and Cigelnik (1997)
<i>RPB2</i>	7cF/11aR	ATGGGYAARCAAGCYATGGG/GCRTGGATCTTRTCRSACC	881	57	O'Donnell <i>et al.</i> (2007)
<i>EF-1<math>\alpha</math></i>	exTEF-F/UFexTEF-R	ACCCGGTTC AAGCATCCGATCTGCGA/AGCTTGCCRGACTTGATCTCAGCTC	1269	64	Vágány (2012)
<i>SIX1</i>	SIX1	GTATCCCTCCGGATTTGAGC/AATAGAGCTGCAAAGCATG	992	59	Lievens <i>et al.</i> (2009)
<i>SIX2</i>	SIX2	CAACGCCGTTTGAATAAGCA/TCTATCCGCTTCTTCTCTC	749	59	Lievens <i>et al.</i> (2009)
<i>SIX3</i>	SIX3	CCAGCCAGAAGGCCAGTTT/GGCAATTAACCACTCTGCC	608	59	Lievens <i>et al.</i> (2009)
<i>SIX4</i>	SIX4	TCAGGCTTCACTTAGCATA/GCCGACCGAAAAACCTAA	967	59	Lievens <i>et al.</i> (2009)
<i>SIX5</i>	SIX5	ACACGCTCTACTACTCTCA/GAAAACTCAACGCGGCAAA	667	59	Lievens <i>et al.</i> (2009)
<i>SIX6</i>	SIX6	CTCTCTGAACCATCAACT/CAAGACCAGGTGATAGGCATT	793	59	Lievens <i>et al.</i> (2009)
<i>SIX7</i>	SIX7	CATCTTTTCGCCGACTTGGT/CTTAGCACCTTGAGTAACT	862	59	Lievens <i>et al.</i> (2009)
<i>SIX8</i>	SIX8	TCGCCGATAAACAAGGTGCCG/TTGTGTAGAACTGGACAGTCGATGC	250	59	Meldrum <i>et al.</i> (2012)
<i>SIX9</i>	FOL SIX9	GGGTGGACATATCAGATGTTTCG/GAATACCTGAGTGGAGTGTGTCTTG	458	69	This study
<i>SIX9</i>	FOC SIX9	GGCCAGCCCTAGTCTAACTCC/AACCTAACATGCTGCCGTC AATCG	347	67	This study
<i>SIX10</i>	SIX10	GTTAGCAACTGCGAGACACTAGAA/AGCAACTTCTCTCTTACTAGC	636	65	This study
<i>SIX11</i>	SIX11	ATTCCGGCTTCGGGTCTCGTTTAC/GAGAGCCTTTTGGTGTATTGTAT	559	61	This study
<i>SIX12</i>	SIX12	CTAACGAAGTGAAGAAGTCTC/GCCTCGCTGGCAAGTATTTGT	449	61	This study
<i>SIX13</i>	SIX13	CCTTCATCATCGACGTACAACG/ATCAAACCCGTAACCTCAGCTCC	1027	61	This study
<i>SIX14</i>	FOL SIX14	ATAAAGTGC GACTGGACTTCTGCC/ACCCCATCCACATTCCTAAGCGA	422	67	This study
<i>SIX14</i>	FOL SIX14 nest	GATCCCAATGGGGCTGTGT/GCTGGTGGCTAGAATCTTTTGA	232	59	This study
<i>SIX14</i>	FOC SIX14	ACAACACCGCGACGCTAAAAT/GCACACTCAGTGCAGCAAGTTC	438	61	This study
<i>C5</i>	C5	AGAGTGTGAAGTGAAGGACGAGGGA/CTACGTTCCGCTCACTCATTGCCT	1064	63	This study
<i>CRX1</i>	CRX1	CACCATGTGTACATAAAGGCCGCC/AAAGTCAAGGACCGGACCGGCCG	1654	69	This study
<i>CRX2</i>	CRX2	TTAGTCGCACATCTACCATCACTG/GGAGTCGATCTAACTTCAGG	856	58	This study
<i>CRX2</i>	CRX2 FP	CCAGTGCAATGGTTGAGACGT/ATGCGCTCGCTTTCTATGTATCTG	902	63	This study
Primers used for real-time reverse transcription-polymerase chain reaction (RT-PCR)					
<i>SIX3</i>	QSIX3	GGCCGTCTTCTACTTCACTTAC/GGGAGAATGTTCTAGCATAACC	69	63	This study
<i>SIX5</i>	QSIX5	TGGGCTCGAAAAGTCCAGCAT/TGTTTCGCCGTC AATGTCGCC	114	63	This study
<i>SIX7</i>	QSIX7	TCGATCTCTTTC AAGACAAGGGCA/GTGGACGCGGGCTTGGTGAAC	130	63	This study
<i>SIX9</i>	QSIX9	GCCGACCCAGACTACGCTTT/GCTGGTTTTGGAAGCCAGTTGT	129	63	This study
<i>SIX10</i>	QSIX10	CCCGAAAAGCCTGCACTGACTA/AGAACAACGTCGGTGGGACCA	53	63	This study
<i>SIX12</i>	QSIX12	TGCTGCTCCAAGTACAACCTT/GCTGATACCTTTGGGTCCAACGC	71	63	This study
<i>SIX14</i>	QSIX14	ATGTCGTATGCCGACGGGAA/TTATCTCGTAGACGCCCTTCT	109	60	This study
<i>C5</i>	QC5	GCCTATGGCAGACTTGTGAC/CCACAGCTTCTGGACTATCTCC	126	63	This study
<i>CRX1</i>	QCRX1	AACTCAGGTACCACATCGGGA/CAGGTCGCTCAGCTCAGT	89	60	This study
<i>CRX2</i>	QCRX2	CAATCAGAAAACACGACGGAA/GGAGTCGATCTAACTTCAGG	89	60	This study
<i>EF-1<math>\alpha</math></i>	QTEF	GGTCAGGTCGGTGTGGTTACG/TGGATCTCGCGGAACCTGCAGG	77	63	This study
<i>TUB2</i>	QTUB	TTCTGCTGTCATGTCGGGTG/TCAGAGGAGCAAAGCCAACCA	134	63	This study

\**EF-1 $\alpha$* , translation elongation factor 1 $\alpha$ ; *RPB2*, RNA polymerase II second largest subunit; *SIX*, Secreted In Xylem; *TUB2*,  $\beta$ -tubulin.

Germany) with minor modifications, whereby the mycelium was first homogenized in a lysing matrix A tube (MP Biomedicals, Santa Ana, CA, USA) placed in a FastPrep-24™ machine (MP Biomedicals) set at 6 m/s for 40 s. The manufacturer's protocol was then followed with the addition of an extra centrifugation step (12 500  $\times$  g for 5 min) after the cell lysis stage. DNA integrity was confirmed by gel electrophoresis.

PCR amplification and sequencing of *EF-1 $\alpha$* , *RPB2* and *TUB2* was carried out for all *Fusarium* isolates using published primers (Table 4) with reactions set up using REDTaq® ReadyMix® (Sigma-Aldrich Gillingham, Dorset, UK) in 20- $\mu$ L volumes containing approximately 50 ng of DNA and a final concentration of 0.5  $\mu$ M of each primer. For *EF-1 $\alpha$* , the thermocycling conditions were as follows: one cycle of 5 min at 94 °C; 40 cycles of 45 s at 94 °C, 30 s at 64 °C and 2 min at 72 °C, followed by one cycle of 10 min at 72 °C. For *RPB2*, the conditions were as follows: one cycle of

1.5 min at 94 °C; 40 cycles of 30 s at 94 °C, 1.5 min at 60 °C and 2 min at 68 °C, followed by one cycle of 10 min at 68 °C. For *TUB2*, the conditions were as follows: one cycle of 3 min at 95 °C; 35 cycles of 1 min at 94 °C, 30 s at 60 °C and 1 min at 72 °C, followed by one cycle of 10 min at 72 °C. All PCR amplicons were purified using a QIAquick PCR Purification Kit (Qiagen), sequenced using forward and reverse primers, and contigs were constructed using the SeqBuilder package of DNASTAR® LaserGene® version 10 (DNASTAR Inc., Madison, WI, USA). Sequences were aligned (CLUSTALW method, Thompson *et al.*, 1994), concatenated using MEGA version 5.1 (Tamura *et al.*, 2011) and a maximum likelihood tree was constructed using the calculated best model, Kimura-2-parameter plus gamma (Kimura, 1980). Bootstrap consensus trees were inferred from 1000 replicates (Felsenstein, 1985). Sequences from published *F. oxysporum* genomes (Broad Institute/MIT, 2007) were also included in this analysis and were identified using a Broad Institute (BI) label.

## Molecular characterization of *Fusarium* isolates: putative effectors

Molecular characterization through the detection of the presence/absence and sequencing of putative effector genes was carried out for all *Fusarium* isolates in Table 1. Isolates were assessed for the presence/absence of *SIX1–SIX14* using published primers where available (for *SIX1–SIX8*) (Lievens *et al.*, 2009; Meldrum *et al.*, 2012), whereas new primers were designed for *SIX9–14* based on the published genome sequences of FOL (isolates MN25/4287, Table 4). Nested primers (FOL *SIX14* nest) were used for *SIX14* because of the very short sequence reads and poor quality sequence data, and PCR was carried out as described below, using 1  $\mu$ L of purified PCR product. FOL isolate MN25 was used as a positive control for all *SIX* genes, with the exception of *SIX4*, where isolate FOL1 was used, as MN25 does not contain *SIX4* (Broad Institute/MIT, 2007). All primer sequences were checked against target *SIX* sequences using a preliminary assembly of the FOC genome (Vágány, 2012) and, following this, new primers were designed for *SIX9* (FOC *SIX9*) and *SIX14* (FOC *SIX14*), because of the large sequence differences between FOC and FOL (Table 4). All PCRs for the *SIX* genes were set up as described for the housekeeping genes with standard thermocycling conditions as follows: one cycle of 2 min at 94 °C; 30 cycles of 45 s at 94 °C, 30 s annealing (see Table 4 for temperatures) and 1 min at 72 °C, followed by one cycle of 5 min at 72 °C. A *de novo* assembly of 70-bp paired-end reads (Illumina, San Diego, CA, USA, GAllx sequencing) was carried out using Velvet to obtain a preliminary FOC genome for isolate FUS2 (Vágány, 2012; Zerbino and Birney, 2008), and resulted in 1511 contigs with an N50 of 184 kb (raw 70-bp paired-end Illumina reads submitted to the NCBI Sequence Read Archive under BioProject PRJNA287483). This assembly was investigated for putative new effectors, including *SIX* genes. Proteins predicted by Augustus (Keller *et al.*, 2011; Stanke and Morgenstern, 2005), which were located on the same contig as any of the *SIX* genes and had at least two of the three main characteristics of effector proteins [short length, presence of a signal peptide as predicted by SignalP version 4 (Petersen *et al.*, 2011), no homology to any other known protein] were considered. This led to the discovery of several putative effectors, including gene *C5*. Then, any predicted protein in the rest of the assembly that met these criteria was considered. This list was refined by screening for the presence of published motifs related to pathogenicity, including RxLR-dEER (Rehmany *et al.*, 2005), which led to the discovery of two very closely related genes, *CRX1* and *CRX2*. Primers were then designed for *CRX1*, *CRX2* and *C5* (Table 4), and PCR was carried out using the thermocycling conditions described for the *SIX* genes. Additional primers were also designed (*CRX2* FP) to amplify *CRX2* from *F. proliferatum* (forward primer based on *F. oxysporum* because of a lack of sequence for *F. proliferatum*; reverse primer based on *F. proliferatum* using sequence from initial PCR with *CRX2* primers) in order to assess the presence/absence of the RxLR domain. All PCR amplicons were purified as described previously, and sequencing was carried out using forward primers or both primers for longer amplicons. Where both primers were used, contigs were constructed as described previously. For *F. proliferatum* (isolates A8, A40 and SP1\_2), contigs were constructed from sequences of both PCRs. Sequences were aligned and maximum likelihood trees were constructed for *SIX7*, *SIX9*, *SIX10* and *SIX12*, as described previously. The models used were Kimura-2-parameter, gamma distributed for *SIX7* and with invariant sites for *SIX9*

(Kimura, 1980), Jukes–Cantor for *SIX10* and Jukes–Cantor, gamma distributed for *SIX12* (Jukes and Cantor, 1969). All available homologous sequences (NCBI) were included and sequences were extracted from all available *Fusarium* genome data following BLAST searches (Broad Institute/MIT, 2007). Trees were not constructed for any of the other *SIX* genes because of the limited number of sequences.

## Expression of putative effectors

For the examination of expression *in planta*, onion seedlings were infected with isolate FUS2 as described in Methods S1 (see Supporting Information). RNA was extracted from the pooled root systems of five seedlings using Trizol® reagent (Life Technologies, Paisley, UK), any DNA was removed using DNase I (Sigma-Aldrich) and first-strand cDNA was synthesized using Superscript II reverse transcriptase (Life Technologies) following the manufacturer's guidelines. Real-time RT-PCR was performed in a Roche Lightcycler 480 using the Lightcycler 480 SYBR Green I Master mix (Roche, Burgess Hill, Sussex, UK), following the manufacturer's instructions. Primers were used at a final concentration of 0.4  $\mu$ M with the annealing temperatures in Table 4. The cycling conditions were as follows: one cycle of 95 °C for 5 min, followed by 45 cycles of 95 °C for 10 s, 60/63 °C for 10 s and 72 °C for 10 s. Melt curve analyses were used to confirm a single PCR product. All samples were run in triplicate, standard curves were plotted for each gene and data were expressed as the quantity of the target gene relative to the geometric mean of *EF-1 $\alpha$*  and *TUB2*.

## ACKNOWLEDGEMENTS

This research was funded through the Biotechnology and Biological Sciences Research Council's Horticulture and Potato Initiative (Project BB/K020870/1), an Agriculture and Horticulture Development Board Fellowship (CP113, to AT) and an Erasmus Fellowship (to AR), and we thank these funding agencies for support. The approaches and preliminary data used in this research (including original *F. oxysporum* f. sp. *cepae* genome sequence) were inspired and developed from the PhD of V. Vagany (as cited) which was funded by the Department for Environment, Food and Rural Affairs (Defra) and originally supervised by the late Dr Dez Barbara (University of Warwick, UK) and Dr Surapareddy (Prasad) Sreenivasaprasad (University of Bedfordshire, UK), with additional support from the Food and Environment Research Agency (FERA). We also thank Jeanette Selby for technical support with genome sequencing, the Warwick Crop Centre Horticultural Services team for their assistance with the glasshouse seedling assays, Reinout de Heer (Hazera Seeds) and Syngenta for providing onion seed, Andy Richardson (Allium and Brassica Centre, Boston, Lincs, UK) for providing onion bulbs and Patrick/Ruth Schäfer (University of Warwick, UK) for advice on growing onions in the sterile system. The authors have no conflicts of interest to declare.

## REFERENCES

- Abawi, G.S. and Lorbeer, J.W. (1972) Several aspects of the ecology and pathology of *Fusarium oxysporum* f. sp. *cepae*. *Phytopathology*, **62**, 870–876.
- Alabouvette, C., Olivain, C., Migheli, Q. and Steinberg, C. (2009) Microbiological control of soil-borne phytopathogenic fungi with special emphasis on wilt-inducing *Fusarium oxysporum*. *New Phytol.* **184**, 529–544.
- Bayraktar, H. and Dolar, F.S. (2011) Molecular identification and genetic diversity of *Fusarium* species associated with onion fields in Turkey. *J. Phytopathol.* **159**, 28–34.

- Bayraktar, H., Türkkan, M. and Dolar, F.S. (2010) Characterization of *Fusarium oxysporum* f.sp. *cepae* from onion in Turkey based on vegetative compatibility and rDNA RFLP analysis. *J. Phytopathol.* **158**, 691–697.
- Boratyn, G.M., Camacho, C., Cooper, P.S., Coulouris, G., Fong, A., Ma, N., Madden, T.L., Matten, W.T., McGinnis, S.D., Merezuk, Y., Raytselis, Y., Sayers, E.W., Tao, T., Ye, J. and Zaretskaya, I. (2013) BLAST: a more efficient report with usability improvements. *Nucleic Acids Res.* **41**, W29–W33.
- Brayford, D. (1996) IMI descriptions of fungi and bacteria set 127. *Mycopathologia*, **133**, 35–63.
- Broad Institute/MIT. (2007) *Fusarium* Comparative Sequencing Project. Available at: <http://www.broadinstitute.org/>. Accessed March 2015.
- Chakrabarti, A., Rep, M., Wang, B., Ashton, A., Dodds, P. and Ellis, J. (2011) Variation in potential effector genes distinguishing Australian and non-Australian isolates of the cotton wilt pathogen *Fusarium oxysporum* f.sp. *vasinfectum*. *Plant Pathol.* **60**, 232–243.
- Covey, P.A., Kuwitzky, B., Hanson, M. and Webb, K.M. (2014) Multilocus analysis using putative fungal effectors to describe a population of *Fusarium oxysporum* from sugar beet. *Phytopathology*, **104**, 886–896.
- Cramer, C. (2000) Breeding and genetics of *Fusarium* basal rot resistance in onion. *Euphytica*, **115**, 159–166.
- Dean, R., Van Kan, J.A., Pretorius, Z.A., Hammond-Kosack, K.E., Di Pietro, A., Spanu, P.D., Rudd, J.J., Dickman, M., Kahmann, R., Ellis, J. and Foster, G.D. (2012) The Top 10 fungal pathogens in molecular plant pathology. *Mol. Plant Pathol.* **13**, 414–430.
- Dissanayake, M., Kashima, R., Tanaka, S. and Ito, S.-I. (2009a) Genetic diversity and pathogenicity of *Fusarium oxysporum* isolated from wilted Welsh onion in Japan. *J. Gen. Plant Pathol.* **75**, 125–130.
- Dissanayake, M., Kashima, R., Tanaka, S. and Ito, S.-I. (2009b) Pathogenic variation and molecular characterization of *Fusarium* species isolated from wilted Welsh onion in Japan. *J. Gen. Plant Pathol.* **75**, 37–45.
- du Toit, L.J., Inglis, D.A. and Pelter, G.Q. (2003) *Fusarium proliferatum* pathogenic on onion bulbs in Washington. *Plant Dis.* **87**, 750–750.
- Entwistle, A.R. (1990) Root diseases. In: *Onions and Allied Crops* (Brewster, J.L., ed.), pp. 103–154. Boca Raton, FL: CRC Press.
- FAOSTAT. (2012) Food and Agricultural Organization of the United Nations – Production Statistics. Available at: <http://faostat3.fao.org/home/E>. Accessed March 2015.
- Felsenstein, J. (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, **39**, 783–791.
- Fraser-Smith, S., Czişlowski, E., Meldrum, R.A., Zander, M., O'Neill, W., Balali, G.R. and Aitken, E.A.B. (2014) Sequence variation in the putative effector gene SIX8 facilitates molecular differentiation of *Fusarium oxysporum* f. sp. *cubense*. *Plant Pathol.* **63**, 1044–1052.
- Galván, G., Koning-Boucoiran, C., Koopman, W., Burger-Meijer, K., González, P., Waalwijk, C., Kik, C. and Scholten, O. (2008) Genetic variation among *Fusarium* isolates from onion, and resistance to *Fusarium* basal rot in related *Allium* species. *Eur. J. Plant Pathol.* **121**, 499–512.
- Gawehns, F., Houterman, P.M., Ichou, F.A., Michielse, C.B., Hijdra, M., Cornelissen, B.J., Rep, M. and Takken, F.L. (2014) The *Fusarium oxysporum* effector *Six6* contributes to virulence and suppresses I-2-mediated cell death. *Mol. Plant-Microbe Interact.* **27**, 336–348.
- Ghanbarzadeh, B., Mohammadi Goltapeh, E. and Safaie, N. (2013) Identification of *Fusarium* species causing basal rot of onion in East Azarbaijan province, Iran and evaluation of their virulence on onion bulbs and seedlings. *Arch. Phytopathol. Plant Protect.* **47**, 1050–1062.
- Houterman, P.M., Speijer, D., Dekker, H.L., De Koster, C.G., Cornelissen, B.J.C. and Rep, M. (2007) The mixed xylem sap proteome of *Fusarium oxysporum*-infected tomato plants. *Mol. Plant Pathol.* **8**, 215–221.
- Houterman, P.M., Cornelissen, B.J.C. and Rep, M. (2008) Suppression of plant resistance gene-based immunity by a fungal effector. *PLoS Pathog.* **4**, e1000061.
- Houterman, P.M., Ma, L., Van Ooijen, G., De Vroomen, M.J., Cornelissen, B.J.C., Takken, F.L.W. and Rep, M. (2009) The effector protein Avr2 of the xylem-colonizing fungus *Fusarium oxysporum* activates the tomato resistance protein I-2 intracellularly. *Plant J.* **58**, 970–978.
- Jiang, R.H.Y., Tripathy, S., Govers, F. and Tyler, B.M. (2008) RXLR effector reservoir in two *Phytophthora* species is dominated by a single rapidly evolving superfamily with more than 700 members. *Proc. Natl. Acad. Sci. USA* **105**, 4874–4879.
- Jukes, T.H. and Cantor, C.R. (1969) Evolution of protein molecules. In: *Mammalian Protein Metabolism* (Munro, H.N., ed.), pp. 21–132. New York: Academic Press.
- Kale, S.D. (2012) Oomycete and fungal effector entry, a microbial Trojan horse. *New Phytol.* **193**, 874–881.
- Kale, S.D., Gu, B., Capelluto, D.G.S., Dou, D., Feldman, E., Rumore, A., Arredondo, F.D., Hanlon, R., Fudal, I., Rouxel, T., Lawrence, C.B., Shan, W. and Tyler, B.M. (2010) External lipid PI3P mediates entry of eukaryotic pathogen effectors into plant and animal host cells. *Cell*, **142**, 284–295.
- Kashiwa, T., Inami, K., Fujinaga, M., Ogiso, H., Yoshida, T., Teraoka, T. and Arie, T. (2013) An avirulence gene homologue in the tomato wilt fungus *Fusarium oxysporum* f. sp. *lycopersici* race 1 functions as a virulence gene in the cabbage yellows fungus *F. oxysporum* f. sp. *conglutinans*. *J. Gen. Plant Pathol.* **79**, 412–421.
- Kehr, A.E., O'Brien, J. and Davis, E.W. (1962) Pathogenicity of *Fusarium oxysporum* f. sp. *cepae* and its interaction with *Pyrenochaeta terrestris* on onion. *Euphytica*, **11**, 197–208.
- Keller, O., Kollmar, M., Stanke, M. and Waack, S. (2011) A novel hybrid gene prediction method employing protein multiple sequence alignments. *Bioinformatics*, **27**, 757–763.
- Kimura, M. (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J. Mol. Evol.* **16**, 111–120.
- Laurence, M.H., Summerell, B.A. and Liew, E.C.Y. (2015) *Fusarium oxysporum* f. sp. *canariensis*: evidence for horizontal gene transfer of putative pathogenicity genes. *Plant Pathol.* **64**, 1068–1075.
- Leslie, J.F. and Summerell, B.A. (2006) *The Fusarium Laboratory Manual*. Oxford: Blackwell Publishing.
- Lieven, B., Houterman, P.M. and Rep, M. (2009) Effector gene screening allows unambiguous identification of *Fusarium oxysporum* f. sp. *lycopersici* races and discrimination from other formae speciales. *FEMS Microbiol. Lett.* **300**, 201–215.
- Lysøe, E., Harris, L.J., Walkowiak, S., Subramanian, R., Divo, H.H., Riiser, E.S., Llorens, C., Gabaldon, T., Kistler, H.C., Jonkers, W., Kolseth, A.K., Nielsen, K.F., Thrane, U. and Frandsen, R.J. (2014) The genome of the generalist plant pathogen *Fusarium avenaceum* is enriched with genes involved in redox, signaling and secondary metabolism. *PLoS One*, **9**, e112703.
- Ma, L.-J., van der Does, H.C., Borkovich, K.A., Coleman, J.J., Daboussi, M.-J., Di Pietro, A., Dufresne, M., Freitag, M., Grabherr, M., Henrissat, B., Houterman, P.M., Kang, S., Shim, W.-B., Woloshuk, C., Xie, X., Xu, J.-R., Antoniw, J., Baker, S.E., Bluhm, B.H., Breakspear, A., Brown, D.W., Butchko, R.A.E., Chapman, S., Coulson, R., Coutinho, P.M., Danchin, E.G.J., Diener, A., Gale, L.R., Gardiner, D.M., Goff, S., Hammond-Kosack, K.E., Hilburn, K., Hua-Van, A., Jonkers, W., Kazan, K., Kodira, C.D., Koehrsen, M., Kumar, L., Lee, Y.-H., Li, L., Manners, J.M., Miranda-Saavedra, D., Mukherjee, M., Park, G., Park, J., Park, S.-Y., Proctor, R.H., Regev, A., Ruiz-Roldan, M.C., Sain, D., Sakthikumar, S., Sykes, S., Schwartz, D.C., Turgeon, B.G., Wapinski, I., Yoder, O., Young, S., Zeng, Q., Zhou, S., Galagan, J., Cuomo, C.A., Kistler, H.C. and Rep, M. (2010) Comparative genomics reveals mobile pathogenicity chromosomes in *Fusarium*. *Nature*, **464**, 367–373.
- Ma, L., Cornelissen, B.J.C. and Takken, F.L.W. (2013) A nuclear localization for Avr2 from *Fusarium oxysporum* is required to activate the tomato resistance protein I-2. *Front. Plant Sci.* **4**, 94.
- Ma, L., Houterman, P.M., Gawehns, F., Cao, L., Sillo, F., Richter, H., Clavijo-Ortiz, M.J., Schmidt, S.M., Boeren, S., Vervoort, J., Cornelissen, B.J., Rep, M. and Takken, F.L. (2015) The AVR2-SIX5 gene pair is required to activate I-2-mediated immunity in tomato. *New Phytol.* **208**, 507–518.
- Meldrum, R.A., Fraser-Smith, S., Tran-Nguyen, L.T.T., Daly, A.M. and Aitken, E.A.B. (2012) Presence of putative pathogenicity genes in isolates of *Fusarium oxysporum* f. sp. *cubense* from Australia. *Australas. Plant Pathol.* **41**, 551–557.
- Michielse, C.B. and Rep, M. (2009) Pathogen profile update: *Fusarium oxysporum*. *Mol. Plant Pathol.* **10**, 311–324.
- O'Donnell, K., Kistler, H.C., Cigelnik, E. and Ploetz, R.C. (1998) Multiple evolutionary origins of the fungus causing Panama disease of banana: concordant evidence from nuclear and mitochondrial gene genealogies. *Proc. Natl. Acad. Sci. USA*, **95**, 2044–2049.
- O'Donnell, K. and Cigelnik, E. (1997) Two divergent dnrtragenomic rDNA ITS2 types within a monophyletic lineage of the fungus *Fusarium* are nonorthologous. *Molecular Phylogenetics and Evolution* **7**, 103–116.
- O'Donnell, K., Sarver, B.A., Brandt, M., Chang, D.C., Noble-Wang, J., Park, B.J., Sutton, D.A., Benjamin, L., Lindsley, M., Padhye, A., Geiser, D.M. and Ward, T.J. (2007) Phylogenetic diversity and microsphere array-based genotyping of human pathogenic *Fusaria*, including isolates from the multistate contact lens-associated U.S. keratitis outbreaks of 2005 and 2006. *J. Clin. Microbiol.* **45**, 2235–248.
- Petersen, T.N., Brunak, S., von Heijne, G. and Nielsen, H. (2011) SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat. Methods*, **8**, 785–786.

- Rehmany, A.P., Gordon, A., Rose, L.E., Allen, R.L., Armstrong, M.R., Whisson, S.C., Kamoun, S., Tyler, B.M., Birch, P.R.J. and Beynon, J.L. (2005) Differential recognition of highly divergent downy mildew avirulence gene alleles by RPP1 resistance genes from two *Arabidopsis* lines. *Plant Cell*, **17**, 1839–1850.
- Rep, M. (2005) Small proteins of plant-pathogenic fungi secreted during host colonization. *FEMS Microbiol. Lett.* **253**, 19–27.
- Rep, M., Van Der Does, H.C., Meijer, M., Van Wijk, R., Houterman, P.M., Dekker, H.L., De Koster, C.G. and Cornelissen, B.J.C. (2004) A small, cysteine-rich protein secreted by *Fusarium oxysporum* during colonization of xylem vessels is required for I-3-mediated resistance in tomato. *Mol. Microbiol.* **53**, 1373–1383.
- Rout, E., Nanda, S., Nayak, S. and Joshi, R.K. (2014) Molecular characterization of NBS encoding resistance genes and induction analysis of a putative candidate gene linked to *Fusarium* basal rot resistance in *Allium sativum*. *Physiol. Mol. Plant Pathol.* **85**, 15–24.
- Sasaki, K., Nakahara, K., Shigyo, M., Tanaka, S. and Ito, S.-i. (2015a) Detection and quantification of onion isolates of *Fusarium oxysporum* f. sp. *cepae* in onion plant. *J. Gen. Plant Pathol.* **81**, 232–236.
- Sasaki, K., Nakahara, K., Tanaka, S., Shigyo, M. and Ito, S.I. (2015b) Genetic and pathogenic variability of *Fusarium oxysporum* f. sp. *cepae* isolated from onion and Welsh onion in Japan. *Phytopathology*, **105**, 525–532.
- Schmidt, S.M., Houterman, P.M., Schreiver, I., Ma, L., Amyotte, S., Chellappan, B., Boeren, S., Takken, F.L. and Rep, M. (2013) MITEs in the promoters of effector genes allow prediction of novel virulence genes in *Fusarium oxysporum*. *BMC Genomics*, **14**, 119.
- Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods*, **9**, 671–675.
- Shimura, A. (2002) Studies on the ecology and control of Welsh onion root rot caused by *Fusarium redolens*. *J. Gen. Plant Pathol.* **68**, 265.
- Southwood, M.J., Viljoen, A., Mostert, G. and McLeod, A. (2012a) Molecular identification of two vegetative compatibility groups of *Fusarium oxysporum* f. sp. *cepae*. *Phytopathology*, **102**, 204–213.
- Southwood, M.J., Viljoen, A., Mostert, L., Rose, L.J. and McLeod, A. (2012b) Phylogenetic and biological characterization of *Fusarium oxysporum* isolates associated with onion in South Africa. *Plant Dis.* **96**, 1250–1261.
- Stanke, M. and Morgenstern, B. (2005) AUGUSTUS: a web server for gene prediction in eukaryotes that allows user-defined constraints. *Nucleic Acids Res.* **33**, W465–W467.
- Stankovic, S., Levic, J., Petrovic, T., Logrieco, A. and Moretti, A. (2007) Pathogenicity and mycotoxin production by *Fusarium proliferatum* isolated from onion and garlic in Serbia. *Eur. J. Plant Pathol.* **118**, 165–172.
- Takken, F. and Rep, M. (2010) The arms race between tomato and *Fusarium oxysporum*. *Mol. Plant Pathol.* **11**, 309–314.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731–2739.
- Taylor, A., Vagany, V., Barbara, D.J., Thomas, B., Pink, D.A.C., Jones, J.E. and Clarkson, J.P. (2013) Identification of differential resistance to six *Fusarium oxysporum* f. sp. *cepae* isolates in commercial onion cultivars through the development of a rapid seedling assay. *Plant Pathol.* **62**, 103–111.
- Thatcher, L.F., Gardiner, D.M., Kazan, K. and Manners, J.M. (2012) A highly conserved effector in *Fusarium oxysporum* is required for full virulence on *Arabidopsis*. *Mol. Plant–Microbe Interact.* **25**, 180–190.
- Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673–4680.
- Vágány, V. (2012) Characterisation of *Fusarium* pathogens in the UK. PhD Thesis. University of Warwick, Warwick: School of Life Sciences.
- Welham, S.J. and Thompson, R. (1997) Likelihood Ratio Tests for Fixed Model Terms using Residual Maximum Likelihood. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **59**, 701–714.
- Yamazaki, M., Morita, Y., Kashiwa, T., Teraoka, T. and Arie, T. (2013) *Fusarium proliferatum*, an additional bulb rot pathogen of Chinese chive. *J. Gen. Plant Pathol.* **79**, 431–434.
- Zerbino, D.R. and Birney, E. (2008) Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* **18**, 821–829.

## SUPPORTING INFORMATION

Additional Supporting information may be found in the online version of this article at the publisher's website:

**Fig. S1** Partial nucleotide alignment of the *SIX5* gene from *Fusarium oxysporum* f. spp. *cepae* and *lycopersici* (isolate MN25, Broad Institute *Fusarium* database). Shaded bases differ from the predominant sequence type.

**Methods S1** Protocol for analysing expression of putative effector genes *in planta*.