

Original citation:

Newbigin, Edward, Rayson, Samantha, Arciga-Reyes, Luis, Wootton, Lucie, De Torres Zabala, Marta, Truman, William, Graham, Neil, Grant, Murray and Davies, Brendan. (2012) A role for nonsense-mediated mRNA decay in plants : pathogen responses are induced in Arabidopsis thaliana NMD mutants. PLoS One, 7 (2). e31917.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/78300>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 3.0 (CC BY 3.0) license and may be reused according to the conditions of the license. For more details see:

<http://creativecommons.org/licenses/by/3.0/>

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

A Role for Nonsense-Mediated mRNA Decay in Plants: Pathogen Responses Are Induced in *Arabidopsis thaliana* NMD Mutants

Samantha Rayson¹, Luis Arciga-Reyes¹, Lucie Wootton¹, Marta De Torres Zabala², William Truman², Neil Graham³, Murray Grant², Brendan Davies^{1*}

1 Centre for Plant Sciences, Faculty of Biological Sciences, University of Leeds, Leeds, United Kingdom, **2** School of Biosciences, University of Exeter, Exeter, United Kingdom, **3** Nottingham Arabidopsis Stock Centre, School of Biosciences, University of Nottingham, Sutton Bonington Campus, Loughborough, United Kingdom

Abstract

Nonsense-mediated mRNA decay (NMD) is a conserved mechanism that targets aberrant mRNAs for destruction. NMD has also been found to regulate the expression of large numbers of genes in diverse organisms, although the biological role for this is unclear and few evolutionarily conserved targets have been identified. Expression analyses of three *Arabidopsis thaliana* lines deficient in NMD reveal that the vast majority of NMD-targeted transcripts are associated with response to pathogens. Congruently, NMD mutants, in which these transcripts are elevated, confer partial resistance to *Pseudomonas syringae*. These findings suggest a biological rationale for the regulation of gene expression by NMD in plants and suggest that manipulation of NMD could offer a new approach for crop protection. Amongst the few non-pathogen responsive NMD-targeted genes, one potential NMD targeted signal, the evolutionarily conserved upstream open reading frame (CuORF), was found to be hugely over-represented, raising the possibility that this feature could be used to target specific physiological mRNAs for control by NMD.

Citation: Rayson S, Arciga-Reyes L, Wootton L, De Torres Zabala M, Truman W, et al. (2012) A Role for Nonsense-Mediated mRNA Decay in Plants: Pathogen Responses Are Induced in *Arabidopsis thaliana* NMD Mutants. PLoS ONE 7(2): e31917. doi:10.1371/journal.pone.0031917

Editor: Edward Newbigin, University of Melbourne, Australia

Received: October 5, 2011; **Accepted:** January 15, 2012; **Published:** February 22, 2012

Copyright: © 2012 Rayson et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was funded by the BBSRC. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: b.h.davies@leeds.ac.uk

Introduction

Nonsense-mediated mRNA-decay (NMD) is a conserved eukaryotic system that performs two major functions. It identifies aberrant mRNA molecules carrying premature termination codons and targets them for destruction, thus preventing the accumulation of potentially harmful truncated proteins [1]. However, NMD also regulates the expression of normal transcripts; microarray analyses of cells and organisms depleted in various NMD effectors have shown that the expression levels of 1–10% of all genes are affected by NMD in yeast, flies, mammals and plants [2–6].

The mRNA features that enable the NMD machinery to identify a termination codon as premature are not yet fully defined and seem to vary between organisms. In mammals, Zebrafish (*Danio rerio*) and *Arabidopsis*, the presence of an exon junction complex (EJC) on the mRNA, downstream of a stop codon, can make it a substrate of NMD [7–9]. In *Saccharomyces cerevisiae*, Hrp1p bound to a downstream sequence element can lead to NMD of the *PGK1* transcript [10,11]. In all organisms, a large distance between a termination codon and the poly(A) tail can also target an mRNA to NMD [9,12–16]. These different mRNA features are targeted by NMD, via a complex containing NMD effector proteins.

Several effectors of NMD have been identified, including the UPFRAMESHIFT1, 2 and 3 and SMG1, 5, 6 and 7 proteins.

Mammalian NMD is dependent upon the cyclic phosphorylation and dephosphorylation of UPF1 [17]. The formation of a complex including UPF1, 2 and 3, bound to target mRNA, enables the phosphorylation of UPF1 by SMG1 [17–20]. The subsequent dephosphorylation of UPF1 is mediated by either SMG5/7 or SMG 6 [21–24]. The UPF proteins are well conserved [25] and UPF1 is indispensable for NMD in all eukaryotes studied [26,27]. In contrast, the SMG1 and SMG5-7 proteins are less well conserved. Some organisms that have a functional NMD pathway lack an SMG1 orthologue and/or homologues of one or two of the SMG5-7 proteins [28]. *Arabidopsis* homologues of *UPF1* [29,30], *UPF2* [9], *UPF3* [31] and *SMG7* [32] have been identified as NMD effectors, but there are no obvious *Arabidopsis* *SMG1*, 5 or 6 homologues.

Despite the finding that NMD regulates the levels of multiple transcripts in every system studied to date, it has been problematic to identify a biological role for such a mechanism of gene regulation. In mammals, rates of NMD can vary in a tissue-specific manner [33] and in response to hypoxia [34]. Other findings suggest an involvement of NMD in the regulation of responses to exogenous change. For example, the elevation of specific transcripts in response to reactive oxygen species in *Saccharomyces pombe* is dependent on a functional UPF1 [35] and amino acid starvation in mammalian and *Drosophila melanogaster* cells leads to the up-regulation of several NMD target mRNAs [3]. More

recently, it has been shown that NMD can be specifically suppressed in differentiating neuronal cells by the expression of a microRNA that targets *UPF1* and a component of the EJC [36]. This local suppression of NMD results in a widespread change in the gene expression profile and is compatible with a role for NMD in normal brain development.

In most model systems it is still unclear why so many genes are subject to post-transcriptional control by NMD. In plants, there are some specific examples of NMD gene regulation either by regulation of alternative splicing in favor of transcripts harboring NMD target features [37–39], or by spatial regulation of factors essential for NMD of a sub-set of *Arabidopsis* targets [40]. Taken together, these findings indicate a potential for additional roles of plant NMD in the dynamic regulation of gene expression. There is, however, currently no known biological rationale for the NMD regulation of non-aberrant transcripts in plants. *Arabidopsis* mutants, impaired in NMD, are available for *UPF1*, *UPF3* and *SMG7* [29,32]. To identify the physiological processes regulated by NMD in *Arabidopsis*, we determined the common transcripts that are enriched in *upf1-5*, *upf3-1* and *smg7-1* mutant plants. Strikingly, the vast majority of such transcripts are associated with pathogen response.

Plants represent a source of nutrients for the microorganisms to which they are exposed. Most potential invaders are kept at bay by innate defences, including both physical and chemical barriers, and inducible defences that respond to pathogen (or microbe) associated molecular patterns (PAMPs/MAMPs), such as flagellin or lipopolysaccharide [41,42]. PAMP-triggered immunity (PTI) in response to bacterial infection involves rapid host transcriptional reprogramming, callose deposition and production of reactive oxygen species [43–46]. Some pathogens have evolved avirulence factors, which confer the ability to evade host pattern recognition receptors or to otherwise suppress PTI [47]. In some cases, host plants have evolved corresponding genes, which allow them to respond to specific avirulence factors by launching the hypersensitive response, in which infected cells and tissues are sacrificed in order to restrict growth and spread of the pathogen [41]. The hypersensitive response commonly precedes systemic acquired resistance (SAR), whereby distal tissues are primed and respond more readily to pathogens. Salicylic acid (SA) is involved with resistance to biotrophic and hemi-biotrophic pathogens and the establishment of SAR [48,49]. Jasmonic acid (JA) and ethylene are associated with resistance to necrotrophic pathogens [48]. ABA, which has a role in plant responses to several abiotic stresses, is a negative regulator of defense responses [50]. Here, we show that *Arabidopsis* NMD mutants constitutively express a set of pathogen response genes, overproduce salicylic acid and consequently show partial resistance to *Pseudomonas syringae* pathovar tomato DC3000 (DC3000). This indicates a role for NMD in the prevention of inappropriate expression of this suite of genes. Furthermore, we identify a smaller subset of NMD targets that are not induced by pathogen infection but which show an overrepresentation of evolutionarily conserved upstream open reading frames. These findings suggest a biological role for NMD in pathogen response and provide evidence for differential regulation of sets of NMD targets in plants.

Results

NMD regulates the abundance of hundreds of *Arabidopsis* transcripts

To determine the effects of NMD on gene expression, microarrays were used to profile transcript abundance in soil-grown *Arabidopsis* wild-type, *upf1-5*, *upf3-1* and *smg7-1* mutant

seedlings. For each mutant line, a volcano plot was used to identify transcripts that differed in expression from wild-type significantly ($p \leq 0.05$) and by at least 1.5-fold. In the *upf1-5*, *upf3-1* and *smg7-1* mutants respectively, 570, 1213 and 756 transcripts are up-regulated (2.5–5.3% of transcripts represented on the array) (Figure 1A) and 414, 1,101 and 716 are down-regulated (1.8–4.8%) (Figure 1B). A condition tree was built using the expression data for these genes (Figure 1C). The *upf1-5* and *upf3-1* expression profiles show no significant differences from each other. The expression profile of *smg7-1*, although more similar to the profiles of the other NMD mutants than to wild-type, shows a greater degree of difference ($p \leq 0.05$), suggesting that SMG7 has additional function(s) distinct from its role in NMD. The genes that are up or down regulated in each mutant were then compared (Figure 1A and B). 206 and 132 of the transcripts (0.9 and 0.6% of transcripts represented on the array) are commonly up or down-regulated in all three NMD mutants. Of transcripts that are differentially regulated in *smg7-1* plants compared to wild-type, 54% are similarly regulated in *upf1-5* or *upf3-1* mutants. A comparable proportion of *UPF3*-regulated transcripts (47%) were commonly regulated by another NMD factor. In contrast, more than 80% of the transcripts regulated by *UPF1* also respond to another NMD factor. For both the up and down-regulated transcripts, the overlap is greater than predicted by chance ($p \leq 0.001$), as expected for independent mutants affecting the same process. Only genes that respond similarly in all three mutants were carried forward for further analyses (listed in Tables S1 and




Figure 1. Gene expression in wild-type and NMD mutant *Arabidopsis* plants. (A and B) Genes that are up-regulated and down-regulated respectively in NMD mutant plants when compared to wild-type ($p < 0.05$, fold change > 1.5). * The overlap of up or down-regulated genes is greater than would be predicted by chance ($p < 0.05$). (C) Condition tree of all genes that are differentially regulated in at least one of the NMD mutants when compared to wild-type ($p < 0.05$, fold change > 1.5). Strongly expressed genes are coloured bright yellow and weakly expressed genes are coloured dark blue. doi:10.1371/journal.pone.0031917.g001

S2). Although this conservative selection of genes will almost certainly exclude some interesting results, its use ensures that genes studied here are differentially regulated by NMD, rather than unrelated processes dependent on individual NMD factors.

There are few evolutionarily conserved targets of NMD

The 206 transcripts that are up-regulated in all three NMD mutants will be referred to as the 'common NMD transcripts'. These include both direct targets of NMD and transcripts that are up-regulated as an indirect effect of compromising the NMD process, for example as a result of the increased stability of an mRNA encoding a transcription factor. The 132 down-regulated transcripts are presumably all indirectly affected, since targets of NMD would be expected to be exclusively up-regulated in NMD-compromised plants. All of the genes that are affected by NMD, directly or indirectly, can provide valuable clues to the biological roles of NMD-regulation of physiological transcripts. By comparing the *Arabidopsis* common NMD transcripts to the lists of genes affected by NMD in other systems, we can determine the extent to which common processes are regulated by NMD across evolution.

Microarray analyses in *Drosophila*, humans and *S. cerevisiae* previously identified only two clusters of orthologous groups (KOGs) of genes that are regulated by NMD across kingdoms [4]. The 206 common NMD genes of *Arabidopsis* map to 165 KOGs [51]. As KOG databases are under constant revision and a further list of NMD-regulated genes is now available for yeast [5], KOGs were reassigned to NMD-regulated genes for each previously studied organism. Forty-two *Arabidopsis* common NMD KOGs overlapped with NMD-regulated KOGs of at least one other organism (Figure 2, listed in Table S3). The re-annotation of KOGs allowed us to identify an additional COG, COG0515 (serine/threonine protein kinase), as NMD-regulated in all four organisms.

One of the previously identified conserved targets, KOG2504 (monocarboxylate transporter) was not represented in the *Arabidopsis* genome. A manual inspection of the TAIR website [52] identified the product of *At2g39210*, which is a common NMD transcript up-regulated in all three NMD mutant *Arabidopsis* lines, as a putative monocarboxylate transporter of unknown function that is located in the plasma membrane. The human NMD-regulated KOG2504 gene is involved in accumulation of pyruvate in hypoxic cells [53], whereas both the yeast gene [54] and *At2g39210* [55] may be associated with nitrogen homeostasis. In summary, it is unclear whether KOG2504 represents a genuine conserved target of NMD, but the regulation of these genes by NMD may fit with the previously suggested conserved functions of NMD in amino acid homeostasis and the response to oxidative stress.

SMG7 belongs to the other previously identified conserved target KOG: KOG2162 (nonsense-mediated decay protein). The *SMG5/7* gene, itself part of the NMD machinery, is targeted by NMD in all organisms studied [2–6,9,32,56], implying an autoregulatory mechanism for NMD. Previously, up-regulation of *SMG7* expression was observed in *Arabidopsis* NMD mutants and in tobacco leaves in which *UPF* genes have been silenced [9]. This study finds that *SMG7* expression is up-regulated in the *upf1-5* and *upf3-1* mutant microarray experiments and this is verified by PCR and real-time PCR (Figure 3). Furthermore, treatment with cycloheximide, which blocks translational elongation and consequently NMD, results in accumulation of *SMG7* mRNA in wild-type plants (Figure 3A). To determine whether *SMG7* is a direct target of NMD, cordycepin was used to halt transcription and the rate of decay of *SMG7* mRNA in wild-type and *upf3-1* mutant seedlings was compared by real-time PCR (Figure 3B). The rate of


decay of *SMG7* mRNA is higher in wild-type plants than in *upf3-1* mutants, showing that the disruption of *UPF3* stabilises *SMG7* mRNA. This validates our microarray and provides compelling evidence that *Arabidopsis* *SMG7* is a direct target of NMD.

It has been noted that a surprising number of NMD-regulated genes in yeast are associated with the dynamics of the plasma membrane and cell wall, particularly with respect to multi-drug resistance [5]. This study identifies a number of similar KOGs that are NMD regulated in *Arabidopsis* (Figure 2 and Table S3). Eight KOGs are represented amongst NMD-regulated genes in both *Arabidopsis* and yeast: COG0697 (permeases of the drug/metabolite transporter (DMT) superfamily); COG0842 (ABC-type multidrug transport system, permease component); COG1132 (ABC-type multidrug transport system, ATPase and permease components); KOG0054 (multidrug resistance-associated protein/mitoxantrone resistance protein, ABC superfamily); KOG0065 (pleiotropic drug resistance proteins (PDR1-15), ABC superfamily), KOG0254 (predicted transporter (major facilitator superfamily)), COG0477 (permeases of the major facilitator superfamily) and COG1131 (ABC-type multidrug transport system, ATPase component). The latter two are also regulated by NMD in *Drosophila*. This suggests that many multi-drug transporters are regulated by NMD in diverse organisms.

Although evolutionarily common NMD target genes have been difficult to identify [4], there is evidence to suggest that certain biological processes are conserved under the control of NMD in different kingdoms. These processes include amino acid homeostasis [3,4,35,57] and protection from oxidative stress [34]. Transcripts encoding proteins involved in amino acid metabolism are overrepresented amongst human NMD targets [3] and present (though not overrepresented) amongst *S. cerevisiae* NMD targets [2], prompting the hypothesis that NMD has a conserved role in regulating the expression of genes involved in amino acid homeostasis. This hypothesis was further supported by similarities between the gene expression profiles of NMD impaired *Drosophila* cells and wild-type starved *Drosophila* [4]. The common NMD transcripts in *Arabidopsis* are enriched for genes involved in the catabolism of amino acids and their derivatives (Figure 4). Of the 16 common NMD genes that have GO terms implicating them in amino-acid homeostasis, ten have features that could target them to NMD; eight have upstream open reading-frames (uORFs), one has a 3'UTR in excess of 300 nt and one has both of these features. This enrichment of NMD target features may imply that this functional sub-set of NMD-regulated genes includes direct targets of NMD. Taken together, these results suggest that in *Arabidopsis*, as in other organisms, NMD is involved in amino-acid homeostasis.

Pathogen response transcripts are up-regulated in *Arabidopsis* NMD mutants

Amongst the common NMD genes, there is a dramatic overrepresentation of transcripts encoding proteins involved in responses to pathogens (Figure 4). Strikingly, all of the functional ontologies that are overrepresented with a p value of <0.01 are related to pathogen response, including defence response (31 genes), systemic acquired resistance (5 genes) and defence response, incompatible interaction (8 genes). The list includes known pathogen induced genes including the key regulators of salicylic acid (SA) mediated plant defence *EDS1*, *EDS5* (*sid1*), *ICS1* and the classical molecular markers of SA activated plant defence *PR1* and *PR5* (reviewed in [58]). Genes involved in indole biosynthesis and more specifically auxin biosynthesis are overrepresented (p<0.05). Notably, other stress-responsive transcripts, such as those involved in the responses to wounding or oxidative

A**B**

KOG I.D.	KOG Description
COG0515:	Serine/threonine protein kinase
KOG2162:	Nonsense-mediated mRNA decay protein
COG0553:	Superfamily II DNA/RNA helicases, SNF2 family
COG4886:	Leucine-rich repeat (LRR) protein
KOG0504:	FOG: Ankyrin repeat
COG0477:	Permeases of the major facilitator superfamily
COG0625:	Glutathione S-transferase
COG1131:	ABC-type multidrug transport system, ATPase component
COG2124:	Cytochrome P450
KOG0867:	Glutathione S-transferase
KOG0157:	Cytochrome P450 CYP4/CYP19/CYP26 subfamilies
COG0457:	FOG: TPR repeat
COG1601:	Translation initiation factor 2, beta subunit (eIF-2beta)/eIF-5 N-terminal domain
KOG1192:	UDP-glucuronosyl and UDP-glucosyl transferase
KOG1565:	Gelatinase A and related matrix metalloproteases
KOG2767:	Translation initiation factor 5 (eIF-5)
KOG4177:	Ankyrin
COG0484:	DnaJ-class molecular chaperone with C-terminal Zn finger domain
COG0500:	SAM-dependent methyltransferases
KOG0583:	Serine/threonine protein kinase
KOG0619:	FOG: Leucine rich repeat
KOG0714:	Molecular chaperone (DnaJ superfamily)
KOG1001:	Helicase-like transcription factor HLTf/DNA helicase RAD5, DEAD-box superfamily
KOG1052:	Glutamate-gated kainate-type ion channel receptor subunit GluR5 and related subunits
COG0443:	Molecular chaperone
COG0446:	Uncharacterized NAD(FAD)-dependent dehydrogenases
COG0524:	Sugar kinases, ribokinase family
COG0697:	Permeases of the drug/metabolite transporter (DMT) superfamily
COG0842:	ABC-type multidrug transport system, permease component
COG1132:	ABC-type multidrug transport system, ATPase and permease components
COG1404:	Subtilisin-like serine proteases
COG1657:	Squalene cyclase
COG1953:	Cytosine/uracil/thiamine/allantoin permeases
COG5147:	Myb superfamily proteins, including transcription factors and mRNA splicing factors
KOG2466:	Uridine permease/thiamine transporter/allantoin transport
KOG0054:	Multidrug resistance-associated protein/mitoxantrone resistance protein, ABC superfamily
KOG0065:	Pleiotropic drug resistance proteins (PDR1-15), ABC superfamily
KOG0254:	Predicted transporter (major facilitator superfamily)
KOG0101:	Molecular chaperones HSP70/HSC70, HSP70 superfamily
KOG0497:	Oxidosqualene-lanosterol cyclase and related proteins
KOG1336:	Monodehydroascorbate/ferredoxin reductase

Figure 2. The overlap of KOGs regulated by NMD between *Arabidopsis* and other organisms. (A) A Venn diagram of the 42 *Arabidopsis* (*At*) NMD-regulated KOGs that are also regulated by NMD in at least one of: *Drosophila* (*Dm*), *Saccharomyces cerevisiae* (Yeast, *Sc*) and Human HeLa cells (*Homo sapiens*, *Hs*). (B) The KOGs represented in the Venn diagram. KOGs are colour coded according to the organisms in which they are NMD-regulated: Blue: *Dm* and *At*, yellow: *Sc* and *At*, red: *Hs* and *At*, green: *Dm*, *Sc* and *At*, purple: *Dm*, *Hs* and *At*, orange *Sc*, *Hs* and *At* and brown: all four organisms.
doi:10.1371/journal.pone.0031917.g002

or osmotic stress, were not overrepresented. Taken together, these observations imply that these transcriptional differences are not a general stress response resulting from a deleterious mutation, but a direct result of deficient NMD leading to the up-regulation of pathogen-responses involving the SA signalling pathway. Many pathogen responses result from co-ordinate regulation by SA, JA and ethylene [59,60], yet no GO terms related to JA or ethylene were over-represented amongst the common NMD genes. This suggests that neither the impairment of NMD, nor the consequent overproduction of salicylic acid, leads to ethylene or JA-dependent responses in *Arabidopsis*.

As there was an overrepresentation of gene ontologies relating to pathogen responses amongst the common NMD genes, publicly available microarray data were mined for expression levels of all of the common NMD genes in response to pathogens, pathogen-associated molecular patterns and SA (Figure 5). The profiles fitted two broad clusters, with 183 of the 206 common NMD genes forming a group of pathogen-responsive genes and 23 common NMD genes being unresponsive (listed in Table S4). All of the pathogen-responsive genes were down-regulated in SA insensitive mutant plants, suggesting a dependency on the SA signalling pathway (Figure 5). Closer inspection of the SA and auxin biosynthetic pathways revealed that five common NMD genes are involved in the

utilisation of chorismate to produce camalexin and SA or to yield tryptophan, the precursor for the synthesis of auxin. Other genes in these pathways were also up-regulated in NMD mutants, but had not passed the rigorous selection criteria (up-regulated at least 1.5 fold at $p \leq 0.05$ in each of the three mutants, Table S5). Genes involved in auxin biosynthesis respond systemically [61] and locally to infection with *P. syringae* [43,62]. Of the 12 genes involved in auxin synthesis that are up-regulated in at least one of the NMD mutant lines, six are also associated with GO terms linked to defence responses. We therefore suspect that the overrepresentation of genes involved in auxin metabolism is part of the SA response rather than a direct result of regulation of auxin metabolism by NMD.

NMD mutants are partially resistant to *Pseudomonas syringae* DC3000

The speed with which plants respond to pathogenic attack by up-regulating inducible defence genes is a key factor in determining resistance or susceptibility [63]. Given the strong correlation to transcriptional reprogramming in response to infection with the hemibiotroph *Pseudomonas syringae* pathovar tomato DC3000 (DC3000) [43,62], we hypothesised that the constitutive up-regulation of pathogen-responsive genes observed in NMD-mutant plants may confer resistance to pathogens. *upf1-5*, *upf3-1* and *upf3-2* mutant plants were challenged with virulent DC3000, however the severe effect of the mutation on the width and shape of the leaf blade prevented us from assaying *smg7-1* plants. Significantly fewer colony-forming units were isolated from leaves of DC3000 challenged *upf1-5*, *upf3-1* and *upf3-2* mutant plants than the wild-type control 3 days post-inoculation (Figure 6). This suggests that NMD mutant plants are more resistant to DC3000.

Pathogen induced hormonal imbalances underpin many pathogen responses [64,65]. SA is primarily associated with defence against biotrophic pathogens and the activation of SAR [48,49]. ABA promotes virulence of diverse pathogens against various host species by down-regulation of *ISOCHORISMATE SYNTHASE 1* (*ICS1*) expression, which is required for SA production, and suppression of PAMP-induced transcripts (reviewed in [66]). Upon infection of *Arabidopsis* with compatible DC3000, coronatine, SA and ABA accumulate rapidly, whereas JA production is induced later in the infection process [50]. The hormone levels of wild-type and mutant plants were profiled before and during infection with DC3000. While levels of JA and abscisic acid (ABA) do not vary significantly between genotypes ($p > 0.05$ in at least 2 of triplicate experiments, Figure S1), levels of SA are significantly higher in NMD mutant plants than in wild-type, even prior to infection ($p < 0.05$ in at least 2 of triplicate experiments, Figure 7). In accordance with the reduced bacterial load, levels of the *Pseudomonad* phytotoxin coronatine [65] are lower in infected NMD mutant plants compared to wild-type (Figure 7). This suggests that the up-regulation of pathogen responsive genes in NMD mutant *Arabidopsis* plants, and their enhanced resistance to DC3000, is a result of constitutive activation of the SA synthesis pathway.

CuORFs are massively enriched amongst the NMD targets that are not pathogen responsive

Several gene features have been associated with NMD substrates in a range of different model systems. In plants, these




Figure 3. *SMG7* mRNA is stabilised in NMD mutant plants. (A) Steady state levels of *SMG7* mRNA in wild-type, *upf1-5* and *upf3-1* mutant plants and plants treated with cycloheximide to disrupt NMD, determined by reverse transcriptase PCR. (B) Degradation of *SMG7* mRNA in wild-type and *upf3-1* mutant plants, determined using real-time qPCR. The error bars represent the standard error of the mean of three technical replicates. The results of two independent treatments are shown. Green bars represent wild-type plants. Blue bars represent *upf3-1* mutant plants.
doi:10.1371/journal.pone.0031917.g003




Figure 4. Biological function of NMD-regulated genes. Biological function gene ontology (GO) terms that are over-represented ($p < 0.05$) amongst the 206 genes that are commonly up-regulated in NMD mutant plants. The horizontal bar indicates the proportion of the 206 NMD genes with the given GO term relative to the proportion of the genome with the same GO term. DRP is defence response to pathogen, SA is salicylic acid and SAR is systemic acquired resistance. GO terms relating to response to stimuli are coloured blue. GO terms relating to auxin metabolism are coloured peach. The remainder are all associated with amino acid catabolism. * p value ≤ 0.01 . ** p value ≤ 0.001 . doi:10.1371/journal.pone.0031917.g004




Figure 5. The expression of the 206 'common NMD genes' in response to biotic stimuli. Green indicates that genes are down-regulated, red indicates up-regulation and black indicates that there was no change. The box surrounds genes that were not pathogen-responsive. doi:10.1371/journal.pone.0031917.g005

include upstream open reading frames (uORFs) [67], UTRs harbouring an intron and 3'UTRs in excess of 300 nt [68]. In common with most NMD target gene analyses in other systems [2–4], there was no convincing enrichment of any of these features amongst the common NMD genes in *Arabidopsis* (Table S6). For example, 34% of the 206 common NMD genes encode transcripts with an upstream open reading frame (uORF), in comparison to 30% of all *Arabidopsis* genes [69]. Limiting these analyses to the 23 non-pathogen responsive genes allows us to exclude the effects of activation of the SA-mediated defence response in up-regulating genes that are not directly targeted by NMD. Amongst these 23 genes, uORFs and 5'UTR introns are over-represented by more than two-fold. Surprisingly, one feature was dramatically over-represented amongst a sub-set of the common NMD genes. Conserved peptide uORFs (CuORFs) are uORFs encoding evolutionarily conserved amino acid sequences. Forty-four *Arabidopsis* genes with CuORFs have been previously identified by comparisons of *Arabidopsis* and rice full length cDNAs [70]. Transcripts with CuORFs are 21-fold over-represented amongst common NMD genes (Table S4, 8 transcripts, $p < 0.0001$). In contrast, no CuORF-containing genes were commonly down-regulated in the NMD mutants. However, when the NMD-regulated CuORFs were mapped onto the cluster analysis, it became clear that all of them belong to the group of 23 common NMD genes that do not respond to pathogens, representing a 190-fold enrichment for this feature.




Figure 6. Reduced growth of *Pseudomonas syringae* DC3000 in NMD mutants. Growth of *Pseudomonas syringae* DC3000 in wild-type and NMD mutant *Arabidopsis* leaves. Numbers of colony forming units were counted 3 days post infection. Bacterial counts are on a \log_{10} scale. Results from independent replicate experiments are shown. doi:10.1371/journal.pone.0031917.g006




Figure 7. Levels of salicylic acid and coronatine in NMD mutants. Salicylic acid (SA) content and coronatine content of wild-type and NMD mutant plants. The salicylic acid and coronatine levels of leaves were measured prior to inoculation (NI) and 10 or 20 hours post-infection with *Pseudomonas syringae* DC3000 (DC10 and DC20 respectively). Wild-type plants are represented by blue bars, *upf1-5* mutants by red bars, *upf3-1* by green bars and *upf3-2* plants by purple bars. doi:10.1371/journal.pone.0031917.g007

Discussion

NMD influences the expression of many *Arabidopsis* genes

The 338 *Arabidopsis* genes (1.6% of transcripts represented on the array) that are either up- or down-regulated in all three mutant backgrounds tested represent a very conservative list of transcripts that are directly or indirectly controlled by NMD in wild type *Arabidopsis* plants. The fact that the NMD mutants used are unlikely to be null, the stringency of the microarray analyses and the use of a developmentally restricted stage, suggests that many more *Arabidopsis* transcripts are affected by NMD. Similar results have been reported in *S. cerevisiae* [2,5], humans [3], *Drosophila* [4] and *Caenorhabditis elegans* [71], where estimates of the proportion of physiological mRNA species affected by NMD range from 3% to greater than 10%. It therefore seems in plants, as in other systems, that the abundance of large numbers of 'normal' physiological transcripts is influenced positively or negatively by NMD. However, it is still not clear what proportion of the transcripts influenced by NMD represent direct targets of the process. A microarray mRNA stability analysis suggested that as many as 45% of the *S. cerevisiae* mRNAs regulated by NMD represent direct targets [5], whereas other analyses in human cell cultures have suggested that the direct targets of NMD represent a smaller proportion of the overall numbers of affected transcripts [72,73]. Further work will be required to identify the true proportion of direct NMD target transcripts in different systems and such analyses should also help to define the typical features that identify direct targets of NMD amongst physiological transcripts.

Our microarray results are compatible with previous analyses of gene expression in other alleles, using different platforms and under different growth conditions [6,74], with overlaps of 29% and 52% respectively. The significant overlaps between different experimental platforms and growth conditions, as well as between different mutant alleles, validate the microarrays and confirm the identification of large numbers of NMD-regulated transcripts in *Arabidopsis*.

Evolutionary conservation of NMD target genes

It has previously been noted that few orthologous genes are commonly regulated by NMD across diverse organisms [4]. The fresh analyses of NMD-regulated genes from *Drosophila*, yeast and human cells, coupled with the *Arabidopsis* NMD-regulated KOGs supports this, with only two KOGs commonly up-regulated across all four organisms. Homologues of the *SMG5-7* genes, represented in KOG2162 (NMD protein), are direct targets of NMD in all organisms tested [2–6,32,56]. It appears that the auto-regulation of these genes is a conserved mechanism for the control of NMD. COG0515 (serine/threonine protein kinase) was also regulated by NMD in all four organisms.

Multidrug resistance may represent another common process affected by NMD. Genes involved in multidrug transport are suppressed by NMD in yeast, *Arabidopsis* and *Drosophila*. Although this study did not identify KOGs containing multidrug transporters amongst the human NMD targets, there is a known example. MRP4 is a mammalian cellular efflux pump. An MRP4 cDNA identified from humans contains two additional exons, which introduce a premature termination codon. These exons are highly conserved across humans, rodents and monkeys and appear to target the mRNAs that harbour them to NMD [75]. Involvement of NMD in the regulation of multi-drug resistance is intriguing, as transporters tend to be expressed specifically in response to their substrates. This implies that the regulation of such genes by NMD would need to be both highly specific and adaptive.

Molecular chaperones were well represented amongst KOGs that are NMD-sensitive in *Arabidopsis* and at least one other organism: COG0443 (yeast, molecular chaperone); KOG0101 (yeast, molecular chaperones HSP70/HSC70, HSP70 superfamily) and KOG0714 (human, molecular chaperone (DnaJ superfamily)). Abrogation of NMD is expected to result in an increased number of mRNAs encoding truncated proteins. If translated, these may induce the unfolded protein response and account for the up-regulation of genes encoding molecular chaperones.

KOGs up-regulated in both *Arabidopsis* and *Drosophila* include COG1601 (translation initiation factor 2, beta subunit (eIF-2beta)/eIF-5 N-terminal domain) and KOG2767 (translation initiation factor 5 (eIF-5)). In human cells, phosphorylation of eIF-2 results in the localisation of UPF1/RENT1 in stress bodies, which inhibits NMD [34]. EIF-5 is involved in recycling of eIF-2 to its active form [76]. Perhaps the suppression of *eIF-2* and *eIF-5* expression by NMD represents another mechanism, in addition to regulation of *SMG5-7* homologues and phosphorylation state of eIF-2alpha, by which NMD is auto-regulated.

NMD regulates gene expression in all organisms tested [2–4,6,71] and it is, therefore, surprising that there appear to be so few evolutionarily conserved targets of this ancient mechanism. It is possible that roles for NMD in addition to transcript quality control, evolved independently in diverse organisms. It seems likely however that the use of different experimental techniques and growth conditions, as well as conservative lists of potential NMD targets and limited KOG annotations, have impeded the identification of conserved targets.

A biological rationale for NMD in plants

NMD-mutant *Arabidopsis* plants constitutively express pathogen-responsive genes and have higher levels of SA than wild-type plants, even in the absence of pathogens. This impairs the virulence of DC3000 on NMD-mutant plants. The possibility remains that the SA pathway is activated in response to the general stress caused by the accumulation of erroneous transcripts, however the specificity of the responsive genes suggests that this is highly unlikely. The phenotype of the NMD mutants mimics that of other *Arabidopsis* lines that overexpress pathogen responsive genes, including plants in which *R* genes are constitutively active [77,78]. As activation of numerous *R* genes causes SA accumulation and numerous *R* genes are up-regulated in response to SA [79,80], it is difficult to conclude whether individual NMD repressed genes are direct or indirect targets of NMD. To understand this further requires the identification of transcripts that are direct targets of plant NMD.

The *R* gene *SNCI* (*SUPPRESSOR OF NPR1-1*, *CONSTITUTIVE 1*) was previously shown to be up-regulated in non-inoculated *upf1-5* and *upf3-1* plants [78]. The abundance of different splice variants, relative to each other, does not appear to vary between NMD mutant and wild-type plants [78]. Unless there is another, previously undetected, splice variant of *SNCI*, this suggests that *SNCI* is unlikely to be a direct target of NMD. Our arrays found a small but reproducible up-regulation of *SNCI* in *smg7-1* plants (fold change = 1.4, $p \leq 0.05$), but not in *upf1-5* or *upf3-1*. As *SNCI* is regulated by a positive amplification loop involving SA, up-regulation of *SNCI* may be a consequence of, rather than the initial stimulus for, the accumulation of SA in NMD mutant plants.

The observation that pathogen responses are de-repressed in NMD mutant plants raises the question of whether this results from a housekeeping function of NMD, preventing the erroneous expression of aberrant but still functional *R* genes, or whether NMD is involved in the response to pathogens. *R* genes are

extensively alternatively spliced and regulation of *R* gene splicing is critical for plant immunity [81–83]. It is probable that NMD targets splice variants that contain premature termination codons. If *R* gene splice variants containing premature termination codons are created in the absence of pathogens, these transcripts could accumulate in NMD mutant plants, leading to mis-expression of *R* genes. The generation of *R* gene splice variants that are targeted by NMD could also provide an opportunity for NMD-mediated regulation of plant-pathogen responses. Levels of NMD activity could change in response to pathogens, perhaps by sequestering one or more NMD factors as in mammalian cells under oxidative stress, or the production of transcripts could shift in favour of splice variants with or without NMD targeted features. This would provide a mechanism by which pathogen-responsive NMD targets could be removed from the sphere of influence of NMD upon infection. There is a precedent for the temporal and spatial regulation of NMD; in mice a brain-specific microRNA, miR-128, represses NMD in differentiating neuronal cells and during brain development leading to the up-regulation of NMD targeted genes that are important for normal neuronal development and function [36]. Global studies of the impact of NMD on relative rates of mRNA decay before and after infection are required to answer these important questions.

Amino-acid homeostasis has previously been suggested as a potential conserved role of NMD [3,4]. In *Arabidopsis*, there is a known link between the regulation of genes involved in amino acid homeostasis and the pathogen response [84,85]. Indeed, of the 16 common NMD genes that are implicated in amino acid homeostasis, 15 are in the pathogen-responsive clade of Figure 5 and seven have GO terms directly related to pathogen responses. It is possible that the role of NMD in regulation of pathogen responses has its evolutionary origin in the NMD-mediated control of homeostasis.

As sessile organisms, plants are arguably more dependent on rapid changes in gene expression in response to biotic and abiotic stressors. This is reflected in the aggressive nature of the hypersensitive response, in which localised programmed cell death restricts pathogen ingress (reviewed in [86]). The tight regulation of such responses is paramount, as the need to thwart pathogens is countered by the detrimental effects of inappropriate expression of pathogen-responsive genes [87]. Indeed, it is possible that many of the deleterious effects of NMD deficiency in *Arabidopsis* are caused by the unwarranted pathogen response. Constitutive expression of pathogen responsive genes often results in reduced fitness [87–89]. Reported phenotypes of *Arabidopsis* NMD mutants include spontaneous necrosis [32], small or narrow leaves that are slightly indented or jagged and impaired flowering [29]. Numerous other *Arabidopsis* mutants that overexpress SA dependent defence responses display similar spontaneous necrotic lesions [90–95] and effects on leaf size [96] and shape [97] have also been identified. SA is also involved in the regulation of flowering [89]. It is tempting to speculate that NMD-mediated repression of pathogen responsive processes evolved to prevent inappropriate expression of genes that have significant fitness costs. Co-ordinate repression of pathogen responses by NMD may also provide the plant with a rapid mechanism to induce pathogen responses by the stabilisation of existing transcripts.

CuORFs are targets of NMD in *Arabidopsis*

It has long been known that the presence of uORFs in naturally-occurring transcripts can target them to NMD [98]. The identification of candidate direct NMD targets in yeast indicated that around 35% could be attributed to the presence of an uORF [5], whereas amongst 17 human direct NMD targets identified

using a proteomic approach, 11 contained an uORF [73]. However, uORFs appear to be more common in the genome than direct targets of NMD, meaning that not all uORFs act as targets for NMD. Indeed, amongst our common NMD genes we did not observe a significant enrichment of uORF-containing transcripts, probably because this list contains a mixture of direct and indirect NMD targets, complicating the identification of specific NMD target signals. Similarly, only a subset of uORF-containing transcripts in *C. elegans* is elevated in an NMD mutant background [71] and no clear defining feature of the uORFs associated with NMD-targeted transcripts could be identified amongst the 11 human NMD target genes [73].

Although uORFs in general did not stand out from the background as a putative NMD targeted feature in our microarray analysis, one particular subset of uORFs is enhanced in the common NMD genes. Evolutionarily conserved uORFs (CuORFs) have been described following a comparison of cDNAs from *Arabidopsis* and rice [70]. Forty-four such genes, belonging to 19 homology groups, were identified from a comparison of *Arabidopsis* and rice cDNA sequences. Eight of these CuORFs are present in the common NMD gene list (8/206 belonging to homology groups 1, 4, 7, 10, 13, 15 and 17 [70]) whereas none are present in the list of transcripts down-regulated in all three mutants (0/132). Even more strikingly, all 8 of the CuORFs found in the common NMD list belong to the sub-group of 23 NMD target genes that are not responsive to pathogens (Table S4), representing an almost 200-fold enrichment. This finding strongly suggests that CuORFs are substrates of NMD in *Arabidopsis* and, given their levels of conservation, potentially in other plant species.

It is not obvious why CuORFs should differ from other uORFs in their ability to target a transcript to NMD. In tobacco, the observation that an uORF of 153 nt was targeted by NMD, whereas those of 108 or less did not, led to the hypothesis that plant NMD targets uORFs that are translatable and encode a peptide of at least 53 amino acids; 331 *Arabidopsis* genes satisfy these criteria [67]. Of the eight NMD responsive CuORF-containing genes identified here, two have CuORFs that encode peptides of less than 35 amino acids, five have CuORFs that encode peptides of 41–52 amino acids and one has a CuORF that encodes a peptide of 57 amino acids. It is unlikely, therefore, that uORF length alone accounts for the overrepresentation of CuORFs amongst the NMD targets. Re-initiation of translation downstream of a premature termination codon can protect an mRNA from NMD [99]. An intriguing possibility is that the conserved peptides produced by these transcripts act, potentially in a condition-dependent fashion, either to decrease the efficiency of the re-initiation of translation at the major (downstream) AUG, or stall the ribosome at the CuORF, thereby activating the NMD pathway to degrade the mRNA. The latter mechanism has already been described in *S. cerevisiae* for *CPAI*, which is targeted by NMD in response to arginine. The uORF of *CPAI* encodes arginine attenuator peptide (AAP). In the presence of arginine, ribosomes are stalled at the uORF. The resultant increased association of ribosomes with the uORF and decreased association of ribosomes with the major ORF targets the transcript to NMD [57,100]. Either of these mechanisms would allow a constant rate of NMD to play a role in the control of individual genes by effectively using the conserved peptides to move specific transcripts into and out of the sphere of influence of NMD. Most of the CuORF genes in the common NMD list are currently uncharacterised, but one of them, *SUPPRESSOR OF ACAULIS51 (SAC51)*, encodes a bHLH transcription factor that has been shown to regulate the *ACL5* gene, which encodes thermospermine synthase [101,102]. Consistent with the role of NMD in suppressing *SAC51* transcript levels using

a CuORF, the *sac* mutant, in which *SAC* expression is elevated, is caused by a point mutation in the CuORF [103]. The group of CuORF-containing transcripts forms a candidate list for individual genes regulated by NMD under diverse conditions and it will be interesting to see whether other CuORF-containing genes, not identified as NMD targets under our conditions, are regulated by NMD at different stages of development or under different conditions. A role for NMD in targeting transcripts with CuORFs could have regulatory effects extending far beyond the CuORF-containing genes, as the overrepresentation of transcription factors amongst CuORF-containing genes suggests that the expression of numerous genes could be affected indirectly by NMD.

In conclusion, we suggest that conserved upstream open reading frames represent specific targets of NMD and may imply a mechanism for the differential regulation of subsets of NMD targets in plants. We also present a biological rationale for the co-ordinate regulation of non-aberrant plant genes by NMD. We show that NMD represses expression of genes involved in the SA-mediated pathogen response and that lifting this repression, by abrogation of NMD, confers partial resistance to *Pseudomonas syringae* pathover tomato DC3000. This raises the possibility that plants may regulate NMD as a means to coordinate pathogen responses. *Pseudomonas syringae* causes a range of diseases to numerous agronomically important crops [100]. Although we have not exposed the NMD mutants to other pathogens, it seems likely that the constitutive up-regulation of pathogen response genes observed may also confer resistance to numerous other pathogens. Perhaps in the future an understanding of the interplay between plant pathogen responses and NMD will facilitate the development of novel approaches in crop protection.

Materials and Methods

Plant materials and growth conditions

Transgenic plants were obtained from the Nottingham *Arabidopsis* Stock Centre (NASC) in the UK [104]. For microarray analyses, wild-type (ecotype Columbia-0) and T-DNA insertion lines SALK_112922 (*upf1-5*), SALK_073354 (*sng7-1*) and SALK_025175 (*upf3-1*) were grown on 3:1 SHL soil:sand (22–24°C, under constant light). For qPCR and experiments involving *Pseudomonas*, plants were grown under 8 hours light (175 $\mu\text{mol m}^{-2} \text{s}^{-1}$)/16 hours dark at 65% relative humidity in SHL compost.

Microarray hybridisation

RNA was isolated from 17 day-old seedlings. Approximately 5 μg of total RNA from each sample was used to produce cDNA using the GeneChip® One-cycle cDNA synthesis kit (Affymetrix), as per manufacturer's instructions. Double stranded cDNA products were purified using the GeneChip® Sample Cleanup Module (Affymetrix). The synthesised cDNAs were in-vitro transcribed by T7 RNA polymerase using biotinylated nucleotides to generate biotinylated complementary RNAs (cRNAs) using the GeneChip® HT IVT labeling kit (Affymetrix), as per manufacturer's instructions. The cRNAs were purified using the GeneChip® Sample Cleanup Module (Affymetrix). The cRNAs were then randomly fragmented at 94°C for 35 minutes in a buffer containing 40 mM Tris-acetate (pH 8.1), 100 mM potassium acetate, and 30 mM magnesium acetate to generate molecules of approximately 35 to 200 bp. Affymetrix *A. thaliana* ATH1-121501 GeneChip® arrays were hybridised with 15 μg of fragmented labelled cRNA for 16 h at 45°C as described in the Affymetrix Technical Analysis Manual using the GeneChip® hybridization control kit and GeneChip® hybridisation, wash and stain kit (Affymetrix). GeneChip® arrays were stained with

streptavidin-phycoerythrin solution and scanned with an Affymetrix 3000 7G GeneArray scanner. Following scanning, non-scaled RNA signal intensity (CEL) files were generated using GeneChip® operating software (GCOS; Affymetrix). All data is MIAME compliant and the raw data has been deposited in both ArrayExpress (E-GEOD-19253 and E-GEOD-32671) and GEO (GSE19253 and GSE32671). The raw data is also available from the NASCArrays database (NASCARRAYS-379 and NASCARRAYS-418, <http://affymetrix.Arabidopsis.info/>; [105]).

Microarray analyses

The non-scaled RNA CEL files were loaded into GeneSpring analysis software (GeneSpring 7.3; Agilent Technologies, USA) using the Robust Multichip Average (RMA) pre-normalisation algorithm [106]. Separate experiments were created for each mutant (*upf1-5*, *upf3-1* and *sng7-1*) using the normalised CEL files for the mutant and corresponding wild-type RNAs. Further normalisations were performed for each experiment using a three step process: (i) probe-sets with a signal value <0.01 were set to 0.01, (ii) per chip normalisation to the 50th percentile, (iii) each gene signal was normalised to the median of that gene. Putative genes with differential hybridisation intensities between a single mutant and corresponding wild-type were identified using a two-step process: (i) genes that were 1.5-fold up- or down-regulated were selected, and (ii) a Welch's t-test was performed ($p < 0.05$). A condition tree was generated for these genes using K-means clustering across conditions (*Arabidopsis* lines) in GeneSpring. In brief, hierarchical clustering was used to compare the similarity in the expression profiles of this group of genes between *Arabidopsis* lines. Genes that were differentially expressed in more than 1 mutant were identified by comparing gene lists of differentially expressed genes in a single mutant using the Venn diagram function of GeneSpring. The DAVID Gene Functional Classification Tool [107,108] was used to search for ontologies that were over-represented amongst the commonly up- or down-regulated NMD genes. Java Treeview [109] was used to profile the expression of the common NMD genes across publicly available microarray data. The STRING database [51] was used to assign KOGs to NMD regulated genes from *Arabidopsis* and other organisms. To create Figure 5 selected publicly available experiments (Table S7) were processed using Bioconductor to yield germa normalised data. Replicates were averaged and the log (base2) ratio to control treatments calculated. The arrays were left as ordered but the genes were organised by SOM and then clustered using hierarchical clustering by uncentered correlation and complete linkage.

Cycloheximide treatment

Leaves of soil-grown wild-type and mutant plants were collected into 2-ml Eppendorf tubes containing MS medium with or without 20 μM cycloheximide and incubated at room temperature for 15 minutes. Following vacuum infiltration for five minutes, the samples were incubated for three hours and then frozen for RNA extraction.

Cordycepin treatment

Wild-type and *upf3-1* plants were grown on MS plates for two weeks. Plants were then transferred to flasks containing incubation buffer (1 mM Pipes, 1 mM sodium citrate, 1 mM KCl, 15 mM sucrose). After 30 minutes, all seedlings except the time = 0 samples were transferred to fresh flasks containing incubation buffer plus cordycepin at 150 $\mu\text{g}/\text{ml}$. All flasks were then subject to a short (1–2 minute) vacuum infiltration. Time = 0 samples were immediately frozen in liquid nitrogen. 4–5 seedlings per

replicate per time-point were taken from the cordycepin-containing flasks and frozen, every hour for four hours.

RNA extraction, RT-PCR and qPCR analyses

Tissue was harvested and snap frozen in liquid nitrogen. Total RNA was extracted from leaves or whole plants using the QIAGEN RNeasy kit as per manufacturer's instructions, including the optional on-column digestion of DNA. mRNA levels were examined by RT-PCR and two micrograms of total, DNA-free RNA were reverse transcribed using Superscript II MMLV Reverse Transcriptase. RNase H was used to destroy any residual RNA and 2 µl of the reaction were then used as a template in a PCR as follows: an initial cycle at 95°C for 3 min followed by 27–37 cycles at 95°C for 1 min, 55–65°C (depending on T_m of primers) for 1 min and 72°C for 3 min, and a final cycle at 72°C for 5 min. The number of cycles was adjusted in each case to avoid over-cycling and all RT-PCR assays were carried out in duplicate. Selected RT-PCR assays were confirmed by qPCR. qPCR was carried out in the presence of the SYBR Green I dye, which binds to double stranded DNA and was monitored with the iCycler thermal cycler (BIO-RAD). Relative values of target expression were calculated as the average of two biological and three technical replicates by normalisation to the EF1α control. Primer sequences are given in Table S8.

Pseudomonas growth assays

A scalpel was used to nick the underside of fully expanded leaves of 5–6 week old plants either side of the mid-vein. A needleless syringe was then used to infiltrate leaves with DC3000 ($A_{600} = 0.0002$) suspended in 10 mM magnesium chloride. 3 days post-infection, a cork borer was used to excise a disc from the infection site. Bacteria were extracted by homogenisation in 10 mM magnesium chloride and cultured on Kings B medium at 28°C. The next day, the numbers of colony forming units were counted. Each experiment consisted of 3 leaves each from 4–6 plants of each genotype.

Hormone profiles

Post-inoculation with DC3000 ($A_{600} = 0.15$), plants were grown under constant light for 10 or 20 hours prior to harvest of inoculated leaves. Each wild-type sample consisted of three leaves from each of two plants. Due to their smaller size, four leaves from each of three mutant plants were pooled/sample. For each time-point, triplicate samples were collected and the mean and SEM were calculated. At harvest, samples were immediately snap frozen in liquid nitrogen. The experiment was performed three times. Hormone measurements were carried out as described in [110].

Supporting Information

Figure S1 Levels of jasmonic acid and abscisic acid in freeze dried tissues of wild type and NMD-deficient *Arabidopsis* leaves. Leaves were harvested from non-inoculated (NI) plants and plants 10 and 20 hours after infection with

Pseudomonas syringae DC3000 (DC10 and DC20). Wild-type plants are represented by blue bars, *upf1-5* mutants by red bars, *upf3-1* by green bars and *upf3-2* plants by purple bars.

(TIFF)

Table S1 Genes that are co-ordinately up-regulated in NMD mutant *Arabidopsis* (common NMD genes).

(XLS)

Table S2 Genes that are co-ordinately down-regulated in NMD mutant *Arabidopsis*.

(XLS)

Table S3 KOGs represented by *Arabidopsis* common NMD genes. The shading indicates the number of species for which genes belonging to the given KOG are up-regulated in response to abrogation of NMD. Darker shades indicate a greater number of species.

(XLS)

Table S4 23 common NMD genes that are not pathogen responsive, those with a CuORF are highlighted.

(XLS)

Table S5 Genes that are both regulated by NMD factors and have a role in auxin synthesis, or in the utilisation of chorismate. Where a gene is commonly up-regulated in all three NMD mutant lines it is shaded red. Orange indicates up-regulation in at least two NMD mutant lines and yellow indicates up-regulation in only one. Genes that are down-regulated in a mutant line are shaded light green. Those genes commonly down-regulated in two NMD mutant lines are coloured dark green.

(XLS)

Table S6 Putative NMD-targeted features of the common NMD genes. The 23 genes that are not in the pathogen-responsive clade are highlighted.

(XLSX)

Table S7 Microarray experiments represented in Figure 5. Four databases were used to compile Figure 5. The identifiers for the experiments used are given beneath the url for the database from which they were retrieved.

(XLSX)

Table S8 Primers used in this study.

(XLS)

Acknowledgments

We thank J. Lloyd and B. Causier for critical reading of the manuscript.

Author Contributions

Conceived and designed the experiments: SR LAR LW MG BD. Performed the experiments: SR LAR LW MDTZ. Analyzed the data: SR WT NG MG BD. Contributed reagents/materials/analysis tools: MG BD. Wrote the paper: SR BD.

References

- Maquat LE (2005) Nonsense-mediated mRNA decay in mammals. *Journal of Cell Science* 118: 1773–1776.
- He F, Li XR, Spatrick P, Casillo R, Dong SY, et al. (2003) Genome-wide analysis of mRNAs regulated by the nonsense-mediated and 5' to 3' mRNA decay pathways in yeast. *Molecular Cell* 12: 1439–1452.
- Mendell JT, Sharifi NA, Meyers JL, Martinez-Murillo F, Dietz HC (2004) Nonsense surveillance regulates expression of diverse classes of mammalian transcripts and mutes genomic noise. *Nature Genetics* 36: 1073–1078.
- Rehwinkel A, Letunic I, Raes J, Bork P, Izaurralde E (2005) Nonsense-mediated mRNA decay factors act in concert to regulate common mRNA targets. *RNA—a Publication of the Rna Society* 11: 1530–1544.
- Culbertson MR, Guan QN, Zheng W, Tang SJ, Liu XS, et al. (2006) Impact of nonsense-mediated mRNA decay on the global expression profile of budding yeast. *PLoS Genetics* 2: 1924–1943.
- Kurihara Y, Matsui A, Hanada K, Kawashima M, Ishida J, et al. (2009) Genome-wide suppression of aberrant mRNA-like noncoding RNAs by NMD in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America* 106: 2453–2458.
- Izaurralde E, Wittkopp N, Huntzinger E, Weiler C, Sauliere J, et al. (2009) Nonsense-mediated mRNA decay effectors are essential for zebrafish embryonic development and survival. *Molecular and Cellular Biology* 29: 3517–3528.

8. Le Hir H, Gatfield D, Izaurralde E, Moore MJ (2001) The exon-exon junction complex provides a binding platform for factors involved in mRNA export and nonsense-mediated mRNA decay. *The EMBO Journal* 20: 4987–4997.
9. Kerenyi Z, Merai Z, Hiriipi L, Benkovics A, Gyula P, et al. (2008) Interkingdom conservation of mechanism of nonsense-mediated mRNA decay. *The EMBO Journal* 27: 1585–1595.
10. Peltz SW, Brown AH, Jacobson A (1993) Messenger-RNA destabilization triggered by premature translational termination depends on at least 3 cis-acting sequence elements and one trans-acting factor. *Genes & Development* 7: 1737–1754.
11. Gonzalez CI, Ruiz-Echevarria MJ, Vasudevan S, Henry MF, Peltz SW (2000) The yeast hnRNP-like protein Hrp1/Nab4 marks a transcript for nonsense-mediated mRNA decay. *Molecular Cell* 5: 489–499.
12. Amrani N, Ganesan R, Kervestin S, Mangus DA, Ghosh S, et al. (2004) A faux 3'-UTR promotes aberrant termination and triggers nonsense-mediated mRNA decay. *Nature* 432: 112–118.
13. Behm-Ansmant I, Gatfield D, Rehwinkel J, Hilgers V, Izaurralde E (2007) A conserved role for cytoplasmic poly(A)-binding protein 1 (PABPC1) in nonsense-mediated mRNA decay. *The EMBO Journal* 26: 1591–1601.
14. Eberle AB, Stalder L, Mathys H, Orozco RZ, Muhlemann O (2008) Posttranscriptional gene regulation by spatial rearrangement of the 3' untranslated region. *PLoS Biology* 6: e92.
15. Ivanov PV, Gehring NH, Kunz JB, Hentze MW, Kulozik AE (2008) Interactions between UPF1, eRFs, PABP and the exon junction complex suggest an integrated model for mammalian NMD pathways. *The EMBO Journal* 27: 736–747.
16. Singh G, Rebbapragada I, Lykke-Andersen J (2008) A competition between stimulators and antagonists of Upf complex recruitment governs human nonsense-mediated mRNA decay. *PLoS Biology* 6: e111.
17. Pal M, Ishigaki Y, Nagy E, Maquat LE (2001) Evidence that phosphorylation of human Upf1 protein varies with intracellular location and is mediated by a wortmannin-sensitive and rapamycin-sensitive PI 3-kinase-related kinase signaling pathway. *RNA* 7: 5–15.
18. Denning G, Jamieson L, Maquat LE, Thompson EA, Fields AP (2001) Cloning of a novel phosphatidylinositol kinase-related kinase: characterization of the human SMG-1 RNA surveillance protein. *The Journal of Biological Chemistry* 276: 22709–22714.
19. Yamashita A, Kashima I, Ohno S (2005) The role of SMG-1 in nonsense-mediated mRNA decay. *Biochimica Et Biophysica Acta-Proteins and Proteomics* 1754: 305–315.
20. Grimson A, O'Connor S, Newman CL, Anderson P (2004) SMG-1 is a phosphatidylinositol kinase-related protein kinase required for nonsense-mediated mRNA decay in *Caenorhabditis elegans*. *Molecular and Cellular Biology* 24: 7483–7490.
21. Fukuhara N, Ebert J, Unterholzner L, Lindner D, Izaurralde E, et al. (2005) SMG7 is a 14-3-3-like adaptor in the nonsense-mediated mRNA decay pathway. *Molecular Cell* 17: 537–547.
22. Anders KR, Grimson A, Anderson P (2003) SMG-5, required for C-elegans nonsense-mediated mRNA decay, associates with SMG-2 and protein phosphatase 2A. *Embo Journal* 22: 641–650.
23. Ohnishi T, Yamashita A, Kashima I, Schell T, Anders KR, et al. (2003) Phosphorylation of hUPF1 induces formation of mRNA surveillance complexes containing hSMG-5 and hSMG-7. *Molecular Cell* 12: 1187–1200.
24. Chiu SY, Serin G, Ohara O, Maquat LE (2003) Characterization of human Smg5/7a: A protein with similarities to *Caenorhabditis elegans* SMG5 and SMG7 that functions in the dephosphorylation of Upf1. *RNA—A Publication of the RNA Society* 9: 77–87.
25. Serin G, Gersappe A, Black JD, Aronoff R, Maquat LE (2001) Identification and characterization of human orthologues to *Saccharomyces cerevisiae* Upf2 protein and Upf3 protein (*Caenorhabditis elegans* SMG-4). *Molecular and Cellular Biology* 21: 209–223.
26. Lejeune F, Maquat LE (2005) Mechanistic links between nonsense-mediated mRNA decay and pre-mRNA splicing in mammalian cells. *Current Opinion in Cell Biology* 17: 309–315.
27. Conti E, Izaurralde E (2005) Nonsense-mediated mRNA decay: molecular insights and mechanistic variations across species. *Current Opinion in Cell Biology* 17: 316–325.
28. Gatfield D, Unterholzner L, Ciccarelli FD, Bork P, Izaurralde E (2003) Nonsense-mediated mRNA decay in *Drosophila*: at the intersection of the yeast and mammalian pathways. *The EMBO Journal* 22: 3960–3970.
29. Arciga-Reyes L, Wootton L, Kieffer M, Davies B (2006) UPF1 is required for nonsense-mediated mRNA decay (NMD) and RNAi in *Arabidopsis*. *Plant Journal* 47: 480–489.
30. Yoine M, Nishii T, Nakamura K (2006) *Arabidopsis* UPF1 RNA helicase for nonsense-mediated mRNA decay is involved in seed size control and is essential for growth. *Plant and Cell Physiology* 47: 572–580.
31. Hori K, Watanabe Y (2005) UPF3 suppresses aberrant spliced mRNA in *Arabidopsis*. *Plant Journal* 43: 530–540.
32. Riha K, Riehs N, Akimcheva S, Puizina J, Bulankova P, et al. (2008) *Arabidopsis* SMG7 protein is required for exit from meiosis. *Journal of Cell Science* 121: 2208–2216.
33. Zetoun AB, Fontaniere S, Magnin D, Anczukow O, Buisson M, et al. (2008) Comparison of nonsense-mediated mRNA decay efficiency in various murine tissues. *BMC Genetics* 9: 83.
34. Gardner LB (2008) Hypoxic inhibition of nonsense-mediated RNA decay regulates gene expression and the integrated stress response. *Molecular and Cellular Biology* 28: 3729–3741.
35. Rodriguez-Gabriel MA, Watt S, Bahler J, Russell P (2006) Upf1, an RNA helicase required for nonsense-mediated mRNA decay, modulates the transcriptional response to oxidative stress in fission yeast. *Molecular and Cellular Biology* 26: 6347–6356.
36. Wilkinson MF, Bruno LG, Karam R, Huang LL, Bhardwaj A, et al. (2011) Identification of a MicroRNA that Activates Gene Expression by Repressing Nonsense-Mediated RNA Decay. *Molecular Cell* 42: 500–510.
37. Palusa SG, Reddy ASN (2010) Extensive coupling of alternative splicing of pre-mRNAs of serine/arginine (SR) genes with nonsense-mediated decay. *New Phytologist* 185: 83–89.
38. Malanin SY, Nikiforova VY (2010) A link between alternative splicing and nonsense-mediated mRNA decay in the control of THIC gene expression in *Arabidopsis*. *Russian Journal of Plant Physiology* 57: 267–272.
39. Schoning JC, Streitner C, Meyer IM, Gao Y, Staiger D (2008) Reciprocal regulation of glycine-rich RNA-binding proteins via an interlocked feedback loop coupling alternative splicing to nonsense-mediated decay in *Arabidopsis*. *Nucleic Acids Research* 36: 6977–6987.
40. Song HR, Song JD, Cho JN, Amasino RM, Noh B, et al. (2009) The RNA binding protein ELF9 directly reduces SUPPRESSOR OF OVEREXPRESSION OF CO1 transcript levels in *Arabidopsis*, possibly via nonsense-mediated mRNA decay. *The Plant Cell* 21: 1195–1211.
41. Jones JD, Dangl JL (2006) The plant immune system. *Nature* 444: 323–329.
42. Ausubel FM (2005) Are innate immune signaling pathways in plants and animals conserved? *Nature Immunology* 6: 973–979.
43. Thilmony R, Underwood W, He SY (2006) Genome-wide transcriptional analysis of the *Arabidopsis thaliana* interaction with the plant pathogen *Pseudomonas syringae* pv. tomato DC3000 and the human pathogen *Escherichia coli* O157:H7. *The Plant Journal: for cell and molecular biology* 46: 34–53.
44. Truman W, de Zabala MT, Grant M (2006) Type III effectors orchestrate a complex interplay between transcriptional networks to modify basal defence responses during pathogenesis and resistance. *The Plant Journal: for cell and molecular biology* 46: 14–33.
45. Kim MG, da Cunha L, McFall AJ, Belkhadir Y, DebRoy S, et al. (2005) Two *Pseudomonas syringae* type III effectors inhibit RIN4-regulated basal defense in *Arabidopsis*. *Cell* 121: 749–759.
46. Nurnberger T, Brunner F, Kemmerling B, Piater L (2004) Innate immunity in plants and animals: striking similarities and obvious differences. *Immunological Reviews* 198: 249–266.
47. da Cunha L, Sreerekha MV, Mackey D (2007) Defense suppression by virulence effectors of bacterial phytopathogens. *Current Opinion in Plant Biology* 10: 349–357.
48. Glazebrook J (2005) Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annual Review of Phytopathology* 43: 205–227.
49. Gaffney T, Friedrich L, Vernooij B, Negroto D, Nye G, et al. (1993) Requirement of salicylic acid for the induction of systemic acquired resistance. *Science* 261: 754–756.
50. de Torres Zabala M, Bennett MH, Truman WH, Grant MR (2009) Antagonism between salicylic and abscisic acid reflects early host-pathogen conflict and moulds plant defence responses. *The Plant Journal: for cell and molecular biology* 59: 375–386.
51. Sklarczyk D, Franceschini A, Kuhn M, Simonovic M, Roth A, et al. (2011) The STRING database in 2011: functional interaction networks of proteins, globally integrated and scored. *Nucleic Acids Research* 39: D561–568.
52. Swarbreck D, Wilks C, Lamesch P, Berardini TZ, Garcia-Hernandez M, et al. (2008) The *Arabidopsis* Information Resource (TAIR): gene structure and function annotation. *Nucleic Acids Research* 36: D1009–1014.
53. Dimmer KS, Friedrich B, Lang F, Deitmer JW, Broer S (2000) The low-affinity monocarboxylate transporter MCT4 is adapted to the export of lactate in highly glycolytic cells. *The Biochemical Journal* 350(Pt 1): 219–227.
54. Godard P, Urrestarazu A, Vissers S, Kontos K, Bontempi G, et al. (2007) Effect of 21 different nitrogen sources on global gene expression in the yeast *Saccharomyces cerevisiae*. *Molecular and Cellular Biology* 27: 3065–3086.
55. Vincill ED, Szczygłowski K, Roberts DM (2005) GmN70 and LjN70. Anion transporters of the symbiosome membrane of nodules with a transport preference for nitrate. *Plant Physiology* 137: 1435–1444.
56. Chan WK, Huang L, Gudikote JP, Chang YF, Imam JS, et al. (2007) An alternative branch of the nonsense-mediated decay pathway. *The EMBO Journal* 26: 1820–1830.
57. Gaba A, Jacobson A, Sachs MS (2005) Ribosome occupancy of the yeast CPA1 upstream open reading frame termination codon modulates nonsense-mediated mRNA decay. *Molecular Cell* 20: 449–460.
58. Vlot AC, Dempsey DA, Klessig DF (2009) Salicylic acid, a multifaceted hormone to combat disease. *Annual Review of Phytopathology* 47: 177–206.
59. Schenk PM, Kazan K, Wilson I, Anderson JP, Richmond T, et al. (2000) Coordinated plant defense responses in *Arabidopsis* revealed by microarray analysis. *Proceedings of the National Academy of Sciences of the United States of America* 97: 11655–11660.
60. Clarke JD, Volko SM, Ledford H, Ausubel FM, Dong X (2000) Roles of salicylic acid, jasmonic acid, and ethylene in cpr-induced resistance in *Arabidopsis*. *The Plant Cell* 12: 2175–2190.

61. Grant MR, Truman WM, Bennett MH, Turnbull CGN (2010) Arabidopsis auxin mutants are compromised in systemic acquired resistance and exhibit aberrant accumulation of various indolic compounds. *Plant Physiology* 152: 1562–1573.
62. Truman W, de Zabala MT, Grant M (2006) Type III effectors orchestrate a complex interplay between transcriptional networks to modify basal defence responses during pathogenesis and resistance. *Plant Journal* 46: 14–33.
63. Biles CL (2001) Plant Disease Resistance: The response of plants to avirulent pathogens. *Bios* 72: 67–69.
64. Grant MR, Jones JD (2009) Hormone (dis)harmony moulds plant health and disease. *Science* 324: 750–752.
65. Bender CL, Stone HE, Sims JJ, Cooksey DA (1987) Reduced pathogen fitness of *Pseudomonas-syringae* pv tomato Tn5 mutants defective in coronatine production. *Physiological and Molecular Plant Pathology* 30: 273–283.
66. Robert-Seilaniantz A, Grant M, Jones JD (2011) Hormone crosstalk in plant disease and defense: more than just jasmonate-salicylate antagonism. *Annual Review of Phytopathology* 49: 317–343.
67. Nyiko T, Sonkoly B, Merai Z, Benkovics AH, Silhavy D (2009) Plant upstream ORFs can trigger nonsense-mediated mRNA decay in a size-dependent manner. *Plant Molecular Biology* 71: 367–378.
68. Kertesz S, Kerenyi Z, Merai Z, Bartos I, Palfy T, et al. (2006) Both introns and long 3'-UTRs operate as cis-acting elements to trigger nonsense-mediated decay in plants. *Nucleic Acids Research* 34: 6147–6157.
69. Shashikanth M, Krishna AR, Ramya G, Devi G, Ulaganathan K (2008) Genome-wide comparative analysis of *Oryza sativa* (japonica) and *Arabidopsis thaliana* 5' UTR sequences for translational regulatory signals. *Plant Biotechnology* 25: 553–563.
70. Hayden CA, Jorgensen RA (2007) Identification of novel conserved peptide uORF homology groups in Arabidopsis and rice reveals ancient eukaryotic origin of select groups and preferential association with transcription factor-encoding genes. *BMC Biology* 5: 32.
71. Ramani AK, Nelson AC, Kapranov P, Bell I, Gingeras TR, et al. (2009) High resolution transcriptome maps for wild-type and nonsense-mediated decay-defective *Caenorhabditis elegans*. *Genome Biology* 10: R101.
72. Viegas MH, Gehring NH, Breit S, Hentze MW, Kulozik AE (2007) The abundance of RNPS1, a protein component of the exon junction complex, can determine the variability in efficiency of the Nonsense Mediated Decay pathway. *Nucleic Acids Research* 35: 4542–4551.
73. McGlinchy NJ, Tan LY, Paul N, Zavolan M, Lilley KS, et al. (2010) Expression proteomics of UPF1 knockdown in HeLa cells reveals autoregulation of hnRNP A2/B1 mediated by alternative splicing resulting in nonsense-mediated mRNA decay. *BMC Genomics* 11: 565.
74. Yoine M, Ohto MA, Onai K, Mita S, Nakamura K (2006) The lba1 mutation of UPF1 RNA helicase involved in nonsense-mediated mRNA decay causes pleiotropic phenotypic changes and altered sugar signalling in Arabidopsis. *Plant Journal* 47: 49–62.
75. Lamba JK, Adachi M, Sun D, Tammur J, Schuetz EG, et al. (2003) Nonsense mediated decay downregulates conserved alternatively spliced ABCC4 transcripts bearing nonsense codons. *Human Molecular Genetics* 12: 99–109.
76. Jennings MD, Pavitt GD (2010) eIF5 has GDI activity necessary for translational control by eIF2 phosphorylation. *Nature* 465: 378–381.
77. Zhang XC, Gassmann W (2007) Alternative splicing and mRNA levels of the disease resistance gene RPS4 are induced during defense responses. *Plant Physiology* 145: 1577–1587.
78. Yi H, Richards EJ (2007) A cluster of disease resistance genes in Arabidopsis is coordinately regulated by transcriptional activation and RNA silencing. *The Plant Cell* 19: 2929–2939.
79. Venugopal SC, Jeong RD, Mandal MK, Zhu S, Chandra-Shekara AC, et al. (2009) Enhanced disease susceptibility 1 and salicylic acid act redundantly to regulate resistance gene-mediated signaling. *PLoS Genetics* 5: e1000545.
80. Mohr TJ, Mammarella ND, Hoff T, Woffenden BJ, Jelesko JG, et al. (2010) The Arabidopsis downy mildew resistance gene RPP8 is induced by pathogens and salicylic acid and is regulated by W box cis elements. *Molecular Plant-Microbe Interactions: MPPI* 23: 1303–1315.
81. Xu S, Zhang Z, Jing B, Gannon P, Ding J, et al. (2011) Transportin-SR is required for proper splicing of resistance genes and plant immunity. *PLoS Genetics* 7: e1002159.
82. Zhang XC, Gassmann W (2003) RPS4-mediated disease resistance requires the combined presence of RPS4 transcripts with full-length and truncated open reading frames. *The Plant Cell* 15: 2333–2342.
83. Dinesh-Kumar SP, Baker BJ (2000) Alternatively spliced N resistance gene transcripts: their possible role in tobacco mosaic virus resistance. *Proceedings of the National Academy of Sciences of the United States of America* 97: 1908–1913.
84. Liu G, Ji Y, Bhuiyan NH, Pilot G, Selvaraj G, et al. (2010) Amino acid homeostasis modulates salicylic acid-associated redox status and defense responses in Arabidopsis. *The Plant Cell* 22: 3845–3863.
85. Grant M, Ward JL, Forcat S, Beckmann M, Bennett M, et al. (2010) The metabolic transition during disease following infection of Arabidopsis thaliana by *Pseudomonas syringae* pv. tomato. *Plant Journal* 63: 443–457.
86. Heath MC (2000) Hypersensitive response-related death. *Plant Molecular Biology* 44: 321–334.
87. Heidel AJ, Clarke JD, Antonovics J, Dong XN (2004) Fitness costs of mutations affecting the systemic acquired resistance pathway in Arabidopsis thaliana. *Genetics* 168: 2197–2206.
88. Jin JB, Jin YH, Lee J, Miura K, Yoo CY, et al. (2008) The SUMO E3 ligase, AtSIZ1, regulates flowering by controlling a salicylic acid-mediated floral promotion pathway and through effects on FLC chromatin structure. *The Plant Journal: for cell and molecular biology* 53: 530–540.
89. Wada KC, Yamada M, Shiraya T, Takeno K (2010) Salicylic acid and the flowering gene FLOWERING LOCUS T homolog are involved in poor-nutrition stress-induced flowering of *Pharbitis nil*. *Journal of Plant Physiology* 167: 447–452.
90. Devadas SK, Enyedi A, Raina R (2002) The Arabidopsis hrl1 mutation reveals novel overlapping roles for salicylic acid, jasmonic acid and ethylene signalling in cell death and defence against pathogens. *The Plant Journal: for cell and molecular biology* 30: 467–480.
91. Wawrzynska A, Rodibaugh NL, Innes RW (2010) Synergistic activation of defense responses in Arabidopsis by simultaneous loss of the GSL5 callose synthase and the EDR1 protein kinase. *Molecular Plant-Microbe Interactions: MPPI* 23: 578–584.
92. Mosher S, Moeder W, Nishimura N, Jikumaru Y, Joo SH, et al. (2010) The lesion-mimic mutant cpr22 shows alterations in abscisic acid signaling and abscisic acid insensitivity in a salicylic acid-dependent manner. *Plant Physiology* 152: 1901–1913.
93. Jing HC, Anderson L, Sturre MJ, Hille J, Dijkwel PP (2007) Arabidopsis CPR5 is a senescence-regulatory gene with pleiotropic functions as predicted by the evolutionary theory of senescence. *Journal of Experimental Botany* 58: 3885–3894.
94. Shirano Y, Kachroo P, Shah J, Klessig DF (2002) A gain-of-function mutation in an Arabidopsis Toll Interleukin1 receptor-nucleotide binding site-leucine-rich repeat type R gene triggers defense responses and results in enhanced disease resistance. *The Plant Cell* 14: 3149–3162.
95. Nibbe M, Hilpert B, Wasternack C, Miersch O, Apel K (2002) Cell death and salicylate- and jasmonate-dependent stress responses in Arabidopsis are controlled by single cdc genes. *Planta* 216: 120–128.
96. Li Y, Tian Z, Liu J, Xie C (2009) Comparative cDNA-AFLP analysis reveals that DL-beta-amino-butyric acid induces resistance through early activation of the host-defense genes in potato. *Physiologia Plantarum* 136: 19–29.
97. Lee J, Nam J, Park HC, Na G, Miura K, et al. (2007) Salicylic acid-mediated innate immunity in Arabidopsis is regulated by SIZ1 SUMO E3 ligase. *The Plant Journal: for cell and molecular biology* 49: 79–90.
98. Ruiz-Echevarria MJ, Peltz SW (2000) The RNA binding protein Pub1 modulates the stability of transcripts containing upstream open reading frames. *Cell* 101: 741–751.
99. Hamid R, Hedges LK, Austin E, Phillips JA, Loyd JE, et al. (2010) Transcripts from a novel BMPR2 termination mutation escape nonsense mediated decay by downstream translation re-initiation: implications for treating pulmonary hypertension. *Clinical Genetics* 77: 280–286.
100. Hirano SS, Upper CD (1990) Population biology and epidemiology of *Pseudomonas-syringae*. *Annual Review of Phytopathology* 28: 155–177.
101. Sumper M, Knott JM, Romer P (2007) Putative spermine synthases from *Thalassiosira pseudonana* and Arabidopsis thaliana synthesize thermospermine rather than spermine. *FEBS Letters* 581: 3081–3086.
102. Imai A, Komura M, Yamamoto K, Takahashi T (2005) Analysis of suppressors of the spermine synthase mutant, ac15, that is defective in stem elongation in Arabidopsis. *Plant and Cell Physiology* 46: S30–S30.
103. Takahashi T, Imai A, Hanzawa Y, Komura M, Yamamoto KT, et al. (2006) The dwarf phenotype of the Arabidopsis ac15 mutant is suppressed by a mutation in an upstream ORF of a bHLH gene. *Development* 133: 3575–3585.
104. Scholl RL, May ST, Ware DH (2000) Seed and molecular resources for Arabidopsis. *Plant Physiology* 124: 1477–1480.
105. May S, Craigon DJ, James N, Okyere J, Higgins J, et al. (2004) NASCArrays: a repository for microarray data generated by NASC's transcriptomics service. *Nucleic Acids Research* 32: D575–D577.
106. Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, et al. (2003) Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249–264.
107. Huang DW, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* 4: 44–57.
108. Dennis G, Sherman BT, Hosack DA, Yang J, Gao W, et al. (2003) DAVID: Database for annotation, visualization, and integrated discovery. *Genome Biology* 4: DOI: 10.1186/gb-2003-4-9-r60.
109. Saldanha AJ (2004) Java Treeview-extensible visualization of microarray data. *Bioinformatics* 20: 3246–3248.
110. Grant M, de Torres-Zabala M, Truman W, Bennett MH, Lafforgue G, et al. (2007) *Pseudomonas syringae* pv. tomato hijacks the Arabidopsis abscisic acid signalling pathway to cause disease. *Embo Journal* 26: 1434–1443.