

**Original citation:**

Parashar, D., Bowden, Jack, Starr, Colin, Wernisch, Lorenz and Mander, Adrian (2015) Type I error control in biomarker-stratified clinical trials. In: 3rd International Clinical Trials Methodology Conference, Glasgow SECC, UK, 16-17 Nov 2015. Published in: *Trials*, 16 (Suppl 2). 084.

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/78665>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the CC0 1.0 Universal (CC0 1.0) Public Domain Dedication license and may be reused according to the conditions of the license. For more details see:

<http://creativecommons.org/publicdomain/zero/1.0/>

**A note on versions:**

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)

ORAL PRESENTATION

Open Access

# Type I error control in biomarker-stratified clinical trials

Deepak Parashar<sup>1\*</sup>, Jack Bowden<sup>2</sup>, Colin Starr<sup>2</sup>, Lorenz Wernisch<sup>2</sup>, Adrian Mander<sup>2</sup>

From 3rd International Clinical Trials Methodology Conference  
Glasgow, UK. 16-17 November 2015

Biomarker-stratified clinical trials assess the biomarker signature of subjects and split them into subgroups so that treatment is of benefit to those who are likely to respond. Since multiple hypotheses are tested, it becomes important to control the type I error. Current methods control the false positive rate where one rejects the null hypothesis while in reality that was true. For two subgroups, the false positive rate is controlled across the two hypotheses as a Family Wise Error Rate (FWER) to an overall predetermined significance level.

The concept of “Wrong Positive” rate is presented where one rejects the null hypothesis in one subgroup while in reality the efficacy would be in the other subgroup. Controlling the wrong positives as well as false positives to an overall significance level constitutes a strong FWER. Further options exist for control of Individual Outcome FWER by assigning weighting to reject the respective null hypotheses.

We apply our methodology to an optimal biomarker-stratified Simon two-stage cancer clinical trial design and illustrate the variation in minimum expected sample sizes for different choices of type I controls for a range of operating characteristics. While the weak FWER would suit a phase II trial, additionally controlling the wrong positives could be more appropriate in a confirmatory setting. The chosen design is an adaptive enrichment that evaluates efficacy in biomarker-positive as well as biomarker-negative subpopulations, and optimal designs are obtained using the four different choices of type I error controls: weak or strong FWER with or without individual outcomes.

## Authors' details

<sup>1</sup>University of Warwick, Coventry, UK. <sup>2</sup>MRC Biostatistics Unit, Cambridge, UK.

<sup>1</sup>University of Warwick, Coventry, UK

Full list of author information is available at the end of the article

Published: 16 November 2015

doi:10.1186/1745-6215-16-S2-O84

Cite this article as: Parashar et al.: Type I error control in biomarker-stratified clinical trials. *Trials* 2015 **16**(Suppl 2):O84.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

