

**Original citation:**

Eyden, Julie, Winsper, Catherine, Wolke, Dieter, Broome, Matthew R. and MacCallum, Fiona. (2016) A systematic review of the parenting and outcomes experienced by offspring of mothers with borderline personality pathology : potential mechanisms and clinical implications. Clinical Psychology Review.

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/78897>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

© 2016, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

**A note on versions:**

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)

Running head: MOTHERS WITH BPD: PARENTING & OFFSPRING OUTCOMES

A systematic review of the parenting and outcomes experienced by offspring of mothers with borderline personality pathology: Potential mechanisms and clinical implications

Julie Eyden<sup>1</sup>, Catherine Winsper<sup>2</sup>, Dieter Wolke<sup>1,2</sup>, Matthew R. Broome<sup>2,3,4</sup>, & Fiona MacCallum<sup>1</sup>

<sup>1</sup>Department of Psychology, University of Warwick, Coventry, UK

<sup>2</sup>Division of Mental Health and Wellbeing, Warwick Medical School, University of Warwick, Coventry, UK

<sup>3</sup>Department of Psychiatry, University of Oxford, Oxford, UK

<sup>4</sup>Warneford Hospital, Oxford Health NHS Foundation Trust, Oxford, UK

Correspondence to Julie Eyden: [j.eyden@warwick.ac.uk](mailto:j.eyden@warwick.ac.uk)

Key words: Borderline personality disorder; mothers; parenting; offspring outcomes; systematic review

### Abstract

There is growing interest in whether the parenting strategies and offspring outcomes of mothers with borderline personality disorder (BPD) differ from those of mothers without BPD. We searched PsychINFO, PubMed, EMBASE, Web of Science, Scopus and ASSIA databases for studies examining parenting skills and attitudes among mothers with BPD/BPD symptoms and/or offspring outcomes. PRISMA reporting guidelines were followed. Of 10,067 abstracts screened, 101 full-text articles were retrieved and 33 met pre-determined criteria for qualitative synthesis. Overall, studies suggest that mothers with BPD/BPD symptoms are more likely to engage in maladaptive interactions with their offspring characterised by insensitive, overprotective, and hostile parenting compared to mothers without BPD/BPD symptoms. Adverse offspring outcomes include BPD symptoms, internalising (including depression) and externalising problems, insecure attachment patterns, and emotional dysregulation. Findings suggest that vulnerability from mother to offspring may be partly transmitted via maladaptive parenting and maternal emotional dysfunction. Conclusions were limited by study heterogeneity in methodology and construct definitions, as well as a paucity of clinical comparison groups. Prospective studies of mothers with BPD and their offspring from pregnancy onwards may further elucidate mechanisms of transmission and identify resilience factors across development. Parenting behaviour awareness, improving attachment behaviours and emotional regulation strategies may be important intervention targets.

A systematic review of the parenting and outcomes experienced by offspring of mothers with borderline personality pathology: Potential mechanisms and clinical implications

Borderline Personality Disorder (BPD) is a complex mental condition characterised by extreme emotional, behavioural, and interpersonal dysregulation (American Psychiatric Association, 2013). Consequently, it is associated with a range of long-term negative sequelae, including relationship dysfunction (Daley, Burge, & Hammen, 2000), unemployment (Skodol et al., 2002), co-morbid psychopathology (Grant et al., 2008; Zanarini et al., 1998), self-harm and suicide (Black, Blum, Pfohl, & Hale, 2004), high levels of treatment utilisation (Bender et al., 2001) and imprisonment (Black et al., 2007).

BPD affects approximately 1-6% of the general population (Grant et al., 2008; Swartz, Blazer, George, & Winfield, 1990; Trull, Jahng, Tomko, Wood, & Sher, 2010). While some studies suggest that BPD may be equally distributed across males and females in community populations (Grant et al., 2008; Sansone & Sansone, 2011), a higher proportion of women with BPD are seen in clinical settings (American Psychiatric Association, 2000). Women with BPD are more likely to have disrupted relationships, engage in self-harm, and have greater overall symptomatology, depression and anxiety (Sansone & Sansone, 2011; Silberschmidt, Lee, Zanarini, & Schulz, 2015), which may partly explain the increased utilisation of psychiatric services (Gunderson & Hoffman, 2005; Sansone & Sansone, 2011; Skodol & Bender, 2003). As BPD is associated with pervasive functional impairment, and insecure attachment patterns (Agrawal, Gunderson, Holmes, & Lyons-Ruth, 2004), parenting may be particularly challenging for women who experience the symptoms of BPD. Individuals with BPD are likely to be separated, divorced or to have never

married (Skodol et al., 2002), and those in relationships are at greater risk of experiencing partner conflict and abuse (Chen et al., 2004). As such, mothers with BPD may have limited levels of emotional, social, and financial support potentially exacerbating parenting difficulties.

A small number of narrative reviews of mothers with BPD have examined aspects of parenting and impacts on offspring (Macfie, 2009; Stepp, Whalen, Pilkonis, Hipwell, & Levine, 2012; Wendland et al., 2014). Stepp et al. (2012) hypothesised that certain parenting strategies exhibited by mothers with BPD (i.e., hostile control and passive aloofness) might be involved in the transmission of psychopathology from mother to child. However, at the time, there were limited studies pertaining to parenting and child outcomes, and only one study (Abela, Skitch, Auerbach, & Adams, 2005) explicitly tested potential mechanisms via which maternal BPD may increase risk of negative child outcomes.

The identification of parenting strategies (and potential associated adverse outcomes for offspring) of mothers with BPD could inform tailored interventions for both mother and child (Stepp et al., 2012). Additionally, an understanding of the mechanisms, beyond predisposition (Crowell, Beauchaine, & Linehan, 2009; Distel et al., 2008; Gratz et al., 2015; Torgersen et al., 2000), underpinning links between maternal BPD and offspring outcome could inform theory regarding aetiological trajectories to BPD and other disorders from infancy onwards.

Recently, Petfield, Startup, Droscher and Cartwright-Hatton (2015) published a systematic review examining parenting behaviors in mothers with BPD and the impact on child outcomes. They identified 17 studies focusing solely on mothers with a diagnosis of BPD and their children and infants. In the current review, we expand on these findings by broadening our systematic search to include offspring of any age

(including adults), and mothers with either a BPD diagnosis or BPD symptoms. We considered it important to include studies assessing sub-clinical BPD symptoms, as many researchers prefer a dimensional approach to the assessment of BPD due to concerns regarding appropriate thresholds and heterogeneity of the disorder (Skodol et al., 2002). We therefore took this approach to ensure coverage of all relevant studies. Furthermore, to move the literature forward we explicitly examined the potential mechanisms underpinning the associations between the parenting behaviors of mothers with BPD and offspring outcomes.

The aim of the current review was to systematically search and narratively synthesise all research examining the parenting behaviours and attitudes of mothers with BPD, mother-offspring interactions, and offspring outcomes. Specifically, we addressed the following four questions:

- 1) What are the characteristic parenting behaviours of mothers with borderline personality pathology (i.e., BPD diagnosis or BPD symptoms)?
- 2) How do mothers with borderline personality pathology and their offspring interact?
- 3) What are the psychopathological and psychosocial outcomes for offspring of mothers with borderline personality pathology?
- 4) What are the mechanisms (parenting or mother/offspring characteristics) underpinning associations between maternal borderline personality pathology and offspring outcomes?

### **Methods**

We used the Preferred Reporting Items for Systematic Reviews and Meta Analyses (PRISMA; Moher, Liberati, Tetzlaff, & Altman, 2009) guidelines throughout this review.

## Data Sources

We searched titles and abstracts in PsychINFO, PubMed, EMBASE, Web of science, Scopus and ASSIA to identify articles examining samples of mothers with BPD pathology (i.e., BPD diagnosis or symptoms), and/or children of mothers with BPD pathology, published between 1980 and the 6<sup>th</sup> July 2015. We selected 1980 as the earliest date for inclusion to reflect when BPD was first delineated as a mental disorder in the DSM III. To enhance search sensitivity manual searches were conducted in the *Journal of Personality Disorders* and *Personality Disorders: Theory, Research and Treatment* from January 2010 to July 2015 and the reference lists of retrieved articles and review papers were inspected to identify additional potentially relevant articles.

## Search Terms

We used the following grouped terms: (borderline\* OR “emotionally unstable personality” OR BPD) AND (mother\* OR parent\* OR maternal\*) AND (child\* OR infant\* OR infancy OR offspring OR bab\* OR adolescen\* OR famil\* OR boy\* OR girl\* OR teenager\* OR youth\* OR young\* OR toddler\* OR daughter\* OR son\*).

## Eligibility Criteria

The review included retrospective, cross-sectional, and prospective quantitative studies meeting the following inclusion criteria:

- 1) The study included mothers with BPD or BPD symptoms (clinical or community samples) and/or offspring (of any age) of mothers with BPD or BPD symptoms (assessed via a standardised measure)
- 2) The study reported on maternal parenting, and/or offspring outcomes using a range of assessment methods (i.e., experimental, observational, interview, self-report questionnaires, and other-report questionnaires)

3) Samples in studies reporting on associations with parenting characteristics or child outcomes consisted mainly of mothers (i.e., at least 70% of the parenting sample).

Studies were excluded if they:

- 1) Were reviews, expert opinion commentaries, or individual case studies
- 2) Considered associations with maternal personality disorders generally rather than BPD specifically
- 3) Were not written in English language or were unpublished
- 4) Reported on extreme outcomes resulting in external intervention (i.e., abuse, filicide, etc.).

### **Screening Procedure**

J.E. and C.W. independently screened all titles and abstracts from the initial search to identify articles for full-text retrieval. If a title appeared relevant but no abstract was available, the full article was retrieved. All full-text articles were independently screened by J.E. and C.W. for final inclusion in the review.

### **Data Extraction and Quality Assessment**

We created a data extraction form based on PRISMA and Cochrane guidelines (Higgins & Green, 2011; Moher et al., 2009). This included information on first author, date and country of study, sample characteristics, BPD assessment tool, study aims, study design and methodology, measurement tool, comparison group and main findings. The quality of each study was assessed by J.E. using the Newcastle-Ottawa Scale (Wells et al., 2000). F.M. independently assessed 50% of the studies as a reliability check. This quality assessment tool comprises a star rating across various quality domains, can be used for case-control and cohort studies, and has been adapted for use with cross-sectional studies (Herzog et al., 2013). The quality domains of group selection, comparability and exposure were assessed for case

control studies (maximum 9 stars awarded). For cohort studies, selection, comparability of cohorts, and outcome quality were assessed (maximum 9 stars). For cross-sectional studies, sample selection, comparability, and outcome were assessed (maximum 10 stars).

### **Data Synthesis**

All studies were qualitatively synthesised. A meta-analysis of the results was not feasible due to the heterogeneity across studies in terms of definitions of parenting and offspring constructs, study design (e.g., experimental, observational, self-report), sample frame (e.g., community, clinical, ranging from infants to adults), and assessment tools. For example, insensitive parenting across studies was variably defined as: “maternal sensitivity,” “insensitivity,” “intrusive sensitivity,” “intrusive negativity,” and within a composite variable (including “sensitivity,” “warmth/hostility,” “acceptance/rejection,” “responsiveness” and “demandingness”).

To address the aims of our review, a narrative overview of the main findings is presented in four major sections: 1) Parenting behaviours (e.g., sensitivity, hostility, overprotection) and perceptions (e.g., maternal rating of parenting efficacy, perceptions of their offspring); 2) Mother-offspring interaction dynamics (e.g., interactions, communications, infant/child behaviour); 3) Offspring outcomes (psychopathological and psychosocial); and 4) Mechanisms underpinning associations between maternal borderline personality pathology and child outcomes. Whilst some outcomes naturally fall into more than one category (particularly regarding maternal parenting behaviours and interactions), to avoid repetition each outcome has been included in one section only.

## Results

### Included Articles

We retrieved 10,067 records: 10,046 from the database search and 21 from hand searching. From abstract screening, 101 articles were identified for full-text retrieval, with an excellent level of agreement between researchers (Cohen's Kappa,  $\kappa = .87$ ;  $p < .001$ ). After screening the full-text articles a total of 33 studies were selected. Inter-rater agreement for final inclusion in the review was again excellent ( $\kappa = .88$ ,  $p < .001$ ). Any disagreements were discussed and resolved between J.E. and C.W. The main reasons for exclusion included: only offspring BPD (not maternal BPD) assessed, studies referred to any relative with BPD and not the mother specifically, maternal personality disorders (not BPD specifically) was assessed, there was no valid measure of BPD (see Figure 1).

### Study Characteristics

We identified 21 case-control, 8 cross-sectional, and 4 cohort studies (see Table 1). Two of the cohort studies were drawn from the same Greifswald Family Study sample (Barnow et al., 2013; Reinelt et al., 2014). Studies of offspring outcomes included a range of age groups: 13 reported infant (aged 2-36 months) outcomes, 11 reported child (aged 3-9 years) outcomes, 13 reported adolescent (aged 10-19 years) outcomes, and 4 reported adult (aged 19+ years) outcomes. Some studies utilised samples encompassing both childhood and adolescence (e.g., Abela et al., 2005; Feldman et al., 1995) or adolescence and adulthood (e.g., Cheng, Huang, Liu, & Liu, 2011; Reinelt et al., 2014). Twenty studies reported on parenting characteristics (i.e., behaviours, perceptions and mother-offspring interaction dynamics) of mothers with borderline personality pathology.

A number of countries were represented (see Table 1). In five of the reviewed studies, English was not the first language of the participants. Three were German in origin (Barnow et al., 2013; Barnow, Spitzer, Grabe, Kessler, & Freyberger, 2006; Reinelt et al., 2014), one was French (Delavenne, Gratier, Devouche, & Apter, 2008), and one was Chinese (Cheng et al., 2011). An inspection of these studies did not indicate any obvious cultural differences (e.g., mothers with BPD from Germany appeared to demonstrate similar levels of overprotection to mothers from other countries), though direct comparisons were difficult due to variations across studies.

### **Quality Assessment**

The quality assessment showed substantial inter-rater agreement (Cohen's Kappa,  $\kappa = .77$ ;  $p < .001$ ), and indicated low risk of bias in sample selection, low risk of comparability bias, and low-moderate risk of exposure/outcome bias (see Supplementary Table 1).

### **Findings**

#### **1) Parenting Behaviours and Perceptions**

Studies assessed several maternal parenting behaviours (i.e., sensitivity and intrusiveness, overprotection, emotional warmth, hostility, rejection) and perceptions (i.e., emotion recognition, parenting stress/distress, parenting efficacy, and representations of child). There were wide variations in the definitions and operationalisation of parenting constructs across studies (see Table 2 for details).

***Sensitivity/intrusiveness.*** Seven studies assessed aspects of maternal sensitivity/intrusiveness. Two studies reported that mothers with BPD (or BPD symptoms) displayed (or reported) significantly lower “sensitivity/non-intrusive sensitivity” with their infants compared to healthy control mothers (HCs) (Crandell, Hobson, & Patrick, 2003; Newman, Stevenson, Bergman, & Boyce, 2007). Howard,

Beckwith, Espinosa, & Tyler (1995) found a negative correlation between BPD symptoms and maternal sensitivity, and Hobson, Patrick, Crandell, Garcia-Perez, and Lee (2005) reported significantly higher “intrusive insensitivity” in mothers with BPD. Controlling for anxiety and depression, Kiel, Gratz, Moore, Lutzman, and Tull (2011) found that mothers with clinically relevant levels of BPD symptoms were significantly more “insensitive” with their infants when infant distress persisted for longer durations. Hobson et al. (2009) reported that mothers with BPD displayed more “intrusiveness/negativity” with their infants than healthy and depressed controls, but these differences did not quite reach significance. The only adolescent offspring community-based study did not find a significant correlation between offspring-reported maternal “intrusiveness” and maternal BPD symptoms (Zalewski et al., 2014).

***Overprotection.*** Four studies converge in indicating that mothers with BPD/BPD symptoms are more overprotective (or overinvolved) with their offspring both in infancy (Elliot et al., 2014) and adolescence (Barnow et al., 2006; Reinelt et al., 2014; Zalewski et al., 2014). Elliot et al. (2014) found that mothers with BPD scored significantly higher on self-reported overprotection than HCs. Barnow et al. (2006) found that adolescents of mothers with BPD perceived them as significantly more overprotective than did adolescents of mothers with depressive disorder, cluster C personality disorder, or no mental health condition. Similarly, Reinelt et al. (2014) found that adolescent-reported overprotection was significantly correlated with mother’s BPD symptoms. Finally, Zalewski et al. (2014) found that maternal BPD symptoms were significantly positively correlated with offspring-reported maternal attempts to “control through guilt” (after controlling for depression and alcohol use) and negatively correlated with “acceptance of individuation.”

**Maternal warmth.** Four studies assessed maternal warmth yielding varying results across age groups. When mothers were observed interacting with their preschool children who had behavioural problems, a significant negative correlation between BPD symptoms and maternal warmth (i.e., enthusiastic, encouraging, cheerful, appropriately responding to needs) was reported (Harvey, Stoessel, & Herbert, 2011). However, maternal BPD symptoms did not uniquely predict low maternal warmth in multiple regression analysis controlling for comorbid psychopathology (including other personality disorders, anxiety, depression, and substance abuse). Relatedly, adolescents did not perceive their mothers with BPD as less “emotionally warm” (e.g., whether parents hugged them or whether they felt their parents wanted to be together with them) than did adolescents of control mothers with depression, cluster C personality disorders, or no mental health condition (Barnow et al., 2006). Similarly, in two community samples, adolescent-perceived maternal “warmth” (no specific description reported) was not significantly correlated with maternal BPD symptoms either in unadjusted associations (Reinelt et al., 2014) or following control for demographic variables (Herr, Hammen, & Brennan, 2008).

**Hostility/negativity/overreactivity/harsh punishment.** Six studies largely converge in suggesting that mothers with BPD may be more hostile or negative than control mothers. Elliot et al. (2014) found that mothers with BPD did not show a significant group difference from HCs regarding hostility towards their infants; however, BPD symptom severity was significantly positively correlated with raised levels of maternal hostile/reactive behaviours. Newman et al. (2007) found that mothers with BPD scored significantly higher on the “hostility” subscale of a general psychopathology measure than HCs; however while they were more likely to display “slight” or “covert” hostility towards their infant, these differences did not reach

significance. Harvey et al. (2011) reported a significant correlation between maternal BPD symptoms and displays of “negative affect” (i.e., irritability, annoyance, frustration, anger) and between maternal BPD symptoms and self-reported “overreactivity” (i.e., noticeably frustrated or angry). However, maternal BPD symptoms no longer predicted negative affect or overreactivity in multiple regression analysis following control for comorbid psychopathology (including other personality disorders, anxiety, depression, and substance abuse). In two studies with adolescent offspring, maternal BPD (Frankel-Waldheter, Macfie, Strimpfel, & Watkins, 2015) and maternal BPD symptoms (Herr et al., 2008) were significantly associated with offspring-perceived “maternal hostility” (Herr et al., 2008), and researcher-observed (Frankel-Waldheter et al., 2015) “maternal hostility,” and this association remained following control for maternal depression (Herr et al., 2008). Similarly, in a high-risk community cohort study of adolescent girls, offspring-perceived “harsh punishment” was significantly associated with maternal BPD symptoms following control for depression and alcohol use (Zalewski et al., 2014).

***Rejection and laxness.*** Two studies assessed maternal rejection. Barnow et al. (2006) found that mothers with BPD did not significantly differ in adolescent offspring-perceived “rejection” from mothers with depression, cluster C personality disorders, or no mental health condition. Conversely, Reinelt et al. (2014) reported a significant positive correlation between maternal BPD symptoms and adolescent-perceived maternal rejection. One study assessed maternal laxness. Harvey et al. (2011) found that maternal BPD symptoms were positively associated with maternal self-reported laxness. This association remained significant in multiple regression analysis controlling for comorbid psychopathology (i.e., other personality disorders, anxiety, depression, and substance abuse).

***Emotion recognition.*** Two studies assessed emotion recognition. Elliot et al. (2014) found that mothers with BPD were significantly poorer than HCs at “infant emotion recognition.” In particular, neutral infant expressions were more often perceived as sad (Elliot et al., 2014). Another study looked at discrepancies between mother-reported and researcher observed infant expressions (Whalen, Kiel, Tull, Latzman, & Gratz, 2015). Findings showed that as maternal BPD symptom severity increased, there was greater convergence between mother-reported and observed infant anger. This relationship increased in strength as observed infant anger increased. As healthy mothers typically under report infant negative affect, the authors suggest mothers with BPD may have heightened sensitivity to infant anger. This pattern of convergence was also seen for mother-reported and researcher observed infant fear expression, but only with behavioural symptoms of BPD (e.g., fear of abandonment, impulsivity).

***Maternal representations and perceptions of offspring.*** Four studies assessing maternal perceptions of their offspring (in fairly divergent ways - see Table 2) yielded inconsistent findings. Elliot et al. (2014) reported that mothers with BPD rated their infants significantly higher on a “difficult child” measure in comparison to HCs. In contrast, Newman et al. (2007) found no significant difference between mothers with BPD and HCs on their rating of their infant as “difficult.” Crittenden and Newman (2010) reported that mothers with BPD had “balanced” representations of their perceptions, feelings, motives and interpretation of their relationship with their 3-36 month-old infants, and did not significantly differ from healthy control mothers. Schacht, Hammond, Marks, Wood, and Conroy (2013) found that mothers with BPD were significantly less likely than HCs to use mind-related descriptors (e.g., interests,

imagination) of their children, and suggested that this might indicate a reduced capacity for mentalisation in mothers with BPD.

***Perceptions of parenthood.*** In two infant studies, mothers with BPD reported significantly higher parenting stress and distress (Elliot et al., 2014; Newman et al., 2007), and significantly lower parenting satisfaction (Newman et al., 2007) and efficacy (Elliot et al., 2014; Newman et al., 2007) than HCs. Increased distress was significantly correlated with increased parental dissatisfaction and efficacy, less sensitive parenting (Newman et al., 2007), and mother's emotional dysregulation (Elliot et al., 2014). One study explored mothers' perinatal experiences (Blankley, Galbally, Snellen, Power, & Lewis, 2015). Of the mothers with BPD, 31% reported pregnancy as traumatic, 12% anticipated the delivery as traumatic, and 31% made requests for an early delivery due to distress associated with the pregnancy. Additionally, 38% were reported as having low levels of antenatal care, by not following care recommendations or having erratic antenatal attendance (<70% attendance).

## **2) Mother-Offspring Interaction Dynamics**

This section includes studies assessing mother-offspring interactions, role-reversal, mother-infant communication, and infant/child behaviour (see Table 3).

***Mother-offspring interactions.*** Eight studies converge in indicating maladaptive interactions between mothers with BPD and their offspring. Three infant studies reported that mothers with BPD had significantly less "satisfying/engaged," or more "difficult" or "dysfunctional" mother-child interactions according to observation (Crandell et al., 2003; Newman et al., 2007) and maternal self-report (Elliot et al., 2014) in comparison to HCs. White, Flanagan, Martin, and Silvermann (2011) found that mothers with BPD (without co-morbid major depressive disorder, MDD)

demonstrated significantly less imitation, and mothers with BPD (and BPD/MDD) engaged in significantly less smiling, touching, and playing with their infants than mothers with MDD and HCs. Hobson et al. (2009) observed that mothers with BPD displayed significantly greater “frightened/disoriented” (i.e., fearful, hesitant) interaction with their infants than mothers with depression or HCs. They also displayed more “withdrawal” (e.g., interacting silently) with their infants than depressed and HC mothers, though these differences did not quite reach significance. Wilson and Durbin (2012) found a significant negative correlation between maternal borderline symptoms and mother’s response to their child’s “bids for attention” (the same was seen for mothers with paranoid, antisocial or histrionic personality disorder but not for other personality disorders). In comparison to children of HCs, children of mothers with BPD had significantly lower (i.e., more negative) mother-child relationship expectations (Macfie & Swan, 2009). In the one study of mother-adolescent interactions, mothers with BPD were found to be lacking in “validation,” “engagement,” and “agreement” regarding their offspring’s opinions; were significantly more likely than HCs to over-personalise in disagreements; and significantly more likely to pressure their child to agree without a rational explanation (Frankel-Waldheter et al., 2015).

***Role-reversal.*** Three studies assessed mother-child role-reversal with mixed results. Macfie, Swan, Fitzpatrick, Watkins, and Rivas (2014) found that maternal preoccupied/unresolved attachment pattern (70% of mothers had a diagnosis of BPD) was significantly correlated with children’s narratives of role-reversal. Similarly, controlling for maternal depression, Macfie and Swan (2009) found that the stories of children of mothers with BPD were significantly more likely to describe mother-child role-reversal than the stories of children of HCs. Hobson et al. (2009), however, found

no significant difference between observations of “role/boundary confusion” with dyads of mothers with BPD, depression or HCs.

***Mother-infant communication.*** Three studies considering communication patterns between mothers with BPD and their infant offspring yielded mixed results (Delavenne, Gratier, Devouche, & Apter, 2008; Hobson et al., 2009; White et al., 2011). White et al. (2011) reported that mothers with BPD did not significantly differ from mothers with depression or HCs in “vocalisations” (i.e., percentage of time spent vocalising) when interacting with their infants. Similarly, Delavenne et al. (2008) found no significant difference in overall frequency of vocalisations, however, mothers with BPD engaged in significantly fewer phrases and longer end-of-phrase pauses in interactions with their infants in comparison to HCs. Hobson et al. (2009) found that mothers with BPD displayed significantly greater “disruptive affective communication” (i.e., conflicting emotional cues, unresponsive to infant emotion) in interactions with their infants in comparison to depressed and HCs.

***Infant/child behaviour.*** Five studies considering infant/child engagement or interaction, whilst not fully converging, on the whole suggest less engaged infant behaviour. Two studies found that infants of mothers with BPD were significantly “less involved” (Newman et al., 2007), “less available for positive engagement” (Hobson et al., 2005), and “less responsive to their mother’s attempts to engage with them” (Hobson et al., 2005) than infants of HC mothers. In contrast, Crandell et al. (2003) report that infants of mothers with BPD were no different in their availability for positive engagement both during and after a still-face paradigm task compared with HCs. Similarly, Wilson and Durbin (2012) found that children of mothers with BPD symptoms were not significantly different to children of healthy mothers in their response to their mother’s bids for attention.

Infants of mothers with BPD appear to demonstrate significantly less eye contact than infants of HC mothers, especially during stressful situations such as the still-face paradigm (Crandell et al., 2003; Hobson et al., 2005). White et al. (2011) only reported greater gaze aversion in infants of mothers with BPD co-morbid with MDD compared with HCs (White et al., 2011). Significantly fewer instances of smiling (White et al., 2011), lowered mood state (Hobson et al., 2005), and lowering of affect (Crandell et al., 2003) have also been observed in infants of mothers with BPD in comparison to HCs.

### **3) Offspring Outcomes**

Studies assessed a range of psychopathological (i.e., BPD and BPD symptoms, depression, internalising and externalising problems) and psychosocial (i.e., self-esteem difficulties, interpersonal difficulties, home difficulties and stability) outcomes of offspring of mothers with borderline personality pathology (see Table 4).

#### **Psychopathology outcomes.**

***Borderline personality disorder.*** Five studies all indicated a significant association between maternal BPD/BPD symptoms and offspring BPD symptoms (Barnow et al., 2013; Cheng et al., 2011; Conway, Hammen, & Brennan, 2015; Stepp, Olino, Klein, Seeley, & Lewinsohn, 2013) and disorder (Weiss et al., 1996). Weiss et al. (1996) reported that children of mothers with BPD were significantly more likely to have a diagnosis of BPD than children of mothers with other personality disorders. In a community-based study, Barnow et al. (2013) found that maternal BPD symptoms (and sub-threshold BPD symptoms) significantly predicted offspring BPD symptoms five years later (at age 20). In a high-risk community cohort spanning 4 time-points (14, 18, 24 & 30 years of age), Stepp et al. (2013) found that maternal history of BPD predicted offspring BPD symptoms at age 30, following control for

offspring gender, early offspring psychopathology, parental education and other parental psychopathology. However, this association no longer reached significance ( $p=.069$ ) following additional control for offspring MDD at Time 3 and Time 4. Also in a high-risk community sample, Conway et al. (2015) found that maternal BPD symptoms were significantly associated with offspring BPD symptoms at age 20 but did not influence offspring symptoms over and above other risk factors (e.g., maternal externalising, offspring internalising). Finally, in a large study of high school students, Cheng et al. (2011) found that maternal BPD traits significantly positively correlated with adolescent offspring BPD traits, following control for family income, parental relationship and parental rearing behaviour.

***BPD symptoms and related features.*** A number of studies assessed associations between maternal BPD and offspring individual BPD symptoms (e.g., emotion dysregulation, unstable self-image and identity, suicide ideation) or related features (i.e., features which have a strong association with BPD, such as insecure attachment).

Four studies suggest that maternal borderline personality pathology is significantly associated with offspring emotional dysregulation across age groups operationalized in various ways (see Table 4). “Low soothability” in infancy (White et al., 2011); “boundary confusion between self/fantasy or reality/fantasy” (Macfie & Swan, 2009) and “self-regulation” (Macfie et al., 2014) in childhood, were all significantly associated with maternal BPD, and “low self-control” and “negative affectivity” in adolescence (Zalewski et al., 2014) were significantly associated with maternal BPD symptoms. Alternatively, Gratz et al. (2014) did not find a main effect of maternal BPD symptoms on infant emotion dysregulation (but see later section on

mechanisms of transmission for explication of the indirect association between maternal BPD and infant emotion dysregulation in this study).

Two studies examined aspects of offspring's self-identity. Macfie and Swan (2009) reported that children of mothers with BPD had significantly poorer self-representations (incongruent and shameful) than children of HCs. In a study examining gender identity disorder (GID), Marantz and Coates (1991) found that mothers of boys with GID were significantly more likely to have a diagnosis of BPD than mothers of boys without GID.

Children of mothers with BPD reported significantly more suicide ideation/plans and death wishes, and more suicide attempts (though this difference was non-significant) than children of HCs (Barnow et al., 2006).

Four studies indicated that maternal BPD/BPD symptoms (or BPD/MDD) is associated with offspring insecure attachment (compared to HCs) in infancy (Gratz et al., 2014; Hobson et al., 2005), childhood/adolescence (Abela et al., 2005) and adolescence (Herr et al., 2008). In particular, children's insecure attachment patterns were categorised as disorganised (Hobson et al., 2005), and fearful after controlling for maternal lifetime depression symptoms (Herr et al., 2008).

***Depression.*** Three studies converge in indicating that children and adolescents of mothers with BPD are at greater risk of developing depression (Abela et al., 2005; Barnow et al., 2006; Herr et al., 2008). The extent to which this association is independent of maternal depression however remains unclear. Abela et al. (2005) found significantly increased depression in offspring of mothers with BPD/MDD compared to offspring of mothers with MDD alone. Barnow et al. (2006) reported offspring of mothers with BPD were more likely to have depressive symptoms than offspring of mothers with depression (although this difference did not reach

significance,  $p < .10$ ). Herr et al. (2008) found that youth depression was significantly associated with maternal BPD symptoms, but not after adjustment for maternal depression.

***Internalising/emotional problems.*** Studies pertaining to offspring internalising/emotional problems suggest an association with maternal BPD (in clinical populations). Barnow et al. (2006) reported significantly increased “emotional problems” in adolescents of mothers with BPD compared to adolescents of mothers with depression or no psychopathology. Jellinek, Bishop, Murphy, Biederman, and Rosenbaum (1991) found that significantly more parents (78% mothers) with BPD had children with high total paediatric symptom checklist scores (representing combined internalising/externalising problems: attention, emotion, behaviour and somatic difficulties) than parents with other axis II disorders or HCs. In contrast, in a sample of children with internalising or externalising disorders, Bertino, Connell, and Lewis (2012) found no significant association between parental (80% mothers) borderline traits and child’s “internalising” problems.

***Externalising/behavioural problems.*** Four studies indicated that offspring of mothers with BPD are at an increased risk of behavioural problems or externalising symptoms (Barnow et al., 2006; Bertino et al., 2012; Jellinek et al., 1991; Weiss et al., 1996). Barnow et al. (2006) reported significantly more attention, delinquency, and aggression problems in adolescents of mothers with BPD in comparison to adolescents of HCs. Bertino et al. (2012) found a significant positive correlation between maternal borderline personality symptoms and both child and adolescent externalising symptoms. In comparison to children of mothers with other personality disorders, Weiss et al. (1996) reported significantly more cases of attention deficit

hyperactivity disorder and disruptive behaviour disorder in children of mothers with BPD.

**General psychopathology.** Barnow et al. (2013) reported that maternal BPD symptoms (and depression), assessed when their offspring were aged 15 years old, significantly predicted offspring general psychopathology (assessed using the Symptom Checklist-Revised- SCL-90-R) 5 years later. Barnow et al. (2006) reported that adolescents of mothers with BPD demonstrate higher levels of physical symptoms compared to children of healthy mothers or mothers with depression.

**Psychosocial outcomes.**

**Difficulties with self-esteem.** Abela et al. (2005) found that maternal BPD/MDD was significantly associated with offspring self-criticism, but not offspring self-esteem in children and adolescents of mothers with BPD. Conversely, Barnow et al. (2006) found significantly lower levels of self-esteem in adolescents of mothers with BPD compared to children of mothers with depression, Cluster C personality disorders, or no mental health condition. One study reported that adolescents of mothers with BPD had significantly higher “harm avoidance” (i.e., fearful, excessive worrying) scores in comparison to children of mothers with depression or HCs (Barnow et al., 2006).

**Interpersonal difficulties.** Three studies indicate that offspring of mothers with BPD have difficulties with mental state understanding and social interactions. Schacht et al. (2013) found that children of mothers with BPD demonstrated significantly poorer emotional labelling and understanding of causes of emotion in comparison to children of HCs, even following adjustment for maternal depression. Barnow et al. (2006) reported that scores on the social problem scale were significantly elevated in adolescents of mothers with a BPD diagnosis in comparison to adolescents of mothers

with depressive disorder, cluster C personality disorders, or no mental health condition. Herr et al. (2008) found that maternal BPD symptoms remained significantly associated with adolescent offspring's poor self-perception of the ability to make "close friendships" and be "social accepted," even after controlling for maternal depression.

***Home difficulties and stability.*** Two studies show some evidence that family dynamics are affected in the families of mothers with BPD. Feldman et al. (1995) found that mothers with BPD reported significantly lower family "cohesion" and "organisation" than mothers with other personality disorders. Both adolescent-reported "family stress" and mother-reported "chronic relationship stress" were significantly correlated with maternal BPD symptoms in a community sample, even after controlling for maternal depression (Herr et al., 2008). Children and adolescents of mothers with BPD experienced significantly greater instability, such as frequent changes in school and household composition, than children of mothers with other personality disorders (Feldman et al., 1995). They were also significantly more likely to witness maternal or paternal suicide attempts and be exposed to parental alcohol/drug abuse (Feldman et al., 1995).

***General impairment.*** Two studies assessed general impairment: one with newborns (Blankley et al., 2015), the other with children/adolescents (Weiss et al., 1996). Blankley et al. (2015) found that newborn infants of mothers with BPD were significantly more likely to have been born preterm (<37 weeks), have required resuscitation at birth, or required referral to special care nursing facilities than control mothers without BPD. They were also significantly more likely to have APGAR scores of less than 7 (an assessment of appearance, pulse, grimace, activity and respiration levels of newborn infants) than infants of control mothers. Weiss et al.

(1996) found that children/adolescents of mothers with BPD had significantly higher general impairment in areas of home, school and social life in comparison to children of mothers with non-borderline personality disorder.

#### **4) Potential Mechanisms Underpinning Transmission of Vulnerability from Mother to Offspring**

Six studies have utilised meditation (and/or moderation) analyses to statistically test potential mechanisms contributing to associations between maternal BPD (or core features of maternal BPD) and negative offspring outcomes.

##### ***Maladaptive parenting as a potential mediator or link in a causal chain.***

Two studies (Macfie et al., 2014; Reinelt et al., 2014) considered maladaptive parenting as a potential mediator of the transmission of vulnerability from mother to child. Using a prospective community-based family cohort, Reinelt et al. (2014) found that maladaptive mother-child interactions (represented by a latent variable comprising perceived overprotective and rejecting parenting style and high mother-child discrepancies regarding child's internalising problems) mediated the longitudinal transmission of BPD symptoms from mother to adolescent. Maladaptive mother-child interactions also mediated the relationship between maternal BPD and individual symptoms of impulsiveness, difficulties identifying and describing feelings, and self-esteem.

In a cross-sectional study of young children, Macfie et al. (2014) examined the relationship between mothers' Adult Attachment Interview (George, Kaplan, & Main, 1984) representations (70% of the study mothers had a diagnosis of BPD), mother's observed parenting, and offspring narratives regarding fear of abandonment. Mother's parenting (i.e., a composite of sensitivity, autonomy support and hostility) significantly mediated the relationship between maternal preoccupied/unresolved

attachment and offspring fear of abandonment (i.e., the association between mothers preoccupied unresolved attachment and offspring's fear of abandonment was partly explained by lower levels of parental sensitivity and autonomy support and higher levels of parental hostility).

Examining parenting as an exogenous (rather than mediating) risk factor in a causal chain, Frankel-Waldheter et al. (2015) investigated whether mother's lack of promotion and inhibition of autonomy (i.e., independence) and relatedness (i.e., close relationships) was associated with adolescent outcomes via maternal borderline pathology. Maternal borderline pathology mediated the relationship between lack of promotion of autonomy and relatedness (and inhibition of autonomy and relatedness) and adolescent affective instability and self-harm. Additionally maternal borderline pathology mediated the relationship between maternal inhibition of (and lack of promotion of) autonomy and relatedness, and adolescent internalising (e.g., anxious depression, withdrawn depression) and externalising symptoms (e.g., aggression). The authors concluded that a lack of maternal promotion of offspring independence and close relationships might underlie maternal borderline features and their effect on offspring outcome. It should be noted, however, that this study was cross-sectional, and whilst difficulties with autonomy and relatedness may contribute towards the development of BPD, it is perhaps more plausible to hypothesise that parenting behaviours mediate the association between maternal BPD and offspring outcome.

***Maternal emotional dysfunction as a mediator.*** In a cross-sectional study, Gratz et al. (2014) reported a significant indirect association between clinically relevant levels of maternal BPD symptoms and infant emotional regulation (ER) difficulties via maternal emotional dysfunction (i.e., mothers with BPD were more likely to experience emotional dysfunction, which in turn increased risk of infant

emotional dysregulation). More specifically, maternal emotion regulation difficulties mediated the association between maternal BPD symptoms and expressivity-related indicators of infant ER difficulties (e.g., intense emotional expressions). Further, maternal emotional intensity/reactivity facilitated an indirect effect of maternal BPD symptoms on lower infant self-focused emotional regulation (e.g., self-stimulation) only in infants with an insecure-resistant attachment relationship.

*Offspring characteristics as mediators or moderators of parenting and offspring outcomes.* In a cross-sectional study of 6-14 year-old offspring, Abela et al. (2005) demonstrated that offspring cognitive and interpersonal vulnerabilities (i.e., ruminative response style, negative attribution style, dysfunctional attitudes, self-criticism, excessive reassurance seeking and insecure attachment style) partially mediated the relationship between parental BPD (84% mothers) and offspring current depressive symptoms. As highlighted by the authors, cognitive vulnerabilities only partly mediated the association, indicating that other factors not assessed in the study, such as emotional dysregulation (Gratz et al., 2014), may play a role in the association between parental BPD and offspring depression.

Within a community sample of adolescent girls, Zalewski et al. (2014) explored whether adolescent temperament (i.e., negative emotionality and low self-control) moderated the association between maternal BPD symptoms and parenting behaviours (i.e., control through guilt, lack of acceptance of individuation, harsh punishment). No significant moderating effect was found (i.e., the parenting patterns of mothers with BPD did not vary according to the temperament of their adolescent offspring).

## Discussion

This systematic review examines the parenting and outcomes experienced by offspring of mothers with borderline personality pathology. It adds to the extant literature by highlighting the difficulties faced by mothers with both subsyndromal and syndromal levels of BPD, by exploring the outcomes of offspring from infancy to young adulthood, and by elucidating the potential mechanisms underpinning the transmission of vulnerability from mother to child. Before we summarise the main findings and contextualise within the extant literature, we consider the methodological limitations of the included studies and hence the limitations of the current review.

First, there was a degree of heterogeneity across studies in the operationalisation of parenting constructs and offspring outcomes. For example, insensitive parenting was referred to variously as: “maternal sensitivity” (Crandell et al., 2003), “insensitivity” (Kiel et al., 2011), “intrusive sensitivity” (Hobson et al., 2005) and “intrusiveness/negativity” (Hobson et al., 2009). Some studies combined constructs in composite variables; for example, Crandell et al. (2003) amalgamated insensitivity, warmth/hostility, and rejection/responsiveness into one construct. In contrast, other studies reported associations with individual construct measures. As such, any observed variance in results across studies may be somewhat attributable to study methodology. There was also heterogeneity in study method design (i.e., observational studies, mother-reports, adolescent-report, experimental designs). Whilst this may account for some variance, there were several instances of consistent findings across study types. For example studies converged in finding “difficult mother-child interactions” when using the still-face paradigm (Crandell et al., 2003), mother-report (Elliot et al., 2014), and mother-child play observation (Newman et al., 2007). Notably, the heterogeneity across studies in terms of definitions of parenting

constructs and study design precluded a meta-analytic synthesis of the results. This highlights the future need for a more systematic approach to the operationalisation of specific parenting behaviours emitted more or less often by mothers with BPD (Zalewski & Lengua, 2012).

Second, participant selection criteria differed across studies. Mothers with BPD were not always the index sample (e.g., Bertino et al., 2012; Marantz & Coates, 1991), making comparisons across some studies difficult. Bertino et al. (2012), for example, selected their sample on the basis of offspring emotional and behavioural difficulties, likely confounding the observed associations with maternal parenting behaviours. In addition, a number of studies used healthy control groups only as comparators. Consequentially, we could not always ascertain whether certain parenting characteristics (and offspring outcomes) were specific to mothers with BPD or reflective of maternal psychopathology in general. However, when clinical comparisons were made, BPD specific associations were often indicated. Furthermore, the method of assessment of mother's BPD diagnosis differed across studies. To ensure that we presented a comprehensive review of the literature, we included studies exploring BPD symptoms in addition to those examining BPD diagnoses. Thus, some of our study findings referred to BPD on a subsyndromal level reducing clinical relevance (e.g., Kiel et al., 2011). Nevertheless, it is noteworthy that findings across clinical and non-clinical samples often converged within constructs, e.g., overprotection (Barnow et al., 2006; Elliot et al., 2014; Reinelt et al., 2014; Zalewski et al., 2014), and offspring insecure attachment (Abela et al., 2005; Gratz et al., 2014; Herr et al., 2008; Hobson et al., 2005). The age of offspring also varied with some samples crossing developmental stages, such as childhood and adolescence (e.g., Abela et al., 2005; Feldman et al., 1995), making it difficult to interpret whether

findings were generalisable or age specific. Future studies with repeated intra-individual assessments may help identify differential age effects.

Third, in some cases there were too few studies in respective domains (e.g., maternal emotion recognition, rejection) to make inter-study comparisons or draw firm conclusions. Given the potential mediating role of poor maternal emotion recognition (Gratz et al., 2014) and parental rejection (Reinelt et al., 2014) in the transmission of vulnerability from mother to offspring, further investigation of these domains is indicated.

Fourth, the quality assessment showed a low-moderate risk of outcome/exposure bias, in particular with respect to researchers and coders not being blind to the mother's diagnosis. Further, our results may have been subject to publication bias due to the "file drawer" problem, however, given that non-significant findings are also of interest in this study population the likelihood of a results bias is potentially reduced.

Fifth, the review excluded child outcomes that required external intervention, such as the child being removed from the home. Whilst some children of mothers with BPD will experience neglect and/or abuse (e.g., Kauppi, Vanamo, Karkola, & Merikanto, 2012; Perepletchikova, Ansell, & Axelrod, 2012) child maltreatment is not restricted to mothers with BPD, or indeed mothers with other psychiatric diagnoses, and additional risk factors may be involved such as partner/family violence, serious marital problem, low socioeconomic status or poor education (Chaffin, Kelleher, & Hollenberg, 1996; Department of Human Services, 2013).

Finally, the majority of the studies, with a few exceptions (Barnow et al., 2013; Reinelt et al., 2014; Stepp et al., 2013), were cross-sectional in design. Whilst some convergence of findings was seen between cross-sectional and longitudinal studies (e.g., offspring BPD), cross-sectional studies cannot take into account intra-individual

development over time (Crone & Elzinga, 2015), making it difficult to draw conclusions regarding temporal precedence and aetiological mechanisms.

### **The Parenting Characteristics of Mothers with BPD/BPD Symptoms**

Accepting some inconsistencies across studies, we found that mothers with BPD/BPD symptoms appear less sensitive, more intrusive, more overprotective, more hostile, show less engagement, and are more likely to have maladaptive interactions (such as role-reversal, boundary confusion, fearful/hesitant behaviour) with their offspring than control mothers. This indicates that mothers with BPD may demonstrate inconsistent parenting characterised by over-involvement such as overprotection and inhibiting autonomy (e.g., Barnow et al., 2006; Elliot et al., 2014; Zalewski et al., 2014) on the one hand, and disengagement and hostility on the other (e.g., Frankel-Waldheter et al., 2015; Herr et al., 2008; White et al., 2011). A similar pattern of under-and over involvement was previously hypothesised to be unique to mothers with BPD (Stepp et al., 2012), and has been observed in subsequent studies in this review (e.g., Frankel-Waldheter et al., 2015; Reinelt et al., 2014). Furthermore, recent studies highlight a reluctance to promote independence as an additional parenting behaviour that may be characteristic of mothers with BPD (Frankel-Waldheter et al., 2015) or mothers with BPD symptoms (Zalewski et al., 2014).

While our review indicates various parenting problems for some mothers with borderline personality pathology, findings do not suggest that these mothers lack a desire to care for their child. Indeed, studies measuring overprotection found that mothers reported a concern for their child's health and safety (Elliot et al., 2014; Reinelt et al., 2014; Zalewski et al., 2014). This suggests that mothers want to parent well but may lack the necessary "tools" to effectuate optimal parenting. The parenting stress and lack of efficacy that mothers with BPD report (Elliot et al., 2014; Newman

et al., 2007) further highlights the parenting difficulties mothers with BPD face. These likely arise from a combination of factors including (but not limited to) individual BPD symptoms such as emotional dysregulation (Gratz et al., 2014), symptom severity (e.g., Elliot et al., 2014), comorbid psychopathology (e.g., Abela et al., 2005), and mothers' own childhood experiences (Bandelow, 2005; Zanarini et al., 1997). Precisely how parenting strategies unravel between mother-offspring dyads (i.e., frequency, duration and magnitude of under/over involvement) requires further explication.

From the limited studies utilising clinical control groups (or statistically controlling for other psychopathology) there is some evidence that maternal borderline personality pathology is specifically associated with parenting behaviours including overprotection (Barnow et al., 2006), control through guilt (Zalewski et al., 2014), maternal hostility (Herr et al., 2008), and fearful/hesitant behaviour (Hobson et al., 2009). The potential ramifications of such parenting strategies are high, including offspring anxiety (van der Bruggen, Stams, Bögels, & Paulussen-Hoogeboom, 2010), behavioural problems (Gere, Villabø, Torgersen, & Kendall, 2012), social anxiety (Spokas & Heimberg, 2009) and BPD (Bezirgianian, Cohen, & Brook, 1993); many of which were observed within the review studies (see below). More studies with clinical control groups (e.g., depression, bipolar disorder) are now needed to further clarify the specificity of these and other parenting behaviours to maternal BPD.

### **Offspring Outcomes Across Developmental Domains**

A range of psychopathological and psychosocial outcomes for offspring of mothers with BPD were observed across several stages of development, tentatively indicating that the negative effects on offspring may be enduring.

Studies, on the whole, reported associations between maternal BPD/BPD symptoms and offspring individual BPD symptoms/features (e.g., emotional dysregulation, insecure attachment, depression, internalising and externalising problems, and interpersonal problems) in infancy (Crandell et al., 2003; Gratz et al., 2014; Hobson et al., 2005; White et al., 2011), childhood/adolescence (Abela et al., 2005; Barnow et al., 2006; Jellinek et al., 1991; Macfie & Swan, 2009), and adolescence (Herr et al., 2008; Zalewski et al., 2014). This indicates that these difficulties may manifest across several stages of development, heightening risk of future psychopathology (Winsper & Wolke, 2014) and BPD in particular (Carlson, Egeland, & Sroufe, 2009; Crowell et al., 2009). Indeed, studies were consistent in reporting a significant positive association between maternal and offspring BPD diagnosis in children (Weiss et al., 1996), and also between maternal BPD/BPD symptoms and offspring BPD symptoms in adolescent (Barnow et al., 2013; Cheng et al., 2011) and adult (Stepp et al., 2013) populations. Despite ongoing controversy regarding the diagnosis of BPD in children and adolescents, recent literature demonstrates that the BPD diagnosis in youth is of comparable reliability and validity to the adult diagnosis (Kaess, Brunner, & Chanen, 2014; Winsper et al., 2016), shows similar levels of stability (Winsper et al., 2015), and is of great clinical relevance (Newton-Howes, Clark, & Chanen, 2015). How psychopathological problems evolve in individual offspring of mothers with BPD, however, requires corroboration with prospective repeated assessment studies.

Collectively, studies demonstrate that offspring of mothers with BPD are a high-risk population who are at increased risk of developing psychosocial and mental health problems across developmental domains. Potential difficulties may even be evident at the perinatal stage (Blankley et al., 2015), though research pertaining to

pregnant mothers with BPD is currently scant. The main findings from the review may be interpreted within a developmental psychopathology framework e.g., the biosocial developmental model (BDM; Crowell et al., 2009). Infants of mothers with BPD may inherit a biological vulnerability to emotionality/negative affectivity (Crandell et al., 2003; Hobson et al., 2005), which is potentiated across development by environmental risk factors (Crowell et al., 2009). Aside from potential heritability and other possible risk factors (e.g., in utero stress or substance exposure, sharing of a toxic environment), parental invalidation has been hypothesised as one mechanistic factor underpinning the transmission of BPD from mother to child (Crowell et al., 2009; Stepp et al., 2012). Recent mediational studies indicate that the transmission of BPD symptoms, either collectively or individually (e.g., emotional dysregulation) may be partly explained by an increased risk of insensitive, rejecting and hostile parenting (Macfie et al., 2014; Reinelt et al., 2014), and maternal emotional dysfunction (Gratz et al., 2014). In this way maladaptive transactions between offspring and mother, and other individuals such as peers (Barnow et al., 2006) and family members (Feldman et al., 1995) may continue. Over time these transactions may give rise to increasing levels of emotional (Barnow et al., 2006), interpersonal (Herr et al., 2008), cognitive (Macfie & Swan, 2009) and behavioural (Bertino et al., 2012) dysregulation, until they finally coalesce into a mental disorder (Crowell et al., 2009). One likely outcome is the development of BPD in childhood or adolescence (Cheng et al., 2011; Reinelt et al., 2014; Stepp et al., 2013; Weiss et al., 1996). As BPD in adolescence is predictive of BPD in adulthood (Winsper et al., 2015), the transmission of BPD (as described above) may continue across successive generations (Stepp et al., 2012). Other outcomes are also likely such as emotional and behavioural dysregulation (Barnow et al., 2006; Bertino et al., 2012), which have been found to

predict multifinial outcomes (Abela et al., 2005; Crowell et al., 2009). Of note, resilience factors (e.g., secure attachment) may prevent transmission of symptoms from mother to child (Gratz et al., 2014). Recent research with adolescents found attachment security served as a buffer by enhancing positive emotion regulation strategies, whilst decreased use of positive emotion regulation strategies mediated the relationship between attachment insecurity and adolescent BPD features (Kim, Sharp, & Carbone, 2014).

### **Research and Clinical Implications**

Not all children of mothers with BPD will go on to develop BPD or other psychopathology. However, offspring of mothers with BPD present an ideal cohort for studying the development of BPD, as these samples will yield a higher proportion of individuals actually developing the disorder (Stepp et al., 2012). This will allow for the prospective examination of processes underpinning the development of clinically relevant levels of BPD from pregnancy onwards (De Genna, Feske, Larkby, Angiolieri, & Gold, 2012; Winsper, Wolke, & Lereya, 2014). By investigating the perinatal period, findings could help elucidate early risk factors e.g., prematurity and poorer health of child at birth (Blankley et al., 2015), which may impact on child outcomes and/or mother-child relationships. Systemic factors, such as poverty, partner aggression and disordered extended family members, and factors that may provide resilience against offspring psychopathology (Bartsch, Roberts, Davies, & Proeve, 2015; Kim et al., 2014) should also be explored. Concurrently assessing various offspring psychopathologies (e.g., BPD, depression, and substance use disorder) may help elucidate the determinants of multifinality of outcome across development.

We only identified 5 studies recruiting participants who did not speak English

as their first language making it difficult to definitively ascertain whether there are cross-cultural variations in the parenting styles of mothers with BPD. In view of the potential impact of culture on what is perceived as “adaptive” or “maladaptive” parenting (Sorkhabi & Mandara, 2013) this may be a fruitful area for future research.

Prospective studies would also provide the opportunity to study intra-dyad transactional dynamics, parenting strategies (including potential frequency, duration and magnitude of over/under involved polarised parenting behaviours) and developmental differences in parenting and child outcomes over time. Observing parenting behaviours within the framework of other factors such as the mother’s specific BPD symptoms, symptom severity and comorbid psychopathology, would help target key areas for intervention.

Our review highlights several prevention and early intervention opportunities. Insecure attachment (Macfie et al., 2014) and emotional dysregulation (Gratz et al., 2014) are important targets for both mothers with BPD and their offspring. Intervention from pregnancy onwards including dyadic infant-parent psychotherapy (Macfie et al., 2014; Wendland et al., 2014), parent skills training (Perrin, Sheldrick, McMenemy, Henson, & Carter, 2014), and mentalisation-based treatment (Bateman & Fonagy, 2010) could help prevent the intergenerational transmission of insecure attachment patterns and self-regulation problems. Intervening early could help prevent offspring from embarking on a maladaptive developmental trajectory. As BPD symptoms often become apparent in adolescence (Chanen & Kaess, 2012) and may be more responsive to treatment than in adulthood (Lenzenweger & Castro, 2005), intervention at this point would be timely. Programmes that have limited exclusions for co-morbid psychopathology (common with BPD) and that combine subsyndromal and syndromal BPD symptoms, e.g., HYPE (Chanen et al., 2009),

could provide both intervention and prevention strategies (Chanen & McCutcheon, 2013). A degree of improvement of symptoms has been previously observed with treatments such as cognitive analytic therapy (Chanen et al., 2008), dialectic behavioural therapy, and mindfulness-based training (e.g., Rossouw & Fonagy, 2012) (for a detailed review see Sharp & Fonagy, 2015). Combined therapy with mother and offspring may also help by targeting dyadic interactions (e.g., joint therapy for managing emotion dysregulation). Similarly, interventions which increase the mother's awareness of how their BPD symptoms impact on parenting behaviours and offspring outcomes (Bartsch et al., 2015; Zalewski, Stepp, Whalen, & Scott, 2015) may help further improve mother-offspring interactions. For example a lack of promotion of offspring independence could be associated with mother's fear of abandonment difficulties (Frankel-Waldheter et al., 2015; Zalewski et al., 2014). There are no current interventions specifically designed for mothers with BPD and their children, therefore, future endeavours could include the design and evaluation of tailored interventions for this cohort of mothers and children.

Whilst this review could not definitively cite many parenting behaviours as unique to mothers with BPD, the differences found between mothers with borderline personality pathology and healthy control mothers indicate many ineffectual parenting characteristics. With greater consistency of parenting constructs and BPD assessment, future research may continue to identify the specific parenting behaviours of mothers with BPD. Considering the poor outcome trajectory for some offspring of mothers with BPD, timely interventions for both mother and offspring to prevent distress and persistent functional impairment are warranted.

## References

- Abela, J. R. Z., Skitch, S. A., Auerbach, R. P., & Adams, P. (2005). The impact of parental borderline personality disorder on vulnerability to depression in children of affectively ill parents. *Journal of Personality Disorders, 19*, 68-83. doi: 10.1521/pedi.19.1.68.62177
- Agrawal, H. R., Gunderson, J., Holmes, B. M., & Lyons-Ruth, K. (2004). Attachment studies with borderline patients: A review. *Harvard Review of Psychiatry, 12*, 94-101. doi: 10.1080/10673220490447218
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders (4th ed., text revision)*. Washington, DC: Author.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (5th ed.)*. Washington, DC: Author.
- Bandelow, B., Krause, J., Wedekind, D., Broocks, A., Hajak, G., & Ruther, E. (2005). Early traumatic life events, parental attitudes, family history, and birth risk factors in patients with borderline personality disorder and healthy controls. *Psychiatry Research, 134*, 169-179. doi: 10.1016/j.psychres.2003.07.008
- Barnow, S., Aldinger, M., Arens, E. A., Ulrich, I., Spitzer, C., Grabe, H. J., & Stopsack, M. (2013). Maternal transmission of borderline personality disorder symptoms in the community-based Greifswald Family Study. *Journal of Personality Disorders, 27*, 806-819. doi: 10.1521/pedi\_2012\_26\_058
- Barnow, S., Spitzer, C., Grabe, H. J., Kessler, C., & Freyberger, H. J. (2006). Individual characteristics, familial experience, and psychopathology in children of mothers with borderline personality disorder. *Journal of the American Academy of Child & Adolescent Psychiatry, 45*, 965-972. doi: 10.1097/01.chi.0000222790.41853.b9
- Bartsch, D. R., Roberts, R. M., Davies, M., & Proeve, M. (2015). The impact of parental diagnosis of borderline personality disorder on offspring: Learning from clinical practice. *Personality and Mental Health, 9*, 33-43. doi: 10.1002/pmh.1274
- Bateman, A., & Fonagy, P. (2010). Mentalization based treatment for borderline personality disorder. *World Psychiatry, 9*, 11-15. doi: 10.1002/j.2051-5545.2010.tb00255.x
- Bender, D. S., Dolan, R. T., Skodol, A. E., Sainslow, C. A., Dyck, I. R., McGlashan, T. H., . . . Gunderson, J. G. (2001). Treatment utilization by patients with personality disorders. *The American Journal of Psychiatry, 158*, 295-302. doi: 10.1176/appi.ajp.158.2.295
- Bertino, M. D., Connell, G., & Lewis, A. J. (2012). The association between parental personality patterns and internalising and externalising behaviour problems in children and adolescents. *Clinical Psychologist, 16*, 110-117. doi: 10.1111/j.1742-9552.2012.00042.x
- Bezirgianian, S., Cohen, P., & Brook, J. S. (1993). The impact of mother-child interaction on the development of borderline personality disorder. *American Journal of Psychiatry, 150*, 1836-1842.
- Black, D. W., Blum, N., Pfohl, B., & Hale, N. (2004). Suicidal behavior in borderline personality disorder: Prevalence, risk factors, prediction, and prevention. *Journal of Personality Disorders, 18*, 226-239. doi: 10.1521/pedi.18.3.226.35445
- Black, D. W., Guntera, T., Allena, J., Bluma, N., Arndta, S., Wenmana, G., & Sieleni, B. (2007). Borderline personality disorder in male and female offenders newly

- committed to prison. *Comprehensive Psychiatry*, 48, 400-405. doi: 10.1016/j.comppsy.2007.04.006
- Blankley, G., Galbally, M., Snellen, M., Power, J., & Lewis, A. J. (2015). Borderline personality disorder in the perinatal period: Early infant and maternal outcomes. *Australasian Psychiatry*, 23, 688-692. doi: 10.1177/1039856215590254
- Carlson, E. A., Egeland, B., & Sroufe, L. A. (2009). A prospective investigation of the development of borderline personality symptoms. *Development and Psychopathology*, 21, 1311-1334. doi: 10.1017/S0954579409990174
- Chaffin, M., Kelleher, K., & Hollenberg, J. (1996). Onset of physical abuse and neglect: Psychiatric, substance abuse, and social risk factors from prospective community data. *Child Abuse and Neglect*, 20, 191-203. doi: 10.1016/S0145-2134(95)00144-1
- Chanen, A. M., Jackson, H. J., McCutcheon, L. K., Jovev, M., Dudgeon, P., Yuen, H. P., . . . McGorry, P. D. (2008). Early intervention for adolescents with borderline personality disorder using cognitive analytic therapy: Randomised controlled trial. *The British Journal of Psychiatry*, 193, 477-484. doi: 10.1080/00048670902817711
- Chanen, A. M., & Kaess, M. (2012). Developmental pathways to borderline personality disorder. *Current Psychiatry Reports*, 14, 45-53. doi: 10.1007/s11920-011-0242-y
- Chanen, A. M., & McCutcheon, L. (2013). Prevention and early intervention for borderline personality disorder: Current status and recent evidence. *The British Journal of Psychiatry*, 202, Special issue s54, 24-29. doi: 10.1192/bjp.bp.112.119180
- Chanen, A. M., McCutcheon, L. K., Germano, D., Nistico, H., Jackson, H. J., & McGorry, P. D. (2009). The HYPE Clinic: An early intervention service for borderline personality disorder. *Journal of Psychiatric Practice*, 15, 163-172. doi: 10.1097/01.pra.0000351876.51098.f0
- Chen, H., Cohen, P., Johnson, J. G., Kasen, S., Sneed, J. R., & Crawford, T. N. (2004). Adolescent personality disorders and conflict with romantic partners during the transition to adulthood. *Journal of Personality Disorders*, 18, 507-525. doi: 10.1521/pedi.18.6.507.54794
- Cheng, H. G., Huang, Y., Liu, Z., & Liu, B. (2011). Associations linking parenting styles and offspring personality disorder are moderated by parental personality disorder, evidence from China. *Psychiatry Research*, 189, 105-109. doi: 10.1016/j.psychres.2010.12.006
- Conway, C. C., Hammen, C., & Brennan, P. A. (2015). Adolescent precursors of adult borderline personality pathology in a high-risk community sample. *Journal of Personality Disorders*, 29, 316-333. doi: 10.1521/pedi\_2014\_28\_158
- Crandell, L. E., Hobson, R. P., & Patrick, M. P. H. (2003). "Still-face" interactions between mothers with borderline personality disorder and their 2-month-old infants. *British Journal of Psychiatry*, 183, 239-247. doi: 10.1192/bjp.183.3.239
- Crittenden, P. M., & Newman, L. (2010). Comparing models of borderline personality disorder: Mothers' experience, self-protective strategies, and dispositional representations. *Clinical Child Psychology and Psychiatry*, 15, 433-451. doi: 10.1177/1359104510368209

- Crone, E. A., & Elzinga, B. M. (2015). Changing brains: How longitudinal functional magnetic resonance imaging studies can inform us about cognitive and social-affective growth trajectories. *Wiley Interdisciplinary Reviews: Cognitive Science*, 6, 53-63. doi: 10.1002/wcs.1327
- Crowell, S. E., Beauchaine, T. P., & Linehan, M. M. (2009). A biosocial developmental model of borderline personality: Elaborating and extending Linehan's theory. *Psychological Bulletin*, 135, 495-510. doi: 10.1037/a0015616
- Daley, S. E., Burge, D., & Hammen, C. (2000). Borderline personality disorder symptoms as predictors of 4-year romantic relationship dysfunction in young women: Addressing issues of specificity. *Journal of Abnormal Psychology*, 109, 451-460. doi: 10.1037//0021-843X.109.3.451
- De Genna, N. M., Feske, U., Larkby, C., Angiolieri, T., & Gold, M. A. (2012). Pregnancies, abortions and births in women with and without borderline personality disorder (BPD). *Womans Health Issues*, 22, 371-377. doi: 10.1016/j.whi.2012.05.002
- Delavenne, A., Gratier, M., Devouche, E., & Apter, G. (2008). Phrasing and fragmented time in "pathological" mother-infant vocal interaction. *Musicae Scientiae, Special issue*, 47-70. doi: 10.1177/1029864908012001031
- Department of Human Services. (2013). What are the causes of child abuse? Retrieved 9/9/15, from <http://www.dhs.vic.gov.au/for-individuals/children,-families-and-young-people/child-protection/about-child-abuse/what-is-child-abuse/what-are-the-causes-of-child-abuse>
- Distel, M. A., Trull, T. J., Derom, C. A., Thiery, E. W., Grimmer, M. A., Martin, N. G., . . . Boomsma, D. I. (2008). Heritability of borderline personality disorder features is similar across three countries. *Psychological Medicine*, 38, 1219-1229. doi: 10.1017/S0033291707002024
- Elliot, R. L., Campbell, L., Hunter, M., Cooper, G., Melville, J., McCabe, K., . . . Loughland, C. (2014). When I look into my baby's eyes: Infant emotion recognition by mothers with borderline personality disorder. *Infant Mental Health Journal*, 35, 21-32. doi: 10.1002/imhj.21426
- Feldman, R. B., Zelkowitz, P., Weiss, M., Vogel, J., Heyman, M., & Paris, J. (1995). A comparison of the families of mothers with borderline and nonborderline personality disorders. *Comprehensive Psychiatry*, 36, 157-163. doi: 10.1016/S0010-440X(95)90110-8
- Frankel-Waldheter, M., Macfie, J., Strimpfel, J. M., & Watkins, C. D. (2015). Effect of maternal autonomy and relatedness and borderline personality disorder on adolescent symptomatology. *Personality Disorders: Theory, Research, and Treatment*, 6, 152-160. doi: 10.1037/per0000109
- George, C., Kaplan, N., & Main, M. (1984). *The attachment interview for adults*. Unpublished manuscript, Berkeley: University of California.
- Gere, M. K., Villabø, M. A., Torgersen, S., & Kendall, P. C. (2012). Overprotective parenting and child anxiety: The role of co-occurring child behavior problems. *Journal of Anxiety Disorders*, 26, 642-649. doi: 10.1016/j.janxdis.2012.04.003
- Grant, B. F., Chou, S. P., Goldstein, R. B., Huang, B., Stinson, F. S., Saha, T. D., . . . Ruan, W. J. (2008). Prevalence, correlates, disability, and comorbidity of DSM-IV borderline personality disorder: Results from the Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions. *Journal of Clinical Psychiatry*, 69, 533-545. doi: 10.4088/JCP.v69n0404

- Gratz, K. L., Kiel, E. J., Latzman, R. D., Elkin, T. D., Moore, S. A., & Tull, M. T. (2014). Maternal borderline personality pathology and infant emotion regulation: Examining the influence of maternal emotion-related difficulties and infant attachment. *Journal of Personality Disorders, 28*, 52-69. doi: 10.1521/pedi.2014.28.1.52
- Gratz, K. L., Kiel, E. J., Latzman, R. D., Moore, S. A., Elkin, T. D., Megason, G. C., & Tull, M. T. (2015). Complex interrelations of trait vulnerabilities in mothers and their infants. *Infancy, 1*–33. doi: 10.1111/infa.12075
- Gunderson, J. G., & Hoffman, P. D. (2005). *Understanding and treating borderline personality disorder: A guide for professionals and families*. Washington, DC: American Psychiatric Publishing inc.
- Harvey, E., Stoessel, B., & Herbert, S. (2011). Psychopathology and parenting practices of parents of preschool children with behavior problems. *Parenting: Science and Practice, 11*, 239-263. doi: 10.1080/15295192.2011.613722
- Herr, N. R., Hammen, C., & Brennan, P. A. (2008). Maternal borderline personality disorder symptoms and adolescent psychosocial functioning. *Journal of Personality Disorders, 22*, 451-465. doi: 10.1521/pedi.2008.22.5.451
- Herzog, R., Álvarez-Pasquin, M. J., Díaz, C., Del Barrio, J. L., Estrada, J. M., & Gil, Á. (2013). Are healthcare workers' intentions to vaccinate related to their knowledge, beliefs and attitudes? A systematic review. *BMC Public Health, 13*, 154. doi: 10.1186/1471-2458-13-154
- Higgins, J. P. T., & Green, S. (2011). Cochrane handbook for systematic reviews of interventions, version 5.1.0. Available from <http://www.cochrane-handbook.org>: The Cochrane Collaboration.
- Hobson, R. P., Patrick, M., Crandell, L., Garcia-Perez, R., & Lee, A. (2005). Personal relatedness and attachment in infants of mothers with borderline personality disorder. *Development and Psychopathology, 17*, 329-347. doi: 10.1017/S0954579405050169
- Hobson, R. P., Patrick, M. P. H., Hobson, J. A., Crandell, L., Bronfman, E., & Lyons-Ruth, K. (2009). How mothers with borderline personality disorder relate to their year-old infants. *British Journal of Psychiatry, 195*, 325-330. doi: 10.1192/bjp.bp.108.060624
- Jellinek, M. S., Bishop, S. J., Murphy, J. M., Biederman, J., & Rosenbaum, J. F. (1991). Screening for dysfunction in the children of outpatients at a psychopharmacology clinic. *American Journal of Psychiatry, 148*, 1031-1036. doi: 10.1176/ajp.148.8.1031
- Kaess, M., Brunner, R., & Chanen, A. (2014). Borderline personality disorder in adolescents. *Pediatrics, 134*, 782-793. doi: 10.1542/peds.2013-3677
- Kauppi, A. L. M., Vanamo, T., Karkola, K., & Merikanto, J. (2012). Fatal child abuse: A study of 13 cases of continuous abuse. *Mental Illness, 4*, 5-9. doi: 10.4081/mi.2012.e2
- Kiel, E. J., Gratz, K. L., Moore, S. A., Latzman, R. D., & Tull, M. T. (2011). The impact of borderline personality pathology on mothers' responses to infant distress. *Journal of Family Psychology, 25*, 907-918. doi: 10.1037/a0025474
- Kim, S., Sharp, C., & Carbone, C. (2014). The protective role of attachment security for adolescent borderline personality disorder features via enhanced positive emotion regulation strategies. *Personality Disorders: Theory, Research, and Treatment, 5*, 125–136. doi: 10.1037/per0000038
- Lenzenweger, M. F., & Castro, D. D. (2005). Predicting change in borderline personality: Using neurobehavioral systems indicators within an individual

- growth curve framework. *Development and Psychopathology*, *17*, 1207–1237. doi: 10.1017/S0954579405050571
- Macfie, J. (2009). Development in children and adolescents whose mothers have borderline personality disorder. *Child Development Perspectives*, *3*, 66-71. doi: 10.1111/j.1750-8606.2008.00079.x
- Macfie, J., & Swan, S. A. (2009). Representations of the caregiver-child relationship and of the self, and emotion regulation in the narratives of young children whose mothers have borderline personality disorder. *Development and Psychopathology*, *21*, 993-1011. doi: 10.1017/S0954579409000534
- Macfie, J., Swan, S. A., Fitzpatrick, K. L., Watkins, C. D., & Rivas, E. M. (2014). Mothers with borderline personality and their young children: Adult attachment interviews, mother-child interactions, and children's narrative representations. *Development and Psychopathology*, *26*, 539-551. doi: 10.1017/s095457941400011x
- Marantz, S., & Coates, S. (1991). Mothers of boys with gender identity disorder: A comparison of matched controls. *Journal of the American Academy of Child and Adolescent Psychiatry*, *30*, 310-315. doi: 10.1097/00004583-199103000-00022
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Annals of Internal Medicine*, *151*, 264–269. doi: 10.1136/bmj.b2535
- Newman, L. K., Stevenson, C. S., Bergman, L. R., & Boyce, P. (2007). Borderline personality disorder, mother-infant interaction and parenting perceptions: Preliminary findings. *Australian and New Zealand Journal of Psychiatry*, *41*, 598-605. doi: 10.1080/00048670701392833
- Newton-Howes, G., Clark, L. A., & Chanen, A. (2015). Personality disorder across the life course. *The Lancet*, *385*, 727–734. doi: 10.1016/S0140-6736(14)61283-6
- Perepletchikova, F., Ansell, E., & Axelrod, S. (2012). Borderline personality disorder features and history of childhood maltreatment in mothers involved with child protective services. *Child Maltreatment*, *17*, 182-190. doi: 10.1177/1077559512448471
- Perrin, E. C., Sheldrick, R. C., McMenamy, J. M., Henson, B. S., & Carter, A. S. (2014). Improving parenting skills for families of young children in pediatric settings: A randomized clinical trial. *JAMA Pediatrics*, *168*, 16-24. doi: 10.1001/jamapediatrics.2013.2919
- Reinelt, E., Stopsack, M., Aldinger, M., Ulrich, I., Grabe, H. J., & Barnow, S. (2014). Longitudinal transmission pathways of borderline personality disorder symptoms: From mother to child? *Psychopathology*, *47*, 10-16. doi: 10.1159/000345857
- Rossouw, T. I., & Fonagy, P. (2012). Mentalization-based treatment for self-harm in adolescents: A randomized controlled trial. *Journal of the American Academy of Child and Adolescent Psychiatry*, *51*, 1304–1313. doi: 10.1016/j.jaac.2012.09.018
- Sansone, R. A., & Sansone, L. A. (2011). Gender patterns in borderline personality disorder. *Innovations in Clinical Neuroscience*, *8*, 16–20.
- Schacht, R., Hammond, L., Marks, M., Wood, B., & Conroy, S. (2013). The relation between mind-mindedness in mothers with borderline personality disorder and mental state understanding in their children. *Infant and Child Development*, *22*, 68-84. doi: 10.1002/icd.1766

- Sharp, C., & Fonagy, P. (2015). Practitioner review: Borderline personality disorder in adolescence - recent conceptualization, intervention and implications for clinical practice. *Journal of Child Psychology & Psychiatry*, Advance online publication. doi: 10.1111/jcpp.12449
- Silberschmidt, A., Lee, S., Zanarini, M. C., & Schulz, S. C. (2015). Gender differences in borderline personality disorder: Results from a multinational clinical trial sample. *Journal of Personality Disorders*, 29, 828-838. doi: 10.1521/pedi\_2014\_28\_175
- Skodol, A. E., & Bender, D. S. (2003). Why are women diagnosed borderline more than men. *Psychiatric Quarterly*, 74, 349-360. doi: 10.1023/A:1026087410516
- Skodol, A. E., Gunderson, J. G., McGlashan, T. H., Dyck, I. R., Stout, R. L., Bender, D. S., . . . Morey, L. C. (2002). Functional impairment in patients with schizotypal, borderline, avoidant, or obsessive-compulsive personality disorder. *American Journal of Psychiatry*, 159, 276-283. doi: 10.1176/appi.ajp.159.2.276
- Sorkhabi, N., & Mandara, J. (2013). Are the effects of Baumrind's parenting styles culturally specific or culturally equivalent? In R. E. Larzelere, A. S. Morris & A. W. Harrist (Eds.), *Authoritative parenting: Synthesizing nurturance and discipline for optimal child development* (pp. 113-135). Washington DC: American Psychological Association.
- Spokas, M., & Heimberg, R. G. (2009). Overprotective parenting, social anxiety and external locus of control: Cross-sectional and longitudinal relationships. *Cognitive Therapy and Research*, 33, 543-551. doi: 10.1007/s10608-008-9227-5
- Stepp, S. D., Olino, T. M., Klein, D. N., Seeley, J. R., & Lewinsohn, P. M. (2013). Unique influences of adolescent antecedents on adult borderline personality disorder features. *Personality Disorders: Theory, Research, and Treatment*, 4, 223-229. doi: 10.1037/per0000015
- Stepp, S. D., Whalen, D. J., Pilkonis, P. A., Hipwell, A. E., & Levine, M. D. (2012). Children of mothers with borderline personality disorder: Identifying parenting behaviors as potential targets for intervention. *Personality Disorders: Theory, Research, and Treatment*, 3, 76-91. doi: 10.1037/a0023081
- Swartz, M., Blazer, D., George, L., & Winfield, I. (1990). Estimating the prevalence of borderline personality disorder in the community. *Journal of Personality Disorders*, 4, 257 - 272. doi: 10.1521/pedi.1990.4.3.257
- Torgersen, S., Lygren, S., Oien, P. A., Skre, I., Onstad, S., Edvardsen, J., . . . Kringlen, E. (2000). A twin study of personality disorders. *Comprehensive Psychiatry*, 41, 416-425. doi: 10.1053/comp.2000.16560
- Trull, T. J., Jahng, S., Tomko, R. L., Wood, P. K., & Sher, K. J. (2010). Revised NESARC personality disorder diagnoses: Gender, prevalence and comorbidity with substance dependent disorders. *Journal of Personality Disorders*, 24, 412-426. doi: 10.1521/pedi.2010.24.4.412
- van der Bruggen, C., Stams, G., Bögels, S., & Paulussen-Hoogeboom, M. (2010). Parenting behaviour as a mediator between young children's negative emotionality and their anxiety/depression. *Infant and Child Development*, 19, 354-365. doi: 10.1002/icd.665

- Weiss, M., Zelkowitz, P., Feldman, R. B., Vogel, J., Heyman, M., & Paris, J. (1996). Psychopathology in offspring of mothers with borderline personality disorder: A pilot study. *Canadian Journal of Psychiatry, 41*, 285-290.
- Wells, G. A., Shea, B., O'Connell, D., Peterson, J., Welch, V., Losos, M., & Tugwell, P. (2000). The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Unpublished manuscript. [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp)
- Wendland, J., Brisson, J., Medeiros, M., Camon-Senechal, L., Aidane, E., David, M., . . . Rabain, D. (2014). Mothers with borderline personality disorder: Transition to parenthood, parent–infant interaction, and preventive/therapeutic approach. *Clinical Psychology Science and Practice, 21*, 139-153. doi: 10.1111/cpsp.12066
- Whalen, D. J., Kiel, E. J., Tull, M. T., Latzman, R. D., & Gratz, K. L. (2015). Maternal borderline personality disorder symptoms and convergence between observed and reported infant negative emotional expressions. *Personality Disorders: Theory, Research, and Treatment, 6*, 229-238. doi: 10.1037/per0000116
- White, H., Flanagan, T. J., Martin, A., & Silvermann, D. (2011). Mother-infant interactions in women with borderline personality disorder, major depressive disorder, their co-occurrence, and healthy controls. *Journal of Reproductive and Infant Psychology, 29*, 223-235. doi: 10.1080/02646838.2011.576425
- Wilson, S., & Durbin, C. E. (2012). Parental personality disorder symptoms are associated with dysfunctional parent-child interactions during early childhood: A multilevel modeling analysis. *Personality Disorders: Theory, Research, and Treatment, 3*, 55-65. doi: 10.1037/a0024245
- Winsper, C., Lereya, S. T., Marwaha, S., Thompson, A., Eyden, J., & Singh, S. P. (2016). The aetiological and psychopathological validity of borderline personality disorder in youth: A systematic review and meta-analysis. *Clinical Psychology Review, 44*, 13-24. doi: 10.1016/j.cpr.2015.12.001
- Winsper, C., Marwaha, S., Lereya, S. T., Thompson, A., Eyden, J., & Singh, S. P. (2015). Clinical and psychosocial outcomes of borderline personality disorder in childhood and adolescence: A systematic review. *Psychological Medicine, 45*, 2237-2251. doi: 10.1017/S0033291715000318
- Winsper, C., & Wolke, D. (2014). Infant and toddler crying, sleeping and feeding problems and trajectories of dysregulated behavior across childhood. *Journal of Abnormal Psychology, 42*, 831-843. doi: 10.1007/s10802-013-9813-1
- Winsper, C., Wolke, D., & Lereya, T. (2014). Prospective associations between prenatal adversities and borderline personality disorder at 11 to 12 years. *Psychological Medicine, 45*, 1025-1037. doi: 10.1017/S0033291714002128
- Zalewski, M., & Lengua, L. J. (2012). Extending research on parenting in mothers diagnosed with BPD: Commentary on Stepp et al. *Personality Disorders: Theory, Research, and Treatment, 3*, 101-103. doi: 10.1037/a0026120
- Zalewski, M., Stepp, S. D., Scott, L. N., Whalen, D. J., Beeney, J. F., & Hipwell, A. E. (2014). Maternal borderline personality disorder symptoms and parenting of adolescent daughters. *Journal of Personality Disorders, 28*, 541-554. doi: 10.1521/pedi\_2014\_28\_131
- Zalewski, M., Stepp, S. D., Whalen, D. J., & Scott, L. N. (2015). A qualitative assessment of the parenting challenges and treatment needs of mothers with borderline personality disorder. *Journal of Psychotherapy Integration, 25*, 71-89. doi: 10.1037/a0038877

- Zanarini, M. C., Frankenburg, F. R., Dubo, E. D., Sickel, A. E., Trikha, A., Leniv, A., & Reynolds, V. (1998). Axis I comorbidity of borderline personality disorder. *American Journal of Psychiatry*, *155*, 1733-1739. doi: 10.1176/ajp.155.12.1733
- Zanarini, M. C., Williams, A. A., Lewis, R. E., Reich, R. B., Vera, S. C., Marino, M. F., . . . Frankenburg, F. R. (1997). Reported pathological childhood experiences associated with the development of borderline personality disorder. *The American Journal of Psychiatry*, *154*, 1101-1106. doi: 10.1176/ajp.154.8.1101

**Figure 1: Flowchart of the article search and selection process**

(Source: PRISMA, Moher et al 2009)



