

Original citation:

de Philippis, Guido, Marchese, Andrea and Rindler, Filip (2017) On a conjecture of Cheeger. In: Gigli, N., (ed.) Measure Theory in Non-Smooth Spaces. Berlin: De Gruyter Open, pp. 145-155.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/81085>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Attribution-NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 3.0) license and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by-nc-nd/3.0/>

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

Guido De Philippis, Andrea Marchese, and Filip Rindler

On a conjecture of Cheeger

4.1 Introduction

In [7] Cheeger, proved that in every doubling metric measure space (X, ρ, μ) satisfying a Poincaré inequality Lipschitz functions are differentiable μ -almost everywhere. More precisely, he showed the existence of a family $\{(U_i, \varphi_i)\}_{i \in \mathbb{N}}$ of Borel charts (that is, $U_i \subset X$ is a Borel set, $X = \bigcup_i U_i$ up to a μ -negligible set, and $\varphi_i: X \rightarrow \mathbb{R}^{d(i)}$ is Lipschitz) such that for every Lipschitz map $f: X \rightarrow \mathbb{R}$ at μ -almost every $x_0 \in U_i$ there exists a unique (co-)vector $df(x_0) \in \mathbb{R}^{d(i)}$ with

$$\limsup_{x \rightarrow x_0} \frac{|f(x) - f(x_0) - df(x_0) \cdot (\varphi(x) - \varphi(x_0))|}{\rho(x, x_0)} = 0.$$

This fact was later axiomatized by Keith [15], leading to the notion of *Lipschitz differentiability space*, see Section 4.2 below.

Cheeger also conjectured that the push-forward of the reference measure μ under every chart φ_i has to be absolutely continuous with respect to the Lebesgue measure, that is,

$$(\varphi_i)_\#(\mu \llcorner U_i) \ll \mathcal{L}^{d(i)},$$

see [7, Conjecture 4.63]. Some consequences of this fact concerning existence of bi-Lipschitz embeddings of X into some \mathbb{R}^N are detailed in [7, Section 14], also see [8, 9].

Let us assume that $(X, \rho, \mu) = (\mathbb{R}^d, \rho_\varepsilon, \nu)$ with ρ_ε the Euclidean distance and ν a positive Radon measure, is a Lipschitz differentiability space when equipped with the (single) identity chart (note that it follows a-posteriori from the validity of Cheeger's conjecture that no mapping into a higher-dimensional space can be a chart in a Lipschitz differentiability structure of \mathbb{R}^d). In this case the validity of Cheeger's conjecture reduces to the validity of the (weak) converse of Rademacher's theorem, which states that a positive Radon measure ν on \mathbb{R}^d with the property that all Lipschitz functions are differentiable ν -almost everywhere must be absolutely continuous with respect to \mathcal{L}^d . Actually, it is well known to experts that this converse of Rademacher's theorem implies Cheeger's conjecture in any metric space, see for instance [15, Section 2.4], [6, Remark 6.11], and [12].

Guido De Philippis: SISSA, Via Bonomea 265, 34136 Trieste, Italy. E-mail: guido.dephilippis@sissa.it

Andrea Marchese: Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland. E-mail: andrea.marchese@math.uzh.ch

Filip Rindler: Mathematics Institute, University of Warwick, Coventry CV4 7AL, UK. E-mail: F.Rindler@warwick.ac.uk

The (strong) converse of Rademacher's theorem has been known to be true in \mathbb{R} since the work of Zahorski [21], where he characterized the sets $E \subset \mathbb{R}$ that are sets of non-differentiability points of some Lipschitz function. In particular, he proved that for every Lebesgue negligible set $E \subset \mathbb{R}$ there exists a Lipschitz function which is nowhere differentiable on E .

The same result for maps $f: \mathbb{R}^d \rightarrow \mathbb{R}^d$ has been proved by Alberti, Csörnyei & Preiss for $d = 2$ as a consequence of a deep structural result for negligible sets in the plane [1, 2]. In 2011, Csörnyei & Jones [14] announced the extension of the above result to every Euclidean space. For Lipschitz maps $f: \mathbb{R}^d \rightarrow \mathbb{R}^m$ with $m < d$ the situation is fundamentally different and there exists a null set such that every Lipschitz function is differentiable at at least one point from that set, see [17, 18]. We finally remark that the weak converse of Rademacher's theorem in \mathbb{R}^2 can also be obtained by combining the results of [4] and [5], see [5, Remark 6.2 (iv)].

Recently, a result concerning the singular structure of measures satisfying a differential constraint was proved in [10]. When combined with the main result of [5] this proves the weak converse of Rademacher's theorem in any dimension, see [10, Theorem 1.14].

In this note we detail how the results in [5, 10] in conjunction with Bate's result on the existence of a sufficient number of independent Alberti representations in a Lipschitz differentiability space [6] imply Cheeger's conjecture; see Section 4.2 for the relevant definitions.

Theorem 4.1.1. *Let (X, ρ, μ) be a Lipschitz differentiability space and let (U, ϕ) be a d -dimensional chart. Then, $\phi_*(\mu \llcorner U) \ll \mathcal{L}^d$.*

Note that by the same arguments of this paper Cheeger's conjecture would also follow from the results announced in [1] and [14].

After we finished writing this note we learned that similar results have been proved by Kell and Mondino [16] and by Gigli and Pasqualetto [13].

4.2 Setup

4.2.1 Lipschitz differentiability spaces

Throughout this chapter, the triple (X, ρ, μ) will always denote a *metric measure space*, that is, (X, ρ) is a separable, complete metric space and $\mu \in \mathcal{M}_+(X)$ is a positive Radon measure on X .

We call a pair (U, φ) such that $U \subset X$ is a Borel set and $\varphi: X \rightarrow \mathbb{R}^d$ is Lipschitz, a *d -dimensional chart* or simply a *d -chart*. A function $f: X \rightarrow \mathbb{R}$ is said to be *differentiable with respect to a d -chart (U, φ)* at $x_0 \in U$ if there exists a unique (co-)vector $df(x_0) \in$

\mathbb{R}^d such that

$$\limsup_{x \rightarrow x_0} \frac{|f(x) - f(x_0) - df(x_0) \cdot (\varphi(x) - \varphi(x_0))|}{\rho(x, x_0)} = 0.$$

We call a metric measure space (X, ρ, μ) a *Lipschitz differentiability space* (also called a metric measure space that admits a *measurable differentiable structure*) if there exists a countable family of $d(i)$ -charts (U_i, φ_i) ($i \in \mathbb{N}$) such that $X = \bigcup_i U_i$ and any Lipschitz map $f: X \rightarrow \mathbb{R}$ is differentiable with respect to every (U_i, φ_i) at μ -almost every point $x_0 \in U_i$.

4.2.2 Alberti representations

We denote by $\Gamma(X)$ the set of *curves* in X , that is, the set of all Lipschitz maps $\gamma: \text{Dom } \gamma \rightarrow X$, for which the domain $\text{Dom } \gamma \subset \mathbb{R}$ is non-empty and compact. Note that we are not requiring $\text{Dom } \gamma$ to be an interval and thus the set $\Gamma(X)$ is sometimes also called the set of *curve fragments* on X . We equip $\Gamma(X)$ with the Hausdorff metric $\text{dist}_{\mathcal{H}}$ on graphs and we consider it as a subspace of the Polish space

$$\mathcal{K} = \{ K \subset \mathbb{R} \times X : K \text{ compact} \}, \tag{4.1}$$

endowed with the Hausdorff metric. Moreover, by arguing as in [19, Lemma 2.20], it is easy to see that $\Gamma(X)$ is an F_σ -subset of \mathcal{K} , i.e. a countable union of closed sets.

The decomposition of a measure into a family of 1-dimensional Hausdorff measures supported on curves leads to the notion of Alberti representation. First introduced in [4] for the study of the rank-one property of BV-derivatives, this decomposition has turned out to be a key tool in the study of differentiability properties of Lipschitz functions, see for instance [1, 2, 5, 6].

Definition 4.2.1. *Let (X, ρ, μ) be a metric measure space. An Alberti representation of μ on a μ -measurable set $A \subset X$ is a parameterized family $(\mu_\gamma)_{\gamma \in \Gamma(X)}$ of positive Borel measures $\mu_\gamma \in \mathcal{M}_+(X)$ with*

$$\mu_\gamma \ll \mathcal{H}^1 \llcorner \text{Im } \gamma,$$

together with a Borel probability measure $\pi \in \mathcal{P}(\Gamma(X))$ such that

$$\mu(B) = \int \mu_\gamma(B) \, d\pi(\gamma) \quad \text{for all Borel sets } B \subset A. \tag{4.2}$$

Here, the measurability of the integrand is part of the requirement of being an Alberti representation.

Remark 4.2.2. *Note that this definition is slightly different from the one in [6, Definition 2.2] since there the set $\Gamma(X)$ consists of bi-Lipschitz curves. Clearly, the existence of a representation in the sense of [6] implies the existence of a representation in our*

sense and this will suffice for our purposes. Let us, however, point out that the converse holds true as well. Indeed, the part of γ that contributes to the integral in (4.2) can be decomposed into countably many bi-Lipschitz pieces, see [19, Remark 2.17].

We will further need the notion of *independent* Alberti representations of a measure. Let $C \subset \mathbb{R}^d$ be a closed, convex, one-sided cone, i.e. a set of the form

$$C := \{ v \in \mathbb{R}^d : v \cdot w \geq (1 - \theta)\|v\| \}$$

for some $w \in \mathbb{S}^{d-1}$ and $\theta \in (0, 1)$. With a Lipschitz map $\phi: X \rightarrow \mathbb{R}^d$, we say that an Alberti representation $\int v_\gamma \, d\pi(\gamma)$ has ϕ -directions in C if

$$(\phi \circ \gamma)'(t) \in C \setminus \{0\} \quad \text{for } \pi\text{-a.e. curve } \gamma \text{ and } \mathcal{H}^1\text{-a.e. } t \in \text{Dom } \gamma.$$

A number of m Alberti representations of μ are ϕ -independent if there are linearly independent cones C_1, \dots, C_m such that the i 'th Alberti representation has ϕ -directions in C_i . Here, linear independence of the cones C_1, \dots, C_m means that any collection of vectors $v_i \in C_i \setminus \{0\}$ is linearly independent. In the case $X = \mathbb{R}^d$ we will always consider $\phi = \text{Id}$.

One of the main results of [6] asserts that a Lipschitz differentiability space necessarily admits many independent Alberti representations, also cf. [5, Theorem 1.1]. Recall that according to Remark 4.2.2 any representation in the sense of [6] is also a representation in the sense of Definition 4.2.1.

Theorem 4.2.3. *Let (X, ρ, μ) be a Lipschitz differentiability space with a d -chart (U, φ) . Then, there exists a countable decomposition*

$$U = \bigcup_{k \in \mathbb{N}} U_k, \quad U_k \subset U \text{ Borel sets,}$$

such that every $\mu \llcorner U_k$ has d φ -independent Alberti representations.

A proof of this theorem can be found in [6, Theorem 6.6].

4.2.3 One-dimensional currents

To use the results of [10] we need a link between Alberti representations and 1-dimensional currents. Recall that a 1-dimensional current T in \mathbb{R}^d is a continuous linear functional on the space of smooth and compactly supported differential 1-forms on \mathbb{R}^d . The boundary of T , ∂T is the distribution (0-current) defined via $\langle \partial T, f \rangle := \langle T, df \rangle$ for every smooth and compactly supported function $f: \mathbb{R}^d \rightarrow \mathbb{R}$. The mass of T , denoted by $\mathbf{M}(T)$, is the supremum of $\langle T, \omega \rangle$ over all 1-forms ω such that $|\omega| \leq 1$ everywhere. A current T is called *normal* if both T and ∂T have finite mass; we denote the set of normal 1-currents by $\mathbf{N}_1(\mathbb{R}^d)$.

By the Radon–Nikodým theorem, a 1-dimensional current T with finite mass can be written in the form $T = \vec{T}\|T\|$ where $\|T\|$ is a finite positive measure and \vec{T} is a vector field in $L^1(\mathbb{R}^d, \|T\|)$ with $|\vec{T}(x)| = 1$ for $\|T\|$ -almost every $x \in \mathbb{R}^d$. In particular, the action of T on a smooth and compactly supported 1-form ω is given by

$$\langle T, \omega \rangle = \int_{\mathbb{R}^d} \langle \omega(x), \vec{T}(x) \rangle \, d\|T\|(x) .$$

An *integer-multiplicity rectifiable 1-current* (in the following called simply *rectifiable 1-current*) $T = \llbracket E, \tau, m \rrbracket$ is a 1-current which acts on 1-forms ω as

$$\langle T, \omega \rangle = \int_E \langle \omega(x), \tau(x) \rangle m(x) \, d\mathcal{H}^1(x) ,$$

where E is a 1-rectifiable set, $\tau(x)$ is a unit vector spanning the approximate tangent space $\text{Tan}(E, x)$ and m is an integer-valued function such that $\int_E m \, d\mathcal{H}^1 < \infty$. More information on currents can be found in [11].

The relation between Alberti representations and normal 1-currents is partially encoded in the following decomposition theorem, due to Smirnov [20].

Theorem 4.2.4. *Let $T = \vec{T}\|T\| \in \mathbf{N}_1(\mathbb{R}^d)$ be a normal 1-current with $|\vec{T}(x)| = 1$ for $\|T\|$ -almost every x . Then, there exists a family of rectifiable 1-currents*

$$T_\gamma = \llbracket E_\gamma, \tau_\gamma, 1 \rrbracket, \quad \gamma \in \Gamma,$$

where Γ is a measure space endowed with a finite positive Borel measure $\pi \in \mathcal{M}_+(\Gamma)$, such that the following assertions hold:

(i) T can be decomposed as

$$T = \int_{\Gamma} T_\gamma \, d\pi(\gamma)$$

and

$$\mathbf{M}(T) = \int_{\Gamma} \mathbf{M}(T_\gamma) \, d\pi(\gamma) = \int_{\Gamma} \mathcal{H}^1(E_\gamma) \, d\pi(\gamma) ;$$

(ii) $\tau_\gamma(x) = \vec{T}(x)$ for \mathcal{H}^1 -almost every $x \in E_\gamma$ and for π -almost every $\gamma \in \Gamma$;

(iii) $\|T\|$ can be decomposed as

$$\|T\| = \int_{\Gamma} \mu_\gamma \, d\pi(\gamma) ,$$

where each μ_γ is the restriction of \mathcal{H}^1 to the 1-rectifiable set E_γ .

An Alberti representation of an Euclidean measure splits it into measures concentrated on “fragments” of curves. In general, these fragments cannot be glued together

to obtain a 1-dimensional normal current since the boundary may have infinite mass. Nevertheless, the “holes” of every curve appearing in an Alberti representation of a measure $\nu \in \mathcal{M}_+(\mathbb{R}^d)$ can be “filled” in such a way as to produce a normal 1-current T with $\nu \ll \|T\|$. Moreover, if the representation has directions in a cone C then the constructed normal current T has orienting vector \vec{T} in $C \setminus \{0\}$ almost everywhere (with respect to $\|T\|$). Indeed, we have the following lemma, which is essentially [5, Corollary 6.5]; it can be interpreted as a partial converse to Theorem 4.2.4:

Lemma 4.2.5. *Let $\nu \in \mathcal{M}_+(\mathbb{R}^d)$ be a finite Radon measure. If there is an Alberti representation $\nu = \int \nu_\gamma \, d\pi(\gamma)$ with directions in a cone C , then there exists a normal 1-current $T \in \mathbf{N}_1(\mathbb{R}^d)$ such that $\vec{T}(x) \in C \setminus \{0\}$ for $\|T\|$ -almost every $x \in \mathbb{R}^d$ and $\nu \ll \|T\|$.*

Proof. For the purpose of illustration we sketch the proof.

Step 1. Given ν as in the statement, we claim that there exists a normal 1-current $T = \vec{T}\|T\|$ with $\mathbf{M}(T) \leq 1$ and $\mathbf{M}(\partial T) \leq 2$ such that $\vec{T}(x) \in C$, for $\|T\|$ -almost every x and that ν is not singular with respect to $\|T\|$.

The claim follows from the proof of [5, Lemma 6.12]. For the sake of completeness let us present the main line of reasoning. By arguing as in Step 1 of the proof of [5, Lemma 6.12], to every $\gamma \in \Gamma(\mathbb{R}^d)$ with $\gamma'(t) \in C$ and a Borel measure $\nu_\gamma \ll \mathcal{H}^1 \llcorner \text{Im } \gamma$ we can associate a 1-Lipschitz map $\psi_{\nu_\gamma} : [0, 1] \rightarrow \mathbb{R}^d$ satisfying

$$\nu_\gamma(\text{Im}(\psi_{\nu_\gamma})) > 0 \quad \text{and} \quad \psi'_{\nu_\gamma}(t) \in C \setminus \{0\} \quad \text{for } \mathcal{H}^1\text{-a.e. } t \in [0, 1].$$

This map can moreover be chosen such that $\gamma \mapsto \psi_{\nu_\gamma}$ coincides with a Borel measurable map π -almost everywhere once we endow the set of curves with the topology of uniform convergence, see Step 3 in the proof of [5, Lemma 6.12].

Let $T_{\nu_\gamma} := \llbracket \text{Im } \psi_{\nu_\gamma}, \tau_{\psi_{\nu_\gamma}}, 1 \rrbracket$ be the rectifiable 1-current associated to ψ_{ν_γ} and set

$$T := \int T_{\nu_\gamma} \, d\pi(\gamma).$$

Since ψ_{ν_γ} is 1-Lipschitz, $\mathcal{H}^1(\text{Im } \psi_{\nu_\gamma}) \leq 1$ and thus $\mathbf{M}(T) \leq 1$. Moreover, for all smooth compactly supported functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ we have

$$\langle \partial T, f \rangle = \langle T, df \rangle = \int f(\psi_{\nu_\gamma}(1)) - f(\psi_{\nu_\gamma}(0)) \, d\pi(\gamma),$$

so that $\mathbf{M}(\partial T) \leq 2$.

By assumption, $\vec{T}(x) \in C \setminus \{0\}$ for $\|T\|$ -almost every $x \in \mathbb{R}^d$. To show that $\|T\|$ and ν are not mutually singular, for π -almost every γ set

$$\nu'_\gamma := \nu_\gamma \llcorner \text{Im } \psi_{\nu_\gamma} \quad \text{and} \quad \nu' := \int \nu'_\gamma \, d\pi(\gamma),$$

so that $\nu' \neq 0$ and $\nu' \leq \nu$. We will now establish that $\nu' \ll \|T\|$, for which we will prove that ν and $\|T\|$ are not mutually singular. Let $E \subset \mathbb{R}^d$ be such that $\|T\|(E) = 0$. Using

$$T = \int \llbracket \text{Im } \psi_{\nu_\gamma}, \tau_{\psi_{\nu_\gamma}}, 1 \rrbracket \, d\pi(\gamma) \quad \text{with} \quad \tau_{\psi_{\nu_\gamma}} = \frac{\psi'_{\nu_\gamma}}{|\psi'_{\nu_\gamma}|} \in C,$$

we get

$$\mathcal{H}^1(\text{Im } \psi_{v_\gamma} \cap E) = 0 \quad \text{for } \pi\text{-a.e. } \gamma.$$

Since by definition $v_\gamma \ll \mathcal{H}^1 \llcorner \text{Im } \gamma$, we have that $v'_\gamma \ll \mathcal{H}^1 \llcorner \text{Im } \psi_{v_\gamma}$. Thus, $v'(E) = 0$.

Step 2. Let us define

$$\mathcal{T} := \{ T \in \mathbf{N}_1(\mathbb{R}^d) : \mathbf{M}(T) \leq 1, \mathbf{M}(\partial T) \leq 2 \text{ and } \bar{T} \in C \text{ } \|\! \| T \|\! \| \text{-a.e.} \}$$

and

$$\mathcal{T}_v := \{ T \in \mathcal{T} : v \text{ and } T \text{ are not singular} \}.$$

Note that if $C = \{ v \in \mathbb{R}^d : v \cdot w \geq (1 - \theta)\|v\| \}$ for some $w \in \mathbb{S}^{d-1}$, $\theta \in (0, 1)$, then $\bar{T} \in C$ almost everywhere implies that

$$\|T\| \geq T \cdot w \geq (1 - \theta)\|T\| \tag{4.3}$$

as measures (here we are identifying T with an \mathbb{R}^d -valued Radon measure and use the pointwise scalar product). Moreover, as a consequence of the Radon–Nikodým theorem, for every $T \in \mathcal{T}_v$ we may write

$$v = g_{\|T\|} \|T\| + v_{\|T\|}^s \quad \text{with} \quad v_{\|T\|}^s \perp \|T\|, \quad \int g_{\|T\|} \, d\|T\| > 0.$$

Let us set $M := \sup_{T \in \mathcal{T}_v} \int g_{\|T\|} \, d\|T\| > 0$ and let $T_k \in \mathcal{T}_v$ be a sequence with

$$\int g_{\|T_k\|} \, d\|T_k\| \rightarrow M.$$

Define

$$T := \sum_k 2^{-k} T_k$$

and note that $T \in \mathcal{T}$. Moreover, by (4.3), $\|T_k\| \ll \|T\|$ for all $k \in \mathbb{N}$, so that there exist $h_k: \mathbb{R}^d \rightarrow \mathbb{R}$ with

$$\int_E h_k \, d\|T\| = \int_E g_{\|T_k\|} \, d\|T_k\| \leq v(E) \quad \text{for all Borel sets } E \subset \mathbb{R}^d.$$

In particular, $T \in \mathcal{T}_v$ and $h_k \leq g_{\|T\|}$. Set $m_k = \max_{1 \leq j \leq k} h_j$. By the monotone convergence theorem, $m_k \rightarrow m_\infty \leq g_{\|T\|}$ in $L^1(\mathbb{R}^d, \|T\|)$ and

$$M \leq \lim_{k \rightarrow \infty} \int m_k \, d\|T\| = \int m_\infty \, d\|T\| \leq \int g_{\|T\|} \, d\|T\| \leq M.$$

Hence, M is actually a maximum and it is attained by T .

We now claim that $v \ll \|T\|$. Indeed, assume by contradiction that $v = g_{\|T\|} \|T\| + v_{\|T\|}^s$ with $v_{\|T\|}^s \neq 0$. Since the Alberti representation of v induces an Alberti representation of $v_{\|T\|}^s$, we can apply Step 1 to find a normal 1-current

$$S \in \mathcal{T}_{v_{\|T\|}^s} \subset \mathcal{T}_v$$

such that $\nu_{\|T\|}^s$ and $\|S\|$ are not mutually singular. In particular, if $\nu = g_{\|S\|} d\|S\| + \nu_{\|S\|}^s$, then there exists a Borel set $F \subset \mathbb{R}^d$ such that

$$\|T\|(F) = 0 \quad \text{and} \quad \int_F g_{\|S\|} d\|S\| > 0. \tag{4.4}$$

Let us define $W := (T + S)/2$ and note that by (4.3) it holds that $\|T\|, \|S\| \ll \|W\|$ so that $W \in \mathcal{T}_\nu$. Moreover, there are functions $h_T, h_S \leq g_{\|W\|}$ such that

$$\int_E h_T d\|W\| = \int_E g_{\|T\|} d\|T\|, \quad \int_E h_S d\|W\| = \int_E g_{\|S\|} d\|S\|$$

for all Borel sets E . However, for F as in (4.4) we obtain

$$M \geq \int_{\mathbb{R}^d} g_{\|W\|} d\|W\| \geq \int_{\mathbb{R}^d} g_{\|T\|} d\|T\| + \int_F g_{\|S\|} d\|S\| > M,$$

a contradiction. □

4.3 Proof of Cheeger’s conjecture

The key tool to prove Cheeger’s conjecture is the following result from [10, Corollary 1.12]:

Theorem 4.3.1. *Let $T_1 = \vec{T}_1\|T_1\|, \dots, T_d = \vec{T}_d\|T_d\| \in \mathbf{N}_1(\mathbb{R}^d)$ be 1-dimensional normal currents. Let $\nu \in \mathcal{M}_+(\mathbb{R}^d)$ be a positive Radon measure such that*

- (i) $\nu \ll \|T_i\|$ for $i = 1, \dots, d$, and
 - (ii) $\text{span}\{\vec{T}_1(x), \dots, \vec{T}_d(x)\} = \mathbb{R}^d$ for ν -almost every x .
- Then, $\nu \ll \mathcal{L}^d$.*

Combining the above result with Lemma 4.2.5 we immediately get the following:

Lemma 4.3.2. *Let $\nu \in \mathcal{M}_+(\mathbb{R}^d)$ have d independent Alberti representations. Then, $\nu \ll \mathcal{L}^d$.*

Proof. Denote by C_1, \dots, C_d independent cones such that there are d Alberti representations having directions in these cones. By Lemma 4.2.5 there are d normal 1-dimensional currents $T_1 = \vec{T}_1\|T_1\|, \dots, T_d = \vec{T}_d\|T_d\| \in \mathbf{N}_1(\mathbb{R}^d)$ such that

$$\nu \ll \|T_i\| \quad \text{for } i = 1, \dots, d,$$

and $\vec{T}_i(x) \in C_i$ for ν -almost every $x \in \mathbb{R}^d$. By the independence of the cones,

$$\text{span}\{\vec{T}_1(x), \dots, \vec{T}_d(x)\} = \mathbb{R}^d \quad \text{for } \nu\text{-a.e. } x \in \mathbb{R}^d.$$

This implies $\nu \ll \mathcal{L}^d$ via Theorem 4.3.1. □

In order to use the above result to prove Theorem 8.1.1 one also needs the following “push-forward lemma”.

Lemma 4.3.3. *Let (X, ρ, μ) be a Lipschitz differentiability space with a d -chart (U, φ) . If $\mu \llcorner U$ has d φ -independent Alberti representations, then also the push-forward $\varphi_#(\mu \llcorner U) \in \mathcal{M}_+(\mathbb{R}^d)$ has d independent Alberti representations.*

Proof. It is enough to show that if there exists a representation of the form $\mu \llcorner U = \int \mu_\gamma \, d\pi(\gamma)$ with φ -directions in a cone C (i.e. such that $(\varphi \circ \gamma)'(t) \in C \setminus \{0\}$ for almost all $t \in \text{Dom } \gamma$ and for π -almost every γ), then we can build an Alberti representation

$$\varphi_#(\mu \llcorner U) = \int v_{\tilde{\gamma}} \, d\tilde{\pi}(\tilde{\gamma}) \quad \text{with} \quad \tilde{\pi} \in \mathcal{P}(\Gamma(\mathbb{R}^d)),$$

with $\tilde{\gamma}'(t) \in C \setminus \{0\}$ for $\tilde{\pi}$ -almost every $\tilde{\gamma}$ and almost every $t \in \text{Dom } \tilde{\gamma}$. To this end consider the map $\Phi: \Gamma(X) \rightarrow \Gamma(\mathbb{R}^d)$ given by $\Phi(\gamma) := \varphi \circ \gamma$ and let $\tilde{\pi} := \Phi_# \pi \in \mathcal{M}_+(\Gamma(\mathbb{R}^d))$. Note that, by the very definition of the push-forward measure, for $\tilde{\pi}$ -almost every $\tilde{\gamma}$ it holds that $\tilde{\gamma} = \varphi \circ \gamma$ for some $\gamma \in \Gamma(X)$.

By considering π as a probability measure defined on the Polish space \mathcal{K} defined in (4.1), and noting that π is concentrated on $\Gamma(X)$, we can apply the disintegration theorem for measures [3, Theorem 5.3.1] to show that for $\tilde{\pi}$ -almost every $\tilde{\gamma}$ there exists a Borel probability measure $\eta_{\tilde{\gamma}}$ concentrated on $\Phi^{-1}(\tilde{\gamma})$ and such that

$$\pi(A) = \int \eta_{\tilde{\gamma}}(A) \, d\tilde{\pi}(\tilde{\gamma}) \quad \text{for all Borel sets } A \subset \Gamma(X).$$

Note also that, by the disintegration theorem, the map $\tilde{\gamma} \mapsto \eta_{\tilde{\gamma}}$ is Borel measurable. Let us now set

$$v_{\tilde{\gamma}} := \int_{\Phi^{-1}(\tilde{\gamma})} \varphi_#(\mu_\gamma) \, d\eta_{\tilde{\gamma}}(\gamma).$$

Clearly, we have the representation

$$\varphi_#(\mu \llcorner U) = \int v_{\tilde{\gamma}} \, d\tilde{\pi}(\tilde{\gamma})$$

and $\tilde{\gamma}'(t) = (\varphi \circ \gamma)'(t) \in C \setminus \{0\}$ for $\tilde{\pi}$ -almost every $\tilde{\gamma}$ and almost every $t \in \text{Dom } \tilde{\gamma}$. Hence, to conclude the proof we only have to show that

$$v_{\tilde{\gamma}} \ll \mathcal{H}^1 \llcorner \text{Im } \tilde{\gamma} \quad \text{for } \tilde{\pi}\text{-a.e. } \tilde{\gamma}.$$

Let E be a set with $\mathcal{H}^1(E \cap \text{Im } \tilde{\gamma}) = 0$. Since $\tilde{\gamma}'(t) \neq 0$ for almost every $t \in \text{Dom } \tilde{\gamma}$, the area formula implies that $\mathcal{L}^1(\tilde{\gamma}^{-1}(E)) = 0$. If $\gamma \in \Phi^{-1}(\tilde{\gamma})$, say $\tilde{\gamma} = \varphi \circ \gamma$, then

$$\mathcal{H}^1(\varphi^{-1}(E) \cap \text{Im } \gamma) \leq \mathcal{H}^1(\gamma(\tilde{\gamma}^{-1}(E))) = 0 \quad \text{for all } \gamma \in \Phi^{-1}(\tilde{\gamma}).$$

Hence, $\mu_\gamma(\varphi^{-1}(E)) = 0$ for all $\gamma \in \Phi^{-1}(\tilde{\gamma})$ which immediately gives

$$v_{\tilde{\gamma}}(E) = \int_{\Phi^{-1}(\tilde{\gamma})} \mu_\gamma(\varphi^{-1}(E)) \, d\eta_{\tilde{\gamma}}(\gamma) = 0.$$

This concludes the proof. \square

Proof of Theorem 8.1.1. Let (U, φ) be a d -chart. By Theorem 4.2.3 there are d φ -independent Alberti representations of $\mu \llcorner U_k$, where $U = \bigcup_{k \in \mathbb{N}} U_k$ is the decomposition from Bate's theorem. Then, via Lemma 4.3.3, the push-forward $\varphi_{\#}(\mu \llcorner U_k)$ also has d independent Alberti representations. Finally, Lemma 4.3.2 yields $\varphi_{\#}(\mu \llcorner U_k) \ll \mathcal{L}^d$ and this concludes the proof. \square

Acknowledgment: The authors would like to thank Nicola Gigli for his suggestion to write this note and Andrea Schioppa for several useful discussions. G. D. P. is supported by the MIUR SIR-grant "Geometric Variational Problems" (RBSI14RVEZ). A. M. is supported by the ERC-grant "Regularity of area-minimizing currents" (306247). F. R. gratefully acknowledges the support from an EPSRC Research Fellowship on "Singularities in Nonlinear PDEs" (EP/L018934/1).

Bibliography

- [1] G. Alberti, M. Csörnyei, and D. Preiss: Structure of null sets in the plane and applications. In Proceedings of the Fourth European Congress of Mathematics (Stockholm, 2004), pages 3–22. European Mathematical Society, 2005.
- [2] G. Alberti, M. Csörnyei, and D. Preiss: Differentiability of Lipschitz functions, structure of null sets, and other problems. In Proceedings of the International Congress of Mathematicians 2010 (Hyderabad 2010), pages 1379–1394. European Mathematical Society, 2010.
- [3] L. Ambrosio, N. Gigli, and G. Savaré: Gradient flows in metric spaces and in the space of probability measures. Lectures in Mathematics ETH Zürich. Birkhäuser, 2005.
- [4] G. Alberti: Rank one property for derivatives of functions with bounded variation. Proc. Roy. Soc. Edinburgh Sect. A, 123:239–274, 1993.
- [5] G. Alberti and A. Marchese: On the differentiability of Lipschitz functions with respect to measures in the Euclidean space. Geom. Funct. Anal., 26:1–66, 2016.
- [6] D. Bate: Structure of measures in Lipschitz differentiability spaces. J. Amer. Math. Soc., 28:421–482, 2015.
- [7] J. Cheeger: Differentiability of Lipschitz functions on metric measure spaces. Geom. Funct. Anal., 9:428–517, 1999.
- [8] J. Cheeger and B. Kleiner: On the differentiability of Lipschitz maps from metric measure spaces to Banach spaces. In Inspired by S. S. Chern, volume 11 of Nankai Tracts Math., pages 129–152. World Scientific, 2006.
- [9] J. Cheeger and B. Kleiner: Differentiability of Lipschitz maps from metric measure spaces to Banach spaces with the Radon-Nikodým property. Geom. Funct. Anal., 19:1017–1028, 2009.
- [10] G. De Philippis and F. Rindler: On the structure of \mathcal{A} -free measures and applications. Ann. of Math., 2016. to appear, arXiv:1601.06543.
- [11] H. Federer: Geometric measure theory. Die Grundlehren der mathematischen Wissenschaften, Band 153. Springer-Verlag New York Inc., New York, 1969.
- [12] J. Gong: Rigidity of derivations in the plane and in metric measure spaces. Illinois J. Math., 56:1109–1147, 2012.

- [13] N. Gigli and E. Pasqualetto: Behaviour of the reference measure on RCD spaces under charts. arXiv:1607.05188, 2016.
- [14] P. Jones: Product formulas for measures and applications to analysis and geometry, 2011. Talk given at the conference “Geometric and algebraic structures in mathematics”, Stony Brook University, May 2011, <http://www.math.sunysb.edu/Videos/dennisfest/>.
- [15] S. Keith: A differentiable structure for metric measure spaces. *Adv. Math.*, 183:271–315, 2004.
- [16] M. Kell and A. Mondino: On the volume measure of non-smooth spaces with Ricci curvature bounded below. arXiv:1607.02036, 2016.
- [17] D. Preiss: Differentiability of lipschitz functions on banach spaces. *J. Funct. Anal.*, 91:312–345, 1990.
- [18] D. Preiss and G. Speight: Differentiability of Lipschitz functions in Lebesgue null sets. *Invent. Math.*, 199:517–559, 2015.
- [19] A. Schioppa: Derivations and Alberti representations. *Adv. Math.*, 293:436–528, 2016.
- [20] S. K. Smirnov: Decomposition of solenoidal vector charges into elementary solenoids, and the structure of normal one-dimensional flows. *Algebra i Analiz*, 5:206–238, 1993. translation in *St. Petersburg Math. J.* 5 (1994), 841–867.
- [21] Z. Zahorski: Sur l’ensemble des points de non-dérivabilité d’une fonction continue. *Bull. Soc. Math. France*, 74:147–178, 1946.