

**Original citation:**

Nachega, Jean B., Adetokunboh, Olatunji, Uthman, Olalekan A., Knowlton, Amy W. , Geng, Elvin, Peltzer, Karl , Chang, Larry W. , Cutsem, Gilles Van, Jaffar, Shabbar S., Ford, Nathan , Mellins, Claude A., Remien, Robert H. and Mills, Edward. (2016) Community-based interventions to improve and sustain antiretroviral therapy adherence, retention in HIV care and clinical outcomes in low- and middle-income countries for achieving the UNAIDS 90-90-90 targets. Current HIV/AIDS Reports.

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/81338>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

"The final publication is available at Springer via <http://dx.doi.org/10.1007/s11904-016-0325-9>

**A note on versions:**

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP url' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)

# **Community-Based Interventions to Improve and Sustain Antiretroviral Therapy Adherence, Retention in HIV Care and Clinical Outcomes in Low- and Middle-Income Countries for Achieving the UNAIDS 90-90-90 Targets**

## **Authors**

Jean B. Nachega<sup>1,2,3</sup>, Olatunji Adetokunboh<sup>2</sup>, Olalekan A. Uthman<sup>4</sup>, Amy W. Knowlton<sup>3</sup>, Elvin Geng<sup>5</sup>; Karl Peltzer<sup>6,7,8</sup>, Larry W. Chang<sup>3</sup>, Gilles Van Cutsem<sup>9</sup>, Shabbar S. Jaffar<sup>10</sup>, Nathan Ford<sup>11</sup>, Claude A. Mellins<sup>12</sup>, Robert H. Remien<sup>13</sup>, Edward J Mills<sup>14</sup>

## **Affiliations**

1. University of Pittsburgh, Pittsburgh, PA, USA
2. Stellenbosch University, Cape Town, South Africa
3. Johns Hopkins University, Baltimore, MD, USA
4. Warwick Medical School, The University of Warwick, Coventry, UK
5. University of California, San Francisco, CA, USA
6. Mahidol University, Salaya, Thailand
7. University of Limpopo, Polokwane, South Africa
8. Human Sciences Research Council, Pretoria, South Africa
9. Medecins Sans Frontieres, Cape Town, South Africa
10. Liverpool School of Tropical Medicine, Liverpool, UK
11. World Health Organization, Geneva, Switzerland
12. HIV Center for Clinical and Behavioral Studies, Columbia University
13. New York State Psychiatric Institute, New York, NY, USA
14. Precision Global Health, Vancouver, Canada

## **Corresponding Author:**

Jean B. Nachega, MD, PhD, MPH, University of Pittsburgh  
Graduate School of Public Health, Department of Epidemiology  
130 DeSoto Street, 503 Parran Hall, Pittsburgh, PA 15261; USA  
E-mail: [JBN16@PITT.EDU](mailto:JBN16@PITT.EDU)

Word counts: Abstract: 318; Main text: 3,755

## Abstract

Little is known about the effect of community versus health facility-based interventions to improve and sustain antiretroviral therapy (ART) adherence, virologic suppression, and retention in care among HIV-infected individuals in low- and middle-income countries (LMICs). We systematically searched four electronic databases for all available randomized controlled trials (RCTs) and comparative cohort studies in LMICs comparing community versus health facility-based interventions. Relative risks (RRs) for pre-defined adherence, treatment engagement (linkage and retention in care), and relevant clinical outcomes were pooled using random effect models. Eleven cohort studies and eleven RCTs ( $N = 97,657$ ) were included. Meta-analysis of the included RCTs comparing community- versus health facility-based interventions found comparable outcomes in terms of ART adherence ( $RR = 1.02$ , 95 % CI 0.99 to 1.04), virologic suppression ( $RR = 1.00$ , 95 % CI 0.98 to 1.03), and all-cause mortality ( $RR = 0.93$ , 95 % CI 0.73 to 1.18). The result of pooled analysis from the RCTs ( $RR = 1.03$ , 95 % CI 1.01 to 1.06) and cohort studies ( $RR = 1.09$ , 95 % CI 1.03 to 1.15) found that participants assigned to community-based interventions had statistically significantly higher rates of treatment engagement. Two studies found community-based ART delivery model either cost-saving or cost-effective. Community- versus facility-based models of ART delivery resulted in at least comparable outcomes for clinically stable HIV-infected patients on treatment in LMICs and are likely to be cost-effective.

## Introduction

The number of people that have started life-saving antiretroviral therapy (ART) has importantly increased in low- and middle-income countries [1]. However, this positive development has led to health facilities' overcrowding, longer waiting times during visits, and reduced time for counseling and clinical care of newly enrolled patients. In most public sector clinics in resource-limited settings, it has also restricted the workforce's capacity to provide ongoing adherence support and track patients lost to follow-up to ensure optimal ART effects on patient health and community HIV prevention. [2]. Further, in July 2014, UNAIDS called for a global scale-up of Treatment as Prevention (TASP) and efforts to meet the following '90-90-90' targets by 2020: (1) 90% of all people living with HIV should know their HIV status (90% diagnosed); (2) 90% of all people diagnosed with HIV infection should receive ART (90% on treatment); and (3) 90% of all people receiving ART should achieve viral suppression (UNAIDS 2014)[3]. This ambitious target is supported by the latest World Health Organization (WHO) guidelines recommending treating all HIV positive individuals irrespective of immune status. It implies that an additional 21 million people are now eligible for treatment [4]. This underscores the importance of expanding health systems' capacity to meet communities' growing health needs.

Emerging data from both developed and developing countries demonstrate that a substantial reduction in patient retention in medical care occurs between each stage of the HIV treatment continuum, or cascade—from diagnosis and linkage to care, assessment of ART readiness to acceptability, receipt of initial ART, adherence and long-term retention in care, and treatment success as reflected by virologic suppression [5-8]. A systematic review by Fox and Rosen reported that retention of HIV-infected adults on ART at 36 months in middle- and low-income countries averages only 65% to 70%[9]. Success along the HIV treatment cascade is even worse in vulnerable populations, namely, pregnant women, children and adolescents, sex

workers, injecting drug users, and men who have sex with men; and they are at high risk of acquiring as well as transmitting the HIV virus, thus experiencing poor clinical and public health outcomes [10-15]. Against this background, it is critical to determine how effective interventions are at every level of the treatment cascade to prevent new infections and promote health outcomes to achieve the goal of an AIDS- free generation [7].

In Africa, selected approaches to reducing loss at every stage of the HIV treatment cascade include decentralization of services and task-shifting aspects of care to nurses and to non-clinical staff, including lay counselors who may be patients themselves. These approaches have been found to be feasible and to result in good clinical outcomes and are now recommended and being scaled up in resource-limited settings [16-19]. However, such facility-based approaches are reaching their limits as increasing numbers of patients initiate ART. Recently, suggestions have been made to expand accessible and flexible community-based ART services delivery, differentiating the needs of clinically ill patients starting ART or in need of significant adherence counseling from the needs of clinically stable ongoing patients with documented good adherence. Such , have been suggested as important strategies for maintaining and improving ART adherence, retention and quality of care [16].

Indeed, community and home-based programs to promote ART adherence and/or retention in HIV care are now increasingly being recognized as an important and sustainable approach that could contribute significantly toward the UNAIDS 90-90-90 target and ultimately an AIDS-Free generation [20-32]. In addition, they are also seen as an essential mechanism of service delivery, including dispensing of ART, and a means of decongesting formal health services, rather than being purely an adherence adjunct. Furthermore, such interventions are likely to be cost-effective by offering a shift of certain tasks from overburdened healthcare settings directly into communities [33,34].

We aimed to conduct a systematic review and meta-analysis to compare the effect of community-based ART delivery on adherence, virologic outcomes, retention in care and all-cause mortality among HIV-infected individuals in middle- and low-income countries against results obtained through standard health facility care.

## Methods

### Protocol

The study background, rationale, and methods were specified in advance and documented in a study protocol registered in the PROSPERO database (CRD42016034114).

### Study Inclusion Criteria

In order to be eligible for inclusion in the review, studies had to report on adherence, and/or virologic suppression, and/or lost-to-follow up outcomes after initiation of ART. The following selection criteria were used to identify potential studies:

- *Study design:* Observational and experimental studies with primary data using cross-sectional, case – control, and cohort (prospective and retrospective) and randomized controlled trials (RCT s) designs
- *Study population:* HIV-infected individuals initiated on ART.
- *Intervention:* Community-based ART delivery. Models could include: 1) home-based interventions (e.g. friends or family-centred approaches); 2) peer- or HIV patients-led interventions; community ART distribution points (with or without involving primary level formal or informal health facilities); 3) community-based ART adherence clubs (with or without involving primary level formal or informal health facilities); 4) community ART groups (CAGs)
- *Comparator:* health-facility (e.g. hospital or clinics)

- *Outcomes:* Primary: 1) Proportion of HIV-infected patients with optimal ART adherence levels\* (>80%); 2) Proportion of HIV-infected patients with virologic suppression (as defined by the studies) at 12 and 24 months' post ART initiation. Secondary: Proportion of patients lost-to-follow up as defined by the studies at 12 and 24 months' post Art initiation; all-cause mortality; and cost to patient and provider.

### **Data Sources and Searches**

We conducted a systematic literature search using the following databases: Medline (PubMed), Scopus, SCI Web of Science and Cochrane Central Register of Controlled Trials (CENTRAL) through January 2016. In addition, abstracts from major HIV/AIDS or infectious diseases conferences such as the Conference on Retrovirus and Opportunistic Infections (CROI), International AIDS Society (IAS), International AIDS Conference and Infectious Diseases Society of America (IDSA) were reviewed for inclusion.

Our search terms included: "community"; "home-based care"; "health facilities"; "adherence"; "adherence club"; "retention in care"; "retention"; "loss to follow up", "attrition", "antiretroviral therapy"; "HIV"; "community volunteers"; "treatment supporter"; "DAART"; "cost".

### **Study Selection and Data Extraction**

Two of the authors (JBN and OA) screened the search outputs using titles and abstracts and independently went through the full text of all potentially eligible studies to assess whether they meet the inclusion criteria. Discrepancies in the choice of included studies between the two authors were resolved through discussion and consensus. For all eligible studies, the same authors reviewed extracted information regarding publication date, study setting, study design, methods, patient population, study intervention, and outcomes.

### **Risk of bias (quality) assessment**

To appraise the risk of bias for included studies, a tool was adapted from the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) (Appendix) [35]. Briefly, the risk of bias was assessed as low risk, unclear risk or high-risk for each of the following domains: selection (sample population), selection (participation rate), performance bias (outcome assessment), performance bias (analytical methods to control for bias) and other forms of bias. We used the Cochrane Collaboration's tool for assessing the risk of bias for quality assessment of the included studies[36]. The studies were graded based on: (i) sequence generation, (ii) blinding of outcome assessor, (iii) incomplete outcome data, (iv) selective outcome reporting, and (v) other sources of bias.

### **Measures of treatment effect and unit of analysis**

We used relative risks (RR) for the calculation of dichotomous data (such as adherence and retention in care) and mean differences for continuous data (such as change in CD4 cell count). All the results are presented with 95 % confidence intervals (CI).

### **Data Synthesis**

In the absence of statistical heterogeneity, we used a fixed-effect model, and we used a random-effects model where we detected moderate heterogeneity and it was deemed still reasonable to combine trials. We assessed the presence of statistical heterogeneity in the meta-analyses by visual inspection of the forest plot and applying a chi-squared test for heterogeneity with a threshold P value of 0.10 to determine statistical significance. Inconsistency was quantified across studies using the  $I^2$  value. We used Review Manager 5.3 [37] to conduct analyses and analyzed results for trials and cohort studies separately and also pooled these data. This review was reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement [38].

## Results

### Study selection and characteristics of included studies

The process of study identification and selection is shown in **Figure 1**. The literature search yielded 7,928 citations after removing duplicates. After review of the title and abstract, 33 full text articles were selected for critical review. A total of eleven trials were included [21,25,26,29,30,36,39-44] and eleven cohort studies [27,36,39,45-54] with a total of 5861 and 89,388 participants, respectively. These studies were conducted in eight different sub-Saharan African countries, including Botswana, Zimbabwe, South Africa, Nigeria, Uganda, Kenya, Tanzania and Zambia. Other studies were conducted in other low and middle income countries such as Brazil, Haiti, Peru and Thailand. See additional details in **Table 1**. We excluded seventeen studies. One study was excluded due to inclusion of a non-African population[44], four studies were excluded due to non-inclusion of community based data [32,55-57] while seven studies were excluded because the studies were non comparative in nature [19,58-62]. A study was also excluded because only baseline data were reported[63] while four did not show data for different arms of the studies[17,64-66].

### Risk of bias in included studies

The risk of bias of included cohort studies is summarised in **Table 2**. All the included studies had low risk of bias with respect to the selection of sample population and explaining the rationale for case and control selection while the included cohort studies were at risk of bias for sample selection ambiguity and having samples that were unlikely to be representative. All the studies had a high participation rate (>70-85%). In terms of outcome assessment, seven of the included studies had objective measures of adherence such as 'pill count', while two had high risk of bias by measuring the outcome using self-reporting format. All the trials and cohort studies used one or more analytical methods to control for bias in individual studies.

The risk of bias in the included studies was highest from other form of bias, followed by selection of sample population, and lowest from participation rates and analytical methods to control for bias.

The risk of bias of included trials is shown in **Table 3**. Allocation sequence generation was adequate in all the eleven trials. Allocation concealment was adequate in nine trials and unclear in the remaining two trials. Masking of outcome assessors was not clear in all the nine trials. Potential risk of bias due selective reporting and other bias was low in all eleven trials.

### **Optimal ART adherence**

Seven RCTs and three cohort studies reported optimal adherence as an outcome. Individual and pooled RRs for optimal adherence are shown in **Figure 2**. The result of pooled analysis from the RCTs showed no statistically significance difference in optimal adherence rates between the two treatment groups (pooled RR = 1.02, 95% CI 0.99 to 1.04,  $I^2 = 68\%$ ), such that among 6358 participants randomised to community-based ART 5827 (91.7%) achieved optimal ART adherence compared with 4083 of 4619 in the facility-based ART group (88.4%). However, three cohort studies provided evidence that participants in community-based ART had statistically significant higher optimal adherence rate compared to those in facility-based ART group (RR = 1.80, 95% CI 1.04 to 3.13), such that among 274 participants randomised to community-based ART 295 (92.9%) achieved optimal ART adherence compared with 196 of 288 in the facility-based ART group (68.1%).

### **Virologic suppression**

Eight RCTs and Eight cohort studies reported virologic suppression as an outcome. Individual and pooled RRs for virologic suppression are shown in **Figure 3**. The result of pooled analysis from the RCTs showed no statistically significance difference in virologic suppression rates between the two treatment groups (pooled RR = 1.00, 95% CI 0.97 to 1.03), with evidence of

no statistically significant heterogeneity between studies ( $I^2 = 0\%$ ,  $p=0.49$ ). Similarly, the result of pooled analysis from the cohort studies showed no statistically significant difference in virologic suppression rates between the two treatment groups (pooled RR = 1.06, 95% CI 0.77 to 1.46), with evidence of statistically significance substantial heterogeneity between studies ( $I^2 = 100\%$ ,  $p<0.00001$ ).

### Retention in care

Seven RCTs and four cohort studies reported retention in care as an outcome. Individual and pooled RRs for retention in care are shown in **Figure 4**. The result of pooled analysis from the RCTs showed that participants assigned to community-based ART (80.3% [3157 of 3931]) had statistically significant higher rates of retention in care than those in facility-based ART (75.9% [2334 of 3074]) at the end of the follow-up period (RR = 1.03, 95% CI 1.01 to 1.06,  $I^2 = 0\%$ ). Similarly, the result of pooled analysis from the cohort studies showed that participants assigned to community-based ART (89.4% [1074 of 1203]) had statistically significant higher rates of retention in care than those in facility-based ART (84.9% [2578 of 3038]) at the end of the follow-up period (RR = 1.09, 95% CI 1.03 to 1.15,  $I^2 = 69\%$ )

### Change in CD4+ T-cell count

Three RCTs reported CD4+ T-cell count as an outcome. The results of pooled analysis from the three RCTs show no significance difference in change in CD4+ T-cell count between the two treatment groups (mean difference = 0.93, 95% CI -11.36 to 13.21,  $I^2 = 0\%$ ) (**Figure 5**).

### All-cause mortality

Ten RCTs and eight cohort studies reported all-cause mortality as an outcome. Individual and pooled RRs for all-cause mortality are shown in **Figure 6**. The result of pooled analysis from the RCTs showed there was no statistically significant difference in rates of all-cause mortality in assigned to community-based ART (9.3% [388 of 4160]) than those assigned to facility-based ART (10.3% [338 of 3272]) at the end of the follow-up period (RR = 0.93, 95% CI 0.73 to 1.18,  $I^2 = 38\%$ ). Similarly, the result of pooled analysis from the cohort studies showed there was no statistically significant difference in rates of all-cause mortality in assigned to community-based ART (4.2% [1,075 of 25,506]) than those assigned to facility-based ART (6.0% [3,299 of 54,708]) at the end of the follow-up period (RR = 0.44, 95% CI 0.19 to 1.02,  $I^2 = 96\%$ ).

### Loss to follow-up

Six RCTs and seven cohort studies reported loss to follow-up as an outcome. Individual and pooled RRs for loss to follow-up are shown in **Figure 7**. The result of pooled analysis from the RCTs showed there was no statistically significant difference in rates of loss to follow-up in assigned to community-based and those assigned to facility-based ART at the end of the follow-up period (RR = 0.88, 95% CI 0.77 to 1.11,  $I^2 = 0\%$ ). However, the result of pooled analysis from the cohort studies showed that the rate of loss to follow-up was statistically significantly lower in those participants in community-based ART (5.1% [1,289 of 25,065]) than those in facility-based ART (9.7% [5,268 of 54,545]) at the end of the follow-up period (RR = 0.34, 95% CI 0.14 to 0.81,  $I^2 = 97\%$ ).

## Cost to the patient and provider

Jaffar et al.[25] reported costs to access care per patient including transport, lunch, child care costs and lost work time. The average total cost per patient in the first year was US \$29 among the community-based participants compared to the US \$60 facility-based patients. In terms of health-service costs, the same study reported average cost per patient per year to be US\$793 among the community-based participants compared to US\$838 among facility-based patients in Jinja, Uganda. Also, Bango and colleagues reported from South Africa, that ART adherence clubs (AAC) were most cost-effective than standard of care (SOC), with a cost per patient year of \$296 for AAC vs. \$374 for SOC. Retention in care at one year was 95% (95% CI: 94.88-95.86) for SOC, and 98% for ACC (95% CI: 97.6-98.3)[67].

## Discussion

This meta-analysis of nine cohort studies and seven RCTs found no statistical difference in optimal ART adherence, virologic suppression, change in CD4+ T-cell count, all-cause mortality and loss to follow-up between those participants assigned to community-based ART and facility-based ART, when analysis was restricted to RCTs. However, in the pooled analysis from both RCTs and cohort studies, we documented that participants assigned to community-based ART had significantly higher rates of retention in care than those in facility-based ART at the end of the follow-up period.

The above results corroborate the fact that providing patient support and education programs at community level are not inferior compared to facility-based ones and may in fact be superior when it comes to selected outcomes such as lost-to-follow up. Of note, our analysis may be underpowered to show superiority on selected outcomes such as virological outcomes and all-cause mortality. Furthermore, it appears that mortality in the observational studies might be

slightly higher in the community-based studies whereas retention and loss to follow-up seems to suggest that these outcomes in community-based are better than facility-based. It is possible that they are both likely artefacts, particularly in the non-randomized studies, of the fact that outcome ascertainment is better in the community than in the facility based studies since mortality in facility-based studies may be under-ascertained, this will make mortality in the community look higher[68]. At the same time, silent and even official community patient transfers in facility based also tend to be under-captured, thus making retention look worse[68]. For these reasons, studies, retention outcome is usually retention in facility and therefore interpretation of such outcome must with caution.

In building decentralized ART delivery, adherence and retention in care support, community-based ART programs encourage patient autonomy, build social networks and minimize the structural barriers, such as cost of transport to the clinic, which in turn appear to result in better outcomes. Such community-based interventions are likely to be impactful since they tend to involve trained community health workers, peers, volunteers, or patient's own social network members (e.g., family and friends) who assist with ART adherence counselling and support. In addition, there is evidence that they may provide material, instrumental and emotional support, as well as promote other healthy behaviours, such as decreased alcohol and drug use, leading to better health outcomes--- including survival [14,24,29,32]. Furthermore, enhancing certain aspects of the patient–supporter relationships—such as trust, supporter availability, communication, reciprocity of support, and medication assistance—in a manner consistent with patients' expectations may help to optimize the relationship, and its positive impact on patient health [14,24,29].

Our study complements the findings of a previous review that assessed the effect of home-based interventions on viral outcomes in sub-Saharan Africa; this review found that there was

insufficient data to be conclusive [69]. Another recent review summarized the evidence supporting different models of community participation for ART care, or community-based ART in sub-Saharan Africa; these community ART programs made treatment readily accessible and affordable[70]. In Uganda and Kenya community health workers or volunteers delivered ART at home [41,62] while in Tete, Mozambique a demonstration project of people living with HIV/AIDS used self-formed community-based ART groups to deliver ART in the community [19]. Also, in South Africa, Medicines Sans Frontiers, piloted ART adherence clubs with promising results[27]. These clubs may provide some adherence counseling and peer support, as well as enable a “fast track” refill mechanisms. Patients are placed in groups of approximately six patients and one member of the group (rotating each month) is permitted to obtain refills for all of the patients in his or her group. These approaches decrease the patient burden on health facilities, reduce transportation costs and waiting times for patients, and help overcome structural barriers. They also reduce treatment fatigue and loss to follow up, increase disclosure and treatment education, and may help patients develop necessary social ties. While supportive of community-based interventions, these evaluations used observational study designs with possibility of selection and observational bias as well as confounding and most of them did not have a valid comparator and could not be included in our meta-analysis.

We also investigated as secondary outcomes two potential concerns related to community-based ART adherence and retention programs, including reported stigma and low quality of care which could result in an increased all-cause mortality. In terms of stigma, an RCT reported that only 3% of patients refused to participate in the home-based ART program due to stigma [25] Furthermore, it has even been suggested that involvement of community-health care workers in HIV care reduced stigma[71] and being part of peer groups has been found to decrease the perception of social stigma [72].

Our results have important clinical and public health implications in the context of reaching UNAIDS 90-90-90 target for an AIDS-free generation: first, community-based interventions aim to deliver a package of essential ART care functions beyond the clinic into the community such as ART refills, monitoring of treatment adherence and outcomes, and detection of sick patients and rapid referral to care. This, in turn, frees up capacity within the clinic-based medical workforce to be able to focus on complicated tasks such as clinical care for sick patients, training and supervision of lower cadres, and management of health care services. Second, community-based ART delivery and adherence monitoring and support models for clinically stable patients with documented virological suppression holds the potential of enabling countries to build sustainable, cost effective and equitable HIV care for populations in countries with a scarce health care workforce. Indeed, a cost-effectiveness study by Marseille and colleagues concluded that a home-based ART programme in rural Africa may be more cost effective than most previous estimates for facility-based HAART programs [73].

Only three cohort studies, Fatti et al. [46], Grimwood et al. [49] and Massavon et al. [52] involved children. The outcomes reported by these studies were virological suppression, mortality and loss to follow up and all of these were not different from what was obtainable in the adult population. These studies were conducted in South Africa and Uganda

Our study has several strengths. We performed a comprehensive search of several databases and sources to identify eligible cohorts and RCTs with the latter providing the highest quality of evidence. Two authors independently evaluated each study for inclusion and data extraction. Regarding limitations, inclusion of cohort study designs may bias the overall estimate of effects due to unmeasured confounding not adjusted for in multivariate analyses. Indeed, the fact that we are observing a difference between RCTs and observational studies for the ART adherence outcome may reflect that in many if not all of these community based interventions, the patients who end up in the intervention, if it is not randomized, are likely to be quite a bit different – selected somehow – for stability even if not measured. However, in the context of implementation science, observational studies often provide strong signals of

important direction of effect. We could not do subgroup analysis for the primary outcomes based on the location of studies ie sub-Saharan African population versus other low and middle income countries. Finally, with only eleven RCTs, we may be underpowered to show superiority of either type interventions.

In summary, we found that community as compared to facility-based ART interventions were effective in achieving optimal ART adherence and virologic suppression. Retention in HIV care was not different in the two groups, but, overall, we found lower all-cause mortality and loss to follow up in favour of community-based interventions, mostly within cohort studies rather than RCTs. As ART rollout expands in Africa and the need of health systems to adjust for such expansion in Africa, community-based ART delivery adherence monitoring and support models for stable patients hold the promise of enabling countries to build sustainable, cost effective and equitable HIV care for populations in countries with a scarce health workforce. Further research with well powered studies may be needed to further explore effectiveness of such community-based ART programs and their cost-effectiveness and the resources needs of families and communities to sustain optimal outcomes.

### **Role of the funding source**

The present study did not receive any specific funding. All authors had full access to all the data in the study and had final responsibility for the decision to submit for publication.

**Authors' contributions:** JBN and EJM conceived the review. JBN, KP and OA drafted the protocol. OA and JBN conducted eligibility of the searches and researched the data. JBN, OA, OAU and AWK drafted the manuscript. The paper was revised critically for intellectual content by all the co-authors and gave final approval for publication.

**Competing interests:** None

Acknowledgments: None

## References

1. Joint United Nations Programme on HIV/AIDS (UNAIDS): *Report on the global AIDS epidemic 2015*. Geneva, Switzerland: UNAIDS; 2015.
2. Ford N, Mills EJ: **Simplified ART delivery models are needed for the next phase of scale up**. *PLoS Med* 2011, **8**:e1001060.
3. Joint United Nations Programme on HIV/AIDS (UNAIDS): *90-90-90: An Ambitious Treatment Target to Help End the AIDS Epidemic*. Geneva, Switzerland: UNAIDS; 2014.
4. World Health Organization: *Guideline on when to start antiretroviral therapy and on pre-exposure prophylaxis for HIV*. [URL: [http://apps.who.int/iris/bitstream/10665/186275/1/9789241509565\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/186275/1/9789241509565_eng.pdf)]. Accessed March 3, 2016. Geneva, Switzerland: WHO; 2015.
5. Gardner EM, McLees MP, Steiner JF, Del Rio C, Burman WJ: **The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection**. *Clin Infect Dis* 2011, **52**:793-800.
6. Micek MA, Gimbel-Sherr K, Baptista AJ, Matediana E, Montoya P, Pfeiffer J, Melo A, Gimbel-Sherr S, Johnson W, Gloyd S: **Loss to follow-up of adults in public HIV care systems in central Mozambique: identifying obstacles to treatment**. *J Acquir Immune Defic Syndr* 2009, **52**:397-405.
7. Nachega JB, Uthman OA, del Rio C, Mugavero MJ, Rees H, Mills EJ: **Addressing the Achilles' heel in the HIV care continuum for the success of a test-and-treat strategy to achieve an AIDS-free generation**. *Clin Infect Dis* 2014, **59** Suppl 1:S21-27.
8. Supervie V, Costagliola D: **The spectrum of engagement in HIV care in France: strengths and gaps [Abstract #: 1030]**. In *20th Conference on Retroviruses and Opportunistic Infections*. Edited by. Atlanta, GA; 2013.
9. Fox MP, Rosen S: **Retention of Adult Patients on Antiretroviral Therapy in Low- and Middle-Income Countries: Systematic Review and Meta-analysis 2008-2013**. *J Acquir Immune Defic Syndr* 2015, **69**:98-108.
10. Adetokunboh O, Oluwasanu M: **Eliminating mother-to-child transmission of the human immunodeficiency virus in sub-Saharan Africa: The journey so far and what remains to be done**. *J Infect Public Health* 2015.
11. Brown AE, Gill ON, Delpech VC: **HIV treatment as prevention among men who have sex with men in the UK: is transmission controlled by universal access to HIV treatment and care?** *HIV Med* 2013, **14**:563-570.
12. Delpech VC, Brown AE, Conti S, Polavarapu V, Yin Z: **Reducing onward transmission: viral suppression among key population groups living with HIV in the United Kingdom [Abstract 018]**. In *19th Annual Conference of the British HIV Association*. Edited by. 19th Annual Conference of the British HIV Association. Manchester, UK; 2013.

13. Kim SH, Gerver SM, Fidler S, Ward H: **Adherence to antiretroviral therapy in adolescents living with HIV: systematic review and meta-analysis.** *Aids* 2014, **28**:1945-1956.
14. Nachega JB, Hislop M, Nguyen H, Dowdy DW, Chaisson RE, Regensberg L, Cotton M, Maartens G: **Antiretroviral therapy adherence, virologic and immunologic outcomes in adolescents compared with adults in southern Africa.** *J Acquir Immune Defic Syndr* 2009, **51**:65-71.
15. Nachega JB, Uthman OA, Anderson J, Peltzer K, Wampold S, Cotton MF, Mills EJ, Ho YS, Stringer JS, McIntyre JA, et al.: **Adherence to antiretroviral therapy during and after pregnancy in low-income, middle-income, and high-income countries: a systematic review and meta-analysis.** *Aids* 2012, **26**:2039-2052.
16. Bedelu M, Ford N, Hilderbrand K, Reuter H: **Implementing antiretroviral therapy in rural communities: the Lusikisiki model of decentralized HIV/AIDS care.** *J Infect Dis* 2007, **196 Suppl 3**:S464-468.
17. Bemelmans M, van den Akker T, Ford N, Philips M, Zachariah R, Harries A, Schouten E, Hermann K, Mwagomba B, Massaquoi M: **Providing universal access to antiretroviral therapy in Thyolo, Malawi through task shifting and decentralization of HIV/AIDS care.** *Trop Med Int Health* 2010, **15**:1413-1420.
18. Cohen R, Lynch S, Bygrave H, Eggers E, Vlahakis N, Hilderbrand K, Knight L, Pillay P, Saranchuk P, Goemaere E, et al.: **Antiretroviral treatment outcomes from a nurse-driven, community-supported HIV/AIDS treatment programme in rural Lesotho: observational cohort assessment at two years.** *J Int AIDS Soc* 2009, **12**:23.
19. Decroo T, Telfer B, Biot M, Maikere J, Dezembro S, Cumba LI, das Dores C, Chu K, Ford N: **Distribution of antiretroviral treatment through self-forming groups of patients in Tete Province, Mozambique.** *J Acquir Immune Defic Syndr* 2011, **56**:e39-44.
20. Chang LW, Alamo S, Guma S, Christopher J, Suntoké T, Omasete R, Montis JP, Quinn TC, Juncker M, Reynolds SJ: **Two-year virologic outcomes of an alternative AIDS care model: evaluation of a peer health worker and nurse-staffed community-based program in Uganda.** *J Acquir Immune Defic Syndr* 2009, **50**:276-282.
21. Chang LW, Kagaayi J, Nakigozi G, Ssempijja V, Packer AH, Serwadda D, Quinn TC, Gray RH, Bollinger RC, Reynolds SJ: **Effect of peer health workers on AIDS care in Rakai, Uganda: a cluster-randomized trial.** *PLoS One* 2010, **5**:e10923.
22. Chang LW, Njie-Carr V, Kalenge S, Kelly JF, Bollinger RC, Alamo-Talisuna S: **Perceptions and acceptability of mHealth interventions for improving patient care at a community-based HIV/AIDS clinic in Uganda: a mixed methods study.** *AIDS Care* 2013, **25**:874-880.
23. Coetzee D, Boulle A, Hildebrand K, Asselman V, Van Cutsem G, Goemaere E: **Promoting adherence to antiretroviral therapy: the experience from a primary care setting in Khayelitsha, South Africa.** *Aids* 2004, **18 Suppl 3**:S27-31.
24. Duwell MM, Knowlton AR, Nachega JB, Efron A, Goliath R, Morroni C, Maartens G, Chaisson RE: **Patient-nominated, community-based HIV treatment supporters: patient perspectives, feasibility, challenges, and factors for success in HIV-infected South African adults.** *AIDS Patient Care STDS* 2013, **27**:96-102.

25. Jaffar S, Amuron B, Foster S, Birungi J, Levin J, Namara G, Nabiryo C, Ndembu N, Kyomuhangi R, Opio A, et al.: **Rates of virological failure in patients treated in a home-based versus a facility-based HIV-care model in Jinja, southeast Uganda: a cluster-randomised equivalence trial.** *Lancet* 2009, **374**:2080-2089.
26. Kunutsor S, Walley J, Katabira E, Muchuro S, Balidawa H, Namagala E, Ikoona E: **Improving clinic attendance and adherence to antiretroviral therapy through a treatment supporter intervention in Uganda: a randomized controlled trial.** *AIDS Behav* 2011, **15**:1795-1802.
27. Luque-Fernandez MA, Van Cutsem G, Goemaere E, Hilderbrand K, Schomaker M, Mantangana N, Mathee S, Dubula V, Ford N, Hernan MA, et al.: **Effectiveness of patient adherence groups as a model of care for stable patients on antiretroviral therapy in Khayelitsha, Cape Town, South Africa.** *PLoS One* 2013, **8**:e56088.
28. Mills EJ, Lester R, Thorlund K, Lorenzi M, Muldoon K, Kanters S, Linnemayr S, Gross R, Calderon Y, Amico KR, et al.: **Interventions to promote adherence to antiretroviral therapy in Africa: a network meta-analysis.** *Lancet HIV* 2014, **1**:e104-111.
29. Nachega JB, Chaisson RE, Goliath R, Efron A, Chaudhary MA, Ram M, Morroni C, Schoeman H, Knowlton AR, Maartens G: **Randomized controlled trial of trained patient-nominated treatment supporters providing partial directly observed antiretroviral therapy.** *Aids* 2010, **24**:1273-1280.
30. Nakigozi G, Makumbi FE, Bwanika JB, Atuyambe L, Reynolds SJ, Kigozi G, Nalugoda F, Chang LW, Kiggundu V, Serwadda D, et al.: **Impact of Patient-Selected Care Buddies on Adherence to HIV Care, Disease Progression, and Conduct of Daily Life Among Pre-antiretroviral HIV-Infected Patients in Rakai, Uganda: A Randomized Controlled Trial.** *J Acquir Immune Defic Syndr* 2015, **70**:75-82.
31. Remien RH, Mellins CA, Robbins RN, Kelsey R, Rowe J, Warne P, Chowdhury J, Lalkhen N, Hoppe L, Abrams EJ, et al.: **Masivukeni: development of a multimedia based antiretroviral therapy adherence intervention for counselors and patients in South Africa.** *AIDS Behav* 2013, **17**:1979-1991.
32. Robbins RN, Mellins CA, Leu CS, Rowe J, Warne P, Abrams EJ, Witte S, Stein DJ, Remien RH: **Enhancing Lay Counselor Capacity to Improve Patient Outcomes with Multimedia Technology.** *AIDS Behav* 2015, **19 Suppl 2**:163-176.
33. Chang LW, Kagaayi J, Nakigozi G, Serwada D, Quinn TC, Gray RH, Bollinger RC, Reynolds SJ, Holtgrave D: **Cost analyses of peer health worker and mHealth support interventions for improving AIDS care in Rakai, Uganda.** *AIDS Care* 2013, **25**:652-656.
34. Thompson MA, Mugavero MJ, Amico KR, Cargill VA, Chang LW, Gross R, Orrell C, Altice FL, Bangsberg DR, Bartlett JG, et al.: **Guidelines for improving entry into and retention in care and antiretroviral adherence for persons with HIV: evidence-based recommendations from an International Association of Physicians in AIDS Care panel.** *Ann Intern Med* 2012, **156**:817-833, w-284, w-285, w-286, w-287, w-288, w-289, w-290, w-291, w-292, w-293, w-294.
35. Vandembroucke JP, von Elm E, Altman DG, Gotzsche PC, Mulrow CD, Pocock SJ, Poole C, Schlesselman JJ, Egger M: **Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration.** *Int J Surg* 2014, **12**:1500-1524.

36. Higgins JPT, Green S: *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]* London: The Cochrane Collaboration; 2011.
37. Centre TNC: *Review Manager (RevMan) [Computer program], version 5.3.* Copenhagen: The Cochrane Collaboration; 2014.
38. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D: **The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration.** *Bmj* 2009, **339**:b2700.
39. Kiweewa FM, Wabwire D, Nakibuuka J, Mubiru M, Bagenda D, Musoke P, Fowler MG, Antelman G: **Noninferiority of a task-shifting HIV care and treatment model using peer counselors and nurses among Ugandan women initiated on ART: evidence from a randomized trial.** *J Acquir Immune Defic Syndr* 2013, **63**:e125-132.
40. Mfinanga S, Chanda D, Kivuyo SL, Guinness L, Bottomley C, Simms V, Chijoka C, Masasi A, Kimaro G, Ngowi B, et al.: **Cryptococcal meningitis screening and community-based early adherence support in people with advanced HIV infection starting antiretroviral therapy in Tanzania and Zambia: an open-label, randomised controlled trial.** *Lancet* 2015, **385**:2173-2182.
41. Selke HM, Kimaiyo S, Sidle JE, Vedanthan R, Tierney WM, Shen C, Denski CD, Katschke AR, Wools-Kaloustian K: **Task-shifting of antiretroviral delivery from health care workers to persons living with HIV/AIDS: clinical outcomes of a community-based program in Kenya.** *J Acquir Immune Defic Syndr* 2010, **55**:483-490.
42. Taiwo BO, Idoko JA, Welty LJ, Otoh I, Job G, Iyaji PG, Agbaji O, Agaba PA, Murphy RL: **Assessing the virologic and adherence benefits of patient-selected HIV treatment partners in a resource-limited setting.** *J Acquir Immune Defic Syndr* 2010, **54**:85-92.
43. Coker M, Etiebet MA, Chang H, Awwal G, Jumare J, Musa BM, Babashani M, Habib AG, Dakum P, Abimiku AG, et al.: **Socio-Demographic and Adherence Factors Associated with Viral Load Suppression in HIV-Infected Adults Initiating Therapy in Northern Nigeria: A Randomized Controlled Trial of a Peer Support Intervention.** *Curr HIV Res* 2015, **13**:279-285.
44. Gross R, Zheng L, La Rosa A, Sun X, Rosenkranz SL, Cardoso SW, Ssali F, Camp R, Godfrey C, Cohn SE, et al.: **Partner-based adherence intervention for second-line antiretroviral therapy (ACTG A5234): a multinational randomised trial.** *Lancet HIV* 2015, **2**:e12-19.
45. Fatti G, Meintjes G, Shea J, Eley B, Grimwood A: **Improved survival and antiretroviral treatment outcomes in adults receiving community-based adherence support: 5-year results from a multicentre cohort study in South Africa.** *J Acquir Immune Defic Syndr* 2012, **61**:e50-58.
46. Fatti G, Shaikh N, Eley B, Grimwood A: **Improved virological suppression in children on antiretroviral treatment receiving community-based adherence support: a multicentre cohort study from South Africa.** *AIDS Care* 2014, **26**:448-453.
47. Franke MF, Kaigamba F, Socci AR, Hakizamungu M, Patel A, Bagiruwigize E, Niyigena P, Walker KD, Epino H, Binagwaho A, et al.: **Improved retention associated with**

**community-based accompaniment for antiretroviral therapy delivery in rural Rwanda.** *Clin Infect Dis* 2013, **56**:1319-1326.

48. Grimsrud A, Lesosky M, Kalombo C, Bekker LG, Myer L: **Implementation and Operational Research: Community-Based Adherence Clubs for the Management of Stable Antiretroviral Therapy Patients in Cape Town, South Africa: A Cohort Study.** *J Acquir Immune Defic Syndr* 2016, **71**:e16-23.

49. Grimwood A, Fatti G, Mothibi E, Malahlela M, Shea J, Eley B: **Community adherence support improves programme retention in children on antiretroviral treatment: a multicentre cohort study in South Africa.** *J Int AIDS Soc* 2012, **15**:17381.

50. Johnston V, Fielding K, Charalambous S, Mampho M, Churchyard G, Phillips A, Grant AD: **Second-line antiretroviral therapy in a workplace and community-based treatment programme in South Africa: determinants of virological outcome.** *PLoS One* 2012, **7**:e36997.

51. Kipp W, Konde-Lule J, Rubaale T, Okech-Ojony J, Alibhai A, Saunders DL: **Comparing antiretroviral treatment outcomes between a prospective community-based and hospital-based cohort of HIV patients in rural Uganda.** *BMC Int Health Hum Rights* 2011, **11 Suppl 2**:S12.

52. Massavon W, Barlow-Mosha L, Mugenyi L, McFarland W, Gray G, Lundin R, Costenaro P, Nannyonga MM, Penazzato M, Bagenda D, et al.: **Factors Determining Survival and Retention among HIV-Infected Children and Adolescents in a Community Home-Based Care and a Facility-Based Family-Centred Approach in Kampala, Uganda: A Cohort Study.** *Isrn aids* 2014, **2014**:852489.

53. Kaihin R, Kasatpibal N, Chitreechuer J, Grimes RM: **Effect of an Empowerment Intervention on Antiretroviral Drug Adherence in Thai Youth.** *Behav Med* 2015, **41**:186-194.

54. Munoz M, Bayona J, Sanchez E, Arevalo J, Sebastian JL, Arteaga F, Guerra D, Zeladita J, Espiritu B, Wong M, et al.: **Matching social support to individual needs: a community-based intervention to improve HIV treatment adherence in a resource-poor setting.** *AIDS Behav* 2011, **15**:1454-1464.

55. Bussmann H, Wester CW, Thomas A, Novitsky V, Okezie R, Muzenda T, Gaolathe T, Ndwapi N, Mawoko N, Widenfelt E, et al.: **Response to zidovudine/didanosine-containing combination antiretroviral therapy among HIV-1 subtype C-infected adults in Botswana: two-year outcomes from a randomized clinical trial.** *J Acquir Immune Defic Syndr* 2009, **51**:37-46.

56. Henderson KC, Hindman J, Johnson SC, Valuck RJ, Kiser JJ: **Assessing the effectiveness of pharmacy-based adherence interventions on antiretroviral adherence in persons with HIV.** *AIDS Patient Care STDS* 2011, **25**:221-228.

57. Peltzer K, Ramlagan S, Jones D, Weiss SM, Fomundam H, Chanetsa L: **Efficacy of a lay health worker led group antiretroviral medication adherence training among non-adherent HIV-positive patients in KwaZulu-Natal, South Africa: results from a randomized trial.** *Sahara j* 2012, **9**:218-226.

58. Mermin J, Ekwaru JP, Were W, Degerman R, Bunnell R, Kaharuza F, Downing R, Coutinho A, Solberg P, Alexander LN, et al.: **Utility of routine viral load, CD4 cell**

**count, and clinical monitoring among adults with HIV receiving antiretroviral therapy in Uganda: randomised trial.** *Bmj* 2011, **343**:d6792.

59. Mermin J, Were W, Ekwaru JP, Moore D, Downing R, Behumbiize P, Lule JR, Coutinho A, Tappero J, Bunnell R: **Mortality in HIV-infected Ugandan adults receiving antiretroviral treatment and survival of their HIV-uninfected children: a prospective cohort study.** *Lancet* 2008, **371**:752-759.

60. Moore DM, Yiannoutsos CT, Musick BS, Tappero J, Degerman R, Campbell J, Were W, Kaharuza F, Alexander LN, Downing R, et al.: **Determinants of early and late mortality among HIV-infected individuals receiving home-based antiretroviral therapy in rural Uganda.** *J Acquir Immune Defic Syndr* 2011, **58**:289-296.

61. Weidle PJ, Wamai N, Solberg P, Liechty C, Sendagala S, Were W, Mermin J, Buchacz K, Behumbiize P, Ransom RL, et al.: **Adherence to antiretroviral therapy in a home-based AIDS care programme in rural Uganda.** *Lancet* 2006, **368**:1587-1594.

62. Wools-Kaloustian KK, Sidle JE, Selke HM, Vedanthan R, Kemboi EK, Boit LJ, Jebet VT, Carroll AE, Tierney WM, Kimaiyo S: **A model for extending antiretroviral care beyond the rural health centre.** *J Int AIDS Soc* 2009, **12**:22.

63. Decroo T, Van Damme W, Kegels G, Remartinez D, Rasschaert F: **Are Expert Patients an Untapped Resource for ART Provision in Sub-Saharan Africa?** *AIDS Res Treat* 2012, **2012**:749718.

64. Achieng L, Musangi H, Ong'uti S, Ombegoh E, Bryant L, Mwiindi J, Smith N, Keiser P: **An observational cohort comparison of facilitators of retention in care and adherence to anti-retroviral therapy at an HIV treatment center in Kenya.** *PLoS One* 2012, **7**:e32727.

65. Amuron B, Coutinho A, Grosskurth H, Nabiryo C, Birungi J, Namara G, Levin J, Smith PG, Jaffar S: **A cluster-randomised trial to compare home-based with health facility-based antiretroviral treatment in Uganda: study design and baseline findings.** *Open AIDS J* 2007, **1**:21-27.

66. Bemelmans M, Baert S, Goemaere E, Wilkinson L, Vandendyck M, van Cutsem G, Silva C, Perry S, Szumilin E, Gerstenhaber R, et al.: **Community-supported models of care for people on HIV treatment in sub-Saharan Africa.** *Trop Med Int Health* 2014, **19**:968-977.

67. Bango F, Wilkinson L, Van Cutsem G, Cleary S: **Cost-effectiveness of ART adherence clubs for long-term management of clinically stable ART patients.** #A2427356. In *International Conference on AIDS and STIs in Africa (ICASA), 7-11 December 2013*. Edited by. Cape Town, South Africa; 2013.

68. Geng EH, Emenyonu N, Bwana MB, Glidden DV, Martin JN: **Sampling-based approach to determining outcomes of patients lost to follow-up in antiretroviral therapy scale-up programs in Africa.** *JAMA* 2008, **300**:506-507.

69. Chishinga N, Godfrey-Faussett P, Fielding K, Ayles H: **Effect of home-based interventions on virologic outcomes in adults receiving antiretroviral therapy in Africa: a meta-analysis.** *BMC Public Health* 2014, **14**:239.

70. Decroo T, Rasschaert F, Telfer B, Remartinez D, Laga M, Ford N: **Community-based antiretroviral therapy programs can overcome barriers to retention of patients**

**and decongest health services in sub-Saharan Africa: a systematic review.** *Int Health* 2013, **5**:169-179.

71. Apondi R, Bunnell R, Awor A, Wamai N, Bikaako-Kajura W, Solberg P, Stall RD, Coutinho A, Mermin J: **Home-based antiretroviral care is associated with positive social outcomes in a prospective cohort in Uganda.** *J Acquir Immune Defic Syndr* 2007, **44**:71-76.

72. Wouters E, Van Damme W, van Rensburg D, Masquillier C, Meulemans H: **Impact of community-based support services on antiretroviral treatment programme delivery and outcomes in resource-limited countries: a synthetic review.** *BMC Health Serv Res* 2012, **12**:194.

73. Marseille E, Kahn JG, Pitter C, Bunnell R, Epalatai W, Jawe E, Were W, Mermin J: **The cost effectiveness of home-based provision of antiretroviral therapy in rural Uganda.** *Appl Health Econ Health Policy* 2009, **7**:229-243.

## Tables

**Table 1 Table of included studies**

Study	Type	Age –median (years)		Countries included	Duration of study	Sample size (n)	Participants / healthcare service description	
		Intervention	Comparator				Intervention	Comparator
Chang et al. [21]	Cluster-randomized trial	35.5	34.0	South Africa	27 months; May 2006 - July 2008	1,336	Patients with Community-based peer health workers	Control group-accessing standard of care
Fatti et al. [45]	Cohort	35.1	34.6	South Africa	69 months ; January 2004 - September 2010	66,953	Patients who received community-based adherence support	Patients without community-based adherence support
Fatti et al. [46]	Cohort	6.3	6.6	South Africa	69 months; January 2004 - September 2010	4,853	Patients who received community-based adherence support	Patients without community-based adherence support
Franke et al. [47]	Prospective cohort	37.0	37.0	Rwanda	15 months; June 2007 to August 2008	610	Community-Based Accompaniment	Clinic-Based Care
Grimsrud et al. [48]	Cohort	33.9	33.2	South Africa	20 months; May 2012 - December 2013	8,150	Community based adherence club	Community Health Centre
Grimwood et al. [49]	Cohort	6.8	6.2	South Africa	57 months; January 2004 - September 2009	3563	Children with patient advocates	Children without patient advocates
Jaffar et al. [25]	Cluster-randomized equivalence trial	37.0	38.0	Uganda	48 months; February 2005 - January 2009	1,453	Home-based care	Facility based care
Johnston et al [50]	Retrospective cohort	36.0	43.0	South Africa	75 months; January 2003 - March 2010	417	Community cohort	Workplace cohort

Kipp et al. [51]	Cohort	36.8	34.8	Uganda	15 months; March 2006 - May 2007	385	Centre/Community-based cohort	Hospital-based cohort
Kiweewa et al. [39]	Randomized controlled trial	27.0	27.8	Uganda	29 months; May 2007 - September 2009	92	Nurse–Peer model	Doctor–Counsellor model
Luque-Fernandez et al. [27]	Cohort	Not reporter	Not reported	South Africa	40 months; November 2007 - February 2011	2,834	Adherence clubs	Traditional clinic-based care
Massavon et al. [52]	Retrospective cohort	91.0 months	45.9 months	Uganda	2003-2010	1,623	Community home-based care approach	Facility-based family-centred approach
Mfinanga et al. [40]	Open-label, randomized controlled trial	38.0 35.0	37.0 35.0	Tanzania Zambia	19 months; February, 2012 - September, 2013	1,999	Clinic plus community support	Standard care
Nachegea et al. [29]	Open-label, randomized controlled trial	35.7	36.7	South Africa	42 months; February 2005 – July 2008	274	Directly observed therapy (DOT-ART) arm	Self-administered ART (Self-ART arm)
Selke et al. [41]	Cluster randomized controlled clinical trial	38.7	37.5	Kenya	26 months; March 2006 - April 2008.	208	Community Care Coordinators arm patients	Standard of Care arm patients
Taiwo et al [42]	Randomized controlled trial	Not reported	Not reported	Nigeria	19 months; June 2006 - December 2007	499	Treatment partner–assisted ART	Patient-administered standard of care ART
Gross et al. [44]	Randomized controlled trial	38	37	Botswana, Brazil, Haiti, Peru, South Africa, Uganda, Zambia, Zimbabwe	30 months; April 2009- September 2011	259	Partner-based modified directly observed therapy	Standard of care
Nakigozi et al [30]	Randomized controlled trial	37.6	37	Uganda	15 months; October 2010 - January 2012	1209	Patient-selected care buddy	Standard of care

Kaihin et al [53]	Cohort	18.2	19.4	Thailand	4 months; April–July 2011	46	Experimental Group (Empowerment Intervention)	Control
Kunutsor et al [26]	Randomized controlled trial	39.1	39.2	Uganda	8 months; March–September 2010	174	Standard adherence intervention package plus treatment supporter intervention	Standard intervention package
Munoz et al [54]	Cohort	31.7	31.9	Peru	17 months; December 2005 - April 2007	120	Community-based accompaniment with supervised antiretrovirals	Control
Coker et al [43]a	Randomized controlled trial	Not reported	Not reported	Nigeria	18 months; August 2006–January 2008	400	Peer educators arm	Standard of care
Coker et al [43]b	Randomized controlled trial	Not reported	Not reported	Nigeria	18 months; August 2006–January 2008	400	Home visits and peer educators arm	Standard of care

---

**Table 2 Risk of bias in included studies of cohort studies**


<b>Study</b>	<b>Selection (sample population)</b>	<b>Selection (participation rate)</b>	<b>Performance bias (outcome assessment)</b>	<b>Performance bias (analytical methods to control for bias)</b>	<b>Other form of bias</b>
Fatti et al. [45]	High	Low	Unclear	Low	High
Fatti et al. [46]	High	Low	Unclear	Low	High
Franke et al. [47]	High	Low	Unclear	Low	High
Grimsrud et al. [48]	High	Low	Low	Low	High
Grimwood et al. [49]	High	Low	Unclear	Low	High
Johnston et al [50]	High	Low	Unclear	Low	High
Kipp et al. [51]	High	Low	Low	Low	High
Luque-Fernandez et al. [27]	High	Low	Unclear	Low	High
Massavon et al. [52]	High	Low	Low	Low	High
Kaihin et al [53]	High	Low	Low	Low	High
Munoz et al. [54]	High	Low	Low	Low	High

**Table 3 Risk of bias in included studies of randomized controlled trials**


<b>Study</b>	<b>Random sequence generation (selection bias)</b>	<b>Allocation concealment (selection bias)</b>	<b>Blinding of outcome assessors (performance bias)</b>	<b>Incomplete outcome data (attrition bias)</b>	<b>Selective reporting (reporting bias)</b>	<b>Other bias</b>
Chang et al. [21]	Low	Unclear	Unclear	Low	Low	Low
Jaffar et al. [25]	Low	Low	Unclear	Low	Low	Low
Kiweewa et al. [39]	Low	Low	Unclear	Low	Low	Low
Mfinanga et al. [40]	Low	Low	Unclear	Low	Low	Low
Nachega et al. [29]	Low	Low	Unclear	Low	Low	Low
Selke et al. [41]	Low	Low	Unclear	Low	Low	Low
Taiwo et al. [42]	Low	Unclear	Unclear	Low	Low	Low
Gross et al. [44]	Low	Low	Low	Low	Low	Low
Nakigozi at el. [30,44]	Low	Low	High	Low	Low	Low
Kunutsor et al. [26]	Low	Low	Unclear	Low	Low	Low
Coker et al. [43]	Low	Low	Unclear	High	Low	Low

## Figure legends


**Figure 1: PRISMA flow for study selection**




**Figure 2: Forest plot of optimal ART adherence comparing community-based ART versus facility-based ART**




**Figure 3: Forest plot of virologic suppression comparing community-based ART versus facility-based ART**




**Figure 4: Forest plot of retention in care comparing community-based ART versus facility-based ART**



**Figure 5: Forest plot of change in CD4 count comparing community-based ART versus facility-based ART**



**Figure 6: Forest plot of all-cause mortality comparing community-based ART versus facility-based ART**



**Figure 7: Forest plot of loss to follow-up comparing community-based ART versus facility-based ART**

