

**Original citation:**

Zambon, P., Palani, S., Kamnev, A. and Balasubramanian, M. K. (2017) Myo2p is the major motor involved in actomyosin ring contraction in fission yeast. *Current Biology*, 27 (3). R99-R100. doi:10.1016/j.cub.2016.12.024

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/85319>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 4.0 International license (CC BY 4.0) and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/4.0/>

**A note on versions:**

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)


Correspondence

# Myo2p is the major motor involved in actomyosin ring contraction in fission yeast

Paola Zamboni<sup>1,2,\*</sup>, Saravanan Palani<sup>1,2,\*</sup>, Anton Kamnev<sup>1</sup>, and Mohan K. Balasubramanian<sup>1,3,\*</sup>

Cytokinesis in many eukaryotes requires an actomyosin-based contractile ring [1]. In fission yeast, cytokinesis involves the type II myosins Myo2p and Myp2p and the type V myosin Myo51p [2]. A recent study by Laplante *et al.* [3], using deletion mutants of *myp2* and *myo51* and the mis-sense mutant *myo2-E1* [4], concluded that each myosin has distinct functions and proposed that Myp2p plays the dominant role in actomyosin ring contraction. Here we present evidence that Myo2p, not Myp2p, is likely to be the major motor driving actomyosin ring contractility. Since the previous work [3] was performed at 25°C, the permissive temperature for *myo2-E1*, we compared cytokinesis timings in *myo2-E1* and *myo2Δ* at 25°C and found that *myo2-E1* is only partially compromised at 25°C. Furthermore, we find that *myp2Δ* and *myp2Δ myo51Δ* double mutants contract actomyosin rings at ~90% of the rate of wild-type cells at 30°C and 36°C, suggesting that Myp2p plays a minimal role in ring contraction at these temperatures. Finally, ring contraction in our *myo2-E1* strain took longer at 25°C than previously reported [3]. Although faster-acting alleles of *myo2* will be required to evaluate its contribution at 25°C, our work establishes that Myo2p is the major motor involved in ring contraction, under most, if not all, conditions.

Work from several laboratories has shown that *myo2-E1* forms healthy colonies at 25°C but fails to do so at increased temperatures [4,5]. Since previous work has shown that Myo2-E1p (the product of *myo2-E1*) neither



**Figure 1. Actomyosin ring kinetics in *S. pombe* myosin mutants.**

(A,C,E) Time taken for various steps in cytokinesis (coalescence of nodes into a ring, dwell time before contraction, and contraction). (B,D,F) Ring contraction rates. In all cases, Rlc1p-3GFP was used as a marker of cytokinetic nodes and the actomyosin ring. Note that improper rings that underwent aberrant contraction were detected in *myo2-E1* mutants (at 30°C and 36°C), *myo2-E1 myp2Δ* mutants (at 25°C, 30°C and 36°C) and *myo2-E1 myo51Δ* mutants (at 25°C, 30°C and 36°C).

binds actin filaments nor has ATPase activity *in vitro* at 25°C [6], Laplante *et al.* [3] inferred that *myo2-E1* should also be severely compromised *in vivo* at 25°C. It is known that *in vitro* activity and *in vivo* function of products of mutant alleles are not necessarily correlated [7]. Furthermore, Laplante *et al.* [3] extrapolate that, since *myo2-E1* is a severely compromised allele of *myo2*, the mGFP-tagged version (mGFP-*myo2-E1*) used in their work would also be severely compromised. The data in Figure 1D of the accompanying response of Laplante and Pollard and in Figure S1A in Laplante *et al.* [3] suggest, however, that the mGFP tag partially rescues *myo2-E1*.

We first investigated whether *myo2-E1* is an appropriate allele to understand Myo2p function at 25°C. We imaged the dynamics of the actomyosin ring component Blt1p, which localizes to precursor nodes and actomyosin rings [8], in germinating *myo2Δ* spores, with the rationale that for its use in comparative studies it should have a phenotype comparable

to *myo2Δ*. Actomyosin rings never assembled in *myo2Δ* cells, even after 4 hours of imaging, whereas germinated wild-type spores assembled normal actomyosin rings (Figure S1A and Movie S1 in Supplemental Information). However, contractile actomyosin rings assembled in *myo2-E1* cells and importantly even in *myo2-E1 myo51Δ* cells [3] (Figure S1B, see 36'–54' time points, and Movie S2, see 35'–55' time points), suggesting that Myo2-E1p should retain some activity for ring assembly through the search, capture, pull and release (SCPR) mechanism [9]. Thus, *myo2-E1* is not a severely compromised allele at 25°C and is inappropriate for investigating the contribution of Myo2p at 25°C.

We then compared the time taken for various aspects of cytokinesis in wild-type cells, the partially active *myo2-E1* mutant, the *myp2Δ myo51Δ* double mutant, and double mutants combining the partially compromised *myo2-E1* and *myp2Δ* or *myo51Δ* at 25°C (as a control), and at 30°C and 36°C. Through DNA sequencing and



genetic crosses, we ensured that all the *myo2-E1* strains used only carried the previously described G345R mutation and did not carry second-site modifiers.

At 25°C, we made similar observations to Laplante *et al.* [3] when ring assembly, maturation, and contraction times were scored in wild-type cells, and in *myp2Δ*, and *myo51Δ* mutants, but not in *myo2-E1* and *myp2Δ myo51Δ* mutants (Figures 1A and S1B, and Movie S2). Ring assembly time doubled in *myo51Δ* compared with wild-type cells, consistent with previous work reporting an ancillary role for Myo51p in ring assembly [10]. Ring contraction was slower in *myp2Δ* mutants (53.9 ± 7.5 minutes). In our analysis, ring contraction was not as severely affected in *myp2Δ myo51Δ* as reported in Laplante *et al.* [3]. Importantly, even the partially compromised *myo2-E1* (containing Myp2p and Myo51p) had a stronger defect than *myp2Δ* in ring contraction and took ~86 mins for ring contraction, suggesting that Myo2p rather than Myp2p played a major role in ring contraction even at 25°C, with Myp2p playing an ancillary role in ring contraction at 25°C (Figures 1A,B and S1B and Movie S2). The strong additive effect in *myo2-E1 myo51Δ* and *myo2-E1 myp2Δ* mutants made it difficult to demarcate different steps in cytokinesis, but the entire process of improper cytokinesis took a comparable amount of time in *myo2-E1*, *myo2-E1 myp2Δ*, and *myo2-E1 myo51Δ* mutants (125–150 minutes).

Ring assembly was defective in *myo2-E1*, *myo2-E1 myo51Δ*, and *myo2-E1 myp2Δ* mutants grown at 30°C or 36°C, confirming the essential role of Myo2p in actomyosin ring assembly (Figures 1C,E and S1C,D and Movie S2). Ring contraction times in *myp2Δ*, *myo51Δ*, and *myp2Δ myo51Δ* mutants grown at 30°C or 36°C were comparable to those in wild-type cells, with ring contraction rates at least 90% of that seen in wild-type cells (Figures 1D,F and S1C,D and Movie S2). These data further reinforced the view that Myp2p is not the major motor involved in ring contraction at 30°C and 36°C.

Laplante and Pollard (accompanying response) have shown that Rlc1p-3GFP shows adverse interactions specifically

with *myo2-E1* and suggest this may have increased the ring contraction time in *myo2-E1 rlc1-3GFP* at 25°C in our work. However, the fact that *rlc1-3GFP* exacerbates the cytokinesis defect of the *myo2-E1* mutant (but not wild-type cells) is itself further evidence that *myo2-E1* is a weak allele that is further weakened by its interaction with Rlc1p-3GFP *in vivo*. This further weakened Myo2-E1p-Rlc1p-3GFP complex causes a significant reduction in ring contraction rate, strengthening our conclusion that Myo2p is the main motor involved in ring contraction.

What then are the specialized functions of Myo2p, Myp2p, and Myo51p? We and others have established that Myo2p is central to ring assembly at all temperatures (current work and [2]), with Myo51p playing an ancillary role [10]. Our analysis has established that ring contraction is independent of Myp2p at 30°C or 36°C. Myp2p may perform an ancillary role in ring contraction at these temperatures. As reported [3], ring contraction is slower in cells lacking Myp2p at 25°C, consistent with previous work that Myp2p is specialized for cytokinesis at lower temperatures [2]. In our work, actomyosin ring contraction in the *myo2-E1* mutant took longer than in the *myp2Δ* mutant at 25°C, suggesting that Myo2p is likely the major motor involved in ring contraction even at 25°C. However, analysis of *myo2-E1* mutants to inform the role of Myo2p in ring contraction should be considered to underestimate Myo2p's function, since *myo2Δ* cells and *myo2-E1 rlc1-3GFP* cells have a stronger phenotype than *myo2-E1* at 25°C. Fast-acting stronger conditional mutant alleles of Myo2p will be essential to investigate the precise role of Myo2p in ring contraction at 25°C.

#### SUPPLEMENTAL INFORMATION

Supplemental Information includes experimental procedures, one figure and two movies and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2016.12.024>.

#### AUTHOR CONTRIBUTIONS

P.Z. and S.P., conception and design, acquisition of data, analysis and

interpretation of data, drafting or revising the article; A.K., analysis tool development and interpretation of data; M.K.B., conceived the project, conception and design, analysis and interpretation of data; S.P. and M.K.B., wrote the manuscript; all authors reviewed the manuscript.

#### ACKNOWLEDGMENTS

We thank Ting Gang Chew and Dan McCollum for critical reading of the manuscript. This work was funded by Wellcome Trust Senior Investigator Award, European Research Council Advanced grant, and a Royal Society Wolfson merit Award to M.K.B.

#### REFERENCES

- Cheffings, T.H., Burroughs, N.J., and Balasubramanian, M.K. (2016). Actomyosin ring formation and tension generation in eukaryotic cytokinesis. *Curr. Biol.* 26, R719–R737.
- Win, T.Z., Mulvihill, D.P., and Hyams, J.S. (2002). Take five: a myosin class act in fission yeast. *Cell Motil. Cytoskeleton* 57, 53–56.
- Laplante, C., Berro, J., Karatekin, E., Hernandez-Leyva, A., Lee, R., and Pollard, T.D. (2015). Three myosins contribute uniquely to the assembly and constriction of the fission yeast cytokinetic contractile ring. *Curr. Biol.* 25, 1955–1965.
- Balasubramanian, M.K., McCollum, D., Chang, L., Wong, K.C., Naqvi, N.I., He, X., Sazer, S., and Gould, K.L. (1998). Isolation and characterization of new fission yeast cytokinesis mutants. *Genetics* 149, 1265–1275.
- Coffman, V.C., Nile, A.H., Lee, I.J., Liu, H., and Wu, J.Q. (2009). Roles of formin nodes and myosin motor activity in Mid1p-dependent contractile-ring assembly during fission yeast cytokinesis. *Mol. Biol. Cell* 20, 5195–5210.
- Lord, M., and Pollard, T.D. (2004). UCS protein Rng3p activates actin filament gliding by fission yeast myosin-II. *J. Cell Biol.* 167, 315–325.
- Reed, S.I., Hadwiger, J.A., and Lorincz, A.T. (1985). Protein kinase activity associated with the product of the yeast cell division cycle gene CDC28. *Proc. Natl. Acad. Sci. USA* 82, 4055–4059.
- Guzman-Vendrell, M., Baldissard, S., Almonacid, M., Mayeux, A., Paoletti, A., and Moseley, J.B. (2013). Bit1 and Mid1 provide overlapping membrane anchors to position the division plane in fission yeast. *Mol. Cell Biol.* 33, 418–428.
- Vavylonis, D., Wu, J.Q., Hao, S., O'Shaughnessy, B., and Pollard, T.D. (2008). Assembly mechanism of the contractile ring for cytokinesis by fission yeast. *Science* 319, 97–100.
- Huang, J., Huang, Y., Yu, H., Subramanian, D., Padmanabhan, A., Thadani, R., Tao, Y., Tang, X., Wedlich-Soldner, R., and Balasubramanian, M.K. (2012). Nonmedially assembled F-actin cables incorporate into the actomyosin ring in fission yeast. *J. Cell Biol.* 199, 831–847.

<sup>1</sup>Division of Biomedical Sciences, Warwick Medical School, University of Warwick, Coventry, CV4 7AL, UK.

<sup>2</sup>P.Z. and S.P. contributed equally to this work.

<sup>3</sup>Lead Contact.

\*E-mail: [M.K.Balasubramanian@warwick.ac.uk](mailto:M.K.Balasubramanian@warwick.ac.uk) (M.K.B.), [S.Palani@warwick.ac.uk](mailto:S.Palani@warwick.ac.uk) (S.P.)