Original citation:

Grothe, Michel J., Scheef, Lukas, Bäuml, Josef, Meng, Chun, Daamen, Marcel, Baumann, Nicole, Zimmer, Claus, Teipel, Stefan, Bartmann, Peter, Boecker, Henning, Wolke, Dieter, Wohlschläger, Afra and Sorg, Christian. (2016) Reduced cholinergic basal forebrain integrity links neonatal complications and adult cognitive deficits after premature birth. *Biological Psychiatry*.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/85390>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

© 2016, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

Reduced cholinergic basal forebrain integrity links neonatal complications and adult cognitive deficits after premature birth

Michel J. Grothe¹, Lukas Scheef², Josef Bäuml^{3,4}, Chun Meng^{3,4}, Marcel Daamen², Nicole Baumann⁵, Claus Zimmer³, Stefan Teipel^{1,6}, Peter Bartmann⁷, Henning Boecker², Dieter Wolke^{5,8}, Afra Wohlschläger^{3,4}, Christian Sorg^{3,4,9}

¹German Center for Neurodegenerative Diseases (DZNE), 18147 Rostock, Germany

²Functional Neuroimaging Group, Department of Radiology, University Hospital Bonn, Bonn, Germany

Department of ³Neuroradiology and ⁹Psychiatry, Technische Universität München TUM, Ismaninger Strasse 22, 81675 Munich, Germany

⁴TUM-Neuroimaging Center of Klinikum rechts der Isar, Technische Universität München TUM, Ismaninger Strasse 22, 81675 Munich, Germany

⁵Department of Psychology, University of Warwick, Coventry, UK

⁶Department of Psychosomatic Medicine, University of Rostock, 18147 Rostock, Germany

⁷Department of Neonatology, University Hospital Bonn, Bonn, Germany

⁸Warwick Medical School, University of Warwick, Coventry, UK

Running Title: Reduced basal forebrain volume in premature-born adults

Corresponding Author:

Michel J. Grothe, Ph.D., German Center for Neurodegenerative Diseases (DZNE), Rostock Gehlsheimer Str. 20, 18147 Rostock, Germany; Tel. 01149-381-494-9479; Fax: 01149-381-494-9472; E-mail: michel.grothe@dzne.de

Manuscript requirements:

Abstract:	244 words
Text:	3989 words
References:	59
Figures:	4
Tables:	1

Abstract

Background: Premature-born individuals have an increased risk for long-term neurocognitive impairments. In animal models, the development of the cholinergic basal forebrain (cBF) is selectively vulnerable to adverse effects of perinatal stressors, and impaired cBF integrity results in lasting cognitive deficits. We hypothesized that in premature-born individuals cBF integrity is impaired and mediates adult cognitive impairments associated with prematurity.

Methods: We used MRI-based volumetric assessments of a cytoarchitectonically defined cBF region-of-interest to determine differences in cBF integrity between 99 adults who were born very preterm and/or with very low birth weight (VP/VLBW) and 106 term born controls from the same birth cohort. MRI-derived cBF volumes were studied in relation to neonatal clinical complications after delivery as well as intelligence measures (IQ) in adulthood.

Results: In VP/VLBW adults, cBF volumes were significantly reduced compared to term-born adults (-4.5% , $F(1, 202) = 11.82$, $p = 0.001$). Lower cBF volume in VP/VLBW adults was specifically associated with both neonatal complications ($r_{\text{part}}(92) = -0.35$, $p < 0.001$) and adult IQ ($r_{\text{part}}(88) = 0.33$, $p = 0.001$) even after controlling for global gray matter and white matter volume. In an additional path analytic model, cBF volume significantly mediated the association between neonatal complications and adult cognitive deficits.

Conclusions: Results provide first time evidence in humans that cBF integrity is impaired after premature birth and links neonatal complications with long-term cognitive outcome. Data suggest that cholinergic system abnormalities may play a relevant role for long-term neurocognitive impairments associated with premature delivery.

Keywords:

preterm birth, perinatal stress, substantia innominata, Nucleus basalis Meynert, intelligence, structural MRI

Introduction

Premature birth has a worldwide prevalence of more than 10% and is associated with an increased risk for birth complication and adverse long-term outcomes (1, 2). Adverse outcomes range from growth failure and higher morbidity to impaired neurocognitive development and lower socioeconomic status (3, 4). Risk for adverse outcomes increases substantially for very premature individuals, i.e. born very preterm (VP; gestational age <32 weeks) and/or with very low birth weight (VLBW; <1500g) (4, 5). Neuronal deficits may arise due to a higher propensity for brain injury caused by a higher risk for perinatal stressors such as hypoxic, ischemic, inflammatory, and/or infectious events in combination with developmental disturbances due to immature birth (2, 6).

However, brain mechanisms linking perinatal brain injury and neurodevelopment with long-term neurocognitive outcome after premature birth are still poorly understood. Previous research has shown that perinatal brain injury affects primarily the forebrain's white matter, characterized by immature and vulnerable glia cells and a relatively poor vascularization before 32 weeks of gestation (7). With increasing severity of injury, white matter impairments extend to gray matter, particularly affecting subcortical structures such as thalamus or basal ganglia (8-12). In this regard, the cholinergic basal forebrain (cBF) may be of special interest, since – at least in animal models – proper development of this system of cortically-projecting nuclei has been shown to be highly sensitive to perinatal stressors, such as glucocorticoid exposure (13), hypoxic-ischemic events (14), or neonatal psychosocial stress (15), resulting in structural and functional impairments that persist into adulthood. Moreover, due to the dependency of cortical circuits on cholinergic inputs, the cBF is highly relevant for cognitive functioning and intellectual capabilities (16), possibly linking perinatal brain injury to lasting effects on neurocognitive development. Thus, due to the higher risk for perinatal stressors in premature-born humans, the hypothesis arises that the cBF may be specifically involved in adverse effects of premature birth and associated with impaired neurocognitive development.

While there is currently little data regarding cBF involvement in human premature delivery, this question is of considerable clinical relevance given that cBF deficits may in principle be amenable to targeted interventions for lowering adverse consequences of premature birth.

Recent developments in stereotactic mapping of the cBF nuclei in the human brain have rendered this region accessible to MRI-based volumetric assessments, providing an in-vivo surrogate marker of cBF structural integrity (17-19). Application of these methods to the field of age-related neurodegenerative diseases has provided initial in-vivo evidence for the prominent role of cBF atrophy in the emergence of cognitive deficits in these conditions (20-24). Here, we used these techniques to assess reductions in cBF volumes in VP/VLBW adults as compared to term-born controls, and to further examine the relation of reduced cBF integrity with both neonatal complications and adult cognitive deficits.

Methods and Materials

Participants

Participants were recruited as part of the prospective Bavarian Longitudinal Study (BLS), which investigates a geographically defined whole-population sample of neonatal at-risk infants born in southern Bavaria (25). The current study sample consisted of 99 VP/VLBW individuals and 106 demographically matched term born individuals (control group) from the same obstetric hospitals who participated in a 26 year follow-up assessment including MRI scanning. More detailed information on the BLS and the selection process for the present study is reported in the supplementary material.

MRI assessments were carried out at the Department of Neuroradiology, Klinikum Rechts der Isar, Technische Universität München, Germany (N = 138), and the Department of Radiology, University Hospital Bonn, Germany (N = 67). The study was approved by the local ethics committees of the Klinikum Rechts der Isar and University Hospital Bonn. All study participants gave written informed consent and received travel expenses and a payment for attendance. Gestational age was determined from maternal reports of the last menstrual period and serial ultrasounds during pregnancy (26).

Assessment of neonatal complications

The intensity of neonatal treatment index (INTI) is a measure of neonatal complications (27). Daily assessments of care level, respiratory support, feeding dependency and neurological status (i.e. mobility, muscle tone, and neurological excitability) were carried out. Each of the 6 variables was scored on a 4-point rating scale (0-3). The INTI was computed as the mean score of daily ratings during the first 10 days of life or until a stable clinical state was reached, depending on which occurred first, ranging from 0 (best state) to 18 (worst state).

Cognitive performance assessment

Prior to and independent from the subsequent MRI examination, subjects from the BLS cohort were asked to take part in an assessment of global cognitive functioning at the age of 26 years by trained psychologists who were blinded to group membership. This included a short version of the German version of the Wechsler Adult Intelligence Scale-III (WAIS-III) (28), allowing the computation of a full-scale (F), verbal (V), and performance (P) IQ.

MRI data acquisition

At both sites, MRI data acquisition was initially performed on Philips Achieva 3T TX systems (Achieva, Philips, the Netherlands), using an 8-channel SENSE head coil. Due to a scanner upgrade, data acquisition in Bonn had to switch to Philips Ingenia 3T system with an 8-channel SENSE head coil after $N = 17$ participants. To account for possible confounds introduced by scanner differences, data analyses included scanner identities as covariates of no interest. Across all scanners, sequence parameters were kept identical. Scanners were checked regularly to provide optimal scanning conditions. Signal-to-noise ratio (SNR) was not significantly different between scanners (one-way ANOVA with factor “scanner-ID” [Bonn 1, Bonn 2, Munich]; $P = 0.811$). A high-resolution T1-weighted 3D-MPRAGE sequence (TI = 1300 ms, TR = 7.7 ms, TE = 3.9 ms, flip angle = 15° ; 180 sagittal slices, FOV = $256 \times 256 \times 180$ mm, reconstruction matrix = 256×256 ; reconstructed voxel size = $1 \times 1 \times 1$ mm³) was acquired.

MRI data processing

MRI data were processed using a fully automated processing pipeline within statistical parametric mapping software (SPM8, Wellcome Trust Center for Neuroimaging) and the VBM8-toolbox (<http://dbm.neuro.uni-jena.de/vbm/>) implemented in MATLAB R2013a (MathWorks, Natick, MA) as described previously (19, 24). First, MRI scans were segmented into gray matter (GM), white matter (WM) and cerebrospinal fluid partitions of 1.5 mm

isotropic voxel-size using the segmentation routine of the VBM8-toolbox. The resulting GM and WM partitions of each subject in native space were then high-dimensionally registered to stereotactic standard space (MNI space). Individual deformation fields resulting from the high-dimensional registration to the reference template were used to warp the GM segments and voxel-values were modulated for volumetric changes introduced by the normalization procedure, such that the total amount of GM volume present before deformation was preserved. All preprocessed GM maps passed a visual inspection for segmentation and registration accuracy.

Individual GM volumes of the cBF were extracted automatically from the warped GM segments by summing up the modulated GM voxel values within a region-of-interest (ROI) defined by a stereotactic atlas of BF cholinergic nuclei in MNI space. This atlas was derived from combined histology and in cranio post-mortem MRI of a 56-year old man without any evidence of cognitive decline or psychiatric illnesses prior to death (18). High-dimensional image registration of the post-mortem MRI data was used to accurately map the histologic delineations of cholinergic forebrain nuclei onto the MNI standard space template. The cBF mask comprises detailed outlines of different cholinergic subdivisions within the basal forebrain, including cell clusters corresponding to the medial septum, diagonal band, nucleus subputaminalis, and nucleus basalis Meynert (18). In the current study we investigated the entire volume of the cBF ROI, including all cholinergic subdivisions, as a proxy for overall BF cholinergic system integrity (Fig. 1).

To study the relative specificity of reduced cBF integrity in VP/VLBW adults and its association with neonatal complications and adult cognitive deficits, we also assessed the respective effects for global GM and WM volume. WM alterations have been described to be the most prevalent form of neuronal damage associated with preterm birth (7) and previous imaging studies found reductions in total WM volume to be a major determinant of IQ in VP/VLBW adolescents (11, 29). To account for differences in head size, cBF as well as

global GM and WM volumes were scaled to the intracranial volume (ICV), calculated as the sum of total segmented volumes of the GM, WM and cerebrospinal fluid partitions.

Statistical analysis

Statistical analyses were carried out using the software package IBM SPSS Statistics version 21. Statistical significance of the differences in ICV-normalized cBF, GM and WM volume between the VP/VLBW and control groups was assessed using analysis of covariance, controlling for MRI acquisition center. In addition to the group-wise volumetric comparisons, partial correlation coefficients of ICV-normalized cBF, GM and WM volume with INTI scores were calculated within the VP/VLBW group, controlling for MRI acquisition center and gender. Partial correlations between cBF volume and INTI score were also assessed when additionally controlling for global GM and WM volume. INTI score was not available for one subject who was excluded from this analysis.

Associations of cBF, GM and WM volume with cognitive test performance were assessed separately for VP/VLBW and controls using partial correlation coefficients controlled for MRI acquisition center and gender. Partial correlations between cBF volume and cognitive test performance were also assessed when additionally controlling for global GM and WM volume. FIQ was used as the primary neuropsychological outcome measure, and secondary correlation analyses assessed domain-specific associations with VIQ and PIQ. Cognitive test performance was not available for five VP/VLBW and three control subjects who were excluded from this analysis.

Mediation analysis

Additional mediation analysis was performed to test whether adult cognitive impairments associated with neonatal complications may be mediated by reduced cBF volume. First, associations between INTI scores and adult IQ scores were established using partial

correlations controlled for gender. In the mediation model, INTI score was entered as the causal variable, FIQ as the outcome variable, cBF volume as the mediator variable, and MRI acquisition center and gender as covariates of no interest. Path coefficients were estimated using (unstandardized) regression coefficients of multiple regression analyses, and statistical significance of the indirect effect was tested using a non-parametric bootstrap approach (with 1000 repetitions) to obtain 95% confidence intervals (30).

Results

Sample characteristics are summarized in Table 1. VP/VLBW and controls differed significantly in gestational age and birth weight. The VP/VLBW group had significantly lower IQ scores compared to the term-born controls.

Cholinergic basal forebrain volumes in VP/VLBW adults and their association with neonatal complications

cBF volumes were reduced in VP/VLBW adults by -4.5% on average compared to controls and this difference was statistically highly significant ($F(1, 202) = 11.82, p = 0.001$). Smaller cBF volumes were significantly correlated with a higher INTI score within the VP/VLBW group (Fig. 2; $r_{\text{part}}(94) = -0.27, p = 0.009$). By contrast, global GM volume did not differ between the two groups ($F(1,202) = 0.39, p = 0.53$) and WM volume was only marginally lower in VP/VLBW adults compared to controls (-1.2%, $F(1, 202) = 2.98, p = 0.086$). Neither global GM ($r_{\text{part}}(94) = 0.07, p = 0.50$) nor WM volumes ($r_{\text{part}}(94) = -0.16, p = 0.13$) were significantly associated with INTI scores, and the correlation between cBF volume and INTI score was increased after additionally controlling for global GM and WM volume ($r_{\text{part}}(92) = -0.35, p < 0.001$).

Association between cholinergic basal forebrain volumes and general cognitive function in VP/VLBW adults and term-born controls

Among VP/VLBW adults, smaller cBF volumes were significantly associated with a lower FIQ (Fig. 3; $r_{\text{part}}(90) = 0.33, p = 0.001$), as well as with a lower VIQ ($r_{\text{part}}(90) = 0.26, p = 0.01$) and PIQ ($r_{\text{part}}(90) = 0.31, p = 0.003$). Neither global GM nor WM volumes were associated with any IQ measure (all $p \geq 0.35$), and controlling for these global volumes did not markedly alter the association between cBF volume and intelligence scores ($r_{\text{part}}(88) = 0.33/0.25/0.33$, for FIQ/VIQ/PIQ, all $p \leq 0.02$). In term-born adults, neither cBF volumes ($r_{\text{part}}(99) = -0.17, p$

= 0.10) nor GM or WM volumes (all $p \geq 0.22$) were significantly associated with FIQ, or any other IQ measure.

Mediational effect of cBF volume on the association between neonatal complications and neurocognitive limitations in VP/VLBW adults

Among VP/VLBW adults, the INTI score was significantly associated with a lower FIQ ($r_{\text{part}}(90) = -0.21$, $p = 0.04$) and VIQ ($r_{\text{part}}(90) = -0.23$, $p = 0.03$), but not with a lower PIQ ($r_{\text{part}}(90) = -0.14$, $p = 0.19$). In a mediation analysis (Fig. 4), the effect of the INTI score on adult FIQ performance (total effect: $c = -0.72 \pm 0.33$, $t(90) = -2.18$, $p = 0.03$) was not significant when controlling for cBF volume (direct effect: $c' = -0.46 \pm 0.33$, $t(89) = -1.39$, $p = 0.17$), and the bootstrapped 95% confidence interval revealed the indirect effect (i.e. mediation: total - direct effect) to be significantly different from zero ($[-0.70, -0.08]$).

Discussion

Structural MRI-based volumetric assessments of a cytoarchitectonically defined cholinergic BF ROI were used to investigate cBF integrity in VP/VLBW adults and to link it with neonatal complications and adult cognition. Volume of the cBF was significantly reduced in VP/VLBW adults in comparison to term-born controls. Among VP/VLBW individuals, reduced cBF volume was associated with both higher neonatal treatment intensity and reduced IQ scores in adulthood. Notably, the association between neonatal complications and adult cognitive problems was found to be mediated by reduced cBF volume. These results provide first evidence in humans that the cBF is involved in long-term adverse effects of premature birth.

Neuropathological studies in premature-born newborns (9) and in-vivo imaging studies in premature-born newborns (10), children (31, 32), adolescents (11, 29, 33), and adults (12, 34-36) reported consistent volume reductions in subcortical brain regions, including brainstem, thalamus, and basal ganglia, but – to the best of our knowledge – volume differences of the cBF have not yet been investigated. By contrast, research in animal models demonstrated a high vulnerability of the developing cBF system against perinatal stressors such as increased glucocorticoids and hypoxic-ischemic events (13-15), which have a high prevalence in premature born infants (2). Our finding of reduced cBF volumes in VP/VLBW adults may reflect an impaired neurodevelopment of this brain system with long-term consequences for cognitive abilities. While we cannot infer developmental trajectories from the examined cross-sectional imaging data acquired in adulthood, a significant association between adult cBF volume reduction and the degree of neonatal complications further supports the idea that development of this brain region is particularly sensitive to perinatal stressors and neonatal treatment. However, more detailed studies in premature newborns and complementary longitudinal studies mapping cBF developmental trajectories are needed to substantiate this finding.

Combining in-vivo cBF volumetry with cognitive performance assessments, we found that reduced cBF volume in VP/VLBW adults was associated with lower intellectual abilities. This is in line with the widely supported role of the cBF system in higher cognitive functions, and may reflect an impaired cholinergic innervation of cognition-relevant cortical circuits (16). Clinicopathologic correlation studies in several neurologic dementing diseases characterized by impaired cBF integrity have demonstrated associations between cBF degeneration and dementia severity (37, 38). Furthermore, reduced cBF integrity has also been associated with more subtle cognitive deficits in conditions with less severe alterations of the cBF system, such as those associated with the normal aging process, developmental disturbances, or mild traumatic brain injury (38-40). Recent neuroimaging studies using MRI-based volumetric assessments of the cBF could replicate associations between reduced cBF integrity and cognitive deficits in neurodegenerative conditions in-vivo (20-24). Our findings in VP/VLBW adults suggest that reduced cBF integrity may be associated with cognitive limitations regardless of the etiology of the deficit.

Importantly, complementary assessments of global GM and WM volume revealed a relatively specific role of reduced cBF integrity for persisting intellectual limitations in VP/VLBW adults. Reductions in cerebral WM volume have been described to be the most prevalent form of neuronal damage associated with premature birth (7) and to be a major determinant of IQ in VP/VLBW adolescents (11, 29). However, in our sample of VP/VLBW adults, total WM volume was only marginally reduced compared to controls and was not related to intellectual impairments. While studies on premature birth-associated cerebral abnormalities in adulthood are still rare, two recent volumetric imaging studies on independent cohorts of VP/VLBW adults (34, 35) reported very similar observations of more subtle between-group differences compared to the data in infants (31, 32) and adolescents (11, 29, 33). Thus, rather than showing global differences, structural abnormalities in VP/VLBW adults were found to be restricted to specific subcortical GM and WM structures and select

cortical limbic and associative regions, indicating that some, but not all, aspects of the adverse effects of premature birth on cerebral structure may normalize during late development. Persistent reductions in cBF volume in VP/VLBW adults are consistent with experimental research in animal models demonstrating a weak potential of the cholinergic forebrain to recover from neonatal lesions (41). While our analyses demonstrate a relatively specific role of the cBF in premature birth-associated cognitive deficits that last into adulthood, one would not expect cBF volume to be the only factor related to variance in intellectual abilities among VP/VLBW adults. This is also reflected in the mild to moderate effect size of the observed associations between cBF volume and IQ scores ($r_{\text{part}} = 0.26 - 0.33$), indicating that impairments in other cortical and/or subcortical regions and their networking through the brain's white matter may affect intellectual abilities in VP/VLBW adults in at least partially independent ways (12, 34, 35). In addition, associations between structural/volumetric brain markers and mental capacities are generally indirect in nature and depend on the degree to which the measured structural abnormalities reflect the functional integrity of the neuronal system under study. In line with these considerations, variance in cBF volume among term-born controls was not found to be linked to differences in intellectual abilities, probably because volumetric differences among normally developed adult subjects are only weak indicators of neuronal functioning. Accordingly, variance in hippocampus volume has been shown to exhibit robust in-vivo associations with memory performance across a wide range of neurologic conditions (24, 42), but not so in healthy adults (43). In addition, the association between cBF volume and cognitive performance may be augmented by comorbid brain lesions in the VP/VLBW group (12, 34-36). Thus, experimental animal research (44) and more recent human in-vivo studies (45) have implicated impaired cBF integrity in diminished neural adaptation to white matter injury, possibly further contributing to worse cognitive outcomes in VP/VLBW adults.

A key finding of our study is that reduced cBF volume in VP/VLBW adults is associated with both neonatal complications and adult cognitive deficits. While our analyses are correlational and do not allow any direct inferences on directionality or causality, additional mediation analysis supports the idea of a causal influence of perinatal stressors on cBF integrity and inter-related cognitive problems. More specifically, factors during the perinatal period may affect development of the cBF; such alterations, in turn, likely influence the brain-behavior developmental trajectory towards worse outcomes; this influence realizes as an iterative process over time, when the child engages in interactions with the environment; interactions which themselves may be altered by the early brain insults. Such a model might open new ways in our thinking about adverse consequences of premature birth complications, both in terms of the prognosis of long-term risk for neurocognitive impairments and the treatment of such complications (4). Due to the lowered capacity of an abnormally developed cortical cholinergic input system to cope with additional disease-related stressors, aberrant development of the cholinergic BF system has been discussed as an important vulnerability factor for adult mental illnesses, such as schizophrenia (46) or late-life cognitive decline and dementia (47). Thus, future studies should carefully examine the possible role of cBF integrity in mediating increased risk profiles for neuropsychiatric disorders in VP/VLBW individuals. Concerning potential treatments, more attention should be paid to interventions which may prevent or reduce the negative impact of perinatal stressors on the cBF. For example, perinatal choline supplementation has been shown to protect cholinergic system integrity in animal models of increased vulnerability to cholinergic deficits, such as modeled by fetal alcohol exposure (48) or genetic predisposition (49). Interestingly, a recent placebo-controlled trial of perinatal dietary choline supplementation in healthy women demonstrated a positive effect on a psychophysiologic outcome marker related to cholinergic system development and schizophrenia risk (46). Early postnatal dietary interventions have already been shown to be able to ameliorate adverse effects of preterm birth on cognitive and brain volumetric outcome

variables in adolescence (50), but specific choline supplementations have not yet been considered. Beyond primary preventive perinatal diet supplementations, pharmacological interventions with pro-cholinergic drugs, such as nicotinic agonists or acetylcholinesterase inhibitors, might be additional options to ameliorate long-standing cognitive limitations in VP/VLBW adults. Cholinergic drugs are generally well tolerated, have demonstrated nootropic properties in healthy adults (51), and have shown beneficial effects on cognitive disturbances that are believed to be of cholinergic origin in diverse neurologic and psychiatric conditions (52-55).

A limitation of our study is that the volumetric cBF measurement is only an indirect marker of cholinergic system integrity. The correspondence of the analyzed brain region with the cholinergic space of the BF was determined based on stereotactic mapping of the forebrain's cholinergic nuclei using combined histology and post-mortem MRI (18). Notably, we found that the results of our study were robust against slight variations in the definition of the cBF ROI, as they could be fully reproduced using a previously published stereotactic mask of the cBF based on a different post-mortem brain (17) (data not shown). Nevertheless, one cannot exclude that the volumetric measurement may also reflect change in other non-cholinergic neuronal populations. For example, cholinergic neurons in the BF are intermingled, and interact, with cortically-projecting GABAergic neurons, which are increasingly being recognized for their critical role in BF-mediated modulation of cortical activity (56). Moreover, GABAergic neurons have also been reported to be particularly prone to preterm hypoxia (7). Thus, while the observed associations with neonatal complications and adult IQ scores underline the clinical and functional relevance of in-vivo detectable volumetric cBF changes after premature birth, the implications of these alterations for cholinergic neurotransmission and cortical function need to be studied in more detail. Pharmacologic challenge paradigms (57, 58) and functional imaging studies (24, 59) may be suitable analytic approaches to further investigate the nature of reduced cBF volume and its

association with neuro-cognitive function in premature-born adults. Secondly, one should note that the current study focuses on high-risk premature birth, as defined by VP and/or VLBW birth. This approach follows previous framings for defining high-risk prematurity, facilitating comparability with a wide spectrum of human prematurity studies (2, 5). While we demonstrated a link between neonatal medical complications, cBF, and IQ for VP/VLBW persons, we cannot disentangle which medical or prematurity factors (or their interactions) may be most determining for cBF volume and IQ in adulthood. This indeterminacy is partly based on our sample definition following combined VP/VLBW criteria for determining a general high-risk prematurity profile (2, 5), as well as on the employed INTI measure (27), which is a summary score of medical complications after birth without specifying the exact nature of these complications or their underlying causes. Thirdly, the sample of this MRI study was biased towards VP/VLBW adults with relatively lower neonatal complications and higher IQ compared to the entire BLS baseline cohort, given that individuals with more severe impairments were more likely to be excluded in initial screening for MRI or to decline to participate in MRI scanning. Thus, the reported differences compared to term-born controls may be considered conservative estimates of the differences in cBF volume in VP/VLBW adults.

In summary, we studied an in-vivo marker of structural cBF integrity in a large sample of VP/VLBW adults, and provide novel evidence for a previously unappreciated aspect of the neuronal changes underlying lasting cognitive limitations in this population. Further studies using complementary pharmacologic and functional imaging approaches for investigating alterations in the cholinergic system after premature birth and its implications for neuro-cognitive function are warranted.

Acknowledgments

We thank all current and former members of the Bavarian Longitudinal Study Group who contributed to general study organization, recruitment, and data collection, management and subsequent analyses, including (in alphabetical order): Stephan Czeschka, Claudia Grunzinger, Christian Koch, Diana Kurze, Sonja Perk, Andrea Schreier, Antje Strasser, Julia Trummer, and Eva van Rossum. We are grateful to the staff of the Department of Neuroradiology in Munich and the Department of Radiology in Bonn for their help in data collection. Most importantly, we thank all our study participants and their families for their efforts to take part in this study. This study was supported by the Chinese Scholar Council (CSC, File No: 2010604026 to C.M.) and the German Federal Ministry of Education and Science (BMBF 01ER0801 to P.B. and D.W., BMBF 01EV0710 to A.M.W., BMBF 01ER0803 to C.S.).

Financial Disclosures

The authors report no biomedical financial interests or potential conflicts of interest.

References

1. Blencowe H, Cousens S, Oestergaard MZ, Chou D, Moller AB, Narwal R, et al. (2012): National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *Lancet*. 379:2162-2172.
2. Volpe JJ (2009): Brain injury in premature infants: a complex amalgam of destructive and developmental disturbances. *Lancet Neurol*. 8:110-124.
3. D'Onofrio BM, Class QA, Rickert ME, Larsson H, Langstrom N, Lichtenstein P (2013): Preterm birth and mortality and morbidity: a population-based quasi-experimental study. *JAMA Psychiatry*. 70:1231-1240.
4. Nosarti C, Reichenberg A, Murray RM, Cnattingius S, Lambe MP, Yin L, et al. (2012): Preterm birth and psychiatric disorders in young adult life. *Arch Gen Psychiatry*. 69:E1-8.
5. Saigal S, Doyle LW (2008): An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet*. 371:261-269.
6. Deng W (2010): Neurobiology of injury to the developing brain. *Nat Rev Neurol*. 6:328-336.
7. Salmaso N, Jablonska B, Scafidi J, Vaccarino FM, Gallo V (2014): Neurobiology of premature brain injury. *Nat Neurosci*. 17:341-346.
8. Kinney HC, Haynes RL, Xu G, Andiman SE, Folkerth RD, Sleeper LA, et al. (2012): Neuron deficit in the white matter and subplate in periventricular leukomalacia. *Ann Neurol*. 71:397-406.
9. Pierson CR, Folkerth RD, Billiards SS, Trachtenberg FL, Drinkwater ME, Volpe JJ, et al. (2007): Gray matter injury associated with periventricular leukomalacia in the premature infant. *Acta Neuropathol*. 114:619-631.

10. Srinivasan L, Dutta R, Counsell SJ, Allsop JM, Boardman JP, Rutherford MA, et al. (2007): Quantification of deep gray matter in preterm infants at term-equivalent age using manual volumetry of 3-tesla magnetic resonance images. *Pediatrics*. 119:759-765.
11. Taylor HG, Filipek PA, Juranek J, Bangert B, Minich N, Hack M (2011): Brain volumes in adolescents with very low birth weight: effects on brain structure and associations with neuropsychological outcomes. *Dev Neuropsychol*. 36:96-117.
12. Meng C, Bauml JG, Daamen M, Jaekel J, Neitzel J, Scheef L, et al. (2016): Extensive and interrelated subcortical white and gray matter alterations in preterm-born adults. *Brain Struct Funct*. 221:2109-2121.
13. Emgard M, Paradisi M, Pirondi S, Fernandez M, Giardino L, Calza L (2007): Prenatal glucocorticoid exposure affects learning and vulnerability of cholinergic neurons. *Neurobiol Aging*. 28:112-121.
14. Johnston MV (1983): Neurotransmitter alterations in a model of perinatal hypoxic-ischemic brain injury. *Ann Neurol*. 13:511-518.
15. Aisa B, Gil-Bea FJ, Marcos B, Tordera R, Lasheras B, Del Rio J, et al. (2009): Neonatal stress affects vulnerability of cholinergic neurons and cognition in the rat: involvement of the HPA axis. *Psychoneuroendocrinology*. 34:1495-1505.
16. Hasselmo ME, Sarter M (2011): Modes and models of forebrain cholinergic neuromodulation of cognition. *Neuropsychopharmacology*. 36:52-73.
17. Teipel SJ, Flatz WH, Heinsen H, Bokde AL, Schoenberg SO, Stockel S, et al. (2005): Measurement of basal forebrain atrophy in Alzheimer's disease using MRI. *Brain*. 128:2626-2644.
18. Kilimann I, Grothe M, Heinsen H, Alho EJ, Grinberg L, Amaro E, Jr., et al. (2014): Subregional Basal forebrain atrophy in Alzheimer's disease: a multicenter study. *J Alzheimers Dis*. 40:687-700.

19. Grothe M, Heinsen H, Teipel SJ (2012): Atrophy of the cholinergic basal forebrain over the adult age range and in early stages of Alzheimer's disease. *Biol Psychiatry*. 71:805-813.
20. Grothe M, Zaborszky L, Atienza M, Gil-Neciga E, Rodriguez-Romero R, Teipel SJ, et al. (2010): Reduction of basal forebrain cholinergic system parallels cognitive impairment in patients at high risk of developing Alzheimer's disease. *Cereb Cortex*. 20:1685-1695.
21. Muth K, Schonmeyer R, Matura S, Haenschel C, Schroder J, Pantel J (2010): Mild cognitive impairment in the elderly is associated with volume loss of the cholinergic basal forebrain region. *Biol Psychiatry*. 67:588-591.
22. Grothe M, Heinsen H, Teipel S (2013): Longitudinal measures of cholinergic forebrain atrophy in the transition from healthy aging to Alzheimer's disease. *Neurobiol Aging*. 34:1210-1220.
23. Choi SH, Jung TM, Lee JE, Lee SK, Sohn YH, Lee PH (2012): Volumetric analysis of the substantia innominata in patients with Parkinson's disease according to cognitive status. *Neurobiol Aging*. 33:1265-1272.
24. Grothe MJ, Heinsen H, Amaro E, Jr., Grinberg LT, Teipel SJ (2016): Cognitive Correlates of Basal Forebrain Atrophy and Associated Cortical Hypometabolism in Mild Cognitive Impairment. *Cereb Cortex*. 26:2411-2426.
25. Wolke D, Meyer R (1999): Cognitive status, language attainment, and prereading skills of 6-year-old very preterm children and their peers: the Bavarian Longitudinal Study. *Dev Med Child Neurol*. 41:94-109.
26. Breeman LD, Jaekel J, Baumann N, Bartmann P, Wolke D (2015): Preterm Cognitive Function Into Adulthood. *Pediatrics*. 136:415-423.
27. Casaer P, Eggermont E (1985): Neonatal clinical neurological assessment. In: Harel S, Anastasiow NJ, editors. *The at-risk infant: Psycho/socio/medical aspects* Baltimore: Brooks, pp 197-220.

28. von Aster M, Neubauer A, Horn R (2006): *Wechsler Intelligenztest für Erwachsene - Deutschsprachige Bearbeitung und Adaptation des WAIS-III von David Wechsler*. Frankfurt/Main (Germany): Pearson.
29. Northam GB, Liegeois F, Chong WK, Wyatt JS, Baldeweg T (2011): Total brain white matter is a major determinant of IQ in adolescents born preterm. *Ann Neurol*. 69:702-711.
30. Preacher KJ, Hayes AF (2004): SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behav Res Methods Instrum Comput*. 36:717-731.
31. Zhang Y, Inder TE, Neil JJ, Dierker DL, Alexopoulos D, Anderson PJ, et al. (2015): Cortical structural abnormalities in very preterm children at 7 years of age. *Neuroimage*. 109:469-479.
32. Peterson BS, Vohr B, Staib LH, Cannistraci CJ, Dolberg A, Schneider KC, et al. (2000): Regional brain volume abnormalities and long-term cognitive outcome in preterm infants. *JAMA*. 284:1939-1947.
33. Nosarti C, Giouroukou E, Healy E, Rifkin L, Walshe M, Reichenberg A, et al. (2008): Grey and white matter distribution in very preterm adolescents mediates neurodevelopmental outcome. *Brain*. 131:205-217.
34. Nosarti C, Nam KW, Walshe M, Murray RM, Cuddy M, Rifkin L, et al. (2014): Preterm birth and structural brain alterations in early adulthood. *Neuroimage Clin*. 6:180-191.
35. Bjuland KJ, Rimol LM, Lohaugen GC, Skranes J (2014): Brain volumes and cognitive function in very-low-birth-weight (VLBW) young adults. *Eur J Paediatr Neurol*. 18:578-590.
36. Bauml JG, Daamen M, Meng C, Neitzel J, Scheef L, Jaekel J, et al. (2015): Correspondence Between Aberrant Intrinsic Network Connectivity and Gray-Matter Volume in the Ventral Brain of Preterm Born Adults. *Cereb Cortex*. 25:4135-4145.
37. Rogers JD, Brogan D, Mirra SS (1985): The nucleus basalis of Meynert in neurological disease: a quantitative morphological study. *Ann Neurol*. 17:163-170.

38. Schliebs R, Arendt T (2011): The cholinergic system in aging and neuronal degeneration. *Behav Brain Res.* 221:555-563.
39. Arciniegas DB (2003): The cholinergic hypothesis of cognitive impairment caused by traumatic brain injury. *Curr Psychiatry Rep.* 5:391-399.
40. Perry EK, Lee ML, Martin-Ruiz CM, Court JA, Volsen SG, Merrit J, et al. (2001): Cholinergic activity in autism: abnormalities in the cerebral cortex and basal forebrain. *Am J Psychiatry.* 158:1058-1066.
41. Pappas BA, Payne KB, Fortin T, Sherren N (2005): Neonatal lesion of forebrain cholinergic neurons: further characterization of behavioral effects and permanency. *Neuroscience.* 133:485-492.
42. Geuze E, Vermetten E, Bremner JD (2005): MR-based in vivo hippocampal volumetrics: 2. Findings in neuropsychiatric disorders. *Mol Psychiatry.* 10:160-184.
43. Van Petten C (2004): Relationship between hippocampal volume and memory ability in healthy individuals across the lifespan: review and meta-analysis. *Neuropsychologia.* 42:1394-1413.
44. Croxson PL, Browning PG, Gaffan D, Baxter MG (2012): Acetylcholine facilitates recovery of episodic memory after brain damage. *J Neurosci.* 32:13787-13795.
45. Ray NJ, Metzler-Baddeley C, Khondoker MR, Grothe MJ, Teipel S, Wright P, et al. (2015): Cholinergic Basal forebrain structure influences the reconfiguration of white matter connections to support residual memory in mild cognitive impairment. *J Neurosci.* 35:739-747.
46. Ross RG, Hunter SK, McCarthy L, Beuler J, Hutchison AK, Wagner BD, et al. (2013): Perinatal choline effects on neonatal pathophysiology related to later schizophrenia risk. *Am J Psychiatry.* 170:290-298.

47. Sarter M, Bruno JP (2004): Developmental origins of the age-related decline in cortical cholinergic function and associated cognitive abilities. *Neurobiol Aging*. 25:1127-1139.
48. Monk BR, Leslie FM, Thomas JD (2012): The effects of perinatal choline supplementation on hippocampal cholinergic development in rats exposed to alcohol during the brain growth spurt. *Hippocampus*. 22:1750-1757.
49. Ash JA, Velazquez R, Kelley CM, Powers BE, Ginsberg SD, Mufson EJ, et al. (2014): Maternal choline supplementation improves spatial mapping and increases basal forebrain cholinergic neuron number and size in aged Ts65Dn mice. *Neurobiol Dis*. 70:32-42.
50. Isaacs EB, Gadian DG, Sabatini S, Chong WK, Quinn BT, Fischl BR, et al. (2008): The effect of early human diet on caudate volumes and IQ. *Pediatr Res*. 63:308-314.
51. Heishman SJ, Kleykamp BA, Singleton EG (2010): Meta-analysis of the acute effects of nicotine and smoking on human performance. *Psychopharmacology (Berl)*. 210:453-469.
52. Singh J, Kour K, Jayaram MB (2012): Acetylcholinesterase inhibitors for schizophrenia. *Cochrane Database Syst Rev*. 1:CD007967.
53. Birks J (2006): Cholinesterase inhibitors for Alzheimer's disease. *Cochrane Database Syst Rev*. CD005593.
54. Rolinski M, Fox C, Maidment I, McShane R (2012): Cholinesterase inhibitors for dementia with Lewy bodies, Parkinson's disease dementia and cognitive impairment in Parkinson's disease. *Cochrane Database Syst Rev*. 3:CD006504.
55. Tenovuo O, Alin J, Helenius H (2009): A randomized controlled trial of rivastigmine for chronic sequels of traumatic brain injury-what it showed and taught? *Brain Inj*. 23:548-558.
56. Anaclet C, Pedersen NP, Ferrari LL, Venner A, Bass CE, Arrigoni E, et al. (2015): Basal forebrain control of wakefulness and cortical rhythms. *Nat Commun*. 6:8744.

57. Snyder PJ, Lim YY, Schindler R, Ott BR, Salloway S, Daiello L, et al. (2014): Microdosing of scopolamine as a "cognitive stress test": rationale and test of a very low dose in an at-risk cohort of older adults. *Alzheimers Dement.* 10:262-267.
58. Teipel SJ, Bruno D, Grothe MJ, Nierenberg J, Pomara N (2015): Hippocampus and basal forebrain volumes modulate effects of anticholinergic treatment on delayed recall in healthy older adults. *Alzheimers Dement: DADM.* 1:216-219.
59. Li CS, Ide JS, Zhang S, Hu S, Chao HH, Zaborszky L (2014): Resting state functional connectivity of the basal nucleus of Meynert in humans: in comparison to the ventral striatum and the effects of age. *Neuroimage.* 97:321-332.

Figure legends

Figure 1. Illustration of the cholinergic basal forebrain region of interest

The cholinergic basal forebrain region of interest as defined by a cytoarchitectonic map of cholinergic nuclei in the basal forebrain is superimposed on transparent renderings of the stereotactic standard space template (MNI space).

Figure 2. Association of intensity of neonatal treatment with cholinergic basal forebrain and global gray and white matter volume in preterm-born adults

Intensity of neonatal treatment (INTI) scores in preterm-born adults are plotted against cholinergic basal forebrain (cBF, top), global gray matter (GM, middle), and global white matter (WM, bottom) volumes, normalized for intracranial volume (head size). For visualization purposes, all plots include linear regression trend lines with their corresponding standardized regression coefficient and associated p-value. Only cBF volume shows a significant negative association with INTI scores, which is corroborated in covariate-controlled partial correlation analyses (see Results).

Figure 3. Association of intellectual abilities with cholinergic basal forebrain and global gray and white matter volume in preterm-born adults

Cholinergic basal forebrain (cBF, top), global gray matter (GM, middle), and global white matter (WM, bottom) volumes, normalized for intracranial volume (head size), are plotted against full-scale IQ scores of the Wechsler Adult Intelligence Scale-III in preterm-born adults. For visualization purposes, all plots include linear regression trend lines with their corresponding standardized regression coefficient and associated p-value. Only cBF volume shows a significant positive association with IQ scores, which is corroborated in covariate-controlled partial correlation analyses (see Results).

Figure 4. Cholinergic basal forebrain volume mediates between neonatal complications and neurocognitive limitations in preterm-born adults

Path diagrams illustrating the bivariate model of neonatal complications (INTI scores) affecting neurocognitive limitations (FIQ score) in preterm-born adults, which is being compared to an alternative model of cholinergic basal forebrain volume mediating the effect of neonatal complications on adult neurocognitive limitations. Paths are labeled with the (unstandardized) regression coefficients of the respective effect ($\pm$ SE). a: effect of causal variable on mediator; b: direct effect of mediator on outcome; c: total effect of causal variable on outcome; c': direct effect of causal variable on outcome. INTI: intensity of neonatal treatment index; FIQ: full-scale intelligence quotient of the Wechsler Adult Intelligence Scale-III.

Tables

Table 1. Sample characteristics of demographic, birth-related, and neuropsychologic variables

	VP/VLBW	Controls
N	99	106
Gender (M/F)	57/42	61/45
Age, yrs (SD)	26.7 (0.6)	26.8 (0.7)
Gestational Age, weeks (SD)	30.4 (2.1)**	39.7 (1.1)
Birth Weight, g (SD)	1325 (314) **	3391 (451)
INTI (SD)	11.6 (3.8)	-
Full-scale IQ (SD)	94.7 (12.4) **	102.7 (11.9)
Verbal IQ (SD)	99.5 (13.8)*	105.9 (14.4)
Performance IQ (SD)	90.2 (13.2)**	98.9 (10.1)

* Significantly different from control group at $p < 0.01$

** Significant different from control group at $p < 0.001$

Statistical significance of group differences was assessed using Fisher's exact test for categorical variables and two-sample t-tests for continuous variables. INTI, intensity of neonatal treatment index (arbitrary units: 0 denotes best, 18 denotes worst state); IQ, intelligence quotient; M/F, male/female; SD, standard deviation