

Original citation:

Ambrozio, Lucas, Carlotto, Alessandro and Sharp, Benjamin G.. (2017) A note on the index of closed minimal hypersurfaces of flat tori. Proceedings of the American Mathematical Society

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/90298>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

First published in Proceedings of the American Mathematical Society published by the American Mathematical Society. © 2017 American Mathematical Society.

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

A NOTE ON THE INDEX OF CLOSED MINIMAL HYPERSURFACES OF FLAT TORI

LUCAS AMBROZIO, ALESSANDRO CARLOTTO AND BEN SHARP

ABSTRACT. Generalizing earlier work by Ros in ambient dimension three, we prove an affine lower bound for the Morse index of closed minimal hypersurfaces inside a flat torus in terms of their first Betti number (with purely dimensional coefficients).

1. INTRODUCTION

Motivated by a variety of recent constructions of closed minimal hypersurfaces in positively curved Riemannian manifolds, and by the associated natural classification questions, we presented in [1] a study of the relation between their Morse index and their Betti numbers, namely those pieces of data respectively encoding the most basic analytic and topological information concerning the hypersurface in question. This relation is in fact the object of a conjecture due to Schoen and Marques-Neves [8, 11] that can be stated as follows: in any closed manifold of positive Ricci curvature there is a linear lower bound of the Morse index of a minimal hypersurface M^n in terms of its first Betti number, that is to say

$$(1.1) \quad \text{index}(M) \geq C b_1(M)$$

for some constant C only depending on the ambient manifold. We refer the reader to the introduction of [1] for a broader contextualization of this problem and for a discussion of the various cases for which we could verify this conjecture.

On the other hand, it is straightforward to observe that inequality (1.1) cannot hold true in the special but fundamental case of flat manifolds, as is seen by considering totally geodesic n -dimensional tori inside $(n + 1)$ -dimensional flat tori (in which case one has $\text{index}(M) = 0$ and $b_1(M) = n$ for any $n \geq 2$). In such a setting, the best one can hope for is instead an affine bound with a negative additive constant on the right-hand side. Up to now, an estimate of that sort was only obtained for $n = 2$ by A. Ros [12]. The scope of this note is to prove the following generalization of such a result:

Theorem 1. *Let M^n be a closed minimal hypersurface in a $(n + 1)$ -dimensional flat torus. Assume there is a point p in M^n where all principal curvatures are distinct. Then*

$$\text{index}(M) \geq \frac{2}{n(n+1)}(b_1(M) - (2n - 1)).$$

If $n = 2$ or $n = 3$, then the above inequality holds true without the assumption on the principal curvatures.

Clearly, there is a natural correspondence between minimal hypersurfaces in flat tori and $(n + 1)$ -periodic minimal hypersurfaces in the Euclidean space \mathbb{R}^{n+1} , a topic that has been

Mathematics Subject Classification (MSC 2010): Primary 53A10; Secondary 53C42, 49Q05.

thoroughly investigated with several interesting results: we refer the reader to the article by W. Meeks [9] and to the survey by D. Hoffman [5] as well as references therein for further details about the classical case $n = 2$. Remarkably, lots of interesting examples in \mathbb{R}^3 are actually known, the most basic ones being provided by the Schwarz P and D surfaces, the latter class ensuring that the estimate of Theorem 1 is actually sharp since in that case one can find, in a fundamental domain, that the Morse index equals one (this follows from the work of M. Ross [13]) while the topology is that of a genus three orientable surface. By contrast, the construction of periodic minimal hypersurfaces in ambient dimension at least four is a fascinating theme of current research: see in particular the recent work by Choe and Hoppe [4] for certain higher-dimensional analogues of the aforementioned classical surfaces and related comments therein.

In applying Theorem 1 to obtain classification results, it is also useful to know that non-trivial minimal hypersurfaces inside a flat torus must have sufficiently large first Betti number:

Theorem 2. *(Cf. Theorem 1 in [6], and Theorem 4.1 in [3]) Let M^n be a closed minimal hypersurface in a $(n + 1)$ -dimensional flat torus. Then $b_1(M) \geq n + 1$ unless M^n is a flat totally geodesic n -dimensional torus.*

This fact follows from a more general statement that goes back at least to E. Kelly [6], but see also Theorem 1 in [10] for an interesting generalization to harmonic maps and Theorem 4.1 in [3] for a broad analysis of topological restrictions imposed by the existence of minimal immersions into manifolds of Ricci curvature bounded from below. For the sake of completeness, a simple and direct proof of Theorem 2 is presented in Subsection 2.2.

In [12], Ros proved that a non-orientable, compact stable minimal surface immersed in a flat three-torus T^3 has the topology of a Klein bottle with a handle (cf. Theorem 7 therein). Analogously, as a simple combined application of the two theorems above one can prove that if $M^3 \subset T^4$ is stable (but not totally geodesic) then either $b_1 = 4$ or $b_1 = 5$ and M^3 is non-orientable as can be checked directly by means of the second variation formula. The question of classifying all such stable minimal hypersurfaces remains a challenging open problem.

Acknowledgments. The authors wish to express their gratitude to André Neves for his interest in this work and for a number of enlightening conversations. During the preparation of this article, L.A. was supported by prof. Neves' ERC Start Grant PSC and LMCF 278940.

2. PROOFS

2.1. Notations and ancillary results.

Throughout this note, we consider the ambient manifold $T^{n+1} := \mathbb{R}^{n+1}/\Gamma$, where Γ is a lattice group of maximal rank, endowed with its (flat) Riemannian metric $\langle \cdot, \cdot \rangle$ and the associated Levi-Civita connection D . Furthermore, we let M^n denote a closed, embedded minimal hypersurface in T^{n+1} , that is to say a smooth, closed hypersurface of vanishing mean curvature. For the sake of simplicity, and in order to streamline our arguments, we shall assume that M^n is orientable or, equivalently, two-sided and we let N denote a choice

of its unit normal vector field. The case when M^n is one-sided, for which the statement of Theorem 1 still holds, is discussed in Remark 6. The induced Levi-Civita connection on the submanifold M^n will always be denoted by ∇ , while Δ stands for the Laplace-Beltrami operator. Lastly, we convene that the second fundamental form of $M^n \subset T^{n+1}$ is defined by the formula $A(X, Y) = \langle D_X Y, N \rangle$ for any pair of smooth vector fields X, Y along M^n .

In our setting the Morse index can be defined as follows: the sections of the normal bundle of $M^n \subset T^{n+1}$ can be identified with the set of smooth functions ϕ on M^n , and the second variation of the area functional is given by the quadratic form

$$Q(\phi, \phi) = \int_M (|\nabla\phi|^2 - |A|^2\phi^2) dM,$$

so that the number of *negative variations* is encoded in the spectrum of the Jacobi operator

$$J_M\phi = \Delta\phi + |A|^2\phi$$

and the index of M^n is by definition the number of (strictly) negative eigenvalues of J_M .

In a closed Riemannian manifold (M^n, g) , the Hodge-Laplace operator is the second order differential operator Δ_p acting on p -forms defined by

$$\Delta_p = dd^* + d^*d$$

where $d : \Omega^p(M) \rightarrow \Omega^{p+1}(M)$ is the exterior differential and $d^* : \Omega^p(M) \rightarrow \Omega^{p-1}(M)$ is the formal adjoint of d , defined with respect to the metric g . A p -form ω is called harmonic when $\Delta_p\omega = 0$ and we let $\mathcal{H}^p(M, g)$ denote the vector space of harmonic p -forms on (M^n, g) . When M^n is closed, ω is harmonic if and only if it is closed and co-closed, that is to say when both $d\omega = 0$ and $d^*\omega = 0$ hold true. Hodge's Theorem asserts that in a closed Riemannian manifold one has the isomorphism $\mathcal{H}^1(M, g) \simeq H^1(M; \mathbb{R})$ so that the dimension of the space of harmonic 1-forms coincides with the first Betti number of the manifold. Also, we will use the (special) Bochner-Weitzenböck formula relating the Hodge-Laplace operator with the usual (rough) Laplacian on 1-forms:

$$(2.1) \quad \Delta_1\omega = -\Delta\omega + Ric^M(\omega^\sharp, \cdot).$$

In this note we employ the usual musical isomorphisms to pass from vectors to 1-forms, see Remark 2.1 in [1] for further details.

The proof of our main result relies on the following proposition:

Proposition 3. *Let M^n be a closed, orientable, minimal hypersurface in T^{n+1} and let ω be a harmonic 1-form on M^n . For every parallel 2-form θ on T^{n+1} one has the identity*

$$(2.2) \quad \Delta\langle N^\flat \wedge \omega, \theta \rangle + |A|^2\langle N^\flat \wedge \omega, \theta \rangle = -2 \sum_{i,j=1}^n A(E_i, E_j) \langle E_j^\flat \wedge \nabla_{E_i}\omega, \theta \rangle,$$

where the expression on the right-hand side is globally defined as it does not depend on the particular choice of local orthonormal frame $\{E_i\}$ on M^n .

Remark 4. The reader may want to compare this assertion with Lemma 1 in [12], where similar computations were performed for the coordinates of ω rather than $N^\flat \wedge \omega$ (the two choices being essentially equivalent only if $n = 2$).

Proof. Let $\{E_i\}$ be a local orthonormal frame on M^n , which is geodesic at a point p in M^n (that is to say $(\nabla_{E_i} E_j)(p) = 0$ for all i, j). We have

$$D_{E_i}(N^\flat \wedge \omega) = D_{E_i} N^\flat \wedge \omega + N^\flat \wedge D_{E_i} \omega = - \sum_{j=1}^n A(E_i, E_j) E_j^\flat \wedge \omega + N^\flat \wedge \nabla_{E_i} \omega.$$

Hence, exploiting the fact that θ is parallel, one has at the point p

$$\begin{aligned} \Delta \langle N^\flat \wedge \omega, \theta \rangle &= \sum_{i=1}^n E_i E_i \langle N^\flat \wedge \omega, \theta \rangle = \sum_{i=1}^n E_i \langle D_{E_i}(N^\flat \wedge \omega), \theta \rangle \\ &= \sum_{i=1}^n E_i \langle - \sum_{j=1}^n A(E_i, E_j) E_j^\flat \wedge \omega + N^\flat \wedge \nabla_{E_i} \omega, \theta \rangle \\ &= - \sum_{i,j=1}^n E_i (A(E_i, E_j)) \langle E_j^\flat \wedge \omega, \theta \rangle - \sum_{i,j=1}^n A(E_i, E_j) \langle D_{E_i} E_j^\flat \wedge \omega, \theta \rangle \\ &\quad - \sum_{i,j=1}^n A(E_i, E_j) \langle E_j^\flat \wedge D_{E_i} \omega, \theta \rangle + \sum_{i=1}^n \langle D_{E_i}(N^\flat \wedge \nabla_{E_i} \omega), \theta \rangle. \end{aligned}$$

Since M^n is minimal, by the Codazzi equation for flat ambient manifolds (evaluating, once again, at p)

$$\sum_{j=1}^n E_j (A(E_i, E_j)) = \sum_{j=1}^n E_i (A(E_j, E_j)) = E_i H = 0.$$

Thus, we can deduce

$$\begin{aligned} \Delta \langle N^\flat \wedge \omega, \theta \rangle &= - \sum_{i,j=1}^n A(E_i, E_j) A(E_i, E_j) \langle N^\flat \wedge \omega, \theta \rangle - \sum_{i,j=1}^n A(E_i, E_j) \langle E_j^\flat \wedge \nabla_{E_i} \omega, \theta \rangle \\ &\quad - \sum_{i,j=1}^n A(E_i, E_j) A(E_i, \omega^\sharp) \langle E_j^\flat \wedge N^\flat, \theta \rangle - \sum_{i,j=1}^n A(E_i, E_j) \langle E_j^\flat \wedge \nabla_{E_i} \omega, \theta \rangle \\ &\quad + \sum_{i=1}^n \langle N^\flat \wedge \nabla_{E_i} \nabla_{E_i} \omega, \theta \rangle. \end{aligned}$$

We can rewrite the above as

$$\Delta \langle N^\flat \wedge \omega, \theta \rangle + |A|^2 \langle N^\flat \wedge \omega, \theta \rangle = \langle \Delta \omega, i_N \theta \rangle + \langle A \circ A(\omega^\sharp), (i_N \theta)^\sharp \rangle - 2 \sum_{i,j=1}^n A(E_i, E_j) \langle E_j^\flat \wedge \nabla_{E_i} \omega, \theta \rangle.$$

Now, the Gauss equation for a minimal hypersurface in flat ambient manifolds yields

$$\text{Ric}^M(\omega^\sharp, (i_N \theta)^\sharp) = - \langle A \circ A(\omega^\sharp), (i_N \theta)^\sharp \rangle,$$

while, on the other hand, the Bochner formula (2.1) for harmonic 1-forms on M^n reads

$$\langle \Delta \omega, i_N \theta \rangle = \text{Ric}^M(\omega^\sharp, (i_N \theta)^\sharp)$$

so that formula (2.2) follows at once. \square

Inspired by Chao Li (see [7], Proposition 5.1), we prove the following result.

Proposition 5. *Let M^n be a closed minimal hypersurface in a $(n+1)$ -dimensional flat torus. Assume there is a point p in M^n where all the principal curvatures are distinct. The set of all harmonic 1-forms ω on M^n such that*

$$(2.3) \quad \sum_{i,j=1}^n A(E_i, E_j) E_j^\flat \wedge \nabla_{E_i} \omega = 0$$

has dimension at most $2n - 1$.

Before proceeding with its proof, we want to remind the reader of a general fact about harmonic forms: in a Riemannian manifold (M^n, g) a 1-form $\omega \in \Omega^1(M^n)$ is closed and co-closed if and only if $\nabla\omega$ is a symmetric trace-free tensor.

Proof. By virtue of our assumption on M^n , we can certainly find a positive number ρ (smaller than the injectivity radius of M^n at p) such that the principal curvatures of M^n are all distinct in the geodesic ball $B_\rho(p)$ and, furthermore, there exists a local orthonormal frame $\{E_1, \dots, E_n\}$ diagonalizing the second fundamental form A at every point of such ball (namely: $A(E_i, E_j) = k_i \delta_{ij}$ for all $i, j = 1, \dots, n$ with $k_1 < k_2 < \dots < k_n$). Now, given a harmonic 1-form ω , equation (2.3) and the fact that the tensor $\nabla\omega$ is symmetric imply

$$(k_i - k_j)(\nabla_{E_i} \omega)(E_j) = 0 \quad \text{for all } i, j = 1, \dots, n,$$

so that

$$(2.4) \quad (\nabla_{E_i} \omega)(E_j) = 0 \quad \text{for all } i \neq j$$

on the whole geodesic ball in question. In particular, since $\nabla\omega$ is also trace-free we deduce that the functions $\nabla\omega(E_1, E_1), \dots, \nabla\omega(E_{n-1}, E_{n-1})$ completely determine the tensor $\nabla\omega$ on $B_\rho(p)$.

Now, let us consider the functions defined on $B_\rho(p)$ by

$$\phi_i(q) = \begin{cases} \omega(E_i)(q) & \text{for } 1 \leq i \leq n \\ \nabla\omega(E_{i-n}, E_{i-n})(q) & \text{for } n+1 \leq i \leq 2n-1. \end{cases}$$

We claim that if ω satisfies (2.3) then for every $q \in B_\rho(p)$ the values $(\phi_1, \dots, \phi_{2n-1})(q)$ are uniquely determined by $(\phi_1, \dots, \phi_{2n-1})(p)$, hence the space of harmonic forms in $B_\rho(p)$ satisfying (2.3) has dimension at most $2n - 1$ and the general statement over M^n follows by unique continuation.

To check the claim, we proceed as follows: given $q \in B_\rho(p)$ let $\gamma : [0, \tau] \rightarrow M^n$ be the only geodesic connecting p to q in $B_\rho(p)$ and consider the functions (of one real variable) obtained by restriction of $(\phi_1, \dots, \phi_{2n-1})$ along γ , namely set

$$f_i(t) = \phi_i(\gamma(t)), \quad \text{for } i = 1, \dots, 2n-1.$$

Then, we claim that (f_1, \dots, f_{2n-1}) solves a linear ODE system (in normal form) so that (by Cauchy-Lipschitz) the value at p uniquely determines the value along the curve γ , which is enough to check the claim above.

Notice that $\gamma'(t) = \sum_{j=1}^n \alpha^j(t) E_j(\gamma(t))$ where the coefficients α^j , $j = 1, \dots, n$ are smooth and bounded (since the frame $\{E_1, \dots, E_n\}$ is orthonormal) hence by linearity

$$(2.5) \quad \frac{d}{dt} f_i(t) = \nabla_{\gamma'(t)} \phi_i(\gamma(t)) = \sum_{j=1}^n \alpha^j(t) \nabla_{E_j} \phi_i(\gamma(t))$$

so that it suffices for our scopes to check that for every choice of the indices $1 \leq i \leq 2n-1$ and $1 \leq j \leq n$ the function $\nabla_{E_j} \phi_i$ can be expressed as a linear combination of $\phi_1, \dots, \phi_{2n-1}$, with smooth coefficients.

First of all, for $i \leq n$ we have

$$(2.6) \quad \nabla_{E_j} \phi_i = \delta_{ij} \phi_{n+i} + \sum_{k=1}^n \Gamma_{ji}^k \phi_k.$$

This can be justified as follows:

$$\nabla_{E_j}(\omega(E_i)) = (\nabla_{E_j} \omega)(E_i) + \omega(\nabla_{E_j} E_i) = \nabla \omega(E_i, E_j) + \sum_{k=1}^n \Gamma_{ji}^k \omega(E_k) = \delta_{ij} \phi_{n+i} + \sum_{k=1}^n \Gamma_{ji}^k \phi_k$$

where the last steps relies on equation (2.4). Also, observe that for $i = n$ the RHS of (2.6) needs to be suitably interpreted, namely with $-\sum_{l=1}^{n-1} \phi_{n+l}$ in lieu of ϕ_{2n} .

On the other hand, the differential equation for ϕ_{n+i} takes for $1 \leq i \leq n-1$ one of the following three forms:

Case 1: $1 \leq j \leq n-1, i \neq j$

$$\nabla_{E_j} \phi_{n+i} = (2\Gamma_{ji}^i - \Gamma_{ij}^i) \phi_{n+i} - \Gamma_{ii}^j \phi_{n+j} - \sum_{k=1}^n R_{ikij} \phi_k.$$

Case 2: $j = n$

$$\nabla_{E_j} \phi_{n+i} = (2\Gamma_{ni}^i - \Gamma_{in}^i) \phi_{n+i} + \sum_{k=1}^{n-1} \Gamma_{ii}^k \phi_{n+k} - \sum_{k=1}^n R_{ikin} \phi_k.$$

Case 3: $j = i$

$$\nabla_{E_j} \phi_{n+i} = \sum_{k=1, k \neq i}^n \Gamma_{kk}^i \phi_{n+i} - \sum_{k=1, k \neq i}^{n-1} (2\Gamma_{ik}^k - \Gamma_{ki}^k) \phi_{n+k} + (2\Gamma_{in}^n - \Gamma_{ni}^n) \sum_{k=1}^{n-1} \phi_{n+k} + \sum_{k=1}^n R_{ki} \phi_k.$$

Indeed, for all $j \neq i$ by the Ricci formula for commuting covariant derivatives (no summation on repeated indexes i and j) we obtain

$$\begin{aligned}
\nabla_{E_j}(\nabla\omega(E_i, E_i)) &= \nabla^2\omega(E_i, E_i, E_j) + 2\nabla\omega(\nabla_{E_j}E_i, E_i) \\
&= \nabla^2\omega(E_i, E_j, E_i) - \sum_{k=1}^n R_{ikij}\omega(E_k) + 2\Gamma_{ji}^i\nabla\omega(E_i, E_i) \\
&= \nabla_{E_i}(\nabla\omega(E_i, E_j)) - \nabla\omega(\nabla_{E_i}E_i, E_j) - \nabla\omega(E_i, \nabla_{E_i}E_j) - \sum_{k=1}^n R_{ikij}\omega(E_k) + 2\Gamma_{ji}^i\nabla\omega(E_i, E_i) \\
&= -\Gamma_{ii}^j\phi_{n+j} + (2\Gamma_{ji}^i - \Gamma_{ij}^i)\phi_{n+i} - \sum_{k=1}^n R_{ikij}\phi_k
\end{aligned}$$

where the very last equality was implied by (2.4). Thereby, the first equation is verified and the second one follows along similar lines exploiting the fact that the tensor $\nabla\omega$ is trace free. Lastly, following the same pattern, one has for all $i = 1, \dots, n-1$

$$\begin{aligned}
\nabla_{E_i}(\nabla\omega(E_i, E_i)) &= \sum_{k=1, k \neq i}^n -E_i(\nabla\omega(E_k, E_k)) = \sum_{k=1, k \neq i}^{n-1} \Gamma_{kk}^i\nabla\omega(E_i, E_i) - (2\Gamma_{ik}^k - \Gamma_{ki}^k)\nabla\omega(E_k, E_k) \\
&\quad + \Gamma_{nn}^i\nabla\omega(E_i, E_i) - (2\Gamma_{in}^n - \Gamma_{ni}^n)\nabla\omega(E_n, E_n) + \sum_{k=1, k \neq i}^{n-1} \sum_{l=1}^n R_{klki}\omega(E_l) + \sum_{k=1}^n R_{nkn i}\omega(E_k) \\
&= \sum_{k=1, k \neq i}^n \Gamma_{kk}^i\phi_{n+i} - \sum_{k=1, k \neq i}^{n-1} (2\Gamma_{ik}^k - \Gamma_{ki}^k)\phi_{n+k} + (2\Gamma_{in}^n - \Gamma_{ni}^n) \sum_{k=1}^{n-1} \phi_{n+k} + \sum_{k=1}^n R_{ki}\phi_k.
\end{aligned}$$

This finishes the proof of the claim and thus the whole argument. \square

2.2. Proofs of Theorem 1 and Theorem 2.

As stated in the introduction, we first present a short proof of Theorem 2.

Proof. Let \mathcal{V} be the space of all parallel 1-forms on the flat $(n+1)$ -dimensional torus. This space consists precisely of the forms df , where f is a linear function on the universal cover of the flat torus (viz. $(n+1)$ -dimensional Euclidean space). In particular, its dimension is $n+1$.

Since M^n is minimal and the elements of \mathcal{V} are parallel, the restriction of any $df \in \mathcal{V}$ is a harmonic 1-form. Moreover, $df = 0$ on M^n if and only if M^n is contained in the quotient of level sets of the linear map f by the action of the lattice subgroup of translations of the Euclidean space that generates the flat torus in question. Our assertion follows at once. \square

We then deduce from Proposition 3 and Proposition 5 our main result, Theorem 1. To that scope, we need to recall a rigidity theorem proved by Do Carmo and Dajczer [2] for minimal hypersurfaces M^n in \mathbb{R}^{n+1} such that some principal curvature has multiplicity at least $n-1$ at all points: any such hypersurface must be part of a higher-dimensional catenoid, or flat (see Corollary 4.4 in their paper). In particular, in a four dimensional torus ($n=3$), the assumption on the principal curvatures given in the statement of Theorem 1 (namely the

assumption that all principal curvatures are pairwise distinct) holds for all closed minimal hypersurfaces that are not totally geodesic.

Proof. Given an orthonormal basis $\{\theta_1, \dots, \theta_{n+1}\}$ of parallel 1-forms on T^{n+1} , let M^n be a closed, orientable minimal hypersurface of a flat $(n+1)$ -dimensional torus whose principal curvatures are all distinct at least at one point. Let us denote by k its Morse index, and by $\{\phi_q\}_{q=1}^k$ an L^2 -orthonormal basis of eigenfunctions of the Jacobi operator $J_M = \Delta + |A|^2$ of M^n generating the eigenspace where the operator is negative definite. Let then Φ denote the linear map defined by

$$\begin{aligned} \Phi : \mathcal{H}^1(M^n) &\rightarrow \mathbb{R}^{n(n+1)k/2} \\ \omega &\mapsto \left[\int_M \langle N^\flat \wedge \omega, \theta_i \wedge \theta_j \rangle \phi_q d\mu \right], \end{aligned}$$

where $1 \leq i < j \leq n+1$ and q varies from 1 to k . Clearly,

$$\dim \mathcal{H}^1(M, g) \leq \dim \text{Ker}(\Phi) + \frac{n(n+1)}{2}k.$$

Since $\mathcal{H}^1(M, g) \simeq H^1(M; \mathbb{R})$, Theorem 1 will follow once we analyse the dimension of the kernel of the map Φ and show that indeed $\dim \text{Ker}(\Phi) \leq 2n-1$.

Let ω be an element of the kernel of the map Φ . This precisely means that every function $u_{ij} = \langle N^\flat \wedge \omega, \theta_i \wedge \theta_j \rangle$ is L^2 -orthogonal to each of the first k eigenfunctions of J_M . Since $\text{index}(M) = k$, we must have

$$Q(u_{ij}, u_{ij}) \geq \lambda_{k+1} \int_M u_{ij}^2 d\mu \geq 0 \quad \text{for all } 1 \leq i < j \leq n+1,$$

by the standard variational characterization of eigenvalues. Hence, thanks to Proposition 3 we have

$$\begin{aligned} 0 &\leq \sum_{1 \leq i < j \leq n+1} Q(u_{ij}, u_{ij}) = - \sum_{1 \leq i < j \leq n+1} \int_M u_{ij} J_M(u_{ij}) d\mu \\ &= 2 \int_M \sum_{1 \leq i < j \leq n+1} \langle N^\flat \wedge \omega, \theta_i \wedge \theta_j \rangle \left\langle \sum_{k,l=1}^n A(E_k, E_l) E_l^\flat \wedge \nabla_{E_k} \omega, \theta_i \wedge \theta_j \right\rangle d\mu \\ &= 2 \int_M \sum_{k,l=1}^n A(E_k, E_l) \langle E_l^\flat \wedge \nabla_{E_k} \omega, N^\flat \wedge \omega \rangle d\mu = 0, \end{aligned}$$

the last equality relying on the fact that trivially, by orthogonality, $i_N(E_l^\flat \wedge \nabla_{E_k} \omega) = 0$ for any choice of the indices k and l . It follows that $\langle N^\flat \wedge \omega, \theta_i \wedge \theta_j \rangle$ are all eigenfunctions of the Jacobi operator J_M , associated to the eigenvalue $\lambda_{k+1} = 0$. By Proposition 3, this implies that ω satisfies equation (2.3). Thus, to complete the proof, it is enough to invoke Proposition 5, which ensures that the dimension of $\text{Ker}(\Phi)$ cannot exceed $2n-1$.

Lastly, to obtain an unconditional result when $n = 2, 3$, we need to observe that when the condition on the principal curvatures of M^n is not fulfilled, then M^n is totally geodesic and therefore a stable n -dimensional torus, in which case our inequality is also satisfied. This claim is clear when $n = 2$ and is a consequence of the result by do Carmo and Dajczer when $n = 3$. Thereby the proof is complete. \square

Remark 6. Let us discuss the modifications needed to obtain this index bound for non-orientable minimal hypersurfaces inside T^{n+1} . First of all, by orientability of T^{n+1} we know that any such M^n is also one-sided. In this case, it is customary to introduce the two-sheeted covering $\pi : \hat{M}^n \rightarrow M^n$ (given by couples (x, N) where $x \in M^n$ and N attains one of the two possible choices for the unit normal of M^n at x) and the associated two-sided immersion $\iota : \hat{M}^n \rightarrow T^{n+1}$ (with a well-defined unit normal field \hat{N} given by $\hat{N}(x, N) = N$). As discussed (for instance) in Subsection 2.3 of [1] one can then consider the restriction of the Jacobi operator of \hat{M}^n to the space of odd functions, namely to all $u : \hat{M}^n \rightarrow \mathbb{R}$ satisfying $u \circ \tau = -u$ for $\tau : \hat{M}^n \rightarrow \hat{M}^n$ the natural deck transformation of the covering in question. Hence, the Morse index of M^n is defined to be the number of negative eigenvalues of $J_{\hat{M}}$ acting on odd functions, in the sense above.

These comments being made, our arguments go through in the non-orientable case as well without any substantial modification once it is checked that for any harmonic form $\omega \in \mathcal{H}^1(M, g)$ one has that the test functions $u_{ij} = \langle \hat{N}^b \wedge \pi^*(\omega), \iota^*(\theta_i) \wedge \iota^*(\theta_j) \rangle$ are indeed odd, which is done in the proof of Theorem A of [1].

REFERENCES

- [1] L. Ambrozio, A. Carlotto and B. Sharp, *Comparing the Morse index and the first Betti number of minimal hypersurfaces*, J. Differential Geom. (to appear).
- [2] M. do Carmo and M. Dajczer, *Rotation hypersurfaces in spaces of constant curvature*, Trans. Amer. Math. Soc., 277 (1983), no. 2, 685-709.
- [3] J. Choe, A. Fraser, *Mean curvature in manifolds with Ricci curvature bounded from below*, preprint (arXiv: 1605.06602).
- [4] J. Choe, J. Hoppe, *Higher dimensional Schwarz's surfaces and Scherk's surfaces*, preprint (arXiv: 1607.07153).
- [5] D. Hoffman, *Some basic facts, old and new, about triply periodic embedded minimal surfaces - International Workshop on Geometry and Interfaces (Aussois, 1990)*. J. Physique 51 (1990), no. 23, Suppl. Colloq. C7, 197-208.
- [6] E. Kelly, *Cohomology of compact minimal submanifolds*, Michigan Math. J. **19** (1972), 133-135.
- [7] C. Li, *Index and topology of minimal hypersurfaces in \mathbb{R}^n* , preprint (arXiv:1605.09693).
- [8] F. Marques, *Minimal surfaces - variational theory and applications*, Proceedings of the International Congress of Mathematicians, Seoul 2014.
- [9] W. H. Meeks III, *The theory of triply periodic minimal surfaces*, Indiana Univ. Math. J. **39** (1990), no. 3, 877-936.
- [10] T. Nagano and B. Smyth, *Minimal varieties and harmonic maps in tori*, Comment. Math. Helv. **50** (1975), 249-265.
- [11] A. Neves, *New applications of Min-max Theory*, Proceedings of the International Congress of Mathematicians, Seoul 2014.
- [12] A. Ros, *One-sided complete stable minimal surfaces*, J. Differential Geom. **74** (2006), no. 1, 69-92.
- [13] M. Ross, *Schwarz' P and D surfaces are stable*, Differential Geom. Appl. **2** (1992), no. 2, 179-195.

L. AMBROZIO: IMPERIAL COLLEGE LONDON, SOUTH KENSINGTON CAMPUS, LONDON SW7 2AZ, UNITED KINGDOM, *E-mail address*: l.ambrozio@imperial.ac.uk

A. CARLOTTO: ETH - DEPARTMENT OF MATHEMATICS, RÄMISTRASSE 101, 8092 ZÜRICH, SWITZERLAND, *E-mail address*: alessandro.carlotto@math.ethz.ch

B. SHARP: UNIVERSITY OF WARWICK, GIBBET HILL RD, COVENTRY CV4 7AL, UNITED KINGDOM *E-mail address*: b.sharp@warwick.ac.uk