

**Original citation:**

Taylor, Andrew, Coventry, Emma, Handy, Claire, West, Jon S., Young, Caroline S. and Clarkson, John P. (2018) *Inoculum potential of Sclerotinia sclerotiorum sclerotia depends on isolate and host plant*. Plant Pathology .doi:[10.1111/ppa.12843](https://doi.org/10.1111/ppa.12843)

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/98266>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

"This is the peer reviewed version of the following Taylor, Andrew, Coventry, Emma, Handy, Claire, West, Jon S., Young, Caroline S. and Clarkson, John P. (2018) *Inoculum potential of Sclerotinia sclerotiorum sclerotia depends on isolate and host plant*. Plant Pathology. which has been published in final form at <https://doi.org/10.1111/ppa.12843>. This article may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#)."

**A note on versions:**

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)

## **Inoculum potential of *Sclerotinia sclerotiorum* sclerotia depends on isolate and host plant**

Andrew Taylor<sup>1\*</sup>, Emma Coventry<sup>1</sup>, Claire Handy<sup>1</sup>, Jon S. West<sup>2</sup>, Caroline S. Young<sup>3</sup> and John P. Clarkson<sup>1</sup>

<sup>1</sup>Warwick Crop Centre, School of Life Sciences, University of Warwick, Wellesbourne Campus, Wellesbourne, Warwick, CV35 9EF, UK.

<sup>2</sup>Rothamsted Research, Harpenden, AL5 2JQ. UK

<sup>3</sup>ADAS, Drayton Farm, Alcester Road, Stratford upon Avon CV37 9RQ, UK

\*andrew.taylor@warwick.ac.uk

Running head: Inoculum potential of *S. sclerotiorum*

Keywords: *Sclerotinia sclerotiorum*, crop rotation, sclerotia, oilseed rape, carrot, lettuce.

Accepted for publication: 31/01/2018

## Abstract

The soilborne fungus *Sclerotinia sclerotiorum* infects many important crop plants. Central to the success of this pathogen is the production of sclerotia, which enables survival in soil and constitutes the primary inoculum. This study aimed to determine how crop plant type and *S. sclerotiorum* isolate impact sclerotial production and germination and hence inoculum potential. Three *S. sclerotiorum* isolates (L6, L17, L44) were used to inoculate plants of bean, carrot, lettuce, oilseed rape (OSR) and potato and the number and weight of sclerotia per plant quantified. Carpogenic germination of sclerotia collected from different hosts was also assessed for L6. Production of sclerotia was dependent on both crop plant type and *S. sclerotiorum* isolate with OSR and lettuce supporting the greatest number (42-122) and weight (1.6-3.0 g) of sclerotia per plant. The largest sclerotia were produced on oilseed rape (33-66 mg). The three *S. sclerotiorum* isolates exhibited a consistent pattern of sclerotial production irrespective of crop type; L6 produced large numbers of small sclerotia while L44 produced smaller numbers of large sclerotia with L17 intermediate. Germination rate and percentage was greatest for larger sclerotia (4-6.7mm) and also varied between host plants. Combining sclerotial production data and typical field crop densities suggested that infected carrot and OSR could produce the greatest number (3944 m<sup>-2</sup>) and weight (73 g m<sup>-2</sup>) of *S. sclerotiorum* sclerotia respectively, suggesting these crops potentially contribute a greater increase in inoculum. This information, once further validated in field trials, could be used to inform future crop rotation decisions.

## Introduction

The cosmopolitan necrotrophic fungus *Sclerotinia sclerotiorum* (Lib) de Bary infects more than 400 species of plants throughout the world (Boland & Hall, 1994) leading to severe economic losses in a wide range of crops, including beans, carrot, lettuce, oilseed rape (OSR) and potatoes. In bean (*Phaseolus vulgaris* L., both dry and green bean), the pathogen causes white mould disease, leading to losses of up to 100% under favourable conditions (Schwartz & Singh, 2013) with the fungus infecting stems, branches, leaves, flowers, pods and seeds at every stage of crop development. In carrot, *S. sclerotiorum* has been reported in over twenty carrot producing countries and affects both foliage and roots causing substantial canopy die-back and subsequent crown rot (Jensen *et al.*, 2008). This can also then lead to post-harvest epidemics in storage (Foster *et al.*, 2008). *S. sclerotiorum* also causes lettuce ‘drop’ where plants quickly wilt in the field due to infection of the stem base and lower leaf axils followed by rapid rotting of the tissue (Young *et al.*, 2004, Clarkson *et al.*, 2014). As in other crops, the disease can be very damaging with up to 50% losses reported in UK lettuce (Young *et al.*, 2004). In OSR, *S. sclerotiorum* causes a stem rot and results in substantial yield losses in all major growing areas including Australia, Canada, UK and USA (Derbyshire & Denton Giles, 2016). In this case, flower petals are initially infected which fall onto leaves or leaf axils resulting in development of the disease in stems (Derbyshire & Denton Giles, 2016). Finally, *S. sclerotiorum* can also be a significant disease in potato where it causes a stem rot, infecting the lower parts of the stems towards the end of the growing season, leading to reduced yields. As for OSR, stem lesions are associated with infected flowers dropping onto stems (Atallah & Johnson, 2004).

Central to the success of *S. sclerotiorum* as a plant pathogen is the production of numerous sclerotia on infected plants. In addition to representing a huge reproductive potential and providing the primary inoculum for subsequent epidemics, these sclerotia enable the fungus to survive overwinter (or for longer periods) in soil (Adams & Ayers, 1979). *S. sclerotiorum* sclerotia can then germinate carpogenically or myceliogenically, depending on environmental conditions with the former resulting in the production of apothecia and airborne ascospores. This often represents the principal source of inoculum and hence allows infection of above-ground plant parts (Bolton *et al.*, 2006). Carpogenic germination of *S. sclerotiorum* occurs only in sclerotia that are brought close to the soil surface and, in temperate climates, a period of cold conditioning is also required to break dormancy (Clarkson *et al.*, 2007). Ascospores are then released from mature apothecia under a wide range of conditions with up to  $7.6 \times 10^5$  spores released over a 20-day period. (Clarkson *et al.*, 2003).

The density of *S. sclerotiorum* sclerotia in the soil is clearly an important factor in disease epidemics although very few studies have attempted to quantify populations and relate this to disease incidence. Surveys in lettuce fields in the USA with a history of *Sclerotinia* disease have reported densities ranging from 0.08-2.9 sclerotia / 100 g soil in Yuma, Arizona (Chitrampalam & Pryor, 2013), while an average of 0.06 sclerotia / 100 cm<sup>3</sup> of soil was reported in the San Joaquin Valley, California (Hao & Subbarao, 2005). Other work has demonstrated a positive correlation between the density of *S. sclerotiorum* sclerotia used to inoculate beds of lettuce (2-100 sclerotia m<sup>-2</sup>) and disease incidence at harvest (5-71%) although the authors indicated that in the desert production area under investigation, mycelial germination of sclerotia was the predominant mode of infection as conditions were not conducive to carpogenic germination and development of apothecia (Chitrampalam *et al.*, 2010). In the many lettuce growing areas, where conditions do allow production of apothecia, small numbers of *S. sclerotiorum* sclerotia can still lead to high levels of disease incidence

(Hao & Subbarao, 2005). This is because the inoculum potential of sclerotia germinating carpogenically is much higher than for those germinating to produce mycelium due to the large number of airborne ascospores released, which dramatically increases the spatial spread from a single sclerotium. Work in other crops such as oilseed rape and sunflower has also demonstrated strong relationships between either sclerotial density or apothecial production and Sclerotinia disease incidence (Gugel, 1986, Lehner *et al.*, 2017).

The number of *S. sclerotiorum* sclerotia that are returned to the soil not only depends on the number of plants infected but also on the host type but very little information exists on sclerotial production on different crops. One study reported variation in the number and weight of *S. sclerotiorum* sclerotia produced by different *S. sclerotiorum* isolates on carrot, celery and selected cultivars of cabbage (Leiner & Winton, 2006). However, this work was carried out using detached plant material, rather than a live host. Other researchers also demonstrated that, *in vitro*, different isolates of both *S. sclerotiorum* and *S. trifoliorum* produced different numbers and weights of sclerotia under the same conditions (Akram *et al.*, 2005, Li *et al.*, 2008, Vleugels *et al.*, 2013). Furthermore, it has been shown that larger *S. sclerotiorum* sclerotia tend to produce more apothecia than smaller ones and are also more likely to germinate (Ben-Yephet *et al.*, 1993, Dillard *et al.*, 1995, Hao *et al.*, 2003, Warmington & Clarkson, 2016). However, these studies used sclerotia produced on a single crop type (lettuce; Ben-Yephet *et al.*, 1993), or on detached potato tubers (Hao *et al.*, 2003) or produced artificially *in vitro* (Dillard *et al.*, 1995, Warmington & Clarkson, 2016). Hence, both crop type and *S. sclerotiorum* isolate potentially have a significant impact on the build-up of inoculum in the field through affecting the number and size of sclerotia produced.

The aim of this study was to quantify sclerotial production by different *S. sclerotiorum* isolates on five different crop hosts using whole plants and hence estimate the risk of inoculum

build-up associated with each. In addition, the effect of sclerotial size on the rate and final level of germination as well as the number of apothecia produced was also investigated.

## Materials and Methods

### ***S. sclerotiorum* isolates and production of sclerotia on agar**

The 17 *S. sclerotiorum* isolates used in this study were obtained from wild buttercup (*Ranunculus acris*, eight isolates), carrot (*Daucus carota*, one isolate), OSR (*B. napus*, one isolate), lettuce (*Lactuca sativa*, four isolates), pea (*Pisum sativum*, two isolates) and celery (*Apium graveolens*, one isolate). A single isolate of *S. subarctica* (isolate HE1 from wild buttercup) was also included for comparison; this related pathogen often produces larger sclerotia than *S. sclerotiorum* isolates (Clarkson *et al.*, 2010). Isolates represented different microsatellite haplotypes assigned through a previous study investigating the population structure of *S. sclerotiorum* and *S. subarctica* in the UK (Clarkson *et al.*, 2017; Table 1). In addition, the relative aggressiveness of the same 18 isolates had also been evaluated previously on three brassica hosts (Taylor *et al.*, 2015). Stocks of each *S. sclerotiorum* isolate were maintained as sclerotia stored at 5°C and new cultures initiated by bisecting a sclerotium, placing it face down on potato dextrose agar (PDA) and incubating at 20°C for 4 days to produce actively growing cultures for use in experiments. To assess the production of sclerotia by the different *S. sclerotiorum* isolates on agar, 5 mm agar plugs from actively growing cultures were placed at the centre of PDA plates (20 ml per plate) and incubated at 20°C for four weeks (five replicate plates per isolate). Sclerotia were then picked off and the number and weight of sclerotia per plate recorded. Data were analysed using a REML analysis in Genstat version 18 (VSN International) with number and weight of sclerotia as the fixed effects and replicate plate as a random effect. Statistical significance was determined using an F test. As all residuals were close to the normal distribution, correlations with aggressiveness were examined using Pearson correlation coefficients (Genstat).

### **Number of *S. sclerotiorum* sclerotia produced on different crop plants**

Three *S. sclerotiorum* isolates (L6, L17 and L44) were selected for determining the number of sclerotia produced on five different crop plants based on differences in sclerotial production *in vitro*. Actively growing cultures of each isolate were produced as described previously and 5 x 5 mm square agar plugs from the leading edge used to inoculate sterile wheat grain placed in Petri dishes. Following incubation at 20°C for 3 days, colonised wheat grain was then used as inoculum for each crop; oilseed rape (cv. Temple, three experiments of 10 replicate plants), lettuce (cv. Montel, eight experiments of 10 replicate plants), dwarf bean (cv. Tendergreen, six experiments of 10 replicate plants), carrot (cv. Nairobi, four experiments of 10 replicate plants) and potato (cv. Estima, three experiments of 10 replicate plants). Plants were inoculated at flowering with the exception of lettuce (30 cm diameter plant) and carrot (4-5 fully opened leaves). For bean and potato inoculations, stems were first cut in two different places with a scalpel and 4-5 wheat grains placed into each wound. Lettuce, carrot and oilseed rape plants were inoculated by placing 10 wheat grains into different axils next to the main stem (2-3 grains per axil). A high level of humidity was maintained to encourage infection and disease development for all inoculated plants using automatic misting sprays in a controlled temperature glasshouse (lettuce, bean and carrot) set at a constant 15°C with supplementary lighting as required to achieve a 16 h photoperiod. Potato and OSR experiments were carried out in polytunnels with overhead irrigation. After plants were fully colonised and necrotic, watering was halted and the tissue allowed to die back and dry out completely before sclerotia were collected, air-dried, weighed and counted. For OSR and bean, the main stem was split open and sclerotia from inside the stem added to the count. Data were analysed using REML in Genstat with number / weight of sclerotia and weight per sclerotium as the fixed effects and replicate experiment / replicate plant as random effects. Statistical significance was determined

using an F test. As all residuals were close to a normal distribution, correlations with sclerotial production on agar were examined using Pearson correlation coefficients (Genstat).

### **Effect of crop type and size of sclerotia on carpogenic germination**

To investigate the effect of size of *S. sclerotiorum* sclerotia on carpogenic germination, isolate L6 was selected due to its ability to consistently produce apothecia in previous work (Taylor *et al.*, 2018). Sclerotia of this isolate collected from plants across multiple experiments were pooled according to the crop type on which they were produced and passed through sieves of increasing diameter to divide them into four size classes; <2 mm, 2-4 mm, 4-6.7 mm and >6.7 mm. Sclerotia were then buried at 1 cm depth in compost (150 g, John Innes No. 1, J Arthur Bowers, UK) placed in 600 ml plastic boxes (Malsar Kest Ltd, UK) and water content adjusted to 30% w/v. There were 30 sclerotia per box, and three replicate boxes per crop/size combination. The sclerotia were initially conditioned by placing the boxes at 5°C for 40 days to promote carpogenic germination (Clarkson *et al.*, 2014) after which they were placed at 15°C and the number of germinating sclerotia and number of apothecia produced for each sclerotium recorded over a 12-week period. For sclerotia in the size class 4-6.7 mm collected from carrot and bean plants, only one box of sclerotia was set up due to the very low numbers obtained in this size class. Sclerotia in the size class >6.7 mm were only obtained from oilseed rape and in this case, due to their size, only 16 sclerotia were placed in each box (three replicate boxes). The germination of *S. sclerotiorum* sclerotia produced on the crop plants was compared with those produced on sterilised wheat grain *in vitro*, as used routinely for production of sclerotia in several previous studies (Mylchreest & Wheeler, 1987, Liu & Paul, 2007, Warmington & Clarkson, 2016). As before, three replicate boxes (30 sclerotia per box) were set up for each size class with the exception of those >6.7 mm where a single box was set up due to low numbers of sclerotia of this size being produced on wheat grain. The number of

sclerotia germinating and the number of apothecia per sclerotium was again recorded over a 12-week period and the time taken to reach one third (33%) germination calculated. Data were analysed using REML in Genstat with germination, number of apothecia and time to 33% germination as the fixed effects and replicate box as a random effect. Statistical significance was determined using an F test.

## Results

### Production of sclerotia on agar

Significant differences in the number of sclerotia produced by the 18 *Sclerotinia* isolates were observed, with mean number ranging from 17 (*S. subarctica* isolate HE1) to 44 (*S. sclerotiorum* isolate L6) per PDA plate ( $P < 0.001$ , Fig. 1a). Significant differences were also observed between isolates for the mean weight of sclerotia produced per plate ( $P < 0.001$ , Fig. 1b) which ranged from 116 mg (*S. sclerotiorum* L44) to 306 mg (*S. sclerotiorum* EV9) and also the mean weight per sclerotium ( $P < 0.001$ , Fig. 1c) which ranged from 5.0 mg (*S. sclerotiorum* L6) to 16.9 mg (*S. subarctica* HE1). There were no significant correlations between aggressiveness on three brassica types (*B. oleracea*, *B. napus*, *B. rapa*) assessed previously (Taylor *et al.*, 2015) and the number of sclerotia ( $r = 0.42$ ,  $P = 0.083$ ), weight of sclerotia ( $r = -0.23$ ,  $P = 0.35$ ) or weight per sclerotium ( $r = -0.46$ ,  $P = 0.055$ ). Based on these results, three *S. sclerotiorum* isolates (all from lettuce) were selected for further experiments on plants based on variation in number of sclerotia produced per plate (L6, L17 and L44 producing 44, 36 and 29 sclerotia per plate respectively).

### Number of *S. sclerotiorum* sclerotia produced on different crop plants

All plants inoculated with *S. sclerotiorum* resulted in infection, complete colonisation and plant death for all crop types with the exception of potato, where in contrast to L6, isolates L17 and L44 failed to completely colonise plants. Significant differences were observed in the mean number of sclerotia produced on the different host plants ( $P < 0.001$ , Fig. 2a), with the largest number produced on lettuce (122 sclerotia per plant for isolate L6) and the smallest number produced on bean (7 sclerotia per plant for isolate L44). Significant differences between the mean weight of sclerotia per plant ( $P < 0.001$ , Fig. 2b) were also observed, ranging from 0.21

g on bean (L44) to 3.09 g on OSR (L6). The mean weight per sclerotium produced by each isolate was generally consistent across the different crops (10-28 mg) with the exception of OSR where significantly heavier sclerotia (33-60 mg) were produced for all three isolates ( $P < 0.001$ ; Fig. 2c). OSR also supported the highest proportion of larger sclerotia; for example, for *S. sclerotiorum* isolate L6, 17% were in the size category 4-6.7 mm compared to 3.2-7.4% in the other crop types (Table 2). It should be noted that pod development was incomplete on bean plants, possibly reducing the potential number of sclerotia produced.

Significant differences in the mean number of sclerotia produced were also observed between the *S. sclerotiorum* isolates when averaged across the different crop types with isolates L6, L17 and L44 producing 75, 54 and 32 sclerotia per plant respectively ( $P < 0.001$ , LSD = 6.87; Fig 2a). However, isolate L44 also produced significantly heavier sclerotia ( $P < 0.001$ ; Fig 2b) across crop type (27 mg) compared to L17 (19 mg) and L6 (16 mg). This therefore resulted in there being no significant differences overall between isolates for the mean total weight of sclerotia produced per plant ( $P = 0.16$ ). Isolate L6 produced a greater mean number and weight of sclerotia on potato compared to the other isolates as this was the only isolate to completely colonise plants. Significant crop x isolate interactions were observed for mean number of sclerotia ( $P < 0.001$ ), mean weight ( $P = 0.004$ ) of sclerotia and mean weight per sclerotium ( $P < 0.001$ ) suggesting that isolates behave differently depending on host. However, crop type was the main driving factor for the variation in these data.

Across the three isolates, the mean number of sclerotia produced on agar was correlated with the number produced on crops ( $r = 0.49$ ,  $P = 0.031$ ). However, there was no correlation between results on agar and on plants for mean total weight of sclerotia ( $r = 0.11$ ,  $P = 0.35$ ), or for the mean weight per sclerotium ( $r = 0.073$ ,  $P = 0.40$ ).

### **Effect of crop type and size of sclerotia on carpogenic germination**

Percentage germination was significantly greater for larger sclerotia of *S. sclerotiorum* L6 produced on crop plants ( $P < 0.001$ ; Table 2) with those  $>6.7$  mm having a mean germination of 95.8% compared with those  $<2$  mm having a mean germination of 62.4% across the different crop types. This was consistent irrespective of crop type ( $P = 0.68$  for crop x size interaction). This trend was also observed for the sclerotia produced on wheat grain where the largest sclerotia ( $>6.7$  mm) resulted in 100% germination compared with 78.9% for the smallest sclerotia ( $<2$  mm). Larger sclerotia of *S. sclerotiorum* isolate L6 also produced significantly more apothecia than smaller ones ( $P < 0.001$ , Table 2) irrespective of crop type ( $P = 0.70$  for crop x size interaction). The largest sclerotia ( $>6.7$  mm) produced on OSR yielded an average of 11.9 apothecia per sclerotium compared to 1.1 apothecia per sclerotium for sclerotia  $<2$  mm, 1.6 apothecia per sclerotium for sclerotia 2-4 mm and 2.5 apothecia per sclerotium for sclerotia 4-6.7 mm (across all crop types). The same trend was again observed for *S. sclerotiorum* sclerotia produced on wheat grain ( $P < 0.001$ ; Table 2) where mean numbers of apothecia ranged from 1.6 ( $<2$  mm) to 15.4 ( $> 6.7$ mm).

Larger sclerotia of *S. sclerotiorum* isolate L6 (produced on crops) also germinated significantly faster ( $P < 0.001$ ; Table 2) with a time to 33% germination of 22 days in the largest sclerotia ( $>6.7$  mm) compared to 60 days in the smallest ( $< 2$  mm). Again, this was consistent across the crop types ( $P = 0.16$  for interaction). For sclerotia artificially produced on wheat grain, there were significant differences ( $P = 0.003$ ) in time to 33% germination but in this case, the time for 33% germination for the largest sclerotia (mean 22 days) was not significantly different from the smallest (mean 24 days), with the mid-range sized sclerotia germinating more rapidly (14-16 days).

Crop type significantly affected the number of apothecia produced ( $P < 0.001$ ; Table 2) and this was particularly evident for OSR where sclerotia generally produced more

apothecia; e.g. for the size class 4.0-6.7 mm across all the crops tested, sclerotia from OSR produced 3.2 apothecia compared to 2.0, 1.9, 2.6 and 1.7 for potato, bean, lettuce and carrot respectively. Crop type also had a significant effect on final percentage germination ( $P < 0.001$ ), with those from OSR exhibiting greater germination; e.g. for size class 2-4 mm, sclerotia from OSR resulted in 80.0% germination compared to 59.6, 66.7, 71.1 and 45.6% for potato, bean, lettuce and carrot respectively. Finally, crop type also had a significant effect on germination rate (time to 33% germination,  $P < 0.001$ ; Table 2). Sclerotia from bean were generally the fastest to germinate; e.g. in the size class 2-4mm the time to 33% was 30 days compared to 42, 41, 35 and 69 days for potato, oilseed rape, lettuce and carrot respectively. Sclerotia from carrot were all particularly slow to germinate (53-71 days).

#### **Estimation of sclerotial production for different crops in the field**

Using field planting densities of 3.3, 8, 28, 40 and 150 plants  $m^{-2}$  for potato, lettuce, OSR, bean and carrot respectively, the number of *S. sclerotiorum* sclerotia that could potentially be returned to the soil for each crop was calculated for each isolate (assuming complete plant colonisation). This resulted in the greatest potential number of sclerotia being returned for carrot (2666-4794  $m^{-2}$ ; mean 3944  $m^{-2}$  across isolates) compared to the other crops with the fewest sclerotial returns from potato (50-298  $m^{-2}$ ; mean 145  $m^{-2}$  across isolates (Fig. 3a). When weight of sclerotia produced by the different crops was considered, OSR (56.3-86.4  $g m^{-2}$ , mean 73.3  $g m^{-2}$  across isolates) and carrot (35.2-82.6  $g m^{-2}$ , mean 60.5  $g m^{-2}$  across isolates) returned the greatest values (Fig. 3b).

## Discussion

The number of sclerotia in the soil increases when successive susceptible crops become infected by *S. sclerotiorum* leading to escalating disease problems in many cropping rotations (Mueller *et al.*, 2002). The situation is exacerbated by short rotations and pressure on land use as *S. sclerotiorum* sclerotia are generally thought to survive for 4-5 years in soil under natural conditions (Adams & Ayers, 1979) although this period can vary depending on factors such as depth, temperature and moisture (Mitchell & Wheeler, 1990, Duncan *et al.*, 2006, Ćosić *et al.*, 2012). However, few researchers have attempted to investigate and quantify the effect of crop type and *S. sclerotiorum* isolate on the production of sclerotia and hence understand how these factors might influence subsequent inoculum pressure.

This is the first study to investigate differences in sclerotial production on whole plants rather than on detached tissue which was the approach used in a previous study by Leiner and Winton (2006). Here, a greater number of sclerotia were produced on carrot pieces compared to celery and these were also larger, with weights ranging from 38-90 mg compared to 14-18 mg (Leiner & Winton, 2006). In the same work, the weight of sclerotia produced by two different *S. sclerotiorum* isolates on three different iceberg lettuce varieties ranged from 14-26 mg. By comparison, an almost identical weight range for *S. sclerotiorum* sclerotia produced on lettuce (13-28 mg) was observed in the current study, but much smaller sclerotia were produced on carrot (12-14 mg) with the largest sclerotia produced on oilseed rape (33-60 mg). Leiner & Winton (2006) also estimated that 250-500 sclerotia might be produced on a single iceberg lettuce plant depending on *S. sclerotiorum* isolate, which is substantially more sclerotia than observed on the infected butterhead lettuce in this work (55-122 per plant). However, this might be explained by iceberg lettuce forming a dense compact head similar to a cabbage, which

would have a much greater amount of biomass and hence sustain a greater level of sclerotial production than the open, leafy butterhead type of lettuce used here.

The *S. sclerotiorum* isolates tested in this study exhibited variation in the number and weight of sclerotia produced both on PDA and on plants. This is consistent with previous reports where sclerotial production was assessed on agar media (Akram *et al.*, 2005, Li *et al.*, 2008, Vleugels *et al.*, 2013). Interestingly, the three *S. sclerotiorum* isolates showed a consistent pattern of sclerotial production irrespective of crop type; L6 produced large numbers of small sclerotia while L44 produced small numbers of large sclerotia with L17 intermediate between the two. Leiner & Winton (2006) also observed that one of the two *S. sclerotiorum* isolates they examined consistently produced more sclerotia on cabbage, carrot, celery and lettuce tissue. As well as being more numerous, the sclerotia from this isolate were also larger. This in contrast to our results where all three isolates produced a comparable total biomass of sclerotia either by producing a large number of small sclerotia (L6) or a small number of large sclerotia (L44). These different findings may be due to different experimental approaches but merits further investigation with a greater range of isolates from different locations. The production of large sclerotia by some *S. sclerotiorum* isolates may also confer a survival advantage as they have been shown to persist for longer periods in the soil (Harvey *et al.*, 1995).

As indicated previously, variation in sclerotial production between different *S. sclerotiorum* isolates has generally only been investigated previously on agar media. In the current study, only a weak correlation was observed between the number of sclerotia produced on PDA *in vitro* and on plants ( $r = 0.49$ ) and there was no correlation between the results on plants and agar for either total weight of sclerotia or mean weight per sclerotium. This highlights the importance of using whole plants to properly assess sclerotial production by different *S. sclerotiorum* isolates. There was also no significant correlation between either

number of sclerotia, weight of sclerotia or weight per sclerotium on PDA and pathogenicity on *Brassica* (Taylor *et al.*, 2015). This is in agreement with previous work for *S. sclerotiorum* isolates from sunflower (Li *et al.*, 2008) but in contrast to work on *S. trifoliorum* from red clover where a negative correlation was observed between the number of sclerotia produced and aggressiveness (Vleugels *et al.*, 2013). However, as the correlations observed in the current study were close to being significant, it is possible that a larger sample size would reveal a weak correlation between number / weight of sclerotia produced and pathogenicity.

It was also clear from the results that as sclerotial size increased from small (< 2.0 mm, mean weight 4.6mg) to medium (2.0-4.0 mm, mean weight 18.4 mg) and large (4.0-6.7 mm, mean weight 43.3mg), percentage germination and number of apothecia produced also increased. This observation was generally consistent across all crop types with the larger sclerotia produced on OSR (> 6.7 mm) resulting in a mean of 12 apothecia per sclerotium. In the few similar studies that have been reported, Ben-Yephet *et al.*, (1993) collected sclerotia from infected lettuce in the field and showed that small sclerotia (1-7 mg) resulted in 11-21% germination compared to 29-31% germination for larger sclerotia (7-40 mg). Dillard *et al.* (1995) produced sclerotia on cornmeal/vermiculite and, following a cold conditioning treatment, observed germination ranging from 76% for small sclerotia (1.68-3.36 mm) to 98% for medium (3.36-4.75 mm) and large (>4.75 mm) sclerotia with mean numbers of apothecia per sclerotium of 1.5, 3.2 and 7.8 respectively, which closely matches the results presented here. Similarly, using sclerotia produced on autoclaved potato pieces, Hao *et al.* (2003) recorded 30% germination for small sclerotia (< 1mm) and 100% for large sclerotia (>4.75 mm) and also reported a significant correlation between size of sclerotia and the number of stipes produced per apothecium. However, none of these previous studies tested *S. sclerotiorum* sclerotia produced on a range of crop plants. Furthermore, the results presented here also indicated that larger sclerotia germinate more quickly which, to our knowledge, has

not been previously reported. The reason for the rapid, high levels of carpogenic germination as well as the production of multiple apothecia associated with larger sclerotia of *S. sclerotiorum* may be due to the accumulation of greater nutrient reserves which are utilised during germination (Willetts & Bullock, 1992). However, little is known about how the composition or quantity of different compounds found in sclerotia such as glycogen, protein, polyphosphate and lipid affect survival or germination (Willetts & Bullock, 1992). The number and weight of *S. sclerotiorum* sclerotia produced on agar has been demonstrated previously to increase linearly with sucrose concentration but subsequent production of apothecia for the larger sclerotia was slower or completely inhibited (Budge & Whipps, 1991). Although this conflicts with the results from this and other reports, no conditioning treatment of the sclerotia was employed which the authors suggest may be required by larger sclerotia if nutrient depletion during the survival phase of the pathogen plays a part in triggering carpogenic germination. It was also observed in the current study that *S. sclerotiorum* sclerotia produced on oilseed rape and those produced artificially on autoclaved wheat grain germinated more rapidly and resulted in a higher level of germination. This could again relate to the nutrient status of the sclerotia and highlights the need to understand how different artificial and natural substrates affect both formation and germination of *S. sclerotiorum* sclerotia.

The data generated for the number of *S. sclerotiorum* sclerotia produced on infected plants in this study allowed estimation of the inoculum potential for different crops. Although infection of OSR and lettuce resulted in greater numbers of sclerotia per plant than the other crops, carrot was identified as potentially supporting the greatest returns of sclerotia to the soil when planting density in the field was taken into account, producing between 2666-4794 sclerotia m<sup>-2</sup> compared with 1026-2754 sclerotia m<sup>-2</sup> in OSR. However, based on the weight of sclerotia produced per plant and crop density, both OSR and carrot constitute the greatest returns with 86 g and 83 g sclerotia m<sup>-2</sup> respectively. These are estimated figures and also

assume that all plants are completely colonised by *S. sclerotiorum* which is unlikely in a field situation. Nonetheless these estimates appear to be consistent with some preliminary field data where sclerotia on the soil surface were counted in quadrats for crops with high levels of Sclerotinia disease. Here, up to 3000 sclerotia m<sup>-2</sup> were recorded for a carrot crop compared to up to 900 m<sup>-2</sup> for OSR and 195 m<sup>-2</sup> for potato (Young *et al.*, 2014). Data from a published field survey in the UK suggested that pea, potato and sunflower crops resulted in high returns of sclerotia to the soil of up to six sclerotia kg<sup>-1</sup> soil with 92 sclerotia m<sup>-2</sup> being recorded on the soil surface of an infected pea crop (Archer *et al.*, 1992). These crops were therefore considered a major threat to subsequent OSR plantings. However, subsequent tillage operations may redistribute sclerotia left on the soil surface throughout the soil profile and many will be killed by microorganisms or adverse environmental conditions (Ćosić *et al.*, 2012). Only a small proportion of sclerotia that remain close to the soil surface can subsequently germinate (Wu & Subbarao, 2008) and these are difficult to quantify. As stated in the introduction, just a few germinating *S. sclerotiorum* sclerotia are required to initiate disease due to the large numbers of ascospores produced by apothecia and as few as two apothecia per m<sup>-2</sup> were shown to lead to 11% disease incidence in soybean (Mueller *et al.*, 2002).

In summary, sclerotial production and germination by *S. sclerotiorum* is dependent on isolate, host plant and size of sclerotia. Both carrot and OSR potentially return more sclerotia by weight or number to the soil compared to the other plants tested here and hence potentially pose a bigger disease risk for subsequent susceptible crops. However, further work is required to extensively quantify sclerotial production in the field in order to validate this. This information could then be used to develop optimum crop rotation strategies to reduce the impact of Sclerotinia disease and maximise financial returns. This has been attempted for *S. sclerotiorum* using a modelling approach based on assumptions relating to potential yield loss for different combinations of susceptible and break crops over time (Ahmadi *et al.*, 2013).

Here, it was concluded that rotations selected to reduce build-up of sclerotia were financially justified.

### **Acknowledgements**

We gratefully acknowledge funding for this work from HGCA (AHDB) and Defra through the Sustainable Arable LINK programme (Defra LINK project SA09130 / LK09130; HGCA project RD-2008-3579). We also thank the Warwick Crop Centre Horticultural Services team for raising and maintaining plants.

## References

- Adams PB, Ayers PB, 1979. Ecology of *Sclerotinia* Species. *Phytopathology* **69**, 896-899.
- Ahmadi, BV, Burnett, FJ, Young, CS, McQuilken, MP, Stott, AW, 2013. Reducing the impact of *Sclerotinia* disease by determining optimum crop rotations using dynamic programming. 87th Annual Conference of the Agricultural Economics Society, University of Warwick, United Kingdom.
- Akram A, Iqbal SM, Ahmed N, Iqbal U, Ghafoor A, 2005. Morphological variability and mycelial compatibility among the isolates of *Sclerotinia sclerotiorum* associated with stem rot of chickpea. *Pakistan Journal of Botany* **40**, 2663-2668.
- Archer S, Mitchell S, Wheeler B, 1992. The effects of rotation and other cultural factors on *Sclerotinia* in oilseed rape, peas and potatoes. In. *Brighton Crop Protection Conference, Pests and Diseases* **1**, 99-108.
- Atallah ZK, Johnson DA, 2004. Development of *Sclerotinia* stem rot in potato fields in South-Central Washington. *Plant Disease* **88**, 419-423.
- Ben-Yephet Y, Genizi A, Siti E, 1993. Sclerotial survival and apothecial production by *Sclerotinia sclerotiorum* following outbreaks of lettuce drop. *Phytopathology* **83**, 509-513.
- Boland GJ, Hall R, 1994. Index of plant hosts of *Sclerotinia sclerotiorum*. *Canadian Journal of Plant Pathology* **16**, 93-108.

Bolton MD, Thomma BP, Nelson BD, 2006. *Sclerotinia sclerotiorum* (Lib.) de Bary: biology and molecular traits of a cosmopolitan pathogen. *Molecular Plant Pathology* **7**, 1-16.

Budge SP, Whipps JM, 1991. Glasshouse trials of *Coniothyrium minitans* and *Trichoderma* species for the biological control of *Sclerotinia sclerotiorum* in celery and lettuce. *Plant Pathology* **40**, 59-66.

Chitrampalam P, Pryor BM, 2013. Population density and spatial pattern of sclerotia of *Sclerotinia sclerotiorum* in desert lettuce production fields. *Canadian Journal of Plant Pathology* **35**, 494-502.

Chitrampalam P, Turini TA, Matheron ME, Pryor BM, 2010. Effect of Sclerotium Density and Irrigation on Disease Incidence and on Efficacy of *Coniothyrium minitans* in Suppressing Lettuce Drop Caused by *Sclerotinia sclerotiorum*. *Plant Disease* **94**, 1118-1124.

Clarkson JP, Carter HE, Coventry E, 2010. First report of *Sclerotinia subarctica* nom. prov. (*Sclerotinia* species 1) in the UK on *Ranunculus acris*. *Plant Pathology* **59**, 1173.

Clarkson JP, Fawcett L, Anthony SG, Young C, 2014. A model for *Sclerotinia sclerotiorum* infection and disease development in lettuce, based on the effects of temperature, relative humidity and ascospore density. *Plos One* **9**, e94049.

Clarkson JP, Phelps K, Whipps JM, Young CS, Smith JA, Watling M, 2007. Forecasting *Sclerotinia* disease on lettuce: a predictive model for carpogenic germination of *Sclerotinia sclerotiorum* sclerotia. *Phytopathology* **97**, 621-31.

Clarkson JP, Staveley J, Phelps K, Young CS, Whipps JM, 2003. Ascospore release and survival in *Sclerotinia sclerotiorum*. *Mycological Research* **107**, 213-22.

Clarkson JP, Warmington RJ, Walley PG, Denton-Giles M, Barbetti MJ, Brodal G, Nordskog B, 2017. Population structure of *Sclerotinia subarctica* and *Sclerotinia sclerotiorum* in England, Scotland and Norway. *Frontiers in Microbiology* **8**, 490.

Ćosić J, Jurković D, Vrandečić K, Kaučić D, 2012. Survival of buried *Sclerotinia sclerotiorum* sclerotia in undisturbed soil. *Helia* **35**, 56-73

Dillard HR, Ludwig JW, Hunter JE, 1995. Conditioning sclerotia of *Sclerotinia sclerotiorum* for carpogenic germination. *Plant Disease* **79**, 411-5.

Duncan RW, Dilantha Fernando WG, Rashid KY, 2006. Time and burial depth influencing the viability and bacterial colonization of sclerotia of *Sclerotinia sclerotiorum*. *Soil Biology and Biochemistry* **38**, 275-284.

Foster AJ, McDonald MR, Boland GJ, 2008. Disease progression of *Sclerotinia* rot of carrot, caused by *Sclerotinia sclerotiorum*, from shoot to root before and after harvest. *Canadian Journal of Plant Pathology* **30**, 206-213.

Gugel RK, 1986. Inoculum disease relationships in *Sclerotinia* stem rot of rapeseed in Saskatchewan. *Canadian Journal of Plant Pathology* **8**, 89-96.

Hao JJ, Subbarao KV, 2005. Comparative analyses of lettuce drop epidemics caused by *Sclerotinia minor* and *S. sclerotiorum*. *Plant Disease* **89**, 717-725.

Hao JJ, Subbarao KV, Duniway JM, 2003. Germination of *Sclerotinia minor* and *S. sclerotiorum* sclerotia under various soil moisture and temperature combinations. *Phytopathology* **93**, 443-450.

Harvey IC, Foley LM, Saville DJ, 1995. Survival and germination of shallow-buried sclerotia of *Sclerotinia sclerotiorum* in pastures in Canterbury. *New Zealand Journal of Agricultural Research* **38**, 279-284.

Jensen BD, Finckh MR, Munk L, Hauser TP, 2008. Susceptibility of wild carrot (*Daucus carota* ssp. *carota*) to *Sclerotinia sclerotiorum*. *European Journal of Plant Pathology* **122**, 359-367.

Lehner MS, Pethybridge SJ, Meyer MC, Del Ponte EM, 2017. Meta-analytic modelling of the incidence-yield and incidence-sclerotial production relationships in soybean white mould epidemics. *Plant Pathology* **66**, 460-468.

Leiner RH, Winton LM, 2006. Differential production of sclerotia by isolates of *Sclerotinia sclerotiorum* from Alaska. *Canadian Journal of Plant Pathology* **28**, 435-440.

Li Z, Zhang M, Wang Y, Li R, Fernando WGD, 2008. Mycelial compatibility group and pathogenicity variation of *Sclerotinia sclerotiorum* populations in sunflower from China, Canada and England. *Plant Pathology Journal* **7**, 131-139.

Liu Y, Paul VH, 2007. Studies on the germination of sclerotia of *Sclerotinia sclerotiorum*. *Journal of Plant Diseases and Protection* **114**, 14-19.

Mitchell SJ, Wheeler BEJ, 1990. Factors affecting the production of apothecia and longevity of sclerotia of *Sclerotinia sclerotiorum*. *Plant Pathology* **39**, 70-76.

Mueller DS, Pedersen WL, Hartman GL, 2002. Effect of crop rotation and tillage system on sclerotinia stem rot on soybean. *Canadian Journal of Plant Pathology* **24**, 450-456.

Mylchreest SJ, Wheeler BEJ, 1987. A method for inducing apothecia from sclerotia of *Sclerotinia sclerotiorum*. *Plant Pathology* **36**, 16-20.

Schwartz H, Singh SP, 2013. Breeding common bean for resistance to white mold: a review. *Crop Science* **53**, 1-13.

Taylor A, Coventry E, Jones JE, Clarkson JP, 2015. Resistance to a highly aggressive isolate of *Sclerotinia sclerotiorum* in a *Brassica napus* diversity set. *Plant Pathology* **64**, 932-940.

Taylor A, Rana K, Handy C, Clarkson JP, 2018. Resistance to *Sclerotinia sclerotiorum* in wild Brassica species and the importance of *Sclerotinia subarctica* as a Brassica pathogen. *Plant Pathology* **67**, 433-444.

Vleugels T, Baert J, Van Bockstaele E, 2013. Morphological and pathogenic characterization of genetically diverse *Sclerotinia* isolates from european red clover crops (*Trifolium pratense* L.). *Journal of Phytopathology* **161**, 254-622.

Warmington R, Clarkson JP, 2016. Volatiles from biofumigant plants have a direct effect on carpogenic germination of sclerotia and mycelial growth of *Sclerotinia sclerotiorum*. *Plant and Soil* **401**, 213-229.

Willettts HJ, Bullock S, 1992. Developmental biology of sclerotia. *Mycological Research* **96**, 801-816.

Wu BM, Subbarao KV, 2008. Effects of soil temperature, moisture, and burial depths on carpogenic germination of *Sclerotinia sclerotiorum* and *S. minor*. *Phytopathology* **98**, 1144-1152.

Young CS, Clarkson JP, Smith JA, Watling M, Phelps K, Whipps JM, 2004. Environmental conditions influencing *Sclerotinia sclerotiorum* infection and disease development in lettuce. *Plant Pathology* **53**, 387-397.

Young C, Clarkson J, West J, Bouda Ahmadi B, Burnett F, 2014. Reducing the impact of *Sclerotinia* disease on arable rotations, vegetable crops and land use. AHDB Final Project Report PR538. <https://cereals.ahdb.org.uk/media/550794/rd-2008-3579-fpr.pdf>.

**Table 1:** *Sclerotinia* isolates evaluated for production of sclerotia *in vitro* or on plants.

<b>Isolate</b>	<b>Host</b>	<b>Location</b>	<b>Year</b>	<b>Microsatellite haplotype<sup>a</sup></b>
<i>S. sclerotiorum</i>				
C28	Carrot	Nottinghamshire	2005	390
CE11	Celery	Norfolk	2009	1
DG4	Buttercup	Warwickshire	2009	391
EV9	Buttercup	Powys	2009	28
HE33	Buttercup	Herefordshire	2009	138
L17	Lettuce	Sussex	2005	1
L44	Lettuce	Sussex	2005	63
L5	Lettuce	Sussex	2005	2
L6	Lettuce	Sussex	2005	3
O92	Oilseed rape	Herefordshire	2005	182
P2	Pea	Herefordshire	2009	31
P7	Pea	Herefordshire	2009	1
R12	Buttercup	Warwickshire	2007	21
R17	Buttercup	Warwickshire	2007	199
R19	Buttercup	Warwickshire	2007	72
R28	Buttercup	Warwickshire	2007	6
R30	Buttercup	Warwickshire	2007	206
<i>S. subarctica</i>				
HE1	Buttercup	Herefordshire	2009	1

<sup>a</sup>Clarkson *et al.*, 2017

**Table 2:** Germination and number of apothecia produced by different sizes of *S. sclerotiorum* sclerotia (isolate L6) produced on different crop plants. Letters (in bold) denote significant differences following REML analysis.

Crop Plant	Size class (mm)	Percentage of sclerotia in size class	Percentage germination	Time to 33% germination (days)	Number of apothecia per sclerotium
Potato	< 2.0	41.7	73.3	51.7	1.1
	2.0-4.0	53.6	59.6	42.3	1.3
	4.0-6.7	4.5	84.4	33.0	2.0
	> 6.7	0.2	-	-	-
	<i>LSD</i>		14.7	13.4	0.46
Bean	< 2.0	43.6	50.0	57.7	1.0
	2.0-4.0	50.2	66.7	29.7	1.6
	4.0-6.7	5.1	60.0	25.0	1.9
	> 6.7	1.1	-	-	-
	<i>LSD</i>		14.7 <sup>a</sup>	13.4 <sup>b</sup>	0.48 <sup>c</sup>
OSR	< 2.0	25.0	64.4	56.3	1.1
	2.0-4.0	52.7	80.0	41.3	2.3
	4.0-6.7	16.6	83.3	32.0	3.2
	> 6.7	5.7	95.8	22.3	11.9
	<i>LSD</i>		14.7	13.4	0.98
Lettuce	< 2.0	37.2	59.2	62.3	1.1
	2.0-4.0	59.1	71.1	35.3	1.5
	4.0-6.7	3.2	86.7	26.0	2.6
	> 6.7	0.5	-	-	-
	<i>LSD</i>		14.7	13.4	0.47
Carrot	< 2.0	42.8	43.3	71.3	1.2
	2.0-4.0	48.6	45.6	69.0	1.5
	4.0-6.7	7.4	46.7	53.0	1.7
	> 6.7	1.1	-	-	-
	<i>LSD</i>		14.7 <sup>a</sup>	13.4 <sup>b</sup>	0.57 <sup>d</sup>
Mean (across all crop types)	< 2.0	38.1	62.4 <b>a</b>	59.9 <b>a</b>	1.1 <b>a</b>
	2.0-4.0	52.9	68.4 <b>a</b>	43.5 <b>b</b>	1.6 <b>a</b>
	4.0-6.7	7.4	84.0 <b>b</b>	31.6 <b>c</b>	2.5 <b>b</b>
	> 6.7	1.7	95.8 <b>b</b>	22.3 <b>c</b>	11.9 <b>c</b>
Wheat grain	< 2.0	46.5	78.9 <b>a</b>	23.7 <b>a</b>	1.6 <b>a</b>
	2.0-4.0	48.5	88.9 <b>b</b>	16.0 <b>b</b>	3.9 <b>b</b>
	4.0-6.7	4.8	97.8 <b>c</b>	14.0 <b>b</b>	4.9 <b>c</b>
	> 6.7	0.3	100.0 <b>c</b>	22.0 <b>a</b>	15.4 <b>d</b>

LSD values as stated except when comparing any treatment to the 4.0-6.7 size class where the replication was lower and hence LSD values were:

<sup>a</sup>20.1

<sup>b</sup>19.2

<sup>c</sup>0.66

<sup>d</sup>0.79

## Figure legends

**Figure 1:** Number (a), total weight (b) and individual weight (c) of sclerotia produced by 18 *Sclerotinia* isolates on PDA. Error bars represent the LSD (5% level).

**Figure 2:** Number (a), total weight (b) and individual weight (c) of sclerotia produced by three *S. sclerotiorum* isolates on five different crop plants following inoculation. Error bars represent the LSD values for comparing either individual means (In), crop means (Cr) or isolate means (Is) or at the 5% level. + indicates incomplete colonisation.

**Figure 3:** Estimated number (a) and weight (b) of sclerotia produced on different crop plants in the field. Error bars represent the LSD values for comparing either individual means (In), crop means (Cr) or isolate means (Is) at the 5% level.

Figure 1




Figure 2




Figure 3

