

**Original citation:**

Pope, Tom, Hough, Gemma, Arbona, Charlotte, Bennison, Jude, Roberts, Harriet, Prince, Gillian and Chandler, Dave (2018) *Investigating the potential of an auto-dissemination system for managing populations of vine weevil, Otiorhynchus sulcatus (Coleoptera : Curculionidae) with entomopathogenic fungi*. Journal of Invertebrate Pathology, 154. pp. 79-84. doi:[10.1016/j.jip.2018.04.002](https://doi.org/10.1016/j.jip.2018.04.002)

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/98544>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

© 2018, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

**A note on versions:**

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)

1 **Investigating the potential of an autodissemination system for**  
2 **managing populations of vine weevil, *Otiorhynchus sulcatus***  
3 **(Coleoptera: Curculionidae) with entomopathogenic fungi**

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

1 **HIGHLIGHTS**

- 2 1. Simple plastic artificial refuges for vine weevil (*Otiorhynchus sulcatus*)  
3 can be used to disseminate an entomopathogenic fungus through vine  
4 weevil populations.
- 5 2. Isolates of *Beauveria bassiana* and *Metarhizium brunneum* cause up to  
6 100% mortality in vine weevil adults under laboratory conditions.
- 7 3. Conidial powders of a *Metarhizium brunneum* isolate placed in artificial  
8 refuges significantly increased vine weevil mortality under polytunnel  
9 conditions.

10

11 **ABSTRACT**

12 Vine weevil, also known as black vine weevil, (*Otiorhynchus sulcatus*) is an  
13 economically important pest affecting soft fruit and nursery stock crops in  
14 temperate regions. We used laboratory and polytunnel experiments to  
15 investigate a novel control system based on autodissemination of spores of an  
16 entomopathogenic fungus used on their own or formulated with talc to  
17 populations of adult vine weevils. The fungus was applied as a conidial powder  
18 to a simple plastic artificial refuge for vine weevils. The potential for adult  
19 weevils to disseminate the fungus was investigated first in polytunnel  
20 experiments using fluorescent powders applied to the refuge in lieu of fungal  
21 conidia. In this system, 88% of adult weevils came in contact with the powder  
22 within 48 hours. When the powder was applied to five adult weevils that were  
23 then placed within a population of 35 potential recipients, it was transmitted on  
24 average to 75% of the recipient population within 7 days. Three isolates of  
25 entomopathogenic fungi (*Beauveria bassiana* 433.99, *B. bassiana* 1749.11,

1 *Metarhizium brunneum* 275.86), selected from a laboratory virulence screen,  
2 were then investigated for efficacy when applied as conidial powders in artificial  
3 refuges placed among populations of adult weevils held in experimental boxes  
4 in the laboratory at 20°C. Under this regime, the fungal isolates caused 70 –  
5 90% mortality over 28 days. A final polytunnel experiment tested the efficacy  
6 of conidial powders of *M. brunneum* 275.86 placed in artificial refuges to  
7 increase vine weevil mortality. Overall weevil mortality was relatively low (26-  
8 41%) but was significantly higher in cages in which the conidial powders were  
9 placed in refuge traps than in cages with control traps. The lower weevil  
10 mortality recorded in the polytunnel experiment compared to the laboratory test  
11 was most likely a consequence of the greater amounts of inoculum required to  
12 kill adult weevils when conditions fluctuate between favourable and  
13 unfavourable temperatures e.g. below 15°C. The potential of an  
14 autodissemination system for entomopathogenic fungi as a means of  
15 controlling vine weevil as part of an integrated pest management programme is  
16 discussed.

17

18 Key words: *Beauveria bassiana*; *Metarhizium brunneum*; autodissemination;  
19 refuge; aggregation

20

21

22

## 23 **INTRODUCTION**

24 Vine weevil, also known as black vine weevil, (*Otiorhynchus sulcatus*) is an  
25 economically damaging pest affecting soft fruit and nursery stock crops

1 (Moorhouse *et al.*, 1992; van Tol *et al.*, 2012). It is widely distributed throughout  
2 temperate regions including northern Europe and North America (Warner &  
3 Negley, 1976; Lundmark, 2010). Damage is caused both by the adults, which  
4 feed on leaves, and larvae, which feed on plant roots, corms and tubers (Smith,  
5 1932; Moorhouse *et al.*, 1992). As the larvae are root pests and the adult  
6 weevils are nocturnal, an infestation may pass unnoticed until leaf notching is  
7 evident or plants show signs of wilting, by which time they will have been  
8 damaged beyond recovery (van Tol *et al.*, 2012).

9

10 Biological control using entomopathogenic nematodes and fungi is used  
11 against vine weevil larvae (e.g. Willmott *et al.*, 2002; Georgis *et al.*, 2006; Shah  
12 *et al.*, 2007; Ansari *et al.*, 2008). At present, control of adult vine weevils is  
13 based on use of broad spectrum chemical insecticides (van Tol *et al.*, 2012  
14 Insecticide sprays are often applied at dusk, when the weevils become active,  
15 which makes it difficult to effectively target applications. Broad-spectrum  
16 insecticides also have a negative impact on biocontrol agents used against  
17 other pests and naturally occurring beneficial insects, such as ground beetles  
18 that prey upon vine weevil adults (Cross *et al.*, 2001). Therefore, more  
19 sustainable solutions for adult vine weevils are needed.

20

21 For this study, we were interested in the potential for biological control of adult  
22 vine weevils using autodissemination of entomopathogenic fungi (EPF) to  
23 complement existing biological control of the larval stage of this pest.  
24 Autodissemination is an application system in which pest insects enter a device  
25 containing a reservoir of an entomopathogen, which they then disseminate to

1 other individuals within their environment (Soper, 1978). It has been developed  
2 to control a range of insect pests with EPF including emerald ash borer (Lyons  
3 *et al.*, 2012), Mediterranean fruit fly (Quesada-Moraga *et al.*, 2008), sweet  
4 potato weevil (Yasuda, 1999) and damson-hop aphid (Hartfield *et al.*, 2001).  
5 However, there is no mention in the available literature of investigations for use  
6 of this approach in the control of vine weevil. Adult vine weevils are known to  
7 be susceptible to EPF infection (Moorhouse *et al.*, 1992), although they die  
8 more slowly than infected weevil larvae (Moorhouse, 1990). They also  
9 aggregate in refuges during the day (Smith, 1932; Moorhouse *et al.*, 1992; van  
10 Tol *et al.*, 2004), which could be used as sources of fungal inoculum. Here we  
11 present results from a series of experiments testing the potential efficacy of  
12 autodisseminating an EPF through the use of artificial refuges.

13

## 14 **EXPERIMENTAL METHODS**

### 15 *Vine weevil culture*

16 Adult vine weevils were collected from commercial soft fruit crops in  
17 Staffordshire, UK, and maintained under laboratory conditions at 21°C in  
18 groups of 25-30 individuals in ventilated plastic boxes (200 L x 100 W x 95 D  
19 mm) lined with damp tissue paper (a source of moisture) and a refuge  
20 (corrugated cardboard – 70 x 50 mm). Weevils were fed leaves of yew, *Taxus*  
21 *baccata*, *ad libitum*.

22

### 23 *EPF culture – storage and production*

24 EPF isolates were taken from the Warwick Crop Centre (WCC) collection of  
25 entomopathogenic fungal cultures. Isolates were stored on porous plastic

1 beads at minus 80°C (Chandler, 1994). Laboratory cultures were grown from  
2 these beads on Sabouraud dextrose agar (SDA) slopes and maintained in a  
3 refrigerator at 4°C for up to six months. To produce conidia for experiments,  
4 subcultures were grown from the slope cultures on SDA Petri plates at 23 ±  
5 1°C for 10-12 days in the dark.

6

7 *Experiment 1: Acquisition of fluorescent marker powders by adult weevils from*  
8 *artificial refuges*

9 Based on results from preliminary experiments, Roguard (BASF plc, Cheadle  
10 Hulme, UK) crawling insect bait stations, were selected for use as simple vine  
11 weevil refuges. These bait stations have a black plastic construction (80 mm  
12 diameter x 15 mm height) with four entrances (20 mm x 5 mm) and were used  
13 without the addition of a bait in these experiments. The Roguard bait stations  
14 were otherwise not modified for use as vine weevil refuges. Any aggregation by  
15 weevils within a bait station was as a result of a strong aggregation behaviour  
16 shown by vine weevil adults (e.g. Smith, 1932; Moorhouse *et al.*, 1992). The  
17 artificial refuges were tested in gauze 'tent' cages (145 x 145 x 152 cm) placed  
18 in a ventilated polytunnel (mean temperatures were 22-26°C (daytime) and 11-  
19 13°C (night time). Sixteen *Euonymus fortunei* (cv. Emerald Gaiety) plants  
20 grown in 1.5 L pots using John Innes No. 2 compost (William Sinclair  
21 Horticulture Ltd., Lincoln, UK) were placed on the floor of the cage. Forty adult  
22 weevils were then released into each cage and left to acclimatise for 24 hours,  
23 after which 12 Roguard refuges were placed into each cage. Six refuges were  
24 spread evenly across the floor of the cage while the other six were placed on  
25 the surface of the compost of six pots. Each refuge contained 0.2 g of a

1 hydrophobic fluorescent powder (Swada, Stalybridge, UK) placed in the central  
2 well. The fluorescent powder was used to quantify the numbers of weevils  
3 entering the refuge. Adult weevils were collected seven days after placing the  
4 refuges in the cages and scored for the presence / absence of fluorescent  
5 powder by examining them under a UV light (Lighting Ever, Birmingham, UK).  
6 There were eight replicate cages.

7

#### 8 *Experiment 2: Dissemination of fluorescent powders among adult weevils*

9 Gauze 'tent' cages were prepared as previously described. Thirty-five weevils  
10 were released into each cage and left to acclimatise for 24 hrs, after which 12  
11 Roguard refuges were placed into each cage and arranged as previously  
12 described but with no fluorescent powder. Five adult vine weevils, marked with  
13 water-based paint and then coated in yellow fluorescent powder by placing the  
14 weevils into a 20 ml specimen tube containing approximately 1 g of fluorescent  
15 powder. The lid of the tube was secured in place before gently rotating the tube  
16 for 30 s to ensure that all of the weevils had become coated in the powder.  
17 Each group of five powder coated weevils were placed into a ventilated plastic  
18 box lined with tissue paper for 30 minutes to allow excess powder to be  
19 dislodged before the weevils were released into each cage. All adult weevils  
20 were collected seven days after the powder coated weevils were released into  
21 the cages. Collected weevils were scored for the presence of fluorescent  
22 powder, excluding those that were coated with powder at the start. There were  
23 eight replicate cages.

24



1 *Experiment 3: Susceptibility of adult vine weevils to EPF isolates in a laboratory*  
2 *bioassay*

3 The susceptibility of adult vine weevils to eight isolates of EPF was measured  
4 in a single dose laboratory bioassay. The isolates (Table 1) were selected  
5 based on their availability as commercial biopesticides and / or their virulence  
6 to vine weevil larvae reported in previous research (Moorhouse,1990). Conidia  
7 were grown as described previously, harvested from SDA plates in sterile  
8 0.05% Triton X-100 and filtered through sintered glass thimbles (40-100 µm  
9 pore). Conidia were then enumerated using an improved Neubauer  
10 haemocytometer and aliquots (10 ml) were prepared at a concentration of  $1 \times$   
11  $10^8$  conidia ml<sup>-1</sup>. Groups of five adult weevils were inoculated by immersion in  
12 suspensions of conidia for 10 seconds. Controls were treated with sterile 0.05%  
13 Triton X-100. Excess suspension was removed by filtration through filter paper  
14 under vacuum. The weevils were left to air dry on the filter paper for one hour,  
15 transferred to a ventilated plastic box as described previously, and maintained  
16 at 20°C, 16:8 light:dark with yew leaves and damp tissue (to maintain > 90%  
17 relative humidity) replaced *ad libitum*. Numbers of living and dead weevils were  
18 counted daily for 28 days. Dead weevils were removed and incubated on damp  
19 filter paper within Petri dishes at 23°C, and the production of fungal conidia on  
20 these cadavers was scored. The viabilities of conidia of the fungal isolates were  
21 measured following incubation for 24 h on SDA at 23°C (Goettel & Inglis, 1997).  
22 All isolates exhibited >87% germination. The experiment was done according  
23 to a block design. Each block comprised of the eight fungal isolates plus a  
24 control. There were three blocks in total, each done on a separate occasion.

25

1 *Experiment 4: Quantifying efficacy of the autodissemination technique in a*  
2 *laboratory bioassay*

3 The susceptibility of adult weevils to EPF applied as a conidial powder within  
4 the Roguard refuge was measured in a replicated laboratory bioassay. The  
5 isolates (*B. bassiana* isolate 433.99, *B. bassiana* 1749.11, *M. brunneum*  
6 275.86; see Table 1) were selected on the basis of their virulence to adult vine  
7 weevils in the previous experiment. Conidia were grown on SDA as described  
8 previously, harvested as a powder using a spatula, and the number of conidia  
9 per g of powder was calculated by counting conidia in suspensions (0.1g  
10 conidia in 10ml of 0.05% Triton X-100) using a haemocytometer. The viabilities  
11 of conidia of the fungal isolates were measured following incubation for 24 h on  
12 SDA at 23°C (Goettel & Inglis, 1997). All isolates exhibited > 91% germination.  
13 The conidia powders were added to Roguard refuges (0.4 g to each trap).  
14 Groups of five adult weevils were placed in ventilated plastic boxes as  
15 described previously but with the addition of a single, fungus treated Roguard  
16 refuge in each box. Boxes were maintained at 20°C, 16:8 light:dark with yew  
17 leaves and damp tissue (to maintain > 90% relative humidity) and numbers of  
18 living and dead weevils were counted daily for a total of 28 days. Dead weevils  
19 were removed and incubated on damp filter paper in Petri dishes at 23°C, and  
20 the production of fungal conidia on these cadavers was again scored as  
21 presence or absence. The experiment was done according to a randomised  
22 block design. Each block comprised three fungal isolates, with three blocks in  
23 total. Each block contained two control chambers (refuge containing talc).

24

1 *Experiment 5: Efficacy evaluation of M. brunneum applied against adult vine*  
2 *weevil under polytunnel conditions*

3 An experiment was done to evaluate the efficacy of *M. brunneum* 275.86  
4 against adult vine weevils when applied in Roguard refuges under polytunnel  
5 conditions. Treatments were established within gauze 'tent' cages (see  
6 Experiment 1) contained within a ventilated polytunnel. Twelve strawberry  
7 plants (cv. Malling Centenary) grown in 1.5 L pots using John Innes No. 2  
8 compost were placed in the centre of each cage. A conidia powder of *M.*  
9 *brunneum* 275.86 was prepared as described in Experiment 4. This was then  
10 added to a 50:50 (w/w) mixture of talc (Sigma, UK) and fluorescent powder (see  
11 Experiment 1) at a ratio of 0.3 g of conidia powder: 0.1g talc / fluorescent  
12 powder. Aliquots of 0.4g of this mixture were then placed in the central well of  
13 Roguard refuges. Mean conidia germination was 84% (SE = 2.69). Six refuges  
14 were placed in each cage equally distributed by placing between every other of  
15 the 12 plant pots. Controls consisted of 0.4 g of the talc / fluorescent powder  
16 mixture added to each Roguard refuge. There were five replicate cages.  
17 Groups of 40 adult vine weevils were placed into each cage on the foliage of  
18 the plants. The weevils were marked on their backs with bright yellow nail  
19 varnish before release so that they were easier to find in subsequent  
20 assessments. After five weeks, the numbers of dead and live adult weevils in  
21 each cage were counted, including the number of weevils coated in fluorescent  
22 powder. The presence of sporulating mycelia on weevil cadavers, visualised by  
23 incubating dead weevils on damp filter paper within Petri dishes at 23°C for  
24 approximately three weeks, was used as an indication of fungus-induced

1 mortality. Samples of powder were collected from refuge traps to evaluate  
2 conidia viability.

3

#### 4 *Analysis*

5 Data from experiments 3 and 4 were analysed using SPSS Statistics Version  
6 24.0 (IBM Corp., 2016). A Cox proportional-hazards regression model (Cox,  
7 1972) was used for analysing the time-mortality responses (i.e. survival) of  
8 weevils in all treatments compared to those in the control over 28 days. The  
9 Cox proportional hazard is expressed as the hazard ratio (relative average daily  
10 risk of death), which is assumed to remain constant over time. The event was  
11 death. Factors were replicate and treatment. The proportional cumulative  
12 survival of 50% of the population (i.e. median survival time (MST), of the weevil  
13 populations of each treatment and their 95% confidence intervals were  
14 calculated and pairwise comparisons were done using a log-rank  $\chi^2$  test  
15 (Bewick *et al.* 2004). Data from experiment 5 were analysed using a  
16 Generalised Linear Model (GLM) with a log link function and negative binomial  
17 error distribution for over dispersed count data (R-3.2.2, R Core Team, 2015).  
18 Wald tests were used to determine the significance of predictor variables.

19

## 20 **RESULTS**

21 *Experiment 1: Acquisition of fluorescent marker powders by adult weevils from*  
22 *artificial refuges*

23 Seven days after introducing the Roguard refuges containing fluorescent  
24 powder, a mean of 37 (range of 32 to 40) of the 40 adult vine weevils were  
25 recovered from each cage. Of these, a mean of 88% (range of 83 to 95%) of

1 the recovered weevils had come into contact with fluorescent powder. Weevils  
2 that had contacted the fluorescent powder were typically heavily coated in  
3 powder, with more than 50% of the body area covered.

4

5 *Experiment 2: Horizontal transmission of fluorescent powders among adult*  
6 *weevils*

7 Seven days after introduction of fluorescent powder-coated weevils to a  
8 recipient population, a mean of 33 (range of 30 to 35) of the 35 unmarked  
9 weevils were recorded from the cages. Of these, a mean of 75% (range of 66  
10 to 93%) of the recipient population had fluorescent powder on their cuticle.

11

12 *Experiment 3: Susceptibility of adult vine weevils to EPF isolates in a laboratory*  
13 *bioassay*

14 All of the EPF isolates caused significantly greater mortality of adult weevils  
15 than controls ( $P < 0.001$ ) (Table 2). After 28 days all except two (*B. bassiana*  
16 isolates 342.92 and 432.99) of the eight isolates tested resulted in 100%  
17 mortality of adult weevils. The median survival time (MST) of weevils treated  
18 with two of the isolates (*B. bassiana* isolates 433.99 and 1749.11) were  
19 significantly ( $P < 0.05$ ) less than the other isolates at 7 and 8 days respectively.

20 All of the isolates tested produced conidia on adult cadavers. The majority of  
21 sporulation occurred between the body segments and leg joints.

22

23 *Experiment 4: Quantifying efficacy of the autodissemination technique in a*  
24 *laboratory bioassay*

1 There was significantly ( $P < 0.001$ ) greater mortality of adult weevils in all of the  
2 treatments with refuges containing EPF than in the controls after 28 days (Table  
3 3). The refuges inoculated with isolates *M. brunneum* 275.86 and *B. bassiana*  
4 433.99 resulted in more than 66% weevil mortality after 28 days. The MST of  
5 weevils exposed to *M. brunneum* 275.86 and *B. bassiana* 433.99 inoculated  
6 refuges was 15 and 17 days respectively. Weevils had visible amounts of  
7 fungal conidia on their cuticles within four hours of starting the experiment and  
8 fungal conidia were seen to be carried out of the traps by weevils entering and  
9 leaving. *Beauveria bassiana* 1749.11 was not as effective in the refuges as had  
10 been expected, based on the data from the previous experiment. There was  
11 little evidence that weevils visited refuges containing this isolate, as indicated  
12 by the amount of weevil frass in the refuges and the amount of conidial powder  
13 on the floor of the bioassay chamber outside the refuge.

14

15 *Experiment 5: Efficacy evaluation of M. brunneum applied against adult vine*  
16 *weevil under polytunnel conditions*

17 Daily average temperatures and daily maximum temperatures in the polytunnel  
18 remained above 15°C during the experiment, while daily minimum  
19 temperatures fell below 15°C on 33 of the 40 days. Results from this  
20 experiment are summarised in Table 4. For both the control treatment and the  
21 *M. brunneum* treatment, over 95% of the weevils were recovered. Of the  
22 recovered weevils, mortality in the *M. brunneum* cages was significantly higher  
23 ( $z = 3.00$ ,  $P = 0.003$ ) than in the control cages. None of the dead weevils  
24 recovered from control cages were found to have become infected with *M.*  
25 *brunneum*, while 34% of the dead weevils recovered from the *M. brunneum*

1 treated cages had fungal mycelium emerging through the cuticle. Similarly  
2 numbers of recovered weevils had fluorescent powder on the cuticle in both the  
3 control and *M. brunneum* treated cages.

4

## 5 **DISCUSSION**

6 In this paper we present results from a series of laboratory and polytunnel  
7 experiments investigating the potential for using autodissemination of an EPF  
8 as a means of controlling adult vine weevil. It is proposed that simple plastic  
9 artificial refuges are effective inoculation chambers and that *B. bassiana* and  
10 *M. brunneum* isolates are effective against vine weevil adults.

11

12 Simple plastic crawling insect bait stations were readily used as refuges by vine  
13 weevil adults and in cage experiments there was effective dissemination of a  
14 hydrophobic fluorescent powder. This was apparent even when weevils had  
15 access to a range of refuges known to be exploited in crop habitats (e.g. Smith,  
16 1932; Moorhouse *et al.*, 1992). This effective dissemination of powders is likely  
17 to be due, at least in part, to the strong aggregation behaviour shown by vine  
18 weevil adults (e.g. Smith, 1932; Moorhouse *et al.*, 1992). Through this  
19 aggregation behaviour weevils are likely to come into contact with the  
20 fluorescent powder either by themselves entering one of the artificial refuges or  
21 by coming into contact with a weevil that has. Indeed, the experiment  
22 investigating the horizontal spread of the fluorescent powder shows that large  
23 numbers of weevils using artificial refuges may not be required for spores of an  
24 EPF to be spread through the weevil population. Further work is required to  
25 investigate the effect of refuge position and density on the spread of EPF

1 spores throughout weevil populations. Finally, there is considerable scope to  
2 optimise the design of artificial refuges. Olfactory lures based on sex (Hartfield  
3 *et al.*, 2001) and aggregation (Tinzaara *et al.*, 2007) pheromones as well as  
4 plant volatiles (Klein & Lacey (1999; Lyons *et al.*, 2012) have, for example,  
5 previously been used to promote the autodissemination of an EPF in damson-  
6 hop aphid and emerald ash borer populations respectively. For vine weevil, it  
7 is already known that responses of weevils to refuges may be enhanced  
8 through the addition of plant volatiles such as (Z)-2-pentenol and methyl  
9 eugenol (van Tol *et al.*, 2012).

10

11 Several studies have shown *M. brunneum* to be an effective control of vine  
12 weevil larvae (e.g. Bruck & Donahue, 2007; Moorhouse *et al.*, 1993; Shah *et*  
13 *al.*, 2007). In contrast, few studies have investigated the potential of *B. bassiana*  
14 for control of vine weevil (e.g. Prado, 1980; Bruck, 2004) or the use of EPFs to  
15 control adults. Moorhouse (1990) does, however, report an LT<sub>50</sub> of 13 days for  
16 *M. brunneum* isolate 275.86 when weevil adults were maintained at 20°C. In  
17 Experiment 3, we tested the same isolate under the same set of conditions as  
18 Moorhouse (1990) and recorded a similar, although slightly faster, LT<sub>50</sub>. The  
19 LT<sub>50</sub> for isolate *M. brunneum* 275.86 was comparable to the other *Metarhizium*  
20 isolates tested but two *B. bassiana* isolates, 433.99 and 1749.11, killed 50%  
21 and 90% of the weevil population significantly (P<0.05) faster than the other  
22 isolates tested.

23

24 In Experiment 4, conidia powders of *M. brunneum* isolate 275.86 and *B.*  
25 *bassiana* isolate 433.99 placed within refuges significantly increased weevil



1 mortality under laboratory conditions. However, *B. bassiana* isolate 1749.11  
2 had no effect on weevil mortality, despite this isolate being virulent to vine  
3 weevil adults when applied as a conidia suspension in the bioassay for  
4 Experiment 3. There was little evidence (e.g. frass inside the refuge or  
5 disturbance of the conidia powder) of weevils entering refuges containing a  
6 conidia powder of this isolate. It is, therefore, possible that weevils avoided  
7 conidia of isolate 1749.11. Insect avoidance of pathogenic fungi has been  
8 described in several other systems e.g. the anthocorid bug, *Anthocoris*  
9 *nemorum*, and the ladybird, *Coccinella septempunctata*, avoiding isolates of *B.*  
10 *bassiana* (Meyling and Pell, 2006; Ormond *et al.*, 2011) as well as the termite,  
11 *Macrotermes michaelseni*, avoiding both *B. bassiana* and *M. brunneum* (Mburu  
12 *et al.*, 2009). If this observation is true, it suggests the possibility of developing  
13 a fungus-based chemical repellent.

14

15 Under polytunnel conditions, conidia powders of *M. brunneum* isolate 275.86  
16 placed within refuges had a small but statistically significant effect on weevil  
17 mortality. Although not directly comparable, this result is similar to previous  
18 studies e.g. field testing autodissemination as a means of controlling emerald  
19 ash borer. In a previous study only 1% of beetles in the field site area were  
20 recorded as being infected with the EPF isolate placed in the inoculation  
21 chambers after a six week trapping period (Lyons *et al.*, 2012). However, as  
22 inoculation chambers may remain within crops throughout the season they are  
23 likely to have a cumulative effect on pest populations.

24

1 A feature of the results from the polytunnel experiment (Experiment 5) - in which  
2 we tested the efficacy of conidia powders of *M. brunneum* placed within refuges  
3 - was that far more weevils (144 weevils) came into contact with the conidial  
4 powder than subsequently died from infection due to this pathogen (26 weevils).  
5 This suggests that around 18% of weevils that came into contact with the  
6 conidial powders placed in the refuges acquired a lethal dose under these  
7 experimental conditions. This is much lower than results for the laboratory  
8 condition, which resulted in over 66% weevil mortality after 28 days. This may  
9 reflect the importance of temperature in determining the efficacy of *M.*  
10 *brunneum* in control of vine weevil (Bruck, 2007), although the time between  
11 infection and the end of this experiment was unknown<sup>[JB1]</sup>. A minimum  
12 temperature of around 15°C is required for effective control of larvae and a  
13 similar temperature requirement is likely to apply for control of adults. However,  
14 in the present study temperatures fluctuated between below 15°C at night and  
15 above 20 or 25°C during the day in the polytunnel experiment. How these  
16 temperature fluctuations affected the efficacy of *M. brunneum* is not known.  
17 However, previous studies of the performance of *B. bassiana* as a biopesticide  
18 against the kissing bug, *Rhodnius prolixus*, have shown that conditions  
19 fluctuating between unfavourable high and low temperatures resulted in a  
20 requirement for 10-20 times more inoculum in order to maintain efficacy  
21 (Fargues & Luz, 2000). This may, at least in part, explain the lower levels of  
22 mortality seen in the polytunnel experiment using *M. brunneum* than in the  
23 laboratory experiments.  
24

1 Autodissemination of an EPF through the use of artificial refuges as inoculation  
2 chambers offers promise for controlling vine weevil as a component of an IPM  
3 programme. For example, EPF targeted against adult weevils could be  
4 deployed alongside entomopathogenic nematodes and fungi used against vine  
5 weevil larvae. This might reduce the need for the use of broad spectrum  
6 chemical insecticides to control vine weevil adults (van Tol *et al.*, 2012), which  
7 may disrupt biological control programmes for other pests.

8

### 9 **ACKNOWLEDGEMENTS**

10 This study was funded by the Chemicals Regulations Directorate (CRD) on  
11 behalf of Defra (project PS2134).

12

### 13 **REFERENCES**

14 Ansari MA, Shah FA, Butt TM (2008) Combined use of entomopathogenic  
15 nematodes and *Metarhizium anisopliae* as a new approach for black vine  
16 weevil, *Otiorhynchus sulcatus* (Coleoptera: Curculionidae) control.  
17 *Entomologia Experimentalis et Applicata*. 129: 340–347.

18 Bewick V, Cheek L, Ball J (2004) Statistics review 12: Survival analysis.  
19 *Critical Care*, 8:389-394.

20 Bruck D. (2004) Natural occurrence of entomopathogens in Pacific Northwest  
21 nursery soils and their virulence to the black vine weevil, *Otiorhynchus*  
22 *sulcatus* (F.) (Coleoptera : Curculionidae). *Environmental Entomology*.  
23 33:1335-1343.

- 1 Bruck DJ. (2007) Efficacy of *Metarhizium anisopliae* as a curative application  
2 for Black Vine Weevil (*Otiorhynchus sulcatus*) infesting container-grown  
3 nursery crops. *Journal of Environmental Horticulture*. 25:150–156.
- 4 Bruck DJ, Donahue KM (2007) Persistence of *Metarhizium anisopliae*  
5 incorporated into soilless potting media for control of the black vine weevil,  
6 *Otiorhynchus sulcatus* in container-grown ornamentals. *Journal of*  
7 *Invertebrate Pathology*. 95:146-150.
- 8 Chandler D (1994) Cryopreservation of fungal spores using porous beads.  
9 *Mycological Research*. 98:525-526.
- 10 Cox DR (1972) Regression models and life-tables. *Journal of the Royal*  
11 *Statistical Society B*. 34:187-220.
- 12 Cross JV, Easterbrook MA, Crook AM, Fitzgerald JD, Innocenzi PJ, Jay CN,  
13 Solomon MG (2001) Review: natural enemies and biocontrol of pests of  
14 strawberry in northern and central Europe. *Biocontrol Science and*  
15 *Technology*. 22:165-216.
- 16 Fargues J, Luz C (2000) Effects of fluctuating moisture and temperature  
17 regimes on the infection potential of *Beauveria bassiana* for *Rhodnius*  
18 *prolixus*. *Journal of Invertebrate Pathology*. 75:202-211.
- 19 Georgis R, Koppenhöfer AM, Lacey LA, Bélair G, Duncan LW, Grewal PS,  
20 Samish M, Tan L, Torr P, van Tol RWHM (2006) Successes and failures in  
21 the use of parasitic nematodes for pest control. *Biological Control*. 38:103-  
22 123.

- 1 Goettel MS, Inglis GD (1997) Ch. V-3 Fungi: Hyphomycetes. *In* "Manual of  
2 Techniques in Insect Pathology", (L. A. Lacey, Ed.), pp.213-248, Academic  
3 Press.
- 4 Hartfield CM, Campbell CAM, Hardie J, Pickett JA, Wadhams LJ (2001)  
5 Pheromone traps for the dissemination of an entomopathogen by the  
6 damson-hop aphid *Phorodon humuli*. *Biocontrol Science and Technology*.  
7 11:401-410.
- 8 Klein MG, Lacey LA (1999) An attractant trap for autodissemination of  
9 entomopathogenic fungi into populations of the Japanese beetle *Popillia*  
10 *japonica* (Coleoptera:Scarabaeidae). *Biocontrol Science and Technology*.  
11 9:151–158.
- 12 Lundmark M (2010) *Otiorhynchus sulcatus*, an autopolyploid general-purpose  
13 genotype species? *Hereditas*. 147:278–282.
- 14 Lyons DB, Lavallee R, George K-P, van Frankenhuyzen K, Johnny S, Guertin  
15 C, Francese JA, Jones GC, Blais M (2012) Towards the development of an  
16 autocontamination trap system to manage populations of emerald ash borer  
17 (Coleoptera: Buprestidae) with the native entomopathogenic fungus,  
18 *Beauveria bassiana*. *Journal of Economic Entomology*. 105:1929-1939.
- 19 Meyling N, Pell JK (2006) Detection and avoidance of an entomopathogenic  
20 fungus by a generalist insect predator. *Ecological Entomology*. 31:162–171.
- 21 Mburu DM, Ochola L, Maniania NK, Njagi PGN, Gitonga LM, Ndung'u MW,  
22 Wanjoya AK, Hassanali A (2009) Relationship between virulence and

- 1 repellency of entomopathogenic isolates of *Metarhizium anisopliae* and  
2 *Beauveria bassiana* to the termite *Macrotermes michaelseni*. *Journal of Insect*  
3 *Physiology*. 55:774–780.
- 4 Moorhouse ER (1990) The potential of the entomopathogenic fungus  
5 *Metarhizium anisopliae* as a microbial control agent of the black vine weevil  
6 *Otiorhynchus sulcatus*. *Phd Thesis, University of Bath*. 533pp.
- 7 Moorhouse ER, Charnley AK, Gillespie AT (1992) A review of the biology and  
8 control of the vine weevil, *Otiorhynchus sulcatus* (Coleoptera: Curculionidae).  
9 *Annals of Applied Biology*. 121:431-454.
- 10 Moorhouse ER, Gillespie AT, Charnley AK (1993) The development of  
11 *Otiorhynchus sulcatus* (Fabricius) (Coleoptera: Curculionidae) larvae on a  
12 range of ornamental pot-plant species and the potential for control using  
13 *Metarhizium anisopliae*. *Journal of Horticultural Science*. 68:627-635.
- 14 Ormond EL, Thomas APM, Pell JK, Freeman SN, Roy HE (2011) Avoidance  
15 of a generalist entomopathogenic fungus by the ladybird, *Coccinella*  
16 *septempunctata*. *FEMS Microbiology Ecology*. 77:229–237.
- 17 Prado E (1980) Control of black vine weevil larvae (*Otiorrhynchus sulcatus*)  
18 using the insect pathogenic fungi *Beauveria bassiana*, *Metarrhizium*  
19 *anisopliae* and *Metarrhizium flavoviride*. *Vaxtskyddsnotiser*. 44:160-167.
- 20 Quesada-Moraga E, Martin-Carballo I, Garrido-Jurado I, Santiago-Álvarez C  
21 (2008) Horizontal transmission of *Metarhizium anisopliae* among laboratory

1 populations of *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae).  
2 *Biological Control*. 47:115-124.

3 Shah FA, Ansari MA, Prasas M, Butt TM (2007) Evaluation of black vine  
4 weevil (*Otiorhynchus sulcatus*) control strategies using *Metarhizium*  
5 *anisopliae* with sublethal doses of insecticides in disparate horticultural  
6 growing media. *Biological Control*. 40:246-252.

7 Smith FF (1932) Biology and control of the black vine weevil. *U.S. Department*  
8 *of Agriculture Technical Bulletin*. 325.

9 Soper R (1978) Autodissemination of entomopathogens: fungi. In: Allen GE,  
10 Ignoffo CM, Jaques RP (eds) Future strategies in pest management systems.  
11 Proceedings of the National Science Foundation, United States Department of  
12 Agriculture and the University of Florida Workshop on Microbial Control of  
13 Insect Pests, Gainesville, FL, pp 63–65.

14 Tinzaara W, Gold CS, Dicke M, Van Huis A, Nankinga CM, Kagezi GH,  
15 Ragamaet RE (2007) The use of aggregation pheromone to enhance  
16 dissemination of *Beauveria bassiana* for the control of the banana weevil in  
17 Uganda. *Biocontrol Science and Technology*. 17:111–124.

18 van Tol RWHM, Visser JH, Sabelis MW (2004) Behavioural responses of the  
19 vine weevil, *Otiorhynchus sulcatus*, to semiochemicals from conspecific,  
20 *Otiorhynchus salicicola*, and host plants. *Entomologia Experimentalis et*  
21 *Applicata*. 110:145-150.

1 van Tol RWHM, Bruck DJ, Griepink FC, De Kogel WJ (2012) Field attraction  
2 of the vine weevil *Otiorhynchus sulcatus* to kairomones. *Journal of Economic*  
3 *Entomology*. 105:169-175.

4 Warner RE, Negley FB (1976) The genus *Otiorhynchus* in America north of  
5 Mexico (Coleoptera: Curculionidae). *Proceedings of the Entomological*  
6 *Society of Washington*. 78:240-262.

7 Willmott DM, Hart AJ, Long SJ, Edmondson RN, Richardson PN (2002) Use  
8 of a cold-active entomopathogenic nematode *Steinernema kraussei* to control  
9 overwintering larvae of the black vine weevil *Otiorhynchus sulcatus*  
10 (Coleoptera: Cucurlionidae) in outdoor strawberry plants. *Nematology*. 4:925-  
11 932.

12 Yasuda K. (1999) Auto-infection system for the sweet potato weevil, *Cylas*  
13 *formicarius* (Fabricius) (Coleoptera: Curculionidae) with entomopathogenic  
14 fungi, *Beauveria bassiana* using a modified sex pheromone trap in the field.  
15 *Applied Entomology and Zoology*. 34:501-505.

16

17

18

19

20

21



1

2

3 Table 1. Fungal isolates used in the initial screen.

4

5 Table 2. Survival analysis results of time-mortality responses of adult black  
6 vine weevil treated directly with EPF isolates 28 days post inoculation.

7

8 Table 3. Survival analysis results of time-mortality responses of adult black  
9 vine weevil to EPF inoculated Roguard refuges 28 days post inoculation.

10

11 Table 4. Results from efficacy evaluation of *M. brunneum* applied against  
12 adult vine weevil under polytunnel conditions.

13

14

Isolate <sup>†</sup>	Species	Host/Substrate	Collection site
342.92	<i>Beauveria bassiana</i>	<i>Otiorhynchus sulcatus</i>	UK
432.99 <sup>a</sup> (ATCC 74040)	<i>B. bassiana</i>	<i>Anthonomus grandis</i>	USA
433.99 <sup>b</sup> (strain GHA)	<i>B. bassiana</i>	<i>Diabrotica undecimpunctata</i>	USA
1749.11	<i>B. bassiana</i>	<i>O. sulcatus</i>	UK
35.79	<i>Metarhizium anisopliae s.l.</i>	<i>O. sulcatus</i>	UK
189.83	<i>M. anisopliae s.l.</i>	<i>O. sulcatus</i>	UK
276.86	<i>M. anisopliae s.l.</i>	<i>O. sulcatus</i>	Germany
275.86 <sup>c</sup> (F52 / BIPESCO5)	<i>Metarhizium brunneum</i>	<i>Cydia pomonella</i>	Germany

1 <sup>†</sup>Isolate number in the WCC culture collection (isolate number from culture collection of origin)

2 (a) Isolate forms the active ingredient in the proprietary mycopesticide 'Naturalis' (Troy Biosciences Inc., 113  
3 South 47<sup>th</sup> Ave., Phoenix, AZ 850433, USA). ATCC = American Type Culture Collection

4 (b) Isolate forms the active ingredient in the proprietary mycopesticide 'Botanigard' (Mycotech Corporation, PO  
5 Box 4109, Butte, MT 59702, USA).

6 (c) Isolate forms the active ingredient in the proprietary mycopesticide 'Met52' (Novozymes, Hallas Allé 4400  
7 Kalundborg, Denmark).

8

9

10

11

12

13

14

1

2

Isolate †	Species	% mortality		Factors	MST <sup>b</sup> (95% CI)	HR <sup>c</sup> (95% CI)	Z (HR)	P (HR)	df	n	
		14 dpi <sup>a</sup>	28 dpi								
				Rep			0.002	0.961	1	2	
				Treatment			72.86	<0.001	8	9	
Control		3	13		-	a				30	
342.92	<i>Beauveria bassiana</i>	47	67		14 (3.9 - 24.1)	bd	10.45 (3.31 - 32.98)	16.02	<0.001	1	15
432.99	<i>Beauveria bassiana</i>	47	67		22 (1.8 - 42.2)	b	8.74 (2.73 - 27.96)	13.35	<0.001	1	15
433.99	<i>Beauveria bassiana</i>	100	100		7 (6.4 - 7.6)	e	104.74 (31.42 - 349.09)	57.35	<0.001	1	15
1749.11	<i>Beauveria bassiana</i>	100	100		8 (6.8 - 9.2)	e	79.15 (23.83 - 262.89)	50.95	<0.001	1	15
35.79	<i>Metarhizium anisopliae</i>	80	100		12 (10.1 - 13.9)	c	32.01 (9.95 - 103.03)	33.78	<0.001	1	15
189.83	<i>Metarhizium anisopliae</i>	67	100		10 (8.8 - 11.2)	c	27.54 (8.74 - 86.79)	32.05	<0.001	1	15
276.86	<i>Metarhizium anisopliae</i>	60	100		13 (11.8 - 14.2)	cd	23.29 (7.33 - 73.99)	28.49	<0.001	1	15
275.86	<i>Metarhizium brunneum</i>	87	100		10 (9.1 - 10.9)	c	31.54 (9.78 - 101.76)	33.36	<0.001	1	15

†Isolate number in the WCC culture collection

The Hazard ratios (HR) indicate the relative average daily risk of death compared to the 0.05% Triton-X treated control. The median survival time (MST) gives the proportional cumulative survival of 50% of the populations. MST values followed by different lower case letters within the column are significantly different (log rank  $\chi^2 \geq 3.841$ ,  $P < 0.05$ ).

a dpi = days post inoculation

<sup>b</sup> MST = median survival time, given in days

<sup>c</sup>HR = hazard ratio, compared to the 0.05% Triton-X treated control

1

2



Isolate <sup>†</sup>	Species	Factors	MST <sup>a</sup> (95% CI)	HR <sup>b</sup> (95% CI)	Z (HR)	P (HR)	df	n
		Rep			0.161	0.689	1	2
		Treatment			35.82	<0.001	3	4
Control			-	a				30
433.99	<i>Beauveria bassiana</i>		17 (15.5 - 18.5)	17.74 (4.474 - 70.322)	16.74	<0.001	1	15
1749.11	<i>Beauveria bassiana</i>		-	a	3.81(0.845 - 17.170)	<0.082	1	15
275.86	<i>Metarhizium brunneum</i>		15 (8.8 - 21.2)	34.3 (9.469 - 124.262)	28.97	<0.001	1	15

1 <sup>†</sup>Isolate number in the WCC culture collection

2 The Hazard ratios (HR) indicate the relative average daily risk of death compared to the 0.05% Triton-X treated control. The median survival time (MST) gives the proportional cumulative survival of  
3 50% of the populations. MST values followed by different lower case letters within the column are significantly different (log rank  $\chi^2 \geq 3.841$ , P < 0.05).

4 <sup>a</sup> MST = median survival time, given in days

5 <sup>b</sup>HR = hazard ratio, compared to the talc control

6

7

8

9

10

11

	<b>Control treatment</b>	<b><i>Metarhizium brunneum</i> treatment</b>
Mean number of weevils covered (+/- SE)	38.20 (1.06)	37.80 (0.86)
Mean number of dead weevils recovered (+/- SE)	10.00 (2.10)	15.40 (1.99)
Mean numbers of <i>M. brunneum</i> infected weevils (+/- SE)	0.00 (0.00)	5.20 (1.39)
Mean numbers of weevils coated in fluorescent powder (+/- SE)	27.60 (2.32)	29.00 (2.02)

1