

Original citation:

Taylor, Celia, Nhlema , B., Wroe , E., Aron , M., Makungwa, H. and Dunbar , E. L.. (2018) Determining whether community health workers are “deployment ready” using standard setting. *Annals of Global Health*.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/98811>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher’s statement:

© 2018, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher’s version. Please see the ‘permanent WRAP URL’ above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

**Determining whether Community Health Workers are “deployment ready” using standard
setting**

Celia Taylor¹, BSocSci, PhD: celia.taylor@warwick.ac.uk*

Basimenye Nhlema², BA, MADA: bnhlema@pih.org

Emily Wroe^{2,3,4}, MD, MPH: ewroe@pih.org

Moses Aron², BSc: maron@pih.org

Henry Makungwa², HDip: hmakungwa@pih.org

Elizabeth L Dunbar², MPH: edunbar@pih.org

1. Division of Health Sciences, University of Warwick, UK
2. Partners In Health, PO Box 56, Neno, Malawi
3. Division of Global Health Equity, Department of Medicine, Brigham & Women’s Hospital,
Boston, MA, USA
4. Department of Global Health & Social Medicine, Harvard Medical School, Boston, MA, USA

* Corresponding author

Competing interests

It is possible that PIH have a conflict of interest, in that the standard set determines the number of CHWs who can be immediately deployed and the number requiring additional training. However, this issue would arise regardless of whether or not the research study was being undertaken. Furthermore, PIH adhere to principles of safe clinical practice and ensuring that their CHWs provide safe care (at the absolute minimum) is clearly critical for their reputation.

Funding

CT's visit to Neno to run the standard setting process was funded by PIH.

CT is supported by the NIHR CLAHRC West Midlands initiative. This paper presents independent research and the views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Authors' contributions

EW and ED conceived the study which CT designed. All authors contributed to test design. CT ran the standard setting process. DH and MA managed test use in the field. MA managed the test data; all authors had access to test and standard setting data. CT analysed the data and drafted the manuscript and all other authors contributed to previous drafts and subsequently read and approved the final manuscript.

Article type

Original research: Global Health Education

Key words

Community Health Workers, Assessment, Standard Setting, Ebel

Running head

Are Community Health Workers "deployment ready"?

Abstract

Background

Community Health Workers (CHWs) provide basic health screening and advice to members of their own communities. Although CHWs are trained, no CHW programmes have used a formal method to identify the level of achievement on post-training assessments that distinguishes “safe” from “unsafe”.

Objectives

The aim of this study was to use Ebel method of standard setting for a post-training written knowledge assessment for CHWs in Neno, Malawi.

Methods

12 participants agreed definitions of a “just-deployment ready” and an “ideal” CHW. Participants rated the importance and difficulty of each question on a three point scale and also indicated the proportion of “just-deployment ready” CHWs expected to answer each of the nine question types correctly. Mean scores were used to determine the passing standard, which was reduced by one standard error of measurement (SEM) as this was the first time any passing standard had been employed.

The level of agreement across participants’ ratings of importance and difficulty was calculated using Krippendorff’s alpha. The assessment results from the first cohort of CHW trainees were analysed using classical test theory.

Findings

There was poor agreement between participants on item ratings of both importance and difficulty (Krippendorff's alphas of 0.064 and 0.074 respectively). The pass mark applied to the assessment, following adjustment using the SEM, was 53.3%. Based on this pass mark, 68% of CHW trainees were 'clear passes', 11% 'borderline passes', 9% 'borderline fails' and 12% 'clear fails'.

Conclusions

Determining whether a CHW is deployment-ready is an important, but difficult exercise, as evidenced by a lack of agreement regarding question importance and difficulty. Future exercises should allow more time for training, discussion and modification of ratings. Based on the assessment, most CHWs trained could be considered deployment-ready but following-up their performance in the field will be vital to validate the pass mark set.

Introduction

Community Health Workers (CHWs) provide a wide range of services including basic health promotion, counselling and care within the communities in which they live, providing a critical link between communities and health systems^{1,2}. CHWs are often seen as part of the solution to the shortage of health workers and lack of universal access to healthcare in low- and middle-income settings^{3,4}. While there is accumulating evidence of the effectiveness of CHW programmes, particularly for single disease areas such as screening for tuberculosis, adherence and retention for patients with HIV, or increasing access to maternal health services⁵⁻⁹, CHW programmes are not always as effective as they could be^{10,11}. There are a number of potential reasons for sub-optimal effectiveness, including insufficient or poor quality training^{10,11}, or because CHWs have difficulties in coping with the volume and complexity of the tasks asked of them¹². Taken to its extreme, and in common with the risks associated with receiving care from any healthcare worker, it is plausible that CHWs cause unintended but preventable harm to their clients, particularly if they are not coherently included in the broader health system¹⁰.

A CHW's initial training should prepare them for their role in service delivery and ensure they have the skills required to provide safe, effective and quality care¹³. Post-training assessments of CHWs' knowledge and skills are recommended to evaluate CHWs' learning during such training¹⁴. However, to determine whether a CHW is ready to be deployed into the field (often to work by themselves), a programme provider needs to know what minimum score is indicative of 'minimum competence' or readiness for safe deployment. The need for such a minimum score, or pass mark, can be implied from the studies of Workman et al.¹⁵ and Kalyango et al.¹⁶. Workman et al.¹⁵ report that CHWs in Brazil achieved a mean score of 41% on an assessment of knowledge about childhood cancer, noting that "without adequate training in and knowledge of the warning signs and symptoms of childhood cancer, it is unlikely that ... CHWs can ... thereby begin to decrease the discrepancies in terms of survival rates for childhood cancer" (p. 183-4). However we do not know what score would have

been required to reverse this conclusion. Kalyango et al.¹⁶, meanwhile, report a similar average knowledge score for pneumonia (median 40%) for CHWs in Uganda, but do not conclude that this is an insufficient level of knowledge for deployment. Anecdotal evidence suggests that post-training assessments appear fairly common, but we were only able to find three studies in which a pre-determined pass mark for a post-training assessment of knowledge was reported¹⁷⁻¹⁹. In two of these¹⁷⁻¹⁹ it is not clear how the pass mark was determined, while in the third²⁰ the authors report that their 50% cut-off to determine competency for their written assessment of maternal, neonatal and child health was arbitrary and not “scientifically driven”. Furthermore, Abrahams-Gessel reports that the initial pass mark of 70% for the knowledge test on cardiovascular risk had to be reduced to 60% because of a very low pass rate amongst the first cohort of CHWs to take the assessment and that there was no agreement between supervisors on “the level of competency that was expected to be achieved” (¹⁸, p. 133). A 70% pass mark was also used in the study of Accredited Social Health Activists (ASHAs) in India with the assessment using true-false style questions, with over 20% of ASHAs failing to meet this standard¹⁹. The authors of this study also emphasise the potential risk to client safety if “valuable information is provided in error or omitted [during] counselling” because “many ASHAs lack the essential knowledge to perform their jobs well” (¹⁹, p. 27-8).

In other healthcare professions such as medicine, attempts are made to establish appropriate pass marks for assessments using a process known as standard setting²¹⁻²³. There is no ‘gold standard’ method of standard setting, although the use of ‘absolute’ standards over fixed or relative standards is generally advocated²⁴. A fixed standard would be a pre-determined pass mark of say, 50%, without consideration of the difficulty of the assessment or whether this score is a good reflection of the standard required for safe practice. With a relative standard, a pre-determined percentage of candidates pass each sitting of the assessment, without consideration of differences in the ability level of different cohorts or whether those that pass are actually safe (and vice versa). With an absolute standard, a group of knowledgeable individuals determine, for each sitting of the assessment, the minimum score that reflects safe practice, with all candidates who achieve this

score passing the assessment. Absolute standards also help to ensure fairness across different sittings of an assessment with different content: a particularly hard exam paper would have a relatively low pass mark and vice versa. Two common methods of setting absolute standards for written assessments, Angoff²⁵ and Ebel²⁶, are described in Box 1.

As CHW programmes become more professionalised and CHWs take on more roles and responsibilities (with ensuing increased potential risks for their clients), it would seem appropriate to extend best-practice methods of standard setting used elsewhere to CHWs. Such an approach would help to maximise the health gains that can be achieved by CHWs. However, to our knowledge, no assessments used to evaluate CHWs' knowledge, skills or attitudes have been standard-set using established methods to date.

Box 1: Common methods of standard setting

Each method requires a panel of appropriately qualified individuals (for example, faculty involved with the design and delivery of teaching) to make decisions on individual test items or the test as a whole. Initial decisions are usually be made independently and the results then averaged across panel members or followed by discussion amongst panel members to agree relevant standards (often known as a 'Modified' approach). Research suggests that at least six panel members would be required to achieve sufficient reliability with discussion and ten without²⁷. For the Angoff and Ebel methods, the panel must begin by agreeing the definition of a 'minimally competent' candidate on the assessment.

Angoff²⁵: The panel provide the proportion of minimally competent candidates who would answer each item correctly. The mean standard across all items in the assessment provides the pass mark.

Ebel²⁶: The panel rate each item in two dimensions: (1) importance (e.g. essential, important, useful to know) and (2) difficulty (e.g. easy, moderate, challenging). This process creates a number of different categories of question (with the examples given there would be nine categories). They then

agree what proportion of minimally competent candidates would answer each category of item correctly and the relevant proportion applied to each item to provide its standard. The mean standard across all items in the assessment provides the pass mark.

Methods

Aims

The aims of this paper are:

1. To describe the application of the Ebel standard setting process used for a post-training written assessment of knowledge within a CHW programme,
2. To provide a short-term evaluation of the standard setting process, and
3. To review CHWs' performance on the first sitting of the standard set assessment and consider the implications of applying the pass mark set.

Study setting

The setting for this study is the rural district of Neno, which is in the south-west zone of Malawi on the border with Mozambique. Partners In Health (PIH), a US-based non-governmental organization, has worked with the Ministry of Health in Neno for the last ten years to strengthen primary and secondary healthcare services, with a strong focus on the community. Their initial work included a 'disease-specific' CHW programme, in which CHWs were assigned to individuals who had specific conditions - primarily HIV and tuberculosis. However, they have recently developed a new, integrated 'household model' or polyvalent approach to the CHW programme. In the new model, CHWs will undertake the following tasks:

1. Timely case finding through education and screening for common, treatable conditions.
2. Linkage to care for symptomatic clients along with those identified through routine screening.

3. Ongoing support and accompaniment of patients in care, including adherence support, psychosocial support, and tracking of missed patient visits.
4. Health education for common health conditions and prevention and management of these conditions to optimize prevention, health services uptake, and health management behaviours in the household.

The new model is being rolled-out across Neno over a two-year period. Once rollout is complete, there will be around 1,100 CHWs; the majority of these CHWs were employed in the disease-specific programme, but there will also be some new recruits. CHWs will attend monthly group meetings led by Senior CHWs (who in turn are supported and managed by Site Supervisors) and will also receive quarterly one-on-one mentorship and supervision visits from their Senior CHW.

As part of implementation of the new model, new and existing CHWs will attend a five day initial training course, which will be followed by shorter refresher training sessions. The purpose of the initial training is to orient CHWs to the following: the new programme and its objectives (specific disease areas), their cross-cutting roles, their specific tasks (as listed above), and data collection and documentation requirements. This will be achieved by developing the knowledge, skills, and attitudes CHWs need to successfully perform the tasks listed above. The training is structured around the key point of interaction between CHWs and their clients, monthly home visits. Following the initial training, CHWs will complete a written knowledge assessment, which has been developed based on a blueprint defined by CHWs' cross-cutting roles and disease areas (Figure 1). The assessment consists of 20 items (questions), including 'fill in the blank' with 1 or 3 answers required and single-best answer (1 from 3 or 5) items (Appendix 1). A total of 30 marks are available, with CHWs given 60 minutes to complete the assessment. The length of the test was determined pragmatically in relation to the length of the training course as a whole. The assessment was originally written in English, at which point all items underwent a quality review process using group discussion amongst the PIH development and implementation team. The assessment was

subsequently translated into Chichewa and back-translated to ensure accuracy to the original English version.

Describing the standard setting process

Standard setting process and panel: The standard setting process was undertaken in February 2017 as part of an orientation programme for those involved in training delivery (i.e. a training the trainer event). It was held at the PIH central office in the district of Neno. The standard setting panel was comprised of 12 individuals, including Malawi Ministry of Health/PIH staff involved in training delivery and PIH staff involved in development and implementation of the household model. All panel members have experience working with CHWs and of the requirements of the household model. All participants were given training about standard setting in general, the Ebel method and how it would work in practice.

The standard setting process followed good practice guidance²³ and used the Ebel method²⁶. The panel firstly discussed their expectations of an “ideal” and a “minimally competent” CHW, in terms of the knowledge required to be exceptional and just safe to deploy, respectively. Each panel member then rated each assessment item independently. Answers were provided to participants. Item importance was rated on a three-point scale: very important, important or nice-to-know. The difficulty of each item – for an ideal CHW - was also rated on a three-point scale: easy, moderate or challenging. For items which carried more than one mark, difficulty was rated for a CHW obtaining full marks. As a result, each panel member classified each item into one of nine possible categories (Table 1). Panel members then individually estimated the expected proportion of *minimally competent* (i.e. just deployment-ready) CHWs that *would* answer an item in each of the nine categories correctly in an assessment.

Determining the pass mark for the assessment:

The mean proportion of minimally competent CHWs who would answer each category of item correctly was calculated across all panel members. These means were applied to each panel member's item importance/difficulty category rating to give an item-level standard for each item for each panel member. The mean standard for each item across all panel members was used as the standard for that item. The weighted mean standard across all items was used as the initial pass mark for the assessment (termed the "Ebel pass mark"), with each item's standard weighted by the number of marks allocated to it.

The group discussed the possible consequences of a CHW failing the assessment. Although the assessment was intended to help ensure client safety (i.e. mitigate false positive test results), there was a concern that CHWs may fail because they are not used to assessments, struggle with written tests or may narrowly miss the pass mark but be safe to be deployed with additional supervision and support (i.e. might be false negatives). It was therefore decided to reduce the Ebel pass mark by one standard error of measurement (SEM) to obtain the pass mark that would be applied for this assessment in practice (termed the "pass mark applied"). The SEM of the assessment was calculated using Cronbach's alpha as the measure of reliability, based on the first cohort of CHW trainees to undertake the assessment (further explanation is provided in Box 2).

It was decided that CHWs scoring between the Ebel pass mark and the pass mark applied (i.e. borderline passes) would be required to meet with their supervisors more frequently for their first few months in practice. CHWs scoring below the pass mark applied would be asked to attend additional training and retake the assessment prior to being deployed.

Short-term evaluation of the standard setting process

Level of agreement between panel members:

The overall level of agreement between panel members for each dimension (importance and difficulty) was estimated using Krippendorff's alpha²⁸ in Stata v14 (see Box 2).

Relationship between item-level standards and candidate performance:

To evaluate item-level standards, the standard set for each item was compared to the mean score for the item across the first cohort of CHW trainees. Ideally, performance in only the sub-group of borderline CHWs would be considered, but the relatively small size of the cohort (and hence the low number of borderline CHWs) means that estimates of performance in this sub-group are likely to be imprecise. We would expect mean cohort performance on each item to exceed the standard set for that item (cf. being approximately equal to it for the sub-group of borderline CHWs), but not so high as to suggest the standard was too low. We therefore calculated the percentage of items where the mean score was between 10 and 30 percentage points higher than the standard (noting there is no empirical 'gold standard' for this range). We would also expect a positive correlation between standards and mean cohort scores across all items and we therefore calculated the Pearson correlation coefficient between item-level standards and mean cohort scores (see Box 2).

Candidate performance and implications of using the pass mark

We summarised the scores of the first cohort of CHW candidates taking the assessment. To consider the implications of using the pass mark applied, the proportion of CHWs in each of four score categories was calculated: clear fail (score < pass mark applied – 1 SEM), borderline fail (pass mark applied – 1 SEM < score < pass mark applied), borderline pass (pass mark applied < score < Ebel pass mark) and clear pass (score > Ebel pass mark).

Box 2: Explanation of statistical concepts

Cronbach's alpha: Measures the internal consistency of an assessment, or the extent to which the individual items are related with each other (and can, therefore, be used as a measure of reliability). The coefficient considers the correlations between candidates' scores on all possible pairs of items and can range from 0 to 1. A coefficient of 0 implies that there are no relationships between item

scores, while a coefficient of 1 implies that scores on one item can perfectly predict scores on all other items. Longer assessments tend to have higher Cronbach's alpha coefficients.

Krippendorff's alpha: Measures the inter-rater reliability of raters' judgements. The method can be used with any number of raters, ordinal data (e.g. the three point rating scales used in the Ebel process) and it can also handle missing data²⁸. The possible range of values is 0 to 1; a coefficient of 0 reflects that any agreement between raters is due to chance alone, while a coefficient of 1 reflects perfect agreement across all raters on all items.

Pearson's correlation coefficient: Measures the strength of the linear relationship between two independent, normally distributed variables. The possible range of values is -1 to +1; a coefficient of -1 reflects a perfect negative linear relationship, a coefficient of +1 a perfect positive linear relationship (for both, if plotted on a scatter diagram, all points would lie exactly on a straight line drawn through the points) and a coefficient of 0 no linear relationship (a random distribution of points on the scatter diagram).

Spearman Brown prophecy formula: Can be used to predict how changing the number of items in an assessment would change its reliability, based on the reliability achieved in the original assessment²⁹ and the general rule that reliability increases as test length increases.

Standard error of measurement (SEM): Estimates the precision with which an individual candidate's score has been measured by the assessment. If the candidate were to take the same assessment 100 more times with no learning effects, we would expect 68 of their scores to be within ± 1 SEM of their original score.

Results

Characteristics of the standard setting panel

The twelve-member standard setting panel consisted of nine members from PIH and three from the Malawian Ministry of Health (25%). One-third of panelists were female, and the mean age was 37 years. The median years' experience working in health in Neno District was 5.5, with a range of 1 year to 30 years. Five members were from the community team, three from the clinical team, and two from each of the environmental health and monitoring & evaluation teams. Regarding education, six had a diploma, four a Bachelors degree, and two a Masters degree.

Describing the standard setting process

Definitions of a “minimally competent” and an “ideal” CHW:

The standard setting panel discussed the expectations of minimally competent (just deployment-ready) and ideal CHWs in broad terms, rather than using specific applications of knowledge. Panel members found this a challenging task, and often cited skills and attributes that, whilst important, would be very difficult to assess in a written test. The broad expectations in relation to knowledge, used to define a minimally competent and an ideal CHW, are shown in Box 3.

Box 3: Broad definitions of a minimally competent and an ideal CHW

Minimally competent

Ideal

Has to refer back to their notes

Consistent knowledge base

Only confident in knowledge of one disease area

Breadth of knowledge

Cannot transfer knowledge to a new situation/

Adapts to change and new situations

Repeats word-for-word from notes

Makes preventable errors in documentation

Thorough

May not ask for help if unsure

Knows when and how to get help

Learns quickly

Determining the pass mark for the assessment:

Across all of the 20 items included in the assessment and all 12 panel members, 4.2% of item importance ratings were in the nice-to-know category, 27.7% were in the important category and 68.1% were in the very important category. These ratings suggest that the content of the items is appropriate for the CHW programme in Neno. Similarly, 59.9% of item difficulty ratings were in the easy category, 24.9% in the moderate category and 15.2% in the challenging category. These ratings suggest that the items are at about the right level for an assessment of deployment-readiness (rather than an assessment used to rank CHWs, which would require a higher proportion of challenging items). Both sets of ratings can be found for each standard setter in the supplementary data file.

Each panel member provided their own estimation of the number out of ten just deployment-ready CHWs who would answer each of the nine categories of item correctly in an assessment. The mean of these estimates, shown in Table 1, was used to convert each panel member's ratings of importance and difficulty into a numerical standard for each item. On average, the panel expected that only 12/100 minimally competent CHWs would answer a challenging, nice-to-know item correctly, but that 80/100 would do so for an easy, very important item.

Individual item standards, prior to weighting, ranged from 51.2% to 75.5% (standards for each item are shown in Appendix 1). Question #2 on the test "If a person tests positive for HIV, when should s/he start ARVs?" was most frequently rated as easy and as very important and therefore had the highest standard (per mark available, prior to adjustment with the SEM) at 0.75. No items were rated as only nice-to-know by more than three panel members. Question #10 on the test "What should a CHW do following detection of a missed period?" (with three answer options) was rated as challenging by six of the 12 panel members and therefore had a particularly low standard of 0.53.

The Ebel pass mark for the assessment, prior to adjustment using the SEM, was calculated as 61.6%. The SEM from the assessment undertaken by 129 CHWs in the first cohort was 8.1%, based on a Cronbach alpha coefficient of 0.78 and a standard deviation of CHWs' scores of 17.5%. Subtracting the SEM from the Ebel pass mark gave 53.4%, or 16.02/30 marks. It was decided to round this down so that the pass mark applied was 16/30 marks (53.3%).

Short-term evaluation of the standard setting process

Level of agreement between panel members:

The level of agreement between panel members for ratings of both item importance and difficulty was barely better than chance, with Krippendorff's alpha coefficients of 0.064 and 0.074 respectively. Had more time been available, the panel would have discussed their ratings and been given the opportunity to modify them, as in a "Modified Ebel" standard setting process, which may have reduced the variability amongst panel members.

Relationship between item-level standards and candidate performance: The mean score of the cohort on an item was between 10-30 percentage points higher than the standard set for the item for 10/20 items (50%). The correlation between standards and mean cohort scores was poor (Pearson's $r=0.13$, $p=0.585$), as shown in Figure 2. One item was much easier than predicted by standard setters (top of Figure 2), which asked about the correct use of ready-to-use therapeutic food to treat malnourished children (Question #15). Two items were particularly challenging compared to the standards set for them (bottom right corner of Figure 2); these asked about CHWs' role in the mother and baby visit three days after birth (Question #11) and the main risk of a home birth (Question #19).

Candidate characteristics

Of the 129 CHWs taking the assessment, 99 (77%) were female. Age and education data were available for 122 CHWs. Of these, the mean age was 39 years (range 23 to 67 years). 54 (44%) had

not completed primary school, 28 (23%) had only completed primary school, 30 (25%) had completed junior secondary school (26 obtaining the Junior Certificate of Education) and the remaining 10 (8%) had completed senior secondary school and obtained the Malawi School Certificate of Education.

Candidate performance and implications of using the pass mark

CHWs' scores were calculated following a post-assessment item review process, in which the items with a low overall facility were checked to ensure papers had been correctly marked. Figure 3 shows a histogram of CHWs' scores on the assessment. 12% of the 129 CHWs were in the "clear fail" category, scoring less than 14 marks/30 and a further 9% were in the "borderline fail" category, scoring 14 or 15 marks/30. 11% were "borderline passes", scoring between 16 and 18 marks/30 and 68% were "clear passes", scoring at least 19 marks/30. Hence the overall pass rate was 79%, with 20% of CHWs considered "borderline", i.e. about whom there is some uncertainty as to whether they should have passed or failed.

Discussion

Summary

Although we have not reported a novel approach to standard setting (the Ebel method itself was first reported in 1954)²⁶, to our knowledge this is the first reported attempt to describe a formal standard setting process for a post-training assessment of CHWs' readiness for deployment (aim 1). Although the standard setting panel was of a reasonable size²⁷, there was very low agreement between panel members as to the importance and difficulty of the items included in the assessment (aim 2). As such, the passing standard itself was not very reliable: as also reported by Abrahams-Gessel¹⁸, there seems to be a lack of agreement amongst trainers and programme managers as to the expected level of competency. For nine of the 20 items, the mean cohort performance was below the standard set, suggesting that these items were harder in practice than rated by the panel

members (aim 2). The assessment itself had good internal consistency (Cronbach's alpha 0.78) given its relatively short length. A reliability of at least 0.8 is advocated as a minimum for 'high stakes' assessments³⁰, and the Spearman Brown prophecy formula (see Box 2) suggests that this could be achieved with around three additional items. One in five CHWs were in the borderline performance category (aim 3).

Practical implications

Additional training, supervision and/or performance monitoring will be provided for the borderline pass CHWs (a policy that will be continued as roll out of the household model occurs in other districts). All CHWs participate in quarterly refresher trainings and the topic areas for such training could be identified by using the aggregate results from the assessment. For example, the two most challenging items in the assessment related to peri- and post-natal health. We might expect those CHWs newly recruited for the household model to have scored below their established peers in the post-training assessment (data were not available to test this hypothesis). If so, then a longer training period (currently five days) might be required for new recruits, although these CHWs may "get up to speed" very quickly once in post. The standards and training curriculum for the items with large discrepancies between standards and performance should also be revisited if a similar pattern is seen across all CHW cohorts.

Limitations

We used the terms "minimally competent" and "just-deployment ready" as the threshold to distinguish pass from fail. In a previous study, Ariff et al.²⁰ defined competency as "having sufficient knowledge and skills to comply with predefined clinical standards". We asked panel members to construct their own definition although they found this challenging. It may also be the case that merely being "safe" may not be sufficient to justify the costs of CHW programmes, because CHWs are usually deployed in resource-constrained settings.

Ideally, more training for panel members would be provided and the process repeated. Such additional training could include facilitated discussion of a large number of practice items to help panel members reach a consensus on item difficulty and importance. The provision of example performance data so that the panel members had some prior knowledge of the type of responses that trainee CHWs might give to the items used would be very helpful for such training (as well as for the actual standard setting). In addition, including a facilitated discussion of all “live” items would have been helpful to reduce the variability amongst panel members²⁷.

Future work

We hope that this work can support others who wish to apply a similar formal approach to setting pass marks for post-training assessments. Similar approaches can also be used for practical or clinical assessments, including for competencies such as communication skills. This is particularly important as post-training assessments become more routinely recommended and utilized², and as policymakers discuss professionalizing cadres of CHWs.

Studying the on-the-job performance of borderline pass CHWs (and contrasting it with that of the highest-scoring CHWs) would also provide a better indication of the knowledge of a “just-deployment-ready” CHW (assuming that the assessment is a good predictor of on-the-job performance). Evaluating on-the-job performance will also help PIH to establish the “learning curve” i.e. the rate at which performance improves with experience.

A real test of the standard would require an evaluation of longer-term predictive validity, following-up all CHWs and independently rating their on-the-job performance. It would then be possible to determine whether the pass mark applied was a good predictor of safe practice (i.e. if all those passing were at least safe and vice versa). Clearly such a study is challenging to implement²³ and raises ethical issues, since it requires those who failed the assessment to be deployed to work with clients. However, the data required for such work could also be obtained from existing CHW

programmes that have used post-training assessments without pass marks and which collect on-the-job performance data.

Conclusions

As CHW programmes become more professionalized, ensuring that CHWs are at least safe to be deployed is an issue of increasing importance for CHW programme providers. This requires the use of post-training assessments that are blueprinted to the programme requirements, include well-written and quality-assessed items and where the borderline between pass and fail has been established using an appropriate method of standard setting. With this in mind, the study reported here outlines how programme providers could start to implement standard setting for their assessments. We found that standard setting was a challenging process for panel members. Obtaining reliable standards requires that panel members gain experience of standard setting and we would therefore advocate the introduction of pilot standard setting exercises at an early stage in CHW programme development. This would also have the advantage of increasing consistency amongst trainers and implementers as to the standards of knowledge and performance expected of CHWs and standard setting in practice can often help to identify problems with individual assessment items.

Acknowledgements

Katie Cundale from PIH assisted with the preparation of materials for the standard setting panel meeting. We would like to thank all panel members for contributing to the process.

List of Figures

Figure 1: Assessment blueprint

Figure 2: Relationship between standards set and item performance; Legend: The green diagonal line shows where the mean cohort score is 10 percentage points higher than the standard set and the red diagonal line 30 percentage points.

Figure 3: Histogram of CHWs' scores; Legend: Red line (14/30) - boundary between clear fail and borderline fail; Yellow line (16/30) – boundary between borderline fail and borderline pass; Green line (19/30) – boundary between borderline pass and clear pass. The lines are not equally spaced as they have been rounded to the nearest whole number.

List of abbreviations

CHW: Community Health Worker

SEM: Standard error of measurement

Ethical approval

This work was reviewed and approved by the Malawi Health Services Research Committee (Ref: 16/11/1694) and The University of Warwick's Biomedical and Scientific Research Ethics Committee (Ref: REGO-2016-1899). All panel members used a symbol on their standard setting score sheets rather than their names, so that all scores were recorded anonymously. All panel members provided informed consent prior to participation.

Appendix 1: Post-training assessment with item-level standards

Supplementary data file: Standard setters' ratings of importance and difficulty by item

References

1. World Health Organization. Strengthening the performance of community health workers in primary health care: report of a WHO Study Group [meeting held in Geneva from 2 to 9 December 1987]. 1989.
2. Bhutta ZA, Lassi ZS, Pariyo G, et al. *Global experience of community health workers for delivery of health related Millennium Development Goals: a systematic review, country case studies and recommendations for scaling up*. Geneva: Global Health Workforce Alliance; 2010.
3. Lehmann U, Sanders D. The state of the evidence on programmes, activities, costs and impact on health outcomes of using community health workers. *Geneva: World Health Organization*. 2007.
4. World Health Organization. *Global strategy on human resources for health: Workforce 2030*. Geneva: World Health Organization; 2016.
5. Perry HB, Zulliger R, Rogers MM. Community health workers in low-, middle-, and high-income countries. *Annu Rev Public Health*. 2014;35:399-421.
6. Rich ML, Miller AC, Niyigena P, et al. Excellent clinical outcomes and high retention in care among adults in a community-based HIV treatment program in rural Rwanda. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2012;59(3):e35-e42.
7. Torpey KE, Kabaso ME, Mutale LN, et al. Adherence support workers: a way to address human resource constraints in antiretroviral treatment programs in the public health setting in Zambia. *PLoS One*. 2008;3(5):e2204.
8. Marcos Y, Phelps BR, Bachman G. Community strategies that improve care and retention along the prevention of mother-to-child transmission of HIV cascade: a review. *Journal of the International AIDS Society*. 2012;15(4).

9. Satti H, Motsamai S, Chetane P, et al. Comprehensive approach to improving maternal health and achieving MDG 5: report from the mountains of Lesotho. *PLoS One*. 2012;7(8):e42700.
10. Kok M, Dieleman M, Taegtmeyer M, et al. Which intervention design factors influence performance of community health workers in low- and middle-income countries? *Health Policy Plan*. 2014:1-21.
11. Glenton C, Colvin CJ, Carlsen B, et al. Barriers and facilitators to the implementation of lay health worker programmes to improve access to maternal and child health: qualitative evidence synthesis. *Cochrane Database Syst Rev*. 2013;10.
12. Schneider H, Okello D, Lehmann U. The global pendulum swing towards community health workers in low-and middle-income countries. *Hum Resources Health*. 2016;14(1):65.
13. Crigler L, Hill K, Furth R, et al. *Community Health Worker Assessment and Improvement Matrix*. Bethesda, MD: USAID Health Care Improvement Project; 2013.
14. Amaya K, Alezuyo C, Bainomugisha B, et al. Community-based family planning best practices manual. 2011.
15. Workman GM, Ribeiro RC, Rai SN, et al. Pediatric cancer knowledge: assessment of knowledge of warning signs and symptoms for pediatric cancer among Brazilian community health workers. *Journal of Cancer Education*. 2007;22(3):181-185.
16. Kalyango JN, Rutebemberwa E, Alfvén T, et al. Performance of community health workers under integrated community case management of childhood illnesses in eastern Uganda. *Malaria Journal*. 2012;11(1):282.
17. Gaziano TA, Abrahams-Gessel S, Denman CA, et al. An assessment of community health workers' ability to screen for cardiovascular disease risk with a simple, non-invasive risk assessment instrument in Bangladesh, Guatemala, Mexico, and South Africa: an observational study. *The Lancet Global Health*. 2015;3(9):e556-e563.


18. Abrahams-Gessel S. *An examination of the ability of community health workers to effectively conduct community-based screening for cardiovascular disease in South Africa, Guatemala, and Mexico*, Boston University; 2016.
19. Bajpai N, Dholakia R. *Improving the performance of Accredited Social Health Activists in India: Working Paper No. 1*. Mumbai: Columbia University; 2011.
20. Ariff S, Soofi SB, Sadiq K, et al. Evaluation of health workforce competence in maternal and neonatal issues in public health sector of Pakistan: an assessment of their training needs. *BMC health services research*. 2010;10(1):319.
21. Cusimano MD. Standard setting in medical education. *Academic Medicine*. 1996;71(10):S112.
22. Bandaranayake RC. Setting and maintaining standards in multiple choice examinations: AMEE Guide No. 37. *Medical teacher*. 2008;30(9-10):836-845.
23. Cizek G, Bunch M. *Standard Setting: A guide to establishing and evaluating performance standards on tests*. Thousand Oaks, CA: Sage Publications; 2007.
24. General Medical Council. Theme 5: Developing and implementing curricula and assessments. 2016; <http://www.gmc-uk.org/education/27394.asp>. Accessed 25/04/2016.
25. Angoff W. Scales, norms and equivalent scores. In: Thorndike R, ed. *Educational Measurement*. Washington, DC: American Council on Education; 1971:508-600.
26. Ebel RL. Procedures for the analysis of classroom tests. *Educ Psychol Meas*. 1954;14:352-364.
27. Fowell S, Fewtrell R, McLaughlin P. Estimating the minimum number of judges required for test-centred standard setting on written assessments. do discussion and iteration have an influence? *Advances in health sciences education*. 2008;13(1):11-24.
28. Krippendorff K. *Computing Krippendorff's Alpha-Reliability (1-25-2011)*. Pennsylvania, PA: University of Pennsylvania; 2011.


29. Wainer H, Thissen D. True score theory: the traditional method. In: Wainer H, Thissen D, eds. *Test Scoring*. Mahwah, NJ: Lawrence Erlbaum; 2001.
30. Roberts C, Newble D, Jolly B, et al. Assuring the quality of high-stakes undergraduate assessments of clinical competence. *Medical teacher*. 2006;28(6):535-543.

Table 1: Standards by item importance and difficulty (/10)

	Very important	Important	Nice-to-Know
Easy	8.0	6.5	4.6
Moderate	5.6	4.6	2.6
Challenging	3.5	2.8	1.2

		Cross-cutting knowledge & skills:						
		Approx. weight:	Process of providing care	Education	Monitoring and screening	Working with colleagues and supervisors	Knowledge of disease	Documentation
	Approx. weight:		25%	23%	18%	8%	15%	11%
Specific disease area:	HIV and STIs	15%						
	TB	18%						
	Family planning	6%						
	Maternal and neonatal health	19%						
	Child health and malnutrition	18%						
	Malaria	5%						
	Non-communicable disease	13%						
	General household health	6%						





Appendix 1: CHW Household Model assessment (with answers and Ebel item standards – prior to adjustment using the Standard Error of Measurement)

Name: _____

Date: _____

Catchment Area: _____

Male or Female

Village: _____

1. Which patients should be prioritized for monthly Home Visits? (1 mark: standard 0.66)

Patients who appear on the TRACE report.

2. If a person tests positive for HIV, when should he/she start ARVs? (1 mark: standard 0.75)

Before the end of 2 weeks.

3. List 3 symptoms of TB. (3 marks: standard 2.07)

1 mark for up to 3 of the following: Chest pain, cough for two weeks or more, fever, night sweats, lack of appetite, loss of weight, coughing up blood.

4. When do CHWs stop visiting a TB patient daily? (1 mark: standard 0.60)

- a. When she has completed 3 months of TB treatment.
- b. When she has completed 1 year of TB treatment.
- c. **When she has completed her TB treatment.**
- d. Never stop. CHWs visit TB patients daily, for life.

5. If a sick patient is referred to the health facility, when should the CHW visit them to follow-up and check on the patient? (1 mark: standard 0.69)

Within 2 days of the referral.

6. List 3 of the non-communicable diseases (NCDs) included in the new Household Model Program. (3 marks: standard 1.80)

1 mark for up to 3 of the following: Hypertension, heart failure, asthma, diabetes, epilepsy, mental illness.

7. What are the signs of malnutrition that you would look for when screening a child under 6 months for nutritional status? (3 marks: standard 1.59)

1 mark for up to 3 of the following: can see outline of ribs. Loose skin on upper arms, loose skin on thighs, can see shoulder bones, flesh missing from buttocks.

8. Until what age should you screen a child for nutritional status using the MUAC tape in the Household Model? (1 mark: standard 0.60)

Until they are 5 years old.

9. **If a woman of childbearing age is not on family planning, what should a CHW do?** (1 mark: standard 0.63)

- a. Give her a supply of contraceptive pills.
- b. Recommend the family planning method you think is most suitable for her.
- c. Refer her to the health provider for more information.**
- d. Teach her about all the family planning methods.

10. **What should a CHW do following detection of a missed period?** (1 mark: standard 0.53)

- a. Re-visit the woman in her home to undertake a pregnancy test
- b. Refer the woman to the health facility for a pregnancy test
- c. Accompany the woman to the health facility for a pregnancy test.**

11. **What should a CHW do when woman and baby are due for their 3rd day postnatal visit?** (1 mark: standard 0.64)

- a. Undertake the visit in the woman's home**
- b. Refer the woman to the health facility
- c. Accompany the woman to the health facility.

12. **What material can you use to educate household members about STI symptoms?** (1 mark: standard 0.64)

EITHER STI Counselling Card OR STI Job Card.

13. **When do Senior CHWs visit CHWs each month for the Supervision, Mentorship and Spot Check Visits?** (1 mark: standard 0.57)

In the 1st week of the month.

14. **Why would a CHW put an 'X' in the ANC column on the Household Register?** (1 mark: standard 0.69)

- a. The visit was cancelled.
- b. The woman attended that ANC visit.
- c. The woman does not have to go to that ANC visit.
- d. The woman missed that ANC visit.**

15. **How should food supplements/Ready-to-use therapeutic food (RUTF) be used to treat a child with malnutrition?** (1 mark: standard 0.59)

- a. Eaten directly as it is only by the malnourished child.**
- b. Eaten directly by all children in the household to prevent them from becoming malnourished.
- c. Mixed with porridge.
- d. Mixed with water and drunk.

16. **A child is confirmed as malnourished by the health facility. How is this indicated in the Household Register?** (1 mark: standard 0.57)

By putting a tick in the 'malnutrition' column against the child's name (must give BOTH the correct column and note the need to specify which child is affected – no half marks).

17. List 3 things that people can do to prevent hypertension. (3 marks: standard 1.54)

1 mark for up to 3 of the following: eat less salt, avoid alcohol, avoid/stop smoking, eat more vegetables, increase physical activity.

18. Which 3 groups of people are considered most vulnerable to malaria? (3 marks: standard 2.01)

1 mark for up to 3 of the following: Children, pregnant women, HIV+ patients (PLWHIV).

19. List 1 risk of a home birth you should educate a pregnant couple about. (1 mark: standard 0.62)

Any 1 of the following: HIV transmission between mother and child, both mother and child can die, mother can lose a lot of blood, both mother and child can get infection due to poor sanitation.

20. Which of the following sections does NOT need to be completed on a Referral and Accompaniment form? (1 mark: standard 0.67)

- a. Bottom half: patient details
- b. Bottom half: reason for referral and accompaniment
- c. Clinical section (section for the provider)**
- d. Top half: reason for referral/accompaniment.