

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/111771>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

© 2018 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>.



Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Manuscript Details

Manuscript number	REDOX_2018_955_R1
Title	PKG1 α oxidation negatively regulates food seeking behaviour and reward
Article type	Research Paper

Abstract

Genes that are highly conserved in food seeking behaviour, such as protein kinase G (PKG), are of interest because of their potential role in the global obesity epidemic. PKG1 α can be activated by binding of cyclic guanosine monophosphate (cGMP) or oxidant-induced interprotein disulfide bond formation between the two subunits of this homodimeric kinase. PKG1 α activation by cGMP plays a role in reward and addiction through its actions in the ventral tegmental area (VTA) of the brain. 'Redox dead' C42S PKG1 α knock-in (KI) mice, which are fully deficient in oxidant-induced disulfide-PKG1 α formation, display increased food seeking and reward behaviour compared to wild-type (WT) littermates. Rewarding monoamines such as dopamine, which are released during feeding, are metabolised by monoamine oxidase to generate hydrogen peroxide that was shown to mediate PKG1 α oxidation. Indeed, inhibition of monoamine oxidase, which prevents it producing hydrogen peroxide, attenuated PKG1 α oxidation and increased sucrose preference in WT, but not KI mice. The deficient reward phenotype of the KI mice was rescued by expressing WT kinase that can form the disulfide state in the VTA using an adeno-associated virus, consistent with PKG1 α oxidation providing a break on feeding behaviour. In conclusion, disulfide-PKG1 α in VTA neurons acts as a negative regulator of feeding and therefore may provide a novel therapeutic target for obesity.

Keywords	food seeking, protein kinase G, obesity
Taxonomy	Molecular Biology, Biochemistry
Corresponding Author	Philip Eaton
Order of Authors	Celine Duraffourd, Robert Huckstepp, Ingke Braren, Cathy Fernandes, Olivier Brock, alessio delogu, Oleksandra Prysyazhna, Joseph Burgoyne, Philip Eaton

Submission Files Included in this PDF

File Name [File Type]

letter accompanying submission of revision.docx [Response to Reviewers]

Duraffourd et al 2018_first revision text without figures.docx [Manuscript File]

Fig 1.tif [Figure]

Fig 2.tif [Figure]

Fig 3.tif [Figure]

Fig 4.tif [Figure]

Fig 5.tif [Figure]

Fig 6.tif [Figure]

Fig 7.tif [Figure]

Fig 8.tif [Figure]

Fig 9.tif [Figure]

Fig 10.tif [Figure]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Dear Dr. Levonen,

Please find a revised version of our manuscript for your consideration.

Although reviewers were very positive overall, Reviewer 2 asked about weight gain in the KI mice and specifically said we “*should at least discuss the issue or tone down their statement to put their findings into perspective*”. This is an important point, and guided by the reviewer’s comment, we have add a significant Discussion text on this issue give its importance. This new text reads,

We have also added molecular weight details to the relevant Figures as requested by reviwer 1. We hope this fully addresses the requirements of this revision, and so will make the manuscript suitable for publication.

Should you have any further requests of requirements, please get in contact.

Yours Sincerely,

Philip Eaton

Ref: REDOX_2018_955

Title: PKG1 α oxidation negatively regulates food seeking behaviour and reward

Journal: Redox Biology

Dear Dr. Eaton,

Thank you for submitting your manuscript to Redox Biology. We have completed the review of your manuscript. A summary is appended below. While revising the paper please consider the reviewers' comments carefully. We look forward to receiving your detailed response and your revised manuscript.

To submit your revised manuscript:

- Log into EVISE® at:
http://www.evise.com/evise/faces/pages/navigation/NavController.jsx?JRNL_ACR=REDOX
- Locate your manuscript under the header 'My Submissions that need Revisions' on your 'My Author Tasks' view
- Click on 'Agree to Revise'
- Make the required edits
- Click on 'Complete Submission' to approve

What happens next?

After approving your submission you will receive a notification that the submission is complete. To track the status of your paper throughout the editorial process, log into EVISE® at:

http://www.evise.com/evise/faces/pages/navigation/NavController.jsx?JRNL_ACR=REDOX

Enrich your article to present your research with maximum impact. This journal supports the following [Content Innovations](#):

I look forward to receiving your revised manuscript as soon as possible.

Kind regards,

Anna-Liisa Levonen
Editor
Redox Biology

Comments from the editors and reviewers:

-Reviewer 1

- The manuscript by Durauffourd and colleagues is timely and of interest. The authors define the role of PKG oxidation and upstream signalling involving food behaviour, which may be important in the control of obesity and associated diseases. The paper is very well written and easy to follow. The protocols are carefully planned and support the conclusions. I only have a minor request for the authors: please add molecular weights in all figures involving representative images for immunoblots.

-Reviewer 2

-

The manuscript by Durauffourd et al. examines the food seeking behavior of “Redox dead” C42S PKG1 alpha knock-in (KI) mice that are deficient in oxidant-induced disulfide-PKG1alpha-formation, in comparison of WT littermates. The authors show increased food seeking and reward behaviour in KI mice compared to WT, and that rewarding monoamines such as dopamine are metabolized by monoamine oxidase (MAO) to generate hydrogen peroxide mediating PKG1alpha oxidation. The deficient reward phenotype of the KI mice was rescued by expressing WT kinase using an adeno-associated virus vector. The authors suggest that disulfide-PKG1alpha acts as a negative regulator of feeding thereby providing a therapeutic target for obesity.

This is a well conducted study with solid results and the overall presentation is very clear. However, the authors have not really addressed whether the observed behavioral changes in KI mice result in changes in weight gain. Do the KI mice gain more weight upon feeding an obesogenic diet? This is an important question, as the authors state in the abstract that disulfide-PKG1alpha could be a potential new target for treating obesity. The authors should at least discuss the issue or tone down their statement to put their findings into perspective.

PKG1 α oxidation negatively regulates food seeking behaviour and reward

Celine Duraffourd¹, Robert T. R. Huckstepp², Ingke Braren³, Cathy Fernandes^{4,5}, Olivier Brock⁶, Alessio Delogu⁶, Oleksandra Prysyazhna¹, Joseph Burgoyne¹, Philip Eaton^{1*}

¹ King's College London, School of Cardiovascular Medicine & Sciences, the Rayne Institute, St. Thomas' Hospital, London, SE1 7EH.

² School of Life Sciences, University of Warwick, Coventry, United Kingdom.

³ University Medical Center Eppendorf, Vector Facility, Inst. for Exp. Pharmacology and Toxikology, N30, Room 09, Martinistr. 52, 20246 Hamburg, Germany

⁴ SGDP Research Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom

⁵MRC Centre for Neurodevelopmental Disorders, King's College London, London, United Kingdom

⁶ Basic and Clinical Neuroscience, Institute of Psychiatry, Psychology and Neuroscience, King's College London, United Kingdom

Running title: PKG1 α is a negative regulator of food-seeking behaviour and reward

Subject Areas: PKG1 α , food-seeking behaviour, reward, Ventral Tegmental Area

Total number of Pages: 30 main text (including title page and figures, Supplemental figures and methods),

Figures: 5 Figures
5 Supplemental figures

Total number of words: 4939 (including references and legends)
2876 (methods)

* Address for correspondence:
King's College London,
School of Cardiovascular Medicine & Sciences,
The Rayne Institute, St. Thomas' Hospital,
London, SE1 7EH. Email: philip.eaton@kcl.ac.uk

Abstract

Genes that are highly conserved in food seeking behaviour, such as protein kinase G (PKG), are of interest because of their potential role in the global obesity epidemic. PKG1 α can be activated by binding of cyclic guanosine monophosphate (cGMP) or oxidant-induced interprotein disulfide bond formation between the two subunits of this homodimeric kinase. PKG1 α activation by cGMP plays a role in reward and addiction through its actions in the ventral tegmental area (VTA) of the brain. 'Redox dead' C42S PKG1 α knock-in (KI) mice, which are fully deficient in oxidant-induced disulfide-PKG1 α formation, display increased food seeking and reward behaviour compared to wild-type (WT) littermates. Rewarding monoamines such as dopamine, which are released during feeding, are metabolised by monoamine oxidase to generate hydrogen peroxide that was shown to mediate PKG1 α oxidation. Indeed, inhibition of monoamine oxidase, which prevents it producing hydrogen peroxide, attenuated PKG1 α oxidation and increased sucrose preference in WT, but not KI mice. The deficient reward phenotype of the KI mice was rescued by expressing WT kinase that can form the disulfide state in the VTA using an adeno-associated virus, consistent with PKG1 α oxidation providing a break on feeding behaviour. In conclusion, disulfide-PKG1 α in VTA neurons acts as a negative regulator of feeding and therefore may provide a novel therapeutic target for obesity.

Introduction

Obesity is a major global health problem with 13% of the world's adult population being affected in 2016. The wide availability of highly-palatable and -rewarding food drives their preferential consumption, increasing the prevalence of obesity. Defining mechanisms that drive us to seek vast amounts of highly calorific food may allow the development of rational interventions to attenuate their consumption, potentially reducing obesity and its associated health risks.

Protein kinase G (PKG) is a phylogenetically well-conserved food-seeking and -reward behavioural gene across insects and nematodes [1]. Whilst PKG expression is seen in mammals [2-4], there appears to be no information as to its effect on feeding behaviour. The PKG1 α isoform is classically activated by binding of cyclic guanosine monophosphate (cGMP) [5]. However, under physiological conditions PKG1 α can also be regulated by oxidation [6-8]. Endogenous oxidants, such as hydrogen peroxide, induce a disulfide bond between cysteine 42 on adjacent chains of 2 PKG1 α monomers, forming a homodimer complex that renders the kinase catalytically active [6-8]. The binding of cGMP to PKG1 α attenuates oxidant-induced interprotein disulfide accumulation, perhaps because it causes conformational changes that move the two cysteine 42 residues away from each other [7, 9]. In conjunction, disulfide-PKG1 α dimerisation through oxidation appears to cause a conformational changes in PKG1 α that decreases cGMP-dependent activity, which could be due to oxidation lowering affinity for the cyclic nucleotide [2, 10]. Therefore, the two pathways appear to be antagonistic regulators of PKG1 α .

In the brain, the ventral tegmental area (VTA) represents a key component of the mesolimbic dopaminergic system and is critical to reward learning and addictive behaviours [11]. Interestingly, PKG regulates dopamine release [12, 13], and altering PKG levels within the VTA increases addiction [14]. Furthermore, alteration of PKG activity through manipulation of cGMP, or its activator nitric oxide synthase (NOS), leads to complementary changes in reward [15, 16]. As cGMP binding appears to increase reward and oxidation of PKG1 α negatively regulates binding of this cyclic nucleotide [12, 13], we speculated that disulfide-PKG1 α in the VTA would therefore suppress food reward. To investigate this hypothesis, we assessed food-seeking and -reward behaviour in C42S PKG1 α knock-in (KI) 'redox dead' mice, which are fully deficient in disulfide-dependent activation, whereas their NO-cGMP-dependent activation remains intact [17]. We found that removal of the disulfide-PKG1 α pathway led to an increase in food seeking behaviour in KI mice compared to wild-type (WT) littermates. Breakdown of the primary neurotransmitter of the reward system, dopamine, by monoamine oxidase-B (MAO-B) leads to production of hydrogen peroxide [18]. Inhibition of MAO-B reduced disulfide-PKG1 α in the VTA and increased sucrose preference in WT but not KI mice. In accordance, inhibition of NOS, which is anticipated to decrease cGMP-dependent activation of PKG1 α , concomitantly increased disulfide-PKG1 α and reduced sucrose preference in WT but not KI mice.

Critically, we could rescue the food-reward phenotype of PKG1 α KI mice by expression of WT PKG1 α in the VTA using an adeno-associated virus (AAV). Therefore disulfide- and cGMP-activation of PKG1 α work antagonistically in the VTA to control hedonic eating, where disulfide-PKG1 α in the VTA acts as a negative regulator, an 'off switch', for food-reward. Therefore, PKG1 α may provide a novel therapeutic target for treatment against sugary addictions.

Results

Disulfide-PKG1 α regulates food-seeking behaviour

We began by investigating food seeking behaviour of C42S PKG1 α KI mice (KI). There was no difference in locomotor activity between genotypes over a 24-hour period (WT: 4.8 ± 0.5 m/h, KI: 4.8 ± 0.7 ; $n = 12$ per group; Figure 1a). However, there was an increase in total distance covered at the beginning of the night, between 19:30 and 21:30, the first 2 hours of their dark period (WT: 16 ± 2 m/h, KI: 20 ± 3 ; $n = 12$ per group; $p = 0.005$; Figure 1a). There was no difference between males and females. Interestingly, there were no differences between KI mice and WT of either sex in overall food consumption ($n = 12$ or 13 per group; Figure 1aii), showing overall drive to feed is not altered by this mutation. Furthermore, there were no differences between KI and WT of either sex in overall weight ($n = 12$ or 13 per group; Figure 1aiii), meaning KI mice do not appear to have altered metabolism.

In a novelty-suppressed feeding test, there was no difference between WT or KI mice of either sex in their approach to food ($n = 12$ - 13 per group, Figure 1bi). However, male KI mice began to eat their food faster than male WT, although there was no difference between females of either genotype (male WT: 190.5 ± 40.6 s, male KI: 73.9 ± 13.9 , female WT: 150.1 ± 53.6 , female KI: 142.9 ± 43.0 ; $n = 12$ or 13 per group; $p = 0.04$; Figure 1bii).

Though we did not observe differences between KI and WT female mice in latency to eat, KI female mice presented a higher weight loss than WT female following 24 hours fasting prior to testing (male WT: 2.1 ± 0.2 g, male KI: 2 ± 0.2 , female WT: 1.8 ± 0.1 , female KI: 2.3 ± 0.2 ; $n = 12$ or 13 per group; $p = 0.006$; Figure 1biii). Consistent with this, female KIs increased their food intake the day following testing compared to WT littermates, possibly to compensate for their weight loss (male WT: 5.5 ± 0.3 g, male KI: 5.7 ± 0.3 , female WT: 5.2 ± 0.3 , female KI: 6.4 ± 0.3 ; $n = 12$ or 13 per group; $p = 0.02$; Figure 1biv). Interestingly, no difference were seen between KI males and their WT littermates in either weight loss or food consumption (Figure 1biii,iv)

Specific effect on feeding behaviour in KI mice

Whilst alterations in food behaviour are prominent in PKG KI mice, this could be due to other confounding factors. However, there were no differences between KI or WT of either sex for tests of: Anxiety, i.e., Open field test and elevated-plus maze (Figure 2); alterations in olfaction, i.e., olfaction habituation/dishabituation test (Figure 3); Exploratory behaviour, i.e., novel object recognition (Figure 4); Memory, i.e., Y-maze (Figure 5), and; general motor deficits, i.e., rotarod and grip and strength (Figure 6). Therefore, changes in feeding behaviours were due to a change in the drive to feed or in the reward gained by eating, and not caused by any other, behavioural deficits.

Disulfide-PKG1 α regulates rewarding behaviour

To distinguish between increased food seeking and reward we performed a sucrose preference test to compare hedonic sensitivity of KI or WT mice. Notably, KI mice had an increased preference for sucrose compared to WT but no differences were observed between males and females (male WT: $63.9 \pm 2\%$, male KI: 74.1 ± 2.1 , female WT: 65.7 ± 2.2 , female KI: 74.5 ± 1.5 ; $n = 12$ or 13 per group; $p = 0.0002$; Figure 1ci). Furthermore, there was no difference in side preference between genotypes of either sex, so excluding this as a

confounding factor (n = 12 or 13 per group; Figure 1cii). Therefore, our data indicate oxidation of PKG1 α to the interprotein disulfide state decreases desire to consume rewarding substances.

L-DOPA or dopamine induces disulfide-PKG1 α in the ventral tegmental area (VTA).

Using non-reducing western immunoblot analysis, we observed that disulfide-PKG1 α was increased in isolated VTA tissue blocks treated with dopamine (Cont: 1.1 ± 0.1 UA, dopamine: 3.8 ± 0.6 ; n = 7 per group; p = 0.01; Figure 7ai,ii). Pre-treatment with deprenyl, an inhibitor of monoamine oxidase B (MAO-B), attenuated formation of disulfide-PKG1 α dimers in response to dopamine in WT mice (dopamine: 3.8 ± 0.6 UA, deprenyl: 2.0 ± 0.1 ; n = 7 per group; p = 0.001; Figure 7ai,ii) to a level indistinguishable from controls (Cont: 1.1 ± 0.1 UA; n = 7; Figure 7a). Similarly, increased PKG1 α was observed in the brains of WT mice following an intraperitoneal injection of L-DOPA, a dopamine precursor (Cont: 1.1 ± 0.1 UA, L-DOPA: 4.6 ± 0.9 ; n = 6 per group; p = 0.0007; Figure 7bi,ii). Pre-treatment with deprenyl also prevented this *in vivo* formation of disulfide-PKG1 α in response to L-DOPA in WT mice (L-DOPA: 4.6 ± 0.9 UA, deprenyl: 2.3 ± 0.3 ; n = 6 per group; p = 0.0007; Figure 7b) to a level indistinguishable from controls (Cont: 1.1 ± 0.1 UA; n = 6 per group, Figure 7b). Auranofin, which pharmacologically inhibits thioredoxin reductase to attenuate reduction of disulfide bonds that form and therefore retain PKG1 α in the oxidised state for detection using western immunoblotting, did not lead to a change in disulfide- PKG1 α levels for either experiment (n = 7 per group, Figure 7). Therefore, there is not a significant amount of PKG1 α oxidation under basal conditions. Treatment with deprenyl or saline controls did not cause any change in disulfide-PKG1 α formation in WT mice in either experiment (n = 7, Figure 7a; n = 6, Figure 7b). Taken together, these results indicate dopamine increases disulfide-PKG1 α in the brain of WT mice.

Inhibition of MAO-B reduces disulfide-PKG1 α and increases sucrose preference in WT, but not KI mice

As one by-product of dopamine degradation by MAO-B is hydrogen peroxide [18], we investigated the effect of pharmacologically inhibiting MAO-B on disulfide-PKG1 α in WT mice. Pre-treatment with 0.5 mg/kg deprenyl increased sucrose preference in WT mice (64.0 ± 0.8 to $72 \pm 1.0\%$; n = 4; p < 0.0001; Figures 8ai), and reduced disulfide-PKG1 α (1.0 ± 0.2 to 0.4 ± 0 UA; n = 4 per group; p = 0.04; Figure 8aii,iii). Higher doses of deprenyl (2.5 or 5 mg/kg) did not increase sucrose preference beyond that of the lowest concentration (Figure 8ai). KI mice unable to form disulfide-PKG1 α (Figure 9b) did not show any alteration in sucrose preference following deprenyl administration (63.0 ± 1.1 to 66.3 ± 1.2 ; Figure 9a). Therefore, blocking degradation of dopamine that generates hydrogen peroxide as a by-product, increased disulfide-PKG1 α dimers and sucrose preference of WT mice.

Inhibition of NO synthase increases disulfide-PKG1 α and decreases sucrose preference in WT, but not KI mice

Nitric oxide (NO) binds to guanylyl cyclase to stimulate production of cGMP [19]. As cGMP attenuates disulfide-PKG1 α accumulation, elevating NO or stimulating guanylyl cyclase to increase cGMP should increase reward. Pre-treatment with an NO synthase inhibitor, L-NAME, decreased sucrose preference of WT mice (66.0 ± 1.2 to $59.2 \pm 1.3\%$; n = 8 per group; P < 0.0001; Figures 8bi), and increased disulfide-PKG1 α (1.0 ± 0.3 to 7 ± 1.3 AU; n = 8 per group; p = 0.004; Figure 8bii,iii). Higher doses of L-NAME (5 and 10 mg/kg) did not decrease sucrose preference beyond that of the lowest concentration (Figure 8bi). KI mice, unable to

form disulfide-PKG1 α (Figure 9b), did not show any alteration in sucrose preference following administration of L-NAME (Figure 9a). Therefore, blocking the elevation of cGMP, decreased sucrose preference in WT mice.

VTA injection of the WT PKG1 α rescues the behaviour phenotype of KI mice

As PKG levels within the VTA are directly associated with addiction [3, 14], and we found a deficit in food reward in KI mice or WT mice following pharmacological inhibition of NOS, we hypothesised that deficiency of disulfide-PKG1 α in the VTA leads to enhanced food reward in KI mice. To test this we used an AAV vector to re-express the wild-type form of the PKG1 α (WT PKG1 α) in the VTA of KI mice (Figure 10a,b).

Expression of WT PKG1 α in the VTA of KI mice reduced their sucrose preference compared with KI mice injected with a control virus expressing solely GFP (GFP: $75.8 \pm 1.4\%$, WT PKG: 62.4 ± 2.0 ; $n = 8$ per group; $p = 0.0007$; Figure 10ci). Importantly, sucrose preference in KI mice injected with WT PKG1 α was not different to that in WT mice ($64.8 \pm 1.5\%$; $n = 13$). Transducing the VTA of KI mice to express C42S KI PKG1 α confirmed that overexpressing PKG1 α in the VTA *per se* does not downregulate food reward ($75.2 \pm 1.7\%$; $n = 8$ per group, Figure 10ci). Overexpression of WT PKG1 α , C42S KI PKG1 α , or GFP did not lead to a change in side preference (Figure 10cii). This indicates disulfide-PKG1 α in the VTA is a key component in food reward, acting as a negative regulator that switches off the rewarding properties of sucrose in mice.

Discussion

Feeding behaviour is a complex decision-making process involving where, when, what, and how much to eat. This isn't only a reflexive response to stimuli or change in energy level, but includes environmental factors and previous experience. Obesity is characterized by excess storage of body fat, which is often associated with other diseases, such as type 2 diabetes, hypertension, heart disease, stroke, cancer and non-alcoholic fatty liver disease. The cost of obesity to the National Health Service in the United Kingdom has increased from an estimated £479.3 in 1998 to £6.1 billion 2014. Thus, identification of mechanisms underlying food preference behaviours, which lead to overeating, are now a priority.

PKG1 α is a cytosolic kinase that is a downstream target of cGMP [20]. PKG has perhaps been most extensively studied in the cardiovascular system [20], though it is found in many tissues including the brain [2]. PKG1 α was found to undergo oxidant-induced targeting and activation, forming a catalytically active dimer [4, 6, 7, 21, 22]. Interestingly, increased cGMP activation of PKG produced by sildenafil, an inhibitor of phosphodiesterase 5-dependent cGMP degradation, increases reward [15]. Furthermore, reduction of PKG activity through administration of phosphodiesterase type 5 inhibitor or by pharmacological inhibition or genetic knockout of NOS, which abrogates soluble guanylyl cyclase-dependent cGMP production, decreases reward [15, 16]. As the NOS-cGMP-PKG1 α pathway is highly conserved in both food seeking [1] and reward [14-16], and disulfide-PKG1 α appears to negatively regulate this pathway [9], we hypothesized regulation of the cGMP/PKG1 α pathway by disulfide-PKG1 α would act as an off switch of food-reward.

We found C42S KI 'redox dead' mice increased locomotor activity during the first 2 hours of their dark period when feeding is at its highest [23], which may indicate an increase in foraging behaviour. Furthermore, KI males began to eat faster than male WT littermates in the novelty suppressed feeding test, whilst female KIs ate more chow following food restriction than female WT littermates, with no differences in any other behavioural tests performed. Therefore, changes in feeding behaviours were due to an altered drive to feed or in reward gained by eating, and not because of: a lack of foraging or exploratory behaviour due to anxiety; general motor discoordination or fatigue; a disinterest in food caused by an inability to smell, or; overeating due to a deficit in learning and memory. Thus, conservation of the role of PKG1 α in food seeking behaviour appears to extend to mammals. Interestingly in KI mice, food intake only increased in females following food deprivation which makes food more rewarding[24] or in both sexes when a rewarding stimulus, i.e., sucrose, was present, with neither sex increasing their food intake of non-palatable food when it was abundant.

Obesity may result from a form of addiction regulated through dopaminergic reward centres of the brain [25]. Eating palatable food is associated with increased activity in the VTA, at least in part due to release of dopamine from its projections to the dorsal striatum [26] and nucleus accumbens [27], increasing pleasure sensation whilst eating rewarding food. Furthermore, when food is plentiful the VTA is inhibited by satiating signals such as leptin or insulin [28], reducing food reward and feeding. Dopamine is so important to feeding that mice unable to synthesize it die of starvation [29]. Whilst PKG is found in reward centres of the brain [3, 12-14], we show here for the first time that rewarding foods leads to increased disulfide-PKG1 α in the VTA.

Surprisingly, the changes observed in foraging behaviour of the KI mouse did not lead to a change in body weight. However, this could be explained by the lack of rewarding value of standard chow, as this only minimally activates the mesolimbic dopaminergic system, which we found to be crucial to the change in foraging behaviour of the KI mice. Furthermore, a high sucrose diet which strongly engages the reward system, doesn't lead to weight gain in mice [30]. Whilst KI mice do gain weight on a high fat diet, they do not do so beyond that of controls [31]. A likely explanation is that high fat diets, whilst leading to weight gain, concomitantly reduces dopaminergic signalling [32-34], which may mask the effects that the C42S PKG1 α KI mutation has on food seeking behaviour. Therefore, an obesogenic diet of high fat to increase caloric intake and high sucrose to increase reward would appear more appropriate than the use of either individually. This is because a diet of high fat high combined with sucrose is more rewarding than a diet containing either component alone [35], which may enhance the effects of PKG1 α on feeding behaviour.

Our data also suggest that food restriction may also provide a way to initiate binge eating in the KI mice, which is a common feature of feeding behaviour in obese humans. Interestingly, restricting sucrose still engages the dopaminergic system [36, 37], and restricting the availability of high sucrose [38] or high fat [39-41] causes binge eating, which could potentially be enhanced in KI mice. Therefore, further investigation into weight gain of the KI mice on a restricted obesogenic diet is warranted, which would more closely resemble a western diet and the feeding behaviour of obese people.

Inhibition of NOS, which diminishes cGMP-dependent activation of PKG1 α , reduces consumption of rewarding food [42] and food intake in general [43]. Consistent with this and the fact that KI mice have potentiated food seeking, is the fact that inhibition of the NOS-cGMP

pathway promotes accumulation of disulfide-PKG1 α [7, 44] in the VTA and increased food reward. Interestingly, KI mice did not show reduced sucrose preference following pre-treatment with L-NAME. These mice have a basal heightened state of reward because they cannot form the disulfide-PKG1 α 'off switch' that delimits sucrose preference. Therefore, inhibition of the NOS-cGMP-PKG1 α reward pathway alone appears insufficient to counteract the loss of negative regulation of reward by disulfide-PKG1 α .

We found elevated disulfide-PKG1 α in the VTA following stimulation of the dopaminergic system with L-DOPA, a dopamine precursor, as well as after administration of dopamine itself. Furthermore, blockade of MAO-B, which will reduce cellular hydrogen peroxide, which is a by-product of dopaminergic breakdown [18], reduced disulfide-PKG1 α in the VTA and increased sucrose preference in WT mice. This may provide some insight as to why antidepressants including MAO inhibitors can lead to weight gain [45, 46]. That deprenyl did not alter food reward in KI mice, shows the effect of altering hydrogen peroxide on food reward was significantly through disulfide-PKG1 α . As a WT phenotype was rescued by introducing WT PKG1 α into the VTA of KI mice, this suggests this is a primary site of action of disulfide-PKG1 α . Therefore, we conclude elevated neuronal activity in the VTA leads to an increase in dopamine production and recycling, and with it cellular hydrogen peroxide and disulfide PKG1 α levels, which reduces reward via phosphorylation of a key (yet unknown) substrate, providing a negative feedback loop.

As MAO-B is a cytosolic enzyme found in dopaminergic neurons and not GABAergic neurons, this suggests dopaminergic neurons are a primary site of action of PKG1 α activity in food reward in reward centres [14]. This does not mean a role for PKG1 α in GABAergic neurons should be excluded [3], but instead that the GABAergic and dopaminergic PKG1 α systems may work in tandem. Interestingly MAO-B inhibition induces reward similar to amphetamines [47], and enhances nicotine [48] and cocaine [49] reward. Conversely, NOS inhibition attenuates reward to other stimulants, such as cocaine [50], methamphetamine [51], MDMA[52], morphine [53], or alcohol [54], as it does for food reward. Taken together with the data presented here, it is possible PKG1 α provides a target for reducing multiple types of addiction.

In summary, the NOS-cGMP-PKG1 α pathway promotes food-seeking and -reward behaviour with oxidation of PKG1 α to its interprotein disulfide state acting as a negative regulator of this process.

Material and methods

Mice

All procedures were performed in accordance with the European Commission Directive 2010/63/EU (European Convention for the Protection of Vertebrate Animals used for Experimental and Other Scientific Purposes) and the United Kingdom Home Office (Scientific Procedures) Act (1986) with project approval from the Animal Welfare and Ethical Review Body of King's College London. Mice constitutively expressing C42S PKG1 α were generated on a pure C57BL/6 background by Taconic Artemis. Age (~8 weeks) and body weight (~25 g) matched PKG1 α C42S knock in (KI) mice or their WT littermates (WTs) of either sex were used

in all studies. All animals had ad libitum access to standard chow (Rat and Mouse No. 1, Special Diet Services, Essex, UK) and water, unless otherwise stated.

AAV Vector particle production and purification

AAV transfer plasmids were derived from pAAV-MCS and scAAV-MCS (Biocat GmbH). The CMV promoter in pAAV-MCS was excised using NcoI/Sall while retaining the CMV enhancer sequence.

Synapsin I proximal promoter (Syn) was amplified from pAAV-Syn-Venus (kindly provided by Rolf Sprengel, Institute for Pharmacy and Molecular Biotechnology, University of Heidelberg, Heidelberg, Germany). For insertion of Synapsin I proximal promoter in pAAV-MCS and to generate a GFP expression cassette as a positive control, Syn was amplified using specific primers 5'- ATCGCTATTACCATG G CTGCAGAGGGCCCTGCGTAT, 5'- ACG GTT CAC TAA ACG GCA GAC TGA GGC AGC GCT G) and another fragment comprising an intronic sequence along with GFP (5'-CGTTTGTAGTGAACCGTCAGATC, 5'-GCT TCT GCA GGT CGA CTT ACT TGT ACA GCT CGT CCA TG) from pscAAV-GFP (Cellbiolabs). Both fragments were inserted simultaneously into pAAV-MCS (digested with NcoI/Sall) using the InFusion HD Cloning Kit (Clontech) according to manufacturer recommendations to generate an AAV transfer plasmid, pAAV-ESyn-GFP.

Subsequently, the pAAV-ESyn-GFP transfer plasmid was digested with EcoRI/Sall. GFP comprising a C-terminal 2A peptide (Thosea Asigna 2A peptide) was amplified (5'-CTTTGGAAGTGAATT C ACC ATGGTGAGCAAGGGCGAG, 5'-GCT AGC CGG TGC AGG GCC GGG) and another fragment encoding PKGI α wildtype and mutant C42s (5'- CCT GCACCG GCTAGC ATGGACTACAAGGACGACGAC, 5'- GCT TCT GCA GGT CGA CTT AGA AGT CTA TAT CCC ATC CTG). Both fragments were inserted in a two-fragment cloning strategy into pAAV-ESyn-GFP (digested with EcoRI/Sall) using the InFusion HD Cloning Kit (Clontech) according to manufacturer recommendations to generate two AAV transfer plasmids pAAV-ESyn-GFP-2A-WT.PKGI α and pAAV-ESyn-GFP-2A- C42S.PKGI α . Integrity of AAV transfer plasmids was confirmed by restriction digestion using SmaI and sequencing.

AAV9 pseudotyped vectors were generated by co-transfection of 293AAV cells (Biocat GmbH) with the respective AAV transfer plasmid containing green fluorescent protein gene expression cassette (eGFP) in frame with the gene of interest (PKGI) flanked by AAV2 inverted terminal repeats (ITR) and the AAV packaging plasmid pAAV2/9 (kindly provided by Julie Johnston, Pennsylvania University, Philadelphia, PA, USA) providing the AAV2 rep and AAV9 cap genes and an additional plasmid encoding adenoviral helper functions (pHelper, Biocat GmbH). Generation of recombinant AAV6 and AAV9 particles was carried out as described previously [55] with some variations: 293 cells were cultivated in DMEM (High Glucose) supplemented with 10% (v/v) heat-inactivated fetal calf serum, 0.1 mM MEM Non-Essential Amino Acids (NEAA), 2 mM L-glutamine, 100 IU/ml penicillin, and 100 μ g/ml streptomycin. Tissue culture reagents were obtained from Life technologies. Briefly, 1.5×10^7 293AAV cells were seeded to 15 cm plates each and transfected directly in suspension.

For production of AAV9, 4.5 μ g AAV transfer plasmid and 7.5 μ g of pAAV2/9 and 10 μ g pHelper (Biocat GmbH) with a molar plasmid-ratio 1:1:1 were diluted in 1.6 ml of 150 mM NaCl. 340 μ l PEI (7.5 mM; Polysciences) were diluted with 1.3 ml 150 mM NaCl. PEI mix was added to the DNA mixture, incubated at room temperature for 15 min and added dropwise to cells. After 72 h cells were harvested, washed three times with phosphate-buffered saline

(PBS) and resuspended in (50 mM Tris base, 150 M NaCl, 5 mM MgCl₂, pH 8.5). After three freeze-thaw cycles, Benzonase (250 U/ml; Merk KGaA) was added and the lysates were incubated for 1 h at 37°C. Cell debris was pelleted and vector containing lysates were purified using iodixanol step gradients.

The genomic titers of DNase resistant recombinant AAV particles are determined by quantitative PCR using the SYBR Green qPCR Master MIX 2 (ThermoScientific) and a 7900 HT cyclor (ABI). Vectors are quantified using GFP specific primers (5'-GCTGCTGCCCCGACAACCACTACCTGAGCAC and 5'-CGGCGGTCACGAAGTCCAGCAGGACCATGT).

Realtime PCR was done in 10 µl with 0.3 µM for each primer. Fluorescence was measured at the end of each annealing phase. AAV transfer plasmid was employed as a copy number standard. A standard curve for quantification was generated by serial dilutions of the respective vector plasmid DNA. Cycling conditions were as follows: 50 °C for 2 min, 95°C for 10 min, followed by 35 cycles of 95°C for 15 sec and 60°C for 60 sec. Calculations were done using SDS 2.4 software (ABI).

Stereotaxic brain injections

Mice were anaesthetized with isoflurane (primal healthcare) and anaesthesia maintained throughout the procedure. Animals were placed prone and the head was fixed in a small animal stereotaxic frame (WPI). The skull was exposed via a 1 cm longitudinal midline skin incision. Mice were injected bilaterally using borosilicate glass pipettes connected to an air injector system (Narishige), with 200-300 nL of ESyn-GFP, ESyn-GFP-2A-WT.PKGI α , ESyn-GFP-2A-C42S.PKGI α , or synapsin-GFP containing AAV. VTA coordinates: 4.0 mm caudal and 480 µm ventral to lambda. The wound was sutured closed (6-0 mersilk; ethicon) and the animals were given postoperative buprenorphine (0.1 mg/kg; intramuscular) and allowed to recover for 3 weeks before behavioural experiments began.

Behavioural testing

Mice were kept on a 12-hour day/night cycle. All mice were allowed to habituate for 10 days before undergoing behavioural tests. Ambient temperature in all rooms was maintained at 21 ± 2 °C with 45% humidity. Sawdust and nesting materials (Datesand Ltd, Manchester) in each cage were changed once every two weeks by the experimenter, but never on the day before or the day of testing to minimize the disruptive effect of cage cleaning on behaviour. All mice undergoing behavioural tests were randomized and the experimenter blinded to genotype, with genotypes revealed only after analysis had been performed. Behaviours were recorded using a camera positioned above the test arenas and movement of each mouse tracked using EthoVision (Noldus Information Technologies bv, Wageningen, The Netherlands). After each individual test, boli and urine were removed from the test arena which was cleaned with 1% Anistel® solution ((high level surface disinfectant, Trisel Solution Ltd, Cambridgeshire, UK) to remove any odours.

1. Home-cage locomotor activity:

Mice were recorded in their home cage (20 × 36 × 14 cm). The test room lighting and cycle were adjusted to match their housing room. Red multi-LED cluster lights were used to track their movement in the dark cycle. Fresh sawdust was placed two days prior to the day of the test to allow better automated tracking of the mice by Ethovision software. Two hours prior to the end of their dark cycle (17:00) on the test day, their home-cage lid was removed and a Plexiglas lid, pierced with holes to allow normal air-exchanges was placed on top of each cage. Wet pellets of food were placed into one corner of the cage, and replenished every 8-10 hours. Video recordings were analysed in detail using EthoVision, from which mean speed (cm/s), movement frequency, duration (s) and number of visits to the feeding/drinking area were calculated. Home-cage behaviour was recorded for 24 hours.

2. Sucrose preference test:

Mice were habituated to a two bottles paradigm in their home cage over 3 days with both bottles containing water, to verify that no side preference existed. On day 4, mice were given a free choice between two bottles, one with 2% (wt/vol) sucrose solution, and another with water for 5 days. In some cases, the sucrose preference test continued for a further 5 days with mice administered with various drugs or saline (0.9%) intraperitoneally (IP) once per day in the last hour of the light cycle for 5 consecutive days. IP drugs included: 50 mg/kg L-DOPA (a dopaminergic agonist); 0.5, 2.5, or 5 mg/kg deprenyl (a monoamine oxidase-B inhibitor) or; 2.5, 5, or 10 mg/kg NG-Nitro-L-arginine methyl ester (L-NAME), a NOS inhibitor. All drugs were purchase from Sigma-Aldrich and dissolved in saline.

Sucrose solutions were changed every 3 days to avoid bacterial growth and changes in taste of the solution. The side of sucrose and water bottles was switched daily to ensure results were not confounded by side preference. Extra care was taken to avoid liquid spillage. Bottles were filled in advance and kept in the housing room to reduce any effects of room temperature and pressure. Intake of water, 2% sucrose solution, and total intake was estimated by weighing bottles before starting the test and every 24 hours, with re-weighing whenever solutions were changed. Side preference was calculated as the percentage of left side solution consumed out of the total amount of liquid consumed (left side solution intake/total intake [left and right side] × 100). Sucrose preference was calculated as the percentage of sucrose solution consumed out of the total amount of liquid consumed (sucrose solution intake/total intake × 100).

3. Novelty-suppressed feeding:

Mice were food restricted 24hrs prior to the novelty-suppressed feeding (NSF) test. On the day of testing, the mouse was weighed and placed wall-facing in a new cage containing only one pellet of food on the opposite side of the cage to the mouse. A red light was evenly distributed across the arena. Latency to approach chow and to begin eating was recorded within a 5-min period.

4. Open field:

The open field arena was a clear circular custom-built light grey acrylic arena (40 cm diameter x 40 cm height). A maximum of 10 lux white light was evenly distributed across the arena during testing. Each mouse was taken from their home-cage and placed into the arena, wall-facing, for the beginning of the test session. The session was video-recorded for 5 mins and analysed using EthoVision. Number of faecal boli and presence of urination were recorded at

the end of the test. In EthoVision, a virtual circle (20 cm diameter) of equal distance from the periphery was defined as the 'central zone' to determine centre locomotor activity (cm), duration (s), speed (cm/s) and frequency of entries into the centre zone.

5. *Elevated-plus maze:*

An elevated plus maze constructed from black opaque acrylic was used with 4 arms (30 cm x 5 cm) and a central platform (5 cm x 5 cm). One set of arms, opposing one another, were enclosed completely by 15 cm high walls of black acrylic, while the other 2 arms were open with an 0.5 cm ledge on either side. The maze was elevated 40 cm from the ground on an acrylic stand. Light intensity around the maze was set at 100 lux on open arms during testing. Mice were placed on the central platform, facing towards a closed arm, and allowed to explore the maze freely for 5 mins. Number of faecal boli and presence of urination were recorded at the end of the test.

EPM was analysed using tracking and hand scoring functions in EthoVision, to extract: distances and speed travelled across closed and open arms; approximate time spent in closed and open arms; approximate total entries into closed or open arms; and approximate latency of the first entry onto an open arm. The exact number of entries onto, time spent on, and latency to enter closed and open arms were manually scored by the experimenter. An arm entry was defined as all 4 paws in the arm; an exit was defined as 2 paws out of the arm.

6. *Olfactory habituation/dishabituation test:*

The test was carried out in the mouse's home-cage with a light of 10 lux inside the cage. During acclimation, a fresh dry cotton-tipped wooden applicator is inserted through the lid and presented to mice for 2 mins, prior to starting the test. Following this habituation, a series of different odours (water, banana, and opposite-sex urine) were presented each in three consecutive trials of 2 mins each. Animal were recorded during the entire test.

7. *Novel object:*

This test was performed using a clean new home-cage. Green rough and black soft cylinders (7 cm height and 3 cm diameter) were introduced into the arena as novel objects. Low white lighting, at approximately 30 lux, was evenly distributed across the arena during testing. The novel object was placed prior to introduction of mice in the cage. Each mouse was taken from their home cage and placed into a corner, wall-facing of the open field arena at the start of the test. The third of the arena in which the object was placed was defined as the 'exploration zone' in EthoVision. Latency (s) to initial exploration of the object, as well as frequency and duration (s) to explore the object, and rearing behaviours during the 3 min were measured.

8. *Y-maze:*

The spontaneous spatial novelty preference test was conducted using a perspex Ymaze. Each arm was 22 cm long, 7 cm wide, with 20-cm-high walls. For this test, one entry into an arm was defined as placement of 2 paws into that arm. For the first trial, one of the arms of the Y-maze was closed, therefore mice could either go left or right according to a pseudorandom sequence (equal numbers of left and right arms were blocked in total sessions), mice could also can move in the central arm for 5 mins. In the second trial one hour later to assess short term memory, the test was repeated with access to both arms. An area was defining for each

arm (right and left) and the main arm in EthoVision: speed (cm/s), distance moved (cm), frequency and duration (s) to explore each arm were measured.

9. Rotarod:

Mice were tested over 3 days, with 24h between sessions; on each day, mice underwent 2 trials, with 1h intervals between them. Mice were placed on an accelerating rotarod (0–40 rpm over 5 min). Latency (s) to fall was the only measure recorded. Parameters analysed included latencies to fall at particular trials, as well as differences between trials. If mice fail to walk and simply grip onto the rotating drum, latency to fall was recorded after 2 complete revolutions without walking.

10. Grip strength:

A Linton grip strength meter (MJS Technology) was used to measure forelimb and hindlimb grip strength as an indicator of neuromuscular function. The grip strength meter was positioned horizontally, and mice were held by the tail and lowered toward the apparatus. Mice were allowed to grasp the metal grid with their forelimbs and were then pulled backward in the horizontal plane. The force in grams applied to the bar at the moment the grasp is released was recorded. This was repeated three times, and average grip strength for each mouse was calculated. The test was repeated for hindlimbs. Mice were not trained prior to testing.

Ex vivo brain removal and dopamine treatment

For experiments testing dopamine, naïve mice were given an IP injection of either auranofin (25 mg/kg; to maintain disulfide bonds by inhibiting thioredoxin reductase; Sigma-Aldrich), or auranofin and deprenyl. Brains were removed 30 mins following IP injections. The VTA was dissected out by eye with a razor blade and placed into artificial cerebrospinal fluid (aCSF: 124 mM NaCl, 3 mM KCl, 1 mM CaCl₂, 26 mM NaHCO₃, 1.25 mM NaH₂PO₄, 1 mM MgSO₄, 10 mM D-glucose (saturated with 95% O₂/5% CO₂; pH 7.5). Isolated VTAs were treated with 1 mM dopamine (Sigma-Aldrich) and either auranofin or auranofin and deprenyl (to match the previous IP injections), for 5 mins before being snap frozen in liquid nitrogen for subsequent analysis by western blotting.

In vivo L-DOPA treatment

Mice that had just completed sucrose preference testing, or naïve mice, were given either one or two IP injections separated by 30 mins. Single IP injections consisted of either saline, auranofin, or deprenyl. Experiments with two IP injections followed 2 protocols; 1) Initial injections consisted of auranofin followed by injections of either L-DOPA, L-NAME, or deprenyl, or 2) Initial injections consisted of auranofin and deprenyl followed by an injection of L-DOPA. Carbidopa was added to all injections containing L-DOPA to stop breakdown of L-DOPA in the periphery so that a sufficient concentration may reach the brain. All drugs were dissolved in saline except auranofin, which was dissolved in DMSO. Brains were removed 30 mins after the final IP injection and snap frozen in liquid nitrogen for subsequent analysis by western blotting.

Western blot

Immunoblotting for PKG1 α disulfide dimer was performed as previously described [6] with maleimide (100 mM/L) used in preparation buffers to alkylate thiols and prevent thiol disulfide exchange. Antibodies used in these studies included cGKI α (Santa Cruz Biotechnology), and GAPDH (Cell Signalling Technology). Horseradish peroxidase-linked secondary antibody (Dako) and ECL reagent (GE Healthcare) were used. Digitized immunoblots were analyzed quantitatively with a Gel-Pro Analyzer 3.1. The percentage of PKG1 α disulfide dimer was normalized by the total GAPDH protein expression.

Statistics

Results are presented as mean \pm SEM. Differences between groups were assessed using a 1-way ANOVA, post-hoc testing was performed as either a t-test for comparison of 2 groups or using a Bonferroni correction for comparison of 3 or more groups. The only exception was home cage locomotor activity which was analysed with a 2-way repeated measures ANOVA with Bonferroni correction for analysing distance moved over 24 hours in 30-minute bins, and by t-test for total distance moved between 19.30 and 21.30. Differences were considered significant at a 95% confidence level.

Acknowledgements

This work was supported by the British Heart Foundation, the European Research Council (ERC Advanced award), the Medical Research Council, and the Department of Health via the NIHR cBRC award to Guy's & St Thomas' NHS Foundation Trust.

Bibliography

- [1] Kaun, K. R.; Sokolowski, M. B. cGMP-dependent protein kinase: linking foraging to energy homeostasis. *Genome* **52**:1-7; 2008.
- [2] Feil, S.; Zimmermann, P.; Knorn, A.; Brummer, S.; Schlossmann, J.; Hofmann, F.; Feil, R. Distribution of cGMP-dependent protein kinase type I and its isoforms in the mouse brain and retina. *Neuroscience* **135**:863-868; 2005.
- [3] Nugent, F. S.; Niehaus, J. L.; Kauer, J. A. PKG and PKA Signaling in LTP at GABAergic Synapses. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* **34**:1829-1842; 2009.
- [4] Nakamura, T.; Ranek, M. J.; Lee, D. I.; Shalkey Hahn, V.; Kim, C.; Eaton, P.; Kass, D. A. Prevention of PKG1 α oxidation augments cardioprotection in the stressed heart. *The Journal of Clinical Investigation* **125**:2468-2472; 2015.
- [5] Alverdi, V.; Mazon, H.; Versluis, C.; Hemrika, W.; Esposito, G.; van den Heuvel, R.; Scholten, A.; Heck, A. J. R. cGMP-binding Prepares PKG for Substrate Binding by Disclosing the C-terminal Domain. *J. Mol. Biol.* **375**:1380-1393; 2008.
- [6] Burgoyne, J. R.; Madhani, M.; Cuello, F.; Charles, R. L.; Brennan, J. P.; Schröder, E.; Browning, D. D.; Eaton, P. Cysteine Redox Sensor in PKG1 α Enables Oxidant-Induced Activation. *Science* **317**:1393-1397; 2007.
- [7] Burgoyne, J. R.; Pryszyzna, O.; Rudyk, O.; Eaton, P. cGMP-Dependent Activation of Protein Kinase G Precludes Disulfide Activation: Implications for Blood Pressure Control. *Hypertension* **60**:1301-1308; 2012.
- [8] Landgraf, W.; Regulla, S.; Meyer, H. E.; Hofmann, F. Oxidation of cysteines activates cGMP-dependent protein kinase. *J Biol Chem* **266**:16305-16311; 1991.
- [9] Müller, P. M.; Gnügge, R.; Dhayade, S.; Thunemann, M.; Krippeit-Drews, P.; Drews, G.; Feil, R. H₂O₂ lowers the cytosolic Ca²⁺ concentration via activation of cGMP-dependent protein kinase 1 α . *Free Radic Biol Med* **53**:1574-1583; 2012.
- [10] Pryszyzna, O.; Burgoyne, J. R.; Scotcher, J.; Grover, S.; Kass, D.; Eaton, P. Phosphodiesterase 5 Inhibition Limits Doxorubicin-induced Heart Failure by Attenuating Protein Kinase G 1 α Oxidation. *Journal of Biological Chemistry* **291**:17427-17436; 2016.
- [11] Pierce, R. C.; Kumaresan, V. The mesolimbic dopamine system: The final common pathway for the reinforcing effect of drugs of abuse? *Neurosci Biobehav Rev* **30**:215-238; 2006.
- [12] Oh, J. H.; Lee, D. K.; Shim, Y.-B.; Ryu, I. S.; Seo, S. Y.; Kim, J.; Yang, J. H.; Cho, H.-W.; Choe, E. S. Dopamine D4 receptors linked to protein kinase G are required for changes in dopamine release followed by locomotor activity after repeated cocaine administration. *Exp Brain Res* **233**:1511-1518; 2015.
- [13] Lee, D. K.; Oh, J. H.; Shim, Y.-B.; Choe, E. S. Protein Kinase G Regulates Dopamine Release, Δ FosB Expression, and Locomotor Activity After Repeated Cocaine Administration: Involvement of Dopamine D2 Receptors. *Neurochem Res* **38**:1424-1433; 2013.
- [14] Jouvert, P.; Revel, M.-O.; Lazaris, A.; Aunis, D.; Langley, K.; Zwiller, J. Activation of the cGMP Pathway in Dopaminergic Structures Reduces Cocaine-Induced EGR-1 Expression and Locomotor Activity. *J Neurosci* **24**:10716-10725; 2004.
- [15] Tahsili-Fahadan, P.; Yahyavi-Firouz-Abadi, N.; Orandi, A.; Esmaeili, B.; Basseda, Z.; Dehpour, A. Rewarding properties of sildenafil citrate in mice: role of the nitric oxide-cyclic GMP pathway. *Psychopharmacology* **185**:201-207; 2006.
- [16] Itzhak, Y.; Roger-Sánchez, C.; Anderson, K. L. Role of the nNOS gene in ethanol-induced conditioned place preference in mice. *Alcohol* **43**:285-291; 2009.
- [17] Pryszyzna, O.; Rudyk, O.; Eaton, P. Single atom substitution in mouse protein kinase G eliminates oxidant sensing to cause hypertension. *Nature Medicine* **18**:286; 2012.
- [18] Hermida-Ameijeiras, Á.; Méndez-Álvarez, E. a.; Sánchez-Iglesias, S. a.; Sanmartín-Suárez, C.; Soto-Otero, R. Autoxidation and MAO-mediated metabolism of dopamine as a potential cause of oxidative stress: role of ferrous and ferric ions. *Neurochemistry International* **45**:103-116; 2004.

- [19] Denninger, J. W.; Marletta, M. A. Guanylate cyclase and the $\cdot\text{NO}$ /cGMP signaling pathway. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1411**:334-350; 1999.
- [20] Rainer, P. P.; Kass, D. A. Old dog, new tricks: novel cardiac targets and stress regulation by protein kinase G. *Cardiovasc Res* **111**:154-162; 2016.
- [21] Burgoyne, J. R.; Oka, S.-i.; Ale-Agha, N.; Eaton, P. Hydrogen Peroxide Sensing and Signaling by Protein Kinases in the Cardiovascular System. *Antioxid Redox Signal*. **18**:1042-1052; 2013.
- [22] Rudyk, O.; Prysyazhna, O.; Burgoyne, J. R.; Eaton, P. Nitroglycerin fails to lower blood pressure in redox-dead Cys42Ser PKG1 α knock-in mouse. *Circulation* **126**:287-295; 2012.
- [23] Sanchez-Alavez, M.; Klein, I.; Brownell, S. E.; Tabarean, I. V.; Davis, C. N.; Conti, B.; Bartfai, T. Night eating and obesity in the EP3R-deficient mouse. *PNAS* **104**:3009-3014; 2007.
- [24] Cameron, J. D.; Goldfield, G. S.; Finlayson, G.; Blundell, J. E.; Doucet, É. Fasting for 24 Hours Heightens Reward from Food and Food-Related Cues. *PLOS ONE* **9**:e85970; 2014.
- [25] Volkow, N. D.; Wise, R. A.; Baler, R. The dopamine motive system: implications for drug and food addiction. *Nat Rev Neurosci* **18**:741; 2017.
- [26] Small, D. M.; Jones-Gotman, M.; Dagher, A. Feeding-induced dopamine release in dorsal striatum correlates with meal pleasantness ratings in healthy human volunteers. *NeuroImage* **19**:1709-1715; 2003.
- [27] Norgren, R.; Hajnal, A.; Mungarndee, S. S. Gustatory Reward and the Nucleus Accumbens. *Physiol. Behav.* **89**:531-535; 2006.
- [28] Figlewicz, D. P.; Evans, S. B.; Murphy, J.; Hoen, M.; Baskin, D. G. Expression of receptors for insulin and leptin in the ventral tegmental area/substantia nigra (VTA/SN) of the rat. *Brain Res* **964**:107-115; 2003.
- [29] Szczypka, M. S.; Kwok, K.; Brot, M. D.; Marck, B. T.; Matsumoto, A. M.; Donahue, B. A.; Palmiter, R. D. Dopamine Production in the Caudate Putamen Restores Feeding in Dopamine-Deficient Mice. *Neuron* **30**:819-828; 2001.
- [30] Avena, N. M.; Rada, P.; Hoebel, B. G. Sugar and fat bingeing have notable differences in addictive-like behavior. *The Journal of nutrition* **139**:623-628; 2009.
- [31] Rudyk, O.; Eaton, P. Examining a role for PKG 1 α oxidation in the pathogenesis of cardiovascular dysfunction during diet-induced obesity. *Free radical biology & medicine* **110**:390-398; 2017.
- [32] Carlin, J.; Hill-Smith, T. E.; Lucki, I.; Reyes, T. M. Reversal of dopamine system dysfunction in response to high-fat diet. *Obesity (Silver Spring, Md.)* **21**:2513-2521; 2013.
- [33] Speed, N.; Saunders, C.; Davis, A. R.; Owens, W. A.; Matthies, H. J. G.; Saadat, S.; Kennedy, J. P.; Vaughan, R. A.; Neve, R. L.; Lindsley, C. W.; Russo, S. J.; Daws, L. C.; Niswender, K. D.; Galli, A. Impaired striatal Akt signaling disrupts dopamine homeostasis and increases feeding. *PloS one* **6**:e25169-e25169; 2011.
- [34] Johnson, P. M.; Kenny, P. J. Dopamine D2 receptors in addiction-like reward dysfunction and compulsive eating in obese rats. *Nature neuroscience* **13**:635-641; 2010.
- [35] DiFeliceantonio, A. G.; Coppin, G.; Rigoux, L.; Edwin Thanarajah, S.; Dagher, A.; Tittgemeyer, M.; Small, D. M. Supra-Additive Effects of Combining Fat and Carbohydrate on Food Reward. *Cell Metabolism* **28**:33-44.e33; 2018.
- [36] Bello, N. T.; Sweigart, K. L.; Lakoski, J. M.; Norgren, R.; Hajnal, A. Restricted feeding with scheduled sucrose access results in an upregulation of the rat dopamine transporter. **284**:R1260-R1268; 2003.
- [37] Colantuoni, C.; Schwenker, J.; McCarthy, J.; Rada, P.; Ladenheim, B.; Cadet, J.-L.; Schwartz, G. J.; Moran, T. H.; Hoebel, B. G. Excessive sugar intake alters binding to dopamine and mu-opioid receptors in the brain. **12**:3549-3552; 2001.
- [38] Wojnicki, F. H. E.; Stine, J. G.; Corwin, R. L. W. Liquid sucrose bingeing in rats depends on the access schedule, concentration and delivery system. *Physiology & Behavior* **92**:566-574; 2007.

- [39] Corwin, R. L.; Wojnicki, F. H. E.; Fisher, J. O.; Dimitriou, S. G.; Rice, H. B.; Young, M. A. Limited Access to a Dietary Fat Option Affects Ingestive Behavior But Not Body Composition in Male Rats. *Physiology & Behavior* **65**:545-553; 1998.
- [40] Wojnicki, F. H. E.; Charny, G.; Corwin, R. L. W. Binge-type behavior in rats consuming trans-fat-free shortening. *Physiology & behavior* **94**:627-629; 2008.
- [41] Davis, J. F.; Melhorn, S. J.; Shurdak, J. D.; Heiman, J. U.; Tschöp, M. H.; Clegg, D. J.; Benoit, S. C. Comparison of hydrogenated vegetable shortening and nutritionally complete high-fat diet on limited access-binge behavior in rats. *Physiology & behavior* **92**:924-930; 2007.
- [42] Czech, D. A. A Nitric Oxide Synthase Inhibitor, L-NAME, Attenuates Saccharin Drinking in a Two-Choice Test in Water-Deprived Rats. *Physiol. Behav.* **67**:161-165; 1999.
- [43] Morley, J. E.; Farr, S. A.; Suarez, M. D.; Flood, J. F. Nitric oxide synthase inhibition and food intake: Effects on motivation to eat and in female mice. *Pharmacol Biochem Behav* **50**:369-373; 1995.
- [44] Müller, P. M.; Gnügge, R.; Dhayade, S.; Thunemann, M.; Krippeit-Drews, P.; Drews, G.; Feil, R. H₂O₂ lowers the cytosolic Ca²⁺ concentration via activation of cGMP-dependent protein kinase Ia. *Free Radical Biology and Medicine* **53**:1574-1583; 2012.
- [45] Cantú, T. G.; Korek, J. S. Monoamine Oxidase Inhibitors and Weight Gain. *Drug Intelligence & Clinical Pharmacy* **22**:755-759; 1988.
- [46] Ranjbar, S.; Pai, N. B.; Deng, C. The Association of Antidepressant Medication and Body Weight Gain. In: Kakkilaya, D. S., ed.: Kakkilaya BS; 2013.
- [47] Wu, W.-R.; Zhu, X.-Z. The amphetamine-like reinforcing effect and mechanism of l-deprenyl on conditioned place preference in mice. *European Journal of Pharmacology* **364**:1-6; 1999.
- [48] Villégier, A.-S.; Salomon, L.; Granon, S.; Changeux, J.-P.; Belluzzi, J. D.; Leslie, F. M.; Tassin, J.-P. Monoamine Oxidase Inhibitors Allow Locomotor and Rewarding Responses to Nicotine. *Neuropsychopharmacology* **31**:1704; 2005.
- [49] Schiffer Wynne, K.; Azmoodeh, M.; Gerasimov, M.; Volkow Nora, D.; Fowler Joanna, S.; Dewey Stephen, L. Selegiline potentiates cocaine-induced increases in rodent nucleus accumbens dopamine. *Synapse* **48**:35-38; 2003.
- [50] Pulvirenti, L.; Balducci, C.; Koob, G. F. Inhibition of Nitric Oxide Synthesis Reduces Intravenous Cocaine Self-administration in the Rat. *Neuropharmacology* **35**:1811-1814; 1996.
- [51] Li, S.-M.; Yin, L.-L.; Shi, J.; Lin, Z.-B.; Zheng, J.-W. The effect of 7-nitroindazole on the acquisition and expression of d-methamphetamine-induced place preference in rats. *Eur J Pharmacol.* **435**:217-223; 2002.
- [52] García-Pardo, M. P.; Rodríguez-Arias, M.; Miñarro, J.; Aguilar, M. A. Role of nitric oxide pathway in the conditioned rewarding effects of MDMA in mice. *Behav Brain Res* **330**:75-77; 2017.
- [53] Gholami, A.; Zarrindast, M.-R.; Sahraei, H.; Haerri-Rohani, A. Nitric oxide within the ventral tegmental area is involved in mediating morphine reward. *Eur J Pharmacol.* **458**:119-128; 2003.
- [54] Itzhak, Y.; Martin, J. L. Blockade of alcohol-induced locomotor sensitization and conditioned place preference in DBA mice by 7-nitroindazole. *Brain Res* **858**:402-407; 2000.
- [55] Grieger, J. C.; Choi, V. W.; Samulski, R. J. Production and characterization of adeno-associated viral vectors. *Nat Protoc* **1**:1412-1428; 2006.

Figure 1: C42S PKG1 α knock-in (KI) ‘redox dead’ mice display increased food reward compared to wild-type littermates (WT). A) Home cage recordings show overall locomotor activity of KI mice is not different to WT. Insert: total distance moved is increased in KIs compared to WTs during the first 2 hours of their awake cycle (19:30-21:30) when food intake is at its highest (ai). However, there was not an overall change in feeding as food intake was similar across groups (aii), or an overall change in metabolism as no differences in weight were found between groups (aiii). B) In a novelty-suppressed feeding test, after 24 hours fasting, all mice approach food in a novel environment with the same delay (bi). However, male KI mice feed faster than male WTs (bii), whilst female KI mice lose more weight during food restriction (biii) and consume more food when the restriction is lifted (biv) than female WTs. Thus, feeding behaviour is altered in KI mice. C) In a sucrose preference test, KIs have a higher preference for 2% sucrose over water compared to WTs (ci), though they do not show any side preference under control conditions (cii). * - $p < 0.05$

Figure 2: KI mice did not show any alterations in anxiety levels. A) In an open-field paradigm, there were no differences between KI mice and WTs of either sex for the distance moved in the centre of the arena (ai), the number of entries in to the centre (aii), the amount of time spent in the centre (aiii), or the speed they moved through the centre of the open field (aiv). B) In an elevated-plus maze paradigm, there were no differences between KI mice and WTs of either sex for the distance moved in the open arms (bi), the number of entries in to the open arms (bii), the amount of time spent in the open arms (biii), or the speed they moved whilst on the open arms of the maze (biv).

Figure 3: KI mice did not show any alterations in olfaction. In an olfaction habituation/dishabituation test, there was no difference between KI mice and WTs of either sex for time spent investigating water, banana odour, or urine from a mouse of the opposite sex with all groups displaying habituation/dishabituation to the different odours. Comparisons between KI and WT male mice (a) and KI and WT female mice (b). * - $p < 0.05$, ** - $p < 0.01$, *** $p < 0.001$

Figure 4: KI mice did not show any alterations in exploratory behaviour. In a novel object recognition paradigm, there were no differences between KI mice and WTs of either sex for the distance moved in the object area (a), the number of entries into the object area (b), the amount of time spent in the object area (c), or the speed they moved through the object area (d).

Figure 5: KI mice did not show any alterations in short-term memory. In a Y-maze paradigm, there were no differences between KI mice and WTs of either sex for the distance moved in the novel and unknown arms (a), the number of entries into the novel and unknown arms (b), the amount of time spent in the novel and unknown arms (c), or the speed they moved through the novel and unknown arms (d). * - $p < 0.05$, ** - $p < 0.01$, *** $p < 0.001$






Figure 6: KI mice did not show any alterations in general motor function. A) In a grip strength test, there were no differences between KI mice and WTs of either sex for the force they could generate with their fore (ai) or hind limbs (aii). B) In a rotarod paradigm, there were no differences between KI mice and WTs of either sex for the time they took to fall from the apparatus. bi) Comparisons between male (bi) and female (bii).

Figure 7: Increased dopaminergic signalling leads to increased disulfide-PKG1 α through production of hydrogen peroxide during dopamine degradation. A) In isolated tissue slices of the VTA, dopamine in the presence of auranofin (added to stabilise dimer formation) led to an increased disulfide-PKG1 α , which was decreased by the MAO-B inhibitor deprenyl. Saline, auranofin, and deprenyl controls did not lead to disulfide-PKG1 α formation. ai) summary data, aii) representative western blots. B) *In vivo* L-DOPA in the presence of auranofin led to increased disulfide-PKG1 α in the VTA, which was decreased by deprenyl. Saline, auranofin, and deprenyl controls did not lead to disulfide-PKG1 α formation. bi) summary data, bii) representative western blots. Therefore, increased dopaminergic activity stimulates disulfide-PKG1 α in the VTA. Furthermore, pharmacological inhibition of dopamine breakdown by MAO-B, which otherwise generates hydrogen peroxide as a by-product, reduced disulfide-PKG1 α in the VTA. * - $p < 0.05$, ** - $p < 0.01$, *** $p < 0.001$


Figure 8: Sucrose preference in WT mice is determined by a balance between NOS/cGMP- and disulfide-PKG1 α . a) Sucrose preference is increased (ai) when disulfide-PKG1 α is reduced by deprenyl (aii: representative western blots, aiii: summary data). b) Sucrose preference is decreased (bi) when disulfide-PKG1 α is increased by L-NAME, an inhibitor of NOS (bii: representative western blots, biii: summary data). All western blots were performed in the presence of auranofin to stabilise dimer formation. Inhibition of NOS with L-NAME, which lowers cGMP levels and so abrogates its ability to attenuate disulfide-PKG1 α accumulation, increased oxidation of the kinase. * - $p < 0.05$, ** - $p < 0.01$

Figure 9: KI mice do not show alterations in sucrose preference following manipulations of the NOS/cGMP- and disulfide- activation pathways of PKG1 α . A) Sucrose preference is unaltered by deprenyl or L-NAME. B) Representative western blots showing that C42S PKG1 α KI mice cannot form disulfide bridges. Therefore, the balance between the NOS/cGMP- and - activation pathways of PKG1 α cannot be altered. As such PKG1 α is only activated through the NOS/cGMP pathway, leaving KI mice in a permanent state of increased reward.


Figure 10: Adeno-associated virus-mediated transduction of VTA neurons of KI mice with WT PKG1 α rescues food reward phenotype of WT mice. A) Cartoon of the location of transduced neurons shown in B (red box). B) Micrograph of transduction of dopaminergic neurons (red: marked by tyrosine hydroxylase) with WT PKG1 α (Green: marked by GFP expression). c) In a sucrose preference test, KI transduced with WT PKG1 α have reduced preference for 2% sucrose compared to WTs transduced with GFP (ci), though they do not show any side preference under control conditions (cii). Importantly, over expression of C42S PKG1 α in KI mice did not show any change in sucrose preference (cii), showing an overall increase in PKG1 α production/activity was not responsible for changes seen following transduction with WT PKG1 α . * - $p < 0.05$, ** - $p < 0.01$




ai




aii




aiii




aiv




bi




bii




biii




biv











a






b












ai




bi










aii



bii





a**b**