

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/129327>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Erdős-Ko-Rado for random hypergraphs: asymptotics and stability

Marcelo M. Gaury¹, Hiệp Hàn², and Igor C. Oliveira³

¹Institute of Theoretical Computer Science, ETH Zürich.*

²Instituto de Matemáticas, Pontificia Universidad Católica de Valparaíso.†

³Faculty of Mathematics and Physics, Charles University in Prague.‡

September 11, 2018

Abstract

We investigate the asymptotic version of the Erdős-Ko-Rado theorem for the random k -uniform hypergraph $\mathcal{H}^k(n, p)$. For $2 \leq k(n) \leq n/2$, let $N = \binom{n}{k}$ and $D = \binom{n-k}{k}$. We show that with probability tending to 1 as $n \rightarrow \infty$, the largest intersecting subhypergraph of $\mathcal{H}^k(n, p)$ has size $(1 + o(1))p \frac{k}{n} N$, for any $p \gg \frac{n}{k} \ln^2 \left(\frac{n}{k} \right) D^{-1}$. This lower bound on p is asymptotically best possible for $k = \Theta(n)$. For this range of k and p , we are able to show stability as well.

A different behavior occurs when $k = o(n)$. In this case, the lower bound on p is almost optimal. Further, for the small interval $D^{-1} \ll p \leq (n/k)^{1-\varepsilon} D^{-1}$, the largest intersecting subhypergraph of $\mathcal{H}^k(n, p)$ has size $\Theta(\ln(pD)ND^{-1})$, provided that $k \gg \sqrt{n \ln n}$.

Together with previous work of Balogh, Bohman and Mubayi, these results settle the asymptotic size of the largest intersecting family in $\mathcal{H}^k(n, p)$, for essentially all values of p and k .

1 Introduction

The Erdős-Ko-Rado theorem [11] is a cornerstone in extremal combinatorics. Let $[n]$ denote the set $\{1, 2, \dots, n\}$, and $\binom{[n]}{k}$ denote the set of all k -element subsets of $[n]$. A family of k -element sets $\mathcal{F} \subset \binom{[n]}{k}$ is called a k -uniform hypergraph on the vertex set $[n]$, and such a hypergraph is called *intersecting* if $A \cap B \neq \emptyset$ holds for every edge $A, B \in \mathcal{F}$. The Erdős-Ko-Rado theorem then states that for $2 \leq k \leq n/2$, an intersecting family $\mathcal{F} \subset \binom{[n]}{k}$ must satisfy $|\mathcal{F}| \leq \frac{k}{n} \binom{[n]}{k}$. This is best possible, as seen by the *principal* hypergraphs \mathcal{F}_i , which consist of all edges containing the fixed element $i \in [n]$.

We investigate a random analogue of the Erdős-Ko-Rado theorem in which the ambient space $\binom{[n]}{k}$ in the theorem is replaced by a random space. Random analogues of extremal results have

*A significant fraction of this work was completed while the author was at University of São Paulo. Supported by CNPq grants 132238/2012-8 and 248952/2013-7.

†The author was supported by the Millennium Nucleus Information and Coordination in Networks ICM/FIC RC130003, the FONDECYT Iniciación grant 11150913, FAPESP (Proc. 2010/16526-3 and 2103/03447-6), BEPE (2013/11353-1) and PROBRAL CAPES/DAAD Proc. 430/15.

‡Parts of this work were completed while the author was at Columbia University and as a visitor at University of São Paulo. Supported in part by CNPq grant 200252/2015-1.

been studied extensively in the last decades, and we refer to [24, 8, 6, 23] for the history of this line of research and recent breakthroughs.

The ambient random space we will work with is $\mathcal{H}^k(n, p)$, the binomial random k -uniform hypergraph on the vertex set $[n]$ in which each edge $e \in \binom{[n]}{k}$ is included in $\mathcal{H}^k(n, p)$ independently with probability p . Further, for a k -uniform \mathcal{H} , let $i(\mathcal{H})$ denote the size of the largest intersecting subhypergraph of \mathcal{H} , i.e., $i(\mathcal{H}) = \max\{|\mathcal{F}|: \mathcal{F} \subset \mathcal{H} \text{ and } \mathcal{F} \text{ is intersecting}\}$. In this notation the Erdős-Ko-Rado theorem states that $i(\mathcal{H}^k(n, 1)) = i\left(\binom{[n]}{k}\right) = \frac{k}{n} \binom{n}{k}$.

Notation. All asymptotic limits in this paper are taken as $n \rightarrow \infty$. If we write $a(n) \ll b(n)$ or $a(n) = o(b(n))$, it means that $a(n)/b(n) \rightarrow 0$. In particular, the notation $o(1)$ represents a function that goes to 0 as $n \rightarrow \infty$, as usual. For simplicity, we omit floor and ceiling functions, whenever they are not essential. We say that a sequence of events \mathcal{E}_n holds asymptotically almost surely if $\Pr[\mathcal{E}_n] \rightarrow 1$ as $n \rightarrow \infty$. By $\ln^d c$ we denote $(\ln c)^d$.

We will be interested in $i(\mathcal{H}^k(n, p))$ for $k = k(n)$ and all $p = p(n) \in (0, 1)$. This question was investigated by Balogh, Bohman and Mubayi [3], which obtained very precise results on the size and on the structure of the largest intersecting family in $\mathcal{H}^k(n, p)$, for $k \leq n^{1/2-o(1)}$. For larger k , they obtained asymptotic tight bounds on $i(\mathcal{H}^k(n, p))$, however, only for rather large values of p . In general, their result highly depends on the range of k and p , and hence it is slightly cumbersome to state. Therefore, we will only partially discuss it here, and refer to [3] for detailed information. Their result concerning the large range of k is given below in Proposition 1.1.

Proposition 1.1 (Proposition 1.3 in [3]). *Let $\delta = \delta(n) > 0$ and $N = \binom{n}{k}$. If $\ln n \ll k < (1 - \delta)n/2$ and $p \gg (1/\delta)((\ln n)/k)^{1/2}$, then almost surely as $n \rightarrow \infty$:*

$$i(\mathcal{H}^k(n, p)) = (1 + o(1))p(k/n)N.$$

In other words, for this range of p , the expected size of the intersection of a principal family \mathcal{F}_i with $\mathcal{H}^k(n, p)$ is very close to the size of a maximum intersecting subfamily of $\mathcal{H}^k(n, p)$. We extend this result, and provide an almost complete description of $i(\mathcal{H}^k(n, p))$ as follows.

Theorem 1.2. *For all $0 < \varepsilon < 1$ there exists a constant $C > 0$ for which the following holds. Let $p = p(n) \in (0, 1)$, $k = k(n)$, where $2 \leq k \leq n/2$, $N = \binom{n}{k}$, and $D = \binom{n-k}{k}$. Then almost surely as $n \rightarrow \infty$:*

- (1) $i(\mathcal{H}^k(n, p)) = (1 \pm \varepsilon)pN$ if $N^{-1} \ll p \ll D^{-1}$,
- (2) $i(\mathcal{H}^k(n, p)) \geq (1 - \varepsilon)\frac{N}{D} \ln(pD)$ if $D^{-1} \ll p \leq (n/k)D^{-1}$ and $k \gg \sqrt{n \ln n}$,
- (3) $i(\mathcal{H}^k(n, p)) \leq C\frac{N}{D} \ln(pD)$ if $D^{-1} \ll p \leq (n/k)^{1-\varepsilon}D^{-1}$,
- (4) $i(\mathcal{H}^k(n, p)) = (1 \pm \varepsilon)p\frac{k}{n}N$ if $p \geq C(n/k) \ln^2(n/k)D^{-1}$.

The first bound follows from a standard deletion argument, and we state it here for completeness. Note also that $i(\mathcal{H}^k(n, p))$ is monotone in p , hence, in the range of p around $(n/k)D^{-1}$ not mentioned in the theorem, we have $i(\mathcal{H}^k(n, p)) = O((N/D) \ln^2(n/k))$ due to (4).

If k is linear in n , the bounds in (1) and (4) determine $i(\mathcal{H}^k(n, p))$ asymptotically for essentially all p . Here, we have a change of behaviour around D^{-1} . Roughly speaking, for p below D^{-1} ,

essentially all of $\mathcal{H}^k(n, p)$ is intersecting. Beyond that point, i.e. for $p \gg D^{-1}$, the largest intersecting subhypergraph of $\mathcal{H}^k(n, p)$ has size very close to the size of the intersection of a principal hypergraph with $\mathcal{H}^k(n, p)$. Observe that cases (2) and (3) are trivial for $k = \Theta(n)$.

For $k = o(n)$ there is a rather short range of p where $i(\mathcal{H}^k(n, p))$ reveals a “flat” behaviour. Indeed, the upper bound (3) shows that $i(\mathcal{H}^k(n, p))$ grows slowly with p , since it appears only in the \ln -term. The corresponding lower bound in (2) shows that for $k \geq n^{1/2+o(1)}$ this bound is tight up to a multiplicative constant. We provide no lower bound for the range $k < n^{1/2-o(1)}$ here, as in this case the result of Balogh et al. is more satisfactory. Again, we refer to [3] for further information.

Although the “flat range” phenomenon might come as a surprise, it has been observed elsewhere. Indeed, in the dense case, i.e. for $p = 1$, and for $k = o(n)$, the size of the largest intersecting family is vanishing compared to the ambient space, that is, $i(\binom{[n]}{k}) = \frac{k}{n} \binom{n}{k} = o\left(\binom{n}{k}\right)$. For these so called “degenerate” problems, the random analogues typically reveal such an intermediate flat behaviour, as observed for example in [19, 20].

The question of for which range of p the largest intersecting family $\mathcal{F} \subset \mathcal{H}^k(n, p)$ is indeed the projection of a principal family has been successfully addressed in [3] for $k < n^{1/2-o(1)}$. For larger k , which we are mainly interested in, the problem seems to be more complicated, and has only been studied recently in [14], for constant p . We make no contribution to this question here. However, besides the bounds on $i(\mathcal{H}^k(n, p))$, we are able to show stability for $k = \Theta(n)$ in the same range for p as in case (4) in Theorem 1.2.

Theorem 1.3. *For every $\beta > 0$ and $\varepsilon > 0$ there exist constants $\delta > 0$ and $C > 0$ for which the following holds. For any $\beta n < k(n) < (1/2 - \beta)n$ and $p \geq C \cdot D^{-1}$, asymptotically almost surely stability holds, i.e., for every intersecting family $\mathcal{F} \subset \mathcal{H}^k(n, p)$ of size $|\mathcal{F}| \geq (1 - \delta)p(k/n)N$, there is an element $i \in [n]$ that is contained in all but at most $\varepsilon p(k/n)N$ elements of \mathcal{F} .*

In the dense case, i.e. for $p = 1$, the result was proven by Friedgut [12]. Indeed, the proof of Theorem 1.3 relies on the result of Friedgut and on a removal lemma for the Kneser graph due to Friedgut and Regev [13].

Further results and organization. In proving Theorems 1.2 and 1.3, it will be convenient for us to work with the Kneser graph $K(n, k)$. The vertex set of this graph is $\binom{[n]}{k}$, and two k -element sets form an edge if and only if they are disjoint. Hence $K(n, k)$ is a $\binom{n-k}{k}$ -regular graph on $\binom{n}{k}$ vertices, and a hypergraph $\mathcal{F} \subset \binom{[n]}{k}$ is intersecting if and only if \mathcal{F} is an independent set in $K(n, k)$. Further, let $K(n, k, p)$ denote the subgraph of $K(n, k)$ induced on the random *vertex* set obtained by including each vertex from $\binom{[n]}{k}$ independently with probability p . Due to the correspondence, all bounds on intersecting subgraphs of $\mathcal{H}^k(n, p)$ will follow from corresponding bounds on the size of largest independent sets in $K(n, k, p)$.

Using this translation, Theorem 1.2 follows from a more general scheme which relies on the technical Proposition 2.6 and Lemma 2.2, to be introduced in the next section. Further, for Theorem 1.3 we will need Lemma 2.5, which together with Lemma 2.2 will be proven in Section 3. Based on these results, we will give the proofs of Theorems 1.2 and 1.3 in Section 2.

In general, the proof scheme based on Proposition 2.6 and Lemma 2.2 can be used to bound the size of the largest independent sets in random subgraphs of any D -regular graph G (actually, a sequence of graphs). Here by random subgraph we mean the graph induced on a binomial random subset of the vertex set. This application yields asymptotically sharp bounds if G has an

independent set of size (close to) $-\lambda_{\min}|V(G)|/(D - \lambda_{\min})$. Indeed, Theorem 1.2 shows such an application to the Kneser graph, and there are many other graphs for which this applies. We refer, e.g., to [2] for a list of such graphs which include the weak product of the complete graph, line graphs of regular graphs which contain a perfect matching, Paley graphs, some strongly regular graphs, and appropriate classes of random regular graphs (see Section 5.1. of [2]).

The proof of Proposition 2.6 will be given in Section 4. It is based on a description of all independent sets in locally dense graphs. This idea can be traced back to the work of Kleitman and Winston [18], and has been exploited in various contexts since their work. Though similar proofs have been given elsewhere, none of them seems to fully fit in our context. This also applies to the powerful extension of the ideas of Kleitman and Winston to hypergraphs due to Balogh, Morris and Samotij in [6] (see also [23]), which only partially suits our needs.

2 Proofs of Theorems 1.2 and 1.3

As mentioned before, the proofs of the main theorems rely on Proposition 2.6. A central notion employed in this proposition which applies to $K(n, k)$ is the following.

Definition 2.1. *Given $\lambda \in (0, 1]$, $\gamma \in (0, 1]$, and a graph G on N vertices, we say that G is (λ, γ) -supersaturated if for any subset $S \subseteq V(G)$ with $|S| \geq \lambda N$, we have*

$$e(S) \geq \gamma \left(\frac{|S|}{N} \right)^2 e(G).$$

In addition, let $\lambda = \lambda(n) > 0$ and $\gamma = \gamma(n) > 0$. A sequence $\{G_n\}_{n \in \mathbb{N}}$ is called $(\lambda(n), \gamma(n))$ -supersaturated if G_n is $(\lambda(n), \gamma(n))$ -supersaturated for each $n \in \mathbb{N}$.

Hence, in a (λ, γ) -supersaturated graph G each set S of size at least λN spans many edges. Indeed, up to the multiplicative factor γ , S spans as many edges as expected from a random subset of $V(G)$ of the same size.

Using an extension of Hoffman's spectral bound [16], one can relate supersaturation to the eigenvalues of a graph. We refer to Section 3 for the proof.

Lemma 2.2. *Let G be a D -regular graph on N vertices, and let λ_{\min} denote the smallest eigenvalue of the adjacency matrix of G . Then every set $S \subset V(G)$ satisfies*

$$e(S) \geq \left(\frac{\lambda_{\min} N}{D |S|} + \frac{D - \lambda_{\min}}{D} \right) \left(\frac{|S|}{N} \right)^2 e(G).$$

As the eigenvalues of the Kneser graph are known due to Lovász [21], we immediately conclude the following supersaturation for the Kneser graph.

Lemma 2.3. *Let $2 \leq k \leq n/2$ and $\tau = \tau(n) > 0$. Then $K(n, k)$ is $\left((1 + \tau) \frac{k}{n}, \frac{\tau}{1 + \tau} \right)$ -supersaturated.*

Proof. The Kneser graph $K(n, k)$ has degree $D = \binom{n-k}{k}$, and the smallest eigenvalue of $K(n, k)$ is given by (see [21]):

$$\lambda_{\min} = - \binom{n-k-1}{k-1} = - \frac{k}{n-k} D.$$

Let $S \subset \binom{[n]}{k}$ be of size at least $(1 + \tau)\frac{k}{n}N$, with $N = \binom{n}{k}$. Lemma 2.2 implies that

$$e(S) \geq \left(-\frac{n}{(n-k)(1+\tau)} + \frac{n}{n-k} \right) \left(\frac{|S|}{N} \right)^2 e(G),$$

and the claim follows. \square

Beyond the notion of supersaturation needed for the proof of Theorem 1.2, we will rely on the following notion of robust stability in the proof of Theorem 1.3 (see also [22]).

Definition 2.4. Let $\lambda, \varepsilon, \delta > 0$. Let G be a graph on N vertices, and let $\mathcal{B}(G) \subseteq \mathcal{P}(V(G))$ be a family of sets. We say that G is $(\lambda, \mathcal{B}(G))$ -stable with respect to (ε, δ) if for every $S \subseteq V(G)$ with $|S| \geq (1 - \delta)\lambda N$, we have either

- $e(S) \geq \delta \left(\frac{|S|}{N} \right)^2 \cdot e(G)$, or
- $|S \setminus B| \leq \varepsilon \lambda N$, for some $B \in \mathcal{B}(G)$.

In addition, let $\lambda = \lambda(n) > 0$, $\{G_n\}_{n \in \mathbb{N}}$ be a sequence of graphs, and $\mathcal{B} = \{\mathcal{B}_n\}_{n \in \mathbb{N}}$ with $\mathcal{B}_n \subset \mathcal{P}(V(G_n))$ be a sequence of families of sets. We say that $\{G_n\}_{n \in \mathbb{N}}$ is (λ, \mathcal{B}) -stable if for any $\varepsilon > 0$ there exists $\delta > 0$ and $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, the graph G_n is $(\lambda(n), \mathcal{B}_n)$ -stable with respect to (ε, δ) .

It is instructive to think of $\mathcal{B}(G)$ as the family of largest independent sets in G , and of λN as the size of each $B \in \mathcal{B}$. The first part of the definition roughly says that if G is robustly stable, then any vertex set S whose size is close to the size of a largest independent set in G must either contain many edges, or be close to a largest independent set in structure.

The Kneser graph satisfies robust stability for k linear in n , as stated in the next lemma. It is a direct consequence of the corresponding stability result proven by Friedgut [12], and the removal lemma proven by Friedgut and Regev [13]. Again, we refer to Section 3 for the details of the proof. In the following, let $\mathcal{F}_i \subset \binom{[n]}{k}$ denote the principal hypergraph centered at i , i.e., the hypergraph consisting of all k -element subsets of $[n]$ containing $i \in [n]$.

Lemma 2.5. Let $\beta > 0$ and $k = k(n)$, where $\beta n \leq k \leq (1/2 - \beta)n$, and let $G_n = K(n, k)$. Further, let $\mathcal{B}_n(G_n) = \{\mathcal{F}_i \mid i \in [n]\} \subset \mathcal{P}(V(G_n))$, and set $\mathcal{B} = \{\mathcal{B}_n\}_{n \in \mathbb{N}}$. Then $G = \{G_n\}_{n \in \mathbb{N}}$ is $(k/n, \mathcal{B})$ -stable.

With supersaturation and robust stability defined, we are now ready to state our main technical result. Given a graph H , we use $\alpha(H)$ to denote the size of the largest independent set in H . Also, for a finite set V , we let V_p be a random subset of V obtained by selecting each element $v \in V$ independently with probability p .

Proposition 2.6. Let $\lambda = \lambda(n)$ and $\gamma = \gamma(n)$ be $(0, 1)$ -valued functions, and let $G = \{G_n\}_{n \in \mathbb{N}}$ be a family of graphs, where each G_n has $N = N(n)$ vertices (with $\lim_{n \rightarrow \infty} N(n) = \infty$) and average degree $D = D(n)$. For any constant $0 < \varepsilon < 1$ there exist constants $C = C(\varepsilon) > 0$ and $\delta = \delta(\varepsilon) > 0$ such that for any probability sequence $p = p(n) \in (0, 1]$, the following holds. For a random spanning subgraph $H_n = G_n[V_p]$, where $V = V(G_n)$, we have:

- (i) If $N^{-1} \ll p \ll D^{-1}$, then $\alpha(H_n) = (1 \pm \varepsilon)pN$ asymptotically almost surely.

(ii) If G is (λ, γ) -supersaturated and $9D^{-1} \leq p \leq \lambda^\varepsilon(\lambda\gamma D)^{-1}$, then

$$\mathbb{P}\left(\alpha(H_n) > \frac{4N}{\varepsilon\gamma D} \ln(pD)\right) \leq \exp\left\{-\frac{N}{\gamma D} \ln(pD)\right\}.$$

(iii) If G is (λ, γ) -supersaturated and $p \geq C(\lambda\gamma D)^{-1} \ln^2(e/\lambda)$, then

$$\mathbb{P}(\alpha(H_n) \geq (1 + \varepsilon)\lambda pN) \leq \exp\{-\varepsilon^2 p\lambda N/24\}.$$

(iv) If G is (λ, \mathcal{B}) -stable and $p \geq C(\lambda D)^{-1} \ln^2(e/\lambda)$, then with probability at least $1 - \exp(-\delta^2 \lambda pN/2)$, the following holds: every independent set I in H_n of size at least $(1 - \delta)\lambda pN$ satisfies $|I \setminus B| \leq \varepsilon \lambda pN$ for some $B \in \mathcal{B}_n$.

In addition, the following result will be needed for the lower bound (2) in Theorem 1.2. It is Shearer's extension [25] of a result due to Ajtai, Komlós and Szemerédi [1].

Proposition 2.7 ([1], [25]). *Let $G = \{G_n\}_{n \in \mathbb{N}}$ be a sequence of graphs on $N = N(n)$ vertices with average degree at most $D = D(n) > 1$. If each G_n is triangle-free, then G_n contains an independent set of size $N(D \ln D - D + 1)/(D - 1)^2 \geq N(-1 + \ln D)/D$. \square*

Finally, we shall repeatedly use Chernoff's bound for binomial random variables, which we state here for reference (see [17, Theorem 2.1]).

Lemma 2.8. *Given integers $m, s > 0$ and $\zeta \in [0, 1]$, we have:*

$$\mathbb{P}(\text{Bin}(m, \zeta) \geq m\zeta + s) \leq e^{-s^2/(2\zeta m + s/3)}. \quad (1)$$

$$\mathbb{P}(\text{Bin}(m, \zeta) \leq m\zeta - s) \leq e^{-s^2/(2\zeta m)}. \quad (2)$$

We are now ready to present the proofs of Theorems 1.2 and 1.3.

Proof of Theorem 1.2. Given $0 < \varepsilon < 1$, apply Proposition 2.6 with $\varepsilon/4$ in order to obtain a corresponding constant C_1 . Let $C = \max\{32/\varepsilon^2, 32C_1/\varepsilon\}$. Further, let $k = k(n)$, and $G_n = K(n, k)$. Recall that G_n is a D -regular graph on N vertices, with $D = D(n) = \binom{n-k}{k}$ and $N = N(n) = \binom{n}{k}$. Let $H_n = G_n[V_p]$, where $V = V(G_n)$, and V_p is the set obtained by including each vertex of V independently with probability p . We apply Proposition 2.6 to $\{G_n\}_{n \in \mathbb{N}}$, with functions $N(n)$ and $D(n)$ as defined above. The first bound of Theorem 1.2 follows immediately from the first case of Proposition 2.6.

For the third and fourth bounds of Theorem 1.2, note that by Lemma 2.3 applied with $\tau = \varepsilon/2$, we know that G_n is (λ, γ) -supersaturated, with $\lambda \leq (1 + \varepsilon/2)k/n$ and $\gamma = \varepsilon/4$. Thus we can apply Proposition 2.6 in both cases. We start with the third bound of Theorem 1.2. Assume that $k = o(n)$, since for k linear in n this range of p is trivial. By the second part of Proposition 2.6 applied with $\varepsilon_1 = \varepsilon/2$, we derive that for $9D^{-1} \leq p \leq (n/k)^{1-\varepsilon/2}(\varepsilon D)^{-1}$, which contains our interval for p in the third case, we have

$$i(\mathcal{H}^k(n, p)) < \frac{8}{\varepsilon_1^2} \frac{N}{D} \ln(pD) \leq C \frac{N}{D} \ln(pD)$$

with probability at least $(1 - \exp(-4N \ln(pD)/(\varepsilon D)))$. As $p \gg D^{-1}$, this probability tends to one as n goes to infinity, which gives the upper bound in the third case.

Next we show the fourth bound of Theorem 1.2. The lower bound follows by considering the subhypergraph of $\mathcal{H}^k(n, p)$ consisting of all hyperedges containing, say, the element n . Using the Chernoff bound (Lemma 2.8), we have with high probability that this (intersecting) subhypergraph has size at least $(1 - \varepsilon)p(k/n)N$. For the upper bound, we apply the third bound of Proposition 2.6 with $\varepsilon/4$ and λ, γ as chosen above. Then, by the choice of C , we have $i(\mathcal{H}^k(n, p)) \leq (1 + \varepsilon)\frac{k}{n}pN$ for $p \geq C(n/k)D^{-1} \ln^2(n/k) \geq C_1(\lambda\gamma D)^{-1} \ln^2(e/\lambda)$, and the claim follows.

Finally, we prove the second bound of Theorem 1.2. Observe that this range of p is nontrivial only if $k \ll n$. By Chernoff's bound, almost surely H_n has at least $(1 - \varepsilon/32)pN$ vertices. Further, $\mathbb{E}[e(H_n)] = p^2ND/2$, and it is not hard to see that $\text{Var}[e(H_n)] \leq 2p^3N^2D + p^2ND$. By Chebyshev's inequality, we derive

$$\mathbb{P}(|e(H_n) - \mathbb{E}(e(H_n))| \geq \varepsilon p^2ND/32) \leq \frac{32^2 \text{Var}(e(H_n))}{\varepsilon^2 p^4 N^2 D^2}$$

which goes to zero by the choice of p .

Claim 2.9. *For $(n \ln n)^{1/2} \ll k \ll n$ and $p \leq (n/k)D^{-1}$, asymptotically almost surely the number of triangles in H_n is at most $\varepsilon pN/32$.*

Proof. The expected number of triangles in H_n is at most $p^3 \binom{n}{k} \binom{n-k}{k} \binom{n-2k}{k}$. Using Markov's inequality and $p \leq (n/k)D^{-1}$, the claim follows if we can show that $(n/k)^2 \binom{n-2k}{k} \ll \binom{n-k}{k}$. Indeed,

$$\binom{n-k}{k} \binom{n-2k}{k}^{-1} = \frac{(n-k) \dots (n-2k+1)}{(n-2k) \dots (n-3k+1)} \geq \left(\frac{n-k}{n-2k} \right)^k \geq \left(1 + \frac{k}{n} \right)^k,$$

and using $(1+x) \geq \exp\{x-x^2\}$ for $0 < x < 1$, we obtain together with our assumption $(n \ln n)^{1/2} \ll k \ll n$ that

$$\binom{n-k}{k} \binom{n-2k}{k}^{-1} \geq \exp\{k^2/n - k^3/n^2\} \gg n^2 \geq (n/k)^2,$$

which completes the proof of the claim. \square

Hence, by removing at most $\varepsilon pN/32$ vertices, we obtain a triangle free graph with at least $(1 - \varepsilon/16)pN$ vertices, and no more than $(1/2 + \varepsilon/32)p^2ND$ edges. Consequently, this graph has average degree at most $(1 + \varepsilon/4)pD$, and due to Proposition 2.7, it contains an independent set of size

$$\frac{(1 - \varepsilon/16)pN}{(1 + \varepsilon/4)pD} (\ln((1 + \varepsilon/4)pD) - 1) \geq (1 - \varepsilon) \frac{N}{D} \ln pD.$$

This completes the proof. \square

Proof of Theorem 1.3. Let $\beta > 0$ be fixed, and $\beta n \leq k \leq (1/2 - \beta)n$. Again, let G_n denote the Kneser graph $K(n, k)$. Set $\lambda = k/n$, and for a given n , let \mathcal{B}_n be the set of all principal hypergraphs \mathcal{F}_i , for $i = 1, \dots, n$. By Lemma 2.5, the family $G = \{G_n\}$ is (λ, \mathcal{B}) -stable, where $\mathcal{B} = \{\mathcal{B}_n\}_{n \in \mathbb{N}}$. For a given $\varepsilon > 0$, we apply Proposition 2.6 in order to obtain constants C' and $\delta > 0$. Since $k = \Theta(n)$, it is possible to choose an appropriate constant C such that δ and C satisfy the conclusion of the theorem, which completes the proof. \square

3 Proofs of Lemmas 2.2 and 2.5

As mentioned before, the proof of Lemma 2.2 is a straightforward extension of Hoffman's bound [16].

Proof of Lemma 2.2. Given a D -regular G with N vertices and smallest eigenvalue λ_{\min} , we need to show that for every non-empty $S \subset V(G)$,

$$e_S = e(S) \geq \left(\frac{\lambda_{\min}}{D} \frac{N}{|S|} + \frac{D - \lambda_{\min}}{D} \right) \left(\frac{|S|}{N} \right)^2 e(G).$$

Let M denote the adjacency matrix of G . For $x, y \in \mathbb{R}^N$, let $\langle x, y \rangle = \sum_{i=1}^N x_i y_i$. Also, let v_S be the 0/1-characteristic vector of S . First note that $\langle v_S, M v_S \rangle = 2e_S$. Since M is a symmetric real matrix, it is diagonalizable by an orthonormal basis. Let u_1, \dots, u_N be normalized eigenvectors of M with corresponding eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N = \lambda_{\min}$, respectively. Since G is a D -regular graph, we have $u_1 = (1/\sqrt{N}, \dots, 1/\sqrt{N})$ and $\lambda_1 = D$. Let $v_S = \sum_{i=1}^N a_i u_i$ be the expansion of v_S by eigenvectors. We have

$$2e_S = \langle v_S, M v_S \rangle = \sum_{i=1}^N \lambda_i a_i^2 \geq \lambda_1 a_1^2 + \lambda_{\min} \sum_{i=2}^N a_i^2.$$

Now observe that $a_1 = \langle v_S, u_1 \rangle = |S|/\sqrt{N}$. In addition, $|S| = \langle v_S, v_S \rangle = \sum_{i=1}^N a_i^2$. Therefore,

$$\begin{aligned} 2e_S &\geq D \frac{|S|^2}{N} + \lambda_{\min} \left(|S| - \frac{|S|^2}{N} \right) \\ &= |S| \left(\lambda_{\min} + \frac{|S|}{N} (D - \lambda_{\min}) \right) \\ &= \left(\frac{|S|}{N} \right)^2 2e(G) \left(\frac{\lambda_{\min}}{D} \frac{N}{|S|} + \left(1 - \frac{\lambda_{\min}}{D} \right) \right), \end{aligned}$$

and the lemma follows. \square

We now proceed to show robust stability for the Kneser graph for $k = \Omega(n)$. The proof is a direct consequence of stability due to Friedgut [12] and a removal lemma for the Kneser graph due to Friedgut and Regev [13], which we state next.

Proposition 3.1 (Friedgut [12]). *Given $\beta > 0$, let $k = k(n)$ be a sequence of integers satisfying $\beta n \leq k \leq (1/2 - \beta)n$. For all $\varepsilon > 0$ there exists $\delta > 0$ and n_0 such that, for all $n \geq n_0$, the following holds. If $\mathcal{F} \subseteq \binom{[n]}{k}$ is an intersecting family of size at least $(1 - \delta) \binom{n-1}{k-1}$, then there is $i \in [n]$ such that $|\mathcal{F} \setminus \mathcal{F}_i| \leq \varepsilon \binom{n-1}{k-1}$.*

Proposition 3.2 (Friedgut and Regev [13]). *Given $\beta > 0$, let $k = k(n)$ be a sequence of integers satisfying $\beta n \leq k \leq (1/2 - \beta)n$. Moreover, let $N = \binom{n}{k}$ and $D = \binom{n-k}{k}$. For all $\varepsilon > 0$ there exists $\delta > 0$ and n_0 such that, for all $n \geq n_0$, the following holds. Every family $\mathcal{F} \subseteq \binom{[n]}{k}$ which spans at most $\delta |\mathcal{F}|^2 (D/N)$ non-intersecting pairs can be made intersecting by removing at most $\varepsilon \binom{n-1}{k-1}$ elements from \mathcal{F} .*

Proof of Lemma 2.5. Given any $\varepsilon > 0$, first let $\varepsilon_2 = \varepsilon/2$, and apply Proposition 3.1 to get a corresponding $\delta_2 = \delta_2(\varepsilon_2) > 0$. Now set $\varepsilon_1 = \min(\varepsilon/2, \delta_2/2)$, and use this time Proposition 3.2 in order to obtain an appropriate $\delta_1 = \delta_1(\varepsilon_1) > 0$. Finally, set $\delta = \min(\delta_1, \delta_2/2) = \delta(\varepsilon) > 0$.

It follows that for any family \mathcal{F} with $|\mathcal{F}| \geq (1 - \delta) \binom{n-1}{k-1}$ and $e(\mathcal{F}) \leq \delta(|\mathcal{F}|/N)^2 (ND/2) \leq \delta_1 |\mathcal{F}|^2 (D/N)$ there exists an intersecting family $\mathcal{F}' \subseteq \mathcal{F}$ obtained from \mathcal{F} by removing at most $\varepsilon_1 \binom{n-1}{k-1}$ of its elements such that

$$|\mathcal{F}'| \geq (1 - \delta - \varepsilon_1) \binom{n-1}{k-1} \geq (1 - \delta_2) \binom{n-1}{k-1}.$$

In addition, Proposition 3.1 implies that for some $i \in [n]$, we have $|\mathcal{F}' \setminus \mathcal{F}_i| \leq \varepsilon_2 \binom{n-1}{k-1}$. Therefore,

$$|\mathcal{F} \setminus \mathcal{F}_i| \leq |\mathcal{F} \setminus \mathcal{F}'| + |\mathcal{F}' \setminus \mathcal{F}_i| \leq \varepsilon_1 \binom{n-1}{k-1} + \varepsilon_2 \binom{n-1}{k-1} \leq \varepsilon \binom{n-1}{k-1},$$

which completes the proof. \square

4 Proof of Proposition 2.6

We begin with the proof of a simple structural result for independent sets in graphs (Lemma 4.1). For a given graph G , let $\mathcal{I}_G(t)$ denote the set of independent sets of G of size exactly t , and \mathcal{I}_G denote the set of all independent sets in G .

Lemma 4.1. *Let G be a graph on N vertices, and $\gamma > 0$ be an arbitrary real number. In addition, let $0 < \ell < t$ be integers. Then, for every independent set $I \subset V(G)$ of size at least t , there is a sequence of vertices $x_1, \dots, x_\ell \in I$ and a sequence of subsets $V(G) \supseteq X_1 \supseteq \dots \supseteq X_\ell$ depending only on x_1, \dots, x_ℓ such that:*

- $x_1, \dots, x_i \notin X_i$ for all $i \leq \ell$,
- $I \setminus \{x_1, \dots, x_i\} \subset X_i$ for all $i \leq \ell$.

Moreover, we have either

(i) $|X_i| \leq \left(1 - 2\gamma \frac{e(G)}{N^2}\right) |X_{i-1}|$ for all $1 \leq i \leq \ell$, or

(ii) $e(G[X_i]) < \gamma \frac{|X_i|^2}{N^2} e(G)$ for some $1 \leq i \leq \ell$.

Proof. Fix an independent set I of size at least t . We need to define the required sequences x_1, \dots, x_ℓ and X_1, \dots, X_ℓ . Assume that we have already chosen elements $x_1, \dots, x_{i-1} \in I$ and sets $V(G) = X_0 \supset X_1 \supset \dots \supset X_{i-1}$ satisfying the conditions of our result. Observe that initially no element has been selected, and for convenience we set $X_0 = V(G)$.

Consider an ordering $(v_1, \dots, v_{|X_{i-1}|})$ of the vertices in X_{i-1} which satisfies

$$|N(v_i) \cap \{v_{i+1}, \dots, v_{|X_{i-1}|}\}| \geq |N(v_j) \cap \{v_{i+1}, \dots, v_{|X_{i-1}|}\}|$$

for all $i < |X_{i-1}|$ and all $j > i$. Such an ordering clearly exists, since one can repeatedly choose (and remove) the vertex with highest degree in the remaining graph. In this case we say that this is a *max-ordering* of the elements in X_{i-1} .

Let j be the smallest index such that the vertex v_j in the max-ordering of X_{i-1} is contained in I . Such index must exist, since $I \setminus \{x_1, \dots, x_{i-1}\} \subseteq X_{i-1}$ and $i-1 < \ell < t \leq |I|$. We define $x_i = v_j$, and set $S = X_{i-1} \setminus \{v_1, \dots, v_j\}$.

If $\deg(v_j, S) < 2\gamma|S|e(G)/N^2$ then we let $X_i = S$. Note that, due to the max-ordering and the definition of v_j , every vertex $v \in X_i = \{v_{j+1}, \dots, v_{|X_{i-1}|}\}$ satisfies $\deg(v, X_i) \leq \deg(v_j, X_i)$. This implies that the number of edges in X_i satisfies $e(X_i) < \gamma|X_i|^2e(G)/N^2$. Otherwise, i.e. for the case $\deg(v_j, S) \geq 2\gamma|S|e(G)/N^2$, we let $X_i = S \setminus N(v_j)$. Then,

$$|X_i| \leq |S| - \deg(v_j, S) = \left(1 - 2\gamma\frac{e(G)}{N^2}\right)|S| \leq \left(1 - 2\gamma\frac{e(G)}{N^2}\right)|X_{i-1}|.$$

Finally, observe that it follows from the definition of v_j that we always have $I \setminus \{x_1, \dots, x_i\} \subset X_i$, which completes the proof. \square

From this lemma we immediately deduce the following corollaries.

Corollary 4.2. *Let $G = (V, E)$ be a fixed (λ, γ) -supersaturated graph on N vertices with average degree D , where $\lambda, \gamma > 0$. Let $t \geq 1$, and ℓ be an integer such that $0 < \ell < t$. Finally, set*

$$\nu = \nu(\ell) = \max \left\{ \left(1 - \gamma\frac{D}{N}\right)^\ell, \lambda \right\}.$$

Then, for every independent set $I \in \mathcal{I}_G(t)$, there exists a subset $L \subset I$ of size ℓ and a set $P(L)$, depending only on L , of size at most νN such that $I \setminus L \subset P(L) \subset V(G)$. Further, we have $L \cap P(L) = \emptyset$. In particular, it follows that $|\mathcal{I}_G(t)| \leq \binom{N}{\ell} \binom{\nu N}{t-\ell}$.

Proof. Given $I \in \mathcal{I}_G(t)$, we apply Lemma 4.1 to obtain a sequence of vertices x_1, \dots, x_ℓ and sets $V = X_0, X_1, \dots, X_\ell$, as stated. Now set $L = \{x_1, \dots, x_\ell\}$ and $P(L) = X_\ell$, and observe that $I \setminus L \subset P(L)$ and $L \cap P(L) = \emptyset$.

If $|X_i| \leq \left(1 - 2\gamma\frac{e(G)}{N^2}\right)|X_{i-1}|$ for all $i \leq \ell$, then $|P(L)| \leq \left(1 - 2\gamma\frac{e(G)}{N^2}\right)^\ell N$. In other words, $|P(L)| \leq \left(1 - \gamma\frac{D}{N}\right)^\ell N$. On the other hand, if $e(X_i) < \gamma\frac{|X_i|^2}{N^2}e(G)$ for some $i \leq \ell$, then $|P(L)| \leq \lambda N$, since by assumption G is (λ, γ) -supersaturated. Altogether, it follows that $|P(L)| \leq \nu N$, which completes the proof. \square

Corollary 4.3. *Let $\lambda, \varepsilon, \delta > 0$ and $G = (V, E)$ be graph on N vertices which is (λ, \mathcal{B}) -stable with respect to (ε, δ) . Let $t > \ell \geq \ln\left(\frac{1}{(1-\delta)\lambda}\right)\frac{N^2}{2\delta e(G)}$. Then, for every independent set $I \in \mathcal{I}_G(t)$, there exists a subset $L \subset I$ of size ℓ and a set $P(L) \subset V(G)$ depending only on L such that $I \setminus L \subset P(L)$ and $L \cap P(L) = \emptyset$. Furthermore, either*

- $|P(L)| \leq (1 - \delta)\lambda N$, or
- $|P(L) \setminus B| \leq \varepsilon\lambda N$ for some $B \in \mathcal{B}$.

Proof. We apply Lemma 4.1 to G using $\gamma = \delta$ in order to obtain a sequence of vertices x_1, \dots, x_ℓ and subsets X_1, \dots, X_ℓ with the desired properties. Let $L = \{x_1, \dots, x_\ell\}$. If $|X_i| \leq \left(1 - 2\delta\frac{e(G)}{N^2}\right)|X_{i-1}|$ for all $i \leq \ell$, then set $P(L) = X_\ell$. Using our assumption on ℓ , we get $|P(L)| \leq \left(1 - 2\delta\frac{e(G)}{N^2}\right)^\ell N \leq$

$(1 - \delta)\lambda N$. Otherwise, pick the smallest index $j \leq \ell$ such that $e(G[X_j]) < \delta \frac{|X_j|^2}{N^2} e(G)$, and let $P(L) = X_j$. Again, if $|P(L)| \leq (1 - \delta)\lambda N$ we are done. On the other hand, if this condition does not hold we deduce from the (λ, \mathcal{B}) -stability of G that there exists some $B \in \mathcal{B}$ for which $|P(L) \setminus B| \leq \varepsilon \lambda N$, which completes the proof. \square

Now that we have all the necessary machinery, we proceed with the proof of Proposition 2.6.

Proof of Proposition 2.6. Let $\lambda = \lambda(n)$, $\gamma = \gamma(n)$, and $G = \{G_n\}_{n \in \mathbb{N}}$ be a sequence of graphs. For a given $0 < \varepsilon < 1$, let $C_1 = 800/\varepsilon^3$. For the proof of case (iv) in Proposition 2.6, suppose that G is (λ, \mathcal{B}) -stable for some \mathcal{B} . Then for $\varepsilon' = \varepsilon/2$, there is a constant $\delta' > 0$ and $n_1 \in \mathbb{N}$ such that, for all $n \geq n_1$, the graph G_n is (λ, \mathcal{B}_n) -stable with respect to (ε', δ') . We choose $\delta = \min\{\delta'/4, \varepsilon'/16, 1/16\}$ and $C_2 = 100/\delta^4$. Finally, set $C = \max\{C_1, C_2\}$, and let $n_0 \geq n_1$ be sufficiently large.

We proceed with the proof of the first case of Proposition 2.6. Assume that $N^{-1} \ll p \ll D^{-1}$. Using the Chernoff bound (Lemma 2.8), we have almost surely $|V_p| = (1 \pm \varepsilon/2)pN$, which proves the upper bound. Further, we have $\mathbb{E}(e(H_n)) = \frac{1}{2}NDp^2$ and by Markov's inequality a.a.s. $e(H_n) \leq \varepsilon pN/2$ holds. By deleting at most this number of vertices from H_n , we obtain an independent set of size at least $(1 - \varepsilon)pN$, which proves the lower bound.

For the second part, assume that $9D^{-1} \leq p \leq \lambda^\varepsilon(\lambda\gamma D)^{-1}$. Further, let $\ell = (1 + \varepsilon)\frac{N}{\gamma D} \ln(pD) > 0$, and $t = \frac{4N}{\varepsilon\gamma D} \ln(pD)$. Let X be the random variable counting the number of independent sets of size exactly t in H_n , i.e., $X = |\mathcal{I}_{H_n}(t)|$. By the choice of our parameters, Corollary 4.2 applies, and we obtain:

$$\mathbb{E}[X] \leq \binom{N}{\ell} \binom{\nu(\ell)N}{t - \ell} p^t,$$

where $\nu(\ell) = \max\left\{\left(1 - \gamma\frac{D}{N}\right)^\ell, \lambda\right\}$. Using $\binom{n}{k} \leq \left(\frac{en}{k}\right)^k$ and the choice of ℓ and t , we get

$$\binom{N}{\ell} \leq \left(\frac{e\gamma D}{\ln(pD)}\right)^\ell \quad \text{and} \quad \binom{\nu(\ell)N}{t - \ell} \leq \left(\frac{e\nu(\ell)\gamma D}{\ln(pD)}\right)^{t - \ell}.$$

Combining both inequalities, and noting that our choice of C guarantees that $\ell \leq \varepsilon t/2$, we get:

$$\mathbb{E}[X] \leq \left(\frac{e\gamma p D \nu^{1 - \varepsilon/2}}{\ln(pD)}\right)^t.$$

In case $\nu(\ell) = \lambda$, we have $\gamma p D \nu^{1 - \varepsilon/2} \leq \lambda^{\varepsilon/2} \leq 1$, since $p \leq \lambda^\varepsilon(\lambda\gamma D)^{-1}$. On the other hand, if $\nu(\ell) \leq e^{-\ell\gamma\frac{D}{N}} \leq (pD)^{-1 - \varepsilon}$, we have $\gamma p D \nu^{1 - \varepsilon/2} \leq \gamma(pD)^{-\varepsilon/2 + \varepsilon^2/2} \leq \gamma \leq 1$ since $\varepsilon < 1$. Hence, $\mathbb{E}(X) \leq (e/\ln(pD))^t$, and the claim follows from Markov's inequality.

For the third part, assume that $p \geq C(\lambda\gamma D)^{-1} \ln^2\left(\frac{e}{\lambda}\right)$. Let $t = (1 + \varepsilon)p\lambda N$, and $\ell = \frac{N}{\gamma D} \ln\left(\frac{e}{\lambda}\right)$. We need to upper bound the following probability:

$$q = \mathbb{P}[\exists I \subset V_p, |I| = t, I \text{ is an independent set in } G_n].$$

It follows from Corollary 4.3 that for any $I \in \mathcal{I}_{G_n}(t)$, there exist $L \subset I$ of size ℓ and $P(L)$ such that $I \setminus L \subset P(L) \subset V$. Therefore,

$$q \leq \sum_L \mathbb{P}[L \subset V_p \text{ and } |V_p \cap P(L)| \geq t - \ell].$$

where the sum is taken over all subsets $L \in \binom{V}{\ell}$ that correspond to some independent as given by Corollary 4.3. Using the fact that L and $P(L)$ are disjoint, we obtain

$$q \leq \sum_L \mathbb{P}[L \subset V_p] \cdot \mathbb{P}[|V_p \cap P(L)| \geq t - \ell]. \quad (3)$$

In addition, by our choice of ℓ , it follows that $\nu(\ell) = \lambda$. Therefore, for any such L , we have $|P(L)| \leq \nu(\ell)N \leq \lambda N$. Further, the choice of ℓ and p implies that $\ell \leq (\varepsilon/2)p\lambda N$. Hence with $X = |V_p \cap P(L)|$, we have due to the Chernoff bound that

$$\mathbb{P}(X \geq t - \ell) \leq \mathbb{P}\left(X \geq p|P(L)| + \frac{\varepsilon p \lambda N}{2}\right) \leq \exp\left(-\frac{\varepsilon^2 p \lambda N}{12}\right).$$

From (3) and $\binom{N}{\ell} \leq \left(\frac{eN}{\ell}\right)^\ell$, it follows that:

$$q \leq \left(\frac{eNp}{\ell}\right)^\ell \exp\left(-\frac{\varepsilon^2 p \lambda N}{12}\right) = \exp\left(\ell \cdot \ln\left(\frac{eNp}{\ell}\right) - \frac{\varepsilon^2 p \lambda N}{12}\right).$$

Recall that we want to prove that $q \leq \exp(-\varepsilon^2 p \lambda N/24)$. With the choice $\ell = \frac{N}{\gamma D} \ln(e/\lambda)$ it is now sufficient to show that $\frac{1}{\gamma D} \ln(e/\lambda) \ln\left(\frac{ep\gamma D}{\ln(e/\lambda)}\right) \leq \frac{\varepsilon^2 p \lambda}{24}$, or equivalently

$$\frac{24}{\varepsilon^2 \gamma \lambda D} \ln(e/\lambda) \leq \frac{p}{\ln\left(\frac{ep\gamma D}{\ln(e/\lambda)}\right)}.$$

As the left hand side is independent of p , and the right hand side is increasing in p , it is sufficient to show the inequality for the endpoint $p = C(\lambda\gamma D)^{-1} \ln^2(e/\lambda)$. In this case the inequality follows from $24/\varepsilon^2 \leq C \ln(e/\lambda) / \ln\left(\frac{eC}{\lambda} \ln(e/\lambda)\right)$. Note that $\ln\left(\frac{eC}{\lambda} \ln(e/\lambda)\right) > \ln(e/\lambda) + \ln C$, since $eC/\lambda > \ln(e/\lambda)$. Therefore the bound follows from $48/\varepsilon^2 \leq C \ln(e/\lambda) / (\ln(e/\lambda) + \ln C)$, or equivalently $\frac{48}{\varepsilon^2} \leq \frac{C}{1 + \ln(C)/\ln(e/\lambda)}$. Since the right-hand side is decreasing in λ , it is sufficient to verify for $\lambda = 1$, which is immediate from the choice of C_1 and C .

For the last part, let $p \geq C(\lambda D)^{-1} \ln^2(e/\lambda)$. Further, let

$$\mathcal{T} = \{I \in \mathcal{I}_{G_n} : |I| > (1 - \delta)\lambda p N \text{ and } |I \setminus B| > \varepsilon \lambda p N \text{ for all } B \in \mathcal{B}_n\}.$$

Our task is to upper bound the value of

$$q_{\mathcal{T}} = \mathbb{P}(\text{There is an independent set } I \subset V_p \text{ with } I \in \mathcal{T}).$$

Recall our choice of ε' , δ' , and n_0 , and that G_n is (λ, \mathcal{B}_n) -stable with respect to (ε', δ') for every $n \geq n_0$. We apply Corollary 4.3 with ε' , δ' , $t = (1 - 4\delta)\lambda p N$, and $\ell = \frac{N}{\delta D} \ln \frac{\varepsilon}{\lambda} \leq \delta \lambda p N$. Note that this is a valid choice of ℓ , since $\frac{N}{\delta D} \ln \frac{\varepsilon}{\lambda} \geq \ln\left(\frac{1}{(1-\delta)\lambda}\right) \frac{N^2}{2\delta e(G)}$. This implies that for every $I \in \mathcal{T}$ there is some $L = L(I) \subset I$ of size ℓ and some $P(L) \subset V(G_n)$, depending only on L and disjoint from L , such that $I \setminus L \subset P(L)$. Hence, if there is an $I \subset V_p$ with $I \in \mathcal{T}$, then there is an L of size ℓ with

(A) $L \subset V_p$, and

(B) $|P(L) \cap V_p| \geq (1 - \delta)p\lambda N - \ell \geq (1 - 2\delta)p\lambda N$ and $|(P(L) \setminus B) \cap V_p| > \varepsilon \lambda p N - \ell \geq \frac{3}{4}\varepsilon \lambda p N$ for all $B \in \mathcal{B}_n$, since $\delta \leq \varepsilon'/16 = \varepsilon/32$.

Let $q_{P(L)}$ be the probability that event (B) holds for the random set V_p . As L and $P(L)$ are disjoint, we have

$$q_{\mathcal{T}} \leq \sum_L \mathbb{P}[L \subset V_p] \cdot q_{P(L)}, \quad (4)$$

where the sum ranges over all $L \in \binom{V}{\ell}$ corresponding to some I as given by Corollary 4.3.

From Corollary 4.3 and the chosen parameters, either $|P(L)| \leq (1 - \delta')\lambda N$, or $|P(L) \setminus B| \leq \varepsilon'\lambda N$ for some $B \in \mathcal{B}_n$. Consider each of the cases separately. If $|P(L)| \leq (1 - \delta')\lambda N \leq (1 - 4\delta)\lambda N$ then Chernoff's bound (Lemma 2.8) yields

$$\mathbb{P}(|P(L) \cap V_p| \geq (1 - 2\delta)p\lambda N) \leq \exp\{-\delta^2\lambda p N\}.$$

Similarly, if $|P(L) \setminus B| \leq \varepsilon'\lambda N = \varepsilon\lambda N/2$ for some $B \in \mathcal{B}_n$, then, together with $\delta \leq \varepsilon/32$, we have

$$\mathbb{P}\left(|(P(L) \setminus B) \cap V_p| > \frac{3}{4}\varepsilon\lambda p N\right) \leq \exp\{-\varepsilon\lambda p N/48\} \leq \exp\{-\delta^2\lambda p N\}.$$

Consequently, for every set L as above we have $q_{P(L)} \leq \exp\{-\delta^2\lambda p N\}$.

Hence (4) combined with $\binom{N}{\ell} \leq \left(\frac{eN}{\ell}\right)^\ell$ and the choice of $\ell = \frac{N}{\delta D} \ln(e/\lambda)$ yields

$$q_{\mathcal{T}} \leq \left(\frac{epN}{\ell}\right)^\ell \exp\{-\delta^2\lambda p N\} \leq \exp\left\{\ell \ln\left(\frac{e\delta p D}{\ln(e/\lambda)}\right) - \delta^2\lambda p N\right\}.$$

To complete the proof it suffices therefore to show that $\ell \ln\left(\frac{e\delta p D}{\ln(e/\lambda)}\right) < \delta^2\lambda p N/2$, or equivalently

$$\frac{2 \ln(e/\lambda)}{\lambda \delta^3 D} < \frac{p}{\ln\left(\frac{e\delta p D}{\ln(e/\lambda)}\right)}.$$

As the left hand side does not depend on p , and the right hand side is monotone increasing in p , it is sufficient to verify this inequality for the endpoint $p = C(\lambda D)^{-1} \ln^2(e/\lambda)$. In this case and noting that $e\delta C/\lambda > \ln(e/\lambda)$ due to our choice of C_2 and C , the claim follows from

$$\frac{2}{\delta^3} < \frac{C \ln(e/\lambda)}{\ln\left(\frac{e\delta C}{\lambda} \ln(e/\lambda)\right)} < \frac{C \ln(e/\lambda)}{2 \ln\left(\frac{e\delta C}{\lambda}\right)} = \frac{C}{2 + 2 \ln(\delta C)/\ln(e/\lambda)}.$$

As the right-hand side is decreasing in λ it is sufficient to verify for $\lambda = 1$ which, however, is immediate from the choice of C and C_2 . This completes the proof. \square

5 Concluding remarks

While this work was under review there has been a vivid interest in questions related to random versions of the Erdős-Ko-Rado theorem (cf. [5, 14, 15, 7, 4, 10, 9]). In particular, besides the results of Balogh, Bohman, and Mubayi [3], the question concerning the structure of the largest intersecting family in the random setting has been addressed in [14, 15, 5] for various ranges of k and p . Moreover, an extension of the robust stability result for intersecting families, Lemma 2.5, has been considered in [9], implying that Theorem 1.3 can be extended to a larger range of k . We refer to these papers for further information.

Acknowledgement

The authors are grateful to Yoshiharu Kohayakawa for mentioning the problem and for several helpful comments. Many thanks go to Oded Regev and Ehud Friedgut for sharing a preliminary version of their paper. Finally, we thank the anonymous referee for carefully reading the paper and for many suggestions that improved its readability.

References

- [1] M. Ajtai, J. Komlós, and E. Szemerédi, *A note on Ramsey numbers*, J. Comb. Theory, Ser. A **29** (1980), no. 3, 354–360.
- [2] N. Alon, I. Dinur, E. Friedgut, and B. Sudakov, *Graph products, Fourier analysis and spectral techniques*, Geometric and Functional Analysis **14** (2004), 913–940.
- [3] J. Balogh, T. Bohman, and D. Mubayi, *Erdős–Ko–Rado in random hypergraphs*, Combinatorics, Probability and Computing **18** (2009), no. 5, 629–646.
- [4] J. Balogh, B. Bollobás, and B. Narayanan, *Transference for the Erdős–Ko–Rado theorem*, Forum of Math. Sigma, to appear.
- [5] J. Balogh, S. Das, M. Delcourt, H. Liu, and M. Sharifzadeh, *Intersecting families of discrete structures are typically trivial.*, J. Comb. Theory, Ser. A **132** (2015), 224–245.
- [6] J. Balogh, R. Morris, and W. Samotij, *Independent sets in hypergraphs*, Journal of the American Mathematical Society **28** (2015), no. 3, 669–709.
- [7] B. Bollobás, B. Narayanan, and A. Raigorodskii, *On the stability of the Erdős–Ko–Rado theorem*, J. Combinatorial Theory Ser. A, to appear.
- [8] D. Conlon and W.T. Gowers, *Combinatorial theorems in sparse random sets*, submitted.
- [9] S. Das and T. Tran, *Removal and stability for Erdős–Ko–Rado*, SIAM J. Discrete Math., to appear.
- [10] P. Devlin and J. Kahn, *On stability in the Erdős–Ko–Rado theorem*, submitted.
- [11] P. Erdős, C. Ko, and R. Rado, *Intersection theorems for systems of finite sets*, The Quarterly Journal of Mathematics **12** (1961), no. 1, 313–320.
- [12] E. Friedgut, *On the measure of intersecting families, uniqueness and stability*, Combinatorica **28** (2008), no. 5, 503–528.
- [13] E. Friedgut and O. Regev, *Manuscript*.
- [14] A. Hamm and J. Kahn, *On Erdős–Ko–Rado for random hypergraphs I*, submitted.
- [15] ———, *On Erdős–Ko–Rado for random hypergraphs II*, submitted.
- [16] A.J. Hoffman, *On eigenvalues and colorings of graphs*, Graph Theory and its Applications (Proc. Advanced Sem., Math. Research Center, Univ. of Wisconsin), 1970, pp. 79–91.

- [17] S. Janson, T. Łuczak, and A. Ruciński, *Random graphs*, Wiley-Interscience Series in Discrete Mathematics and Optimization, Wiley-Interscience, New York, 2000.
- [18] D.J. Kleitman and K.J. Winston, *On the number of graphs without 4-cycles*, *Discrete Mathematics* **41** (1982), no. 2, 167–172.
- [19] Y. Kohayakawa, B. Kreuter, and A. Steger, *An extremal problem for random graphs and the number of graphs with large even-girth*, *Combinatorica* **18** (1998), no. 1, 101–120.
- [20] Y. Kohayakawa, S.J. Lee, V. Rödl, and W. Samotij, *The number of sidon sets and the maximum size of sidon sets contained in a sparse random set of integers*, *Random Structures & Algorithms* (2013).
- [21] L. Lovász, *On the Shannon capacity of a graph*, *IEEE Transactions on Information Theory* **25** (1979), no. 1, 1–7.
- [22] W. Samotij, *Stability results for random discrete structures*, *Random Structures & Algorithms* **44** (2014), no. 3, 269–289.
- [23] D. Saxton and A. Thomason, *Hypergraph containers*, *Invent. Math.* **201** (2015), no. 3, 925–992.
- [24] M. Schacht, *Extremal results for random discrete structures*, submitted.
- [25] J.B. Shearer, *A note on the independence number of triangle-free graphs*, *Discrete Mathematics* **46** (1983), no. 1, 83–87.