

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/134047>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Polar relaxation by dynein-mediated removal of cortical myosin II

Bernardo Chapa-y-Lazo,¹ Motonari Hamanaka,^{1,2} Alexander Wray,^{1,3} Mohan K. Balasubramanian,^{1*} and Masanori Mishima^{1*}

¹Centre for Mechanochemical Cell Biology, Warwick Medical School, Gibbet Hill, Coventry CV4 7AL, UK. ²Hokkaido University, Sapporo, Japan. ³University of Nottingham, UK.

*Author for correspondence: M.K.B. (m.k.balasubramanian@warwick.ac.uk) and M.M. (m.mishima@warwick.ac.uk)

Summary/eTOC blurb

Chapa-y-Lazo et al. report that, during anaphase, astral microtubules and dynein remove myosin from the non-equatorial cell cortex and trigger a bidirectional cortical flow towards the equator, which facilitates assembly of the actomyosin contractile ring.

Abstract

Nearly 6 decades ago, Lewis Wolpert proposed the relaxation of the polar cell cortex by the radial arrays of astral microtubules as a mechanism for cleavage furrow induction. While this mechanism has remained controversial, recent work has provided evidence for polar relaxation by astral microtubules, although its molecular mechanisms remain elusive. Here, using *C. elegans* embryos, we show that polar relaxation is achieved through dynein-mediated removal of myosin II from the polar cortexes. Mutants that position centrosomes closer to the polar cortex accelerated

furrow induction whereas suppression of dynein activity delayed furrowing. We show that dynein-mediated removal of myosin II from the polar cortexes triggers a bidirectional cortical flow towards the cell equator, which induces the assembly of the actomyosin contractile ring. These results provide a molecular mechanism for the aster-dependent polar relaxation, which works in parallel with equatorial stimulation to promote robust cytokinesis.

Introduction

During animal cell cytokinesis, cleavage by constriction of an actomyosin contractile ring is spatially coupled to chromosome segregation by the mitotic apparatus (Reymann et al., 2016; Werner et al., 2007; Su et al., 2014; Green et al., 2012; Turlier et al., 2014; Khaliullin et al., 2018; Chen et al., 2008; D'Avino et al., 2015; Salbreux et al., 2009; Dechant and Glotzer, 2003; D'Avino et al., 2006; DeBiasio et al., 1996; Uehara et al., 2016; Zhou and Wang, 2008; Nishimura and Yonemura, 2006; Argiros et al., 2012; Nguyen et al., 2014). The best-understood mechanism for spatial coupling is chemical signaling via activation of the RhoA GTPase at the cell equator by the central spindle, which is mediated by centralspindlin, a microtubule-bundling signaling hub, and ECT2 RhoGEF under the regulation of mitotic kinases (Sedzinski et al., 2011; White and Glotzer, 2012; Wolpert, 1960; Mishima, 2016; Murthy and Wadsworth, 2008; White and Borisy, 1983). A phosphatase-mediated negative signal from the kinetochores has also been reported to reduce the F-actin levels at the polar cortexes (Grill et al., 2001; Rodrigues et al., 2015; Dan, 1954; Pecreaux et al., 2006; Werner et al., 2007; Saunders et al., 2007; Chen et al., 2008; Panbianco et al., 2008; Murthy and Wadsworth, 2008; Nahaboo et al., 2015; Khaliullin et al., 2018; Lee et al., 2015; Maton et al., 2015; Kiyomitsu and

Cheeseman, 2013; Kotak et al., 2012; Schmidt et al., 2017; Fielmich et al., 2018). However, in cells with a relatively small mitotic spindle such as the *Caenorhabditis elegans* one-cell stage embryo, neither positive nor negative signals from the spindle or chromosomes alone can effectively work on the distant cell cortex at the initial stage of cytokinesis before the cleavage furrow deepens (Saunders et al., 2007; Dassow, 2009; Reymann et al., 2016; Mishima, 2016; Lee et al., 2015; Turlier et al., 2014; Nahaboo et al., 2015; Salbreux et al., 2009; Maton et al., 2015; DeBiasio et al., 1996; Zhou and Wang, 2008). In these cells, signals from the astral microtubules, both positive ones to the cell equator and negative ones to the polar/non-equatorial cortexes, play key roles.

Recent studies suggest that the positive signaling to the equator (“equatorial stimulation”) via astral microtubules uses a similar mechanism to the one from the central spindle, mediated by centralspindlin (Verbrugghe and White, 2007; Su et al., 2014; Sedzinski et al., 2011; D’Avino et al., 2006; Werner et al., 2007; Uehara et al., 2016; Bringmann et al., 2007; Nishimura and Yonemura, 2006; Dechant and Glotzer, 2003; Argiros et al., 2012; Nguyen et al., 2014). As another mechanism, the idea of furrow induction by relaxation of the polar cortexes via a signal from the astral microtubules (“astral relaxation/polar relaxation”) has a longer history (Carvalho et al., 2009; Wolpert, 1960; Grill et al., 2001; White and Borisy, 1983; Jordan et al., 2016; Pecreaux et al., 2006; Khaliullin et al., 2018; Saunders et al., 2007; Panbianco et al., 2008; Nahaboo et al., 2015; Lee et al., 2015; Maton et al., 2015; Kiyomitsu and Cheeseman, 2013; Kotak et al., 2012; Schmidt et al., 2017; Fielmich et al., 2018) although early experimental work failed to support this theory (Werner et al., 2007; White and Glotzer, 2012; Saunders et al., 2007; Rappaport, 1996; Reymann et al.,

2016; Mishima, 2016; Lee et al., 2015; Nahaboo et al., 2015; Maton et al., 2015). However, more recently, evidence has been accumulating for the suppression of the contractility of the polar cortexes by radial arrays of astral microtubules (polar relaxation) in various cell types, including *C. elegans* embryos (Wolpert, 1960; Dickinson et al., 2013; Rodrigues et al., 2015; Murthy and Wadsworth, 2008; Verbrugghe and White, 2007; Werner et al., 2007; White and Borisy, 1983; Khaliullin et al., 2018; Bringmann et al., 2007; Dechant and Glotzer, 2003), echinoderm embryos (Dan, 1954; Singh et al., 2019; Dassow, 2009; Werner et al., 2007; Carvalho et al., 2009; Foe and Dassow, 2008; Chen et al., 2008; Jordan et al., 2016; Mishima, 2016; Murthy and Wadsworth, 2008; Khaliullin et al., 2018; Dassow et al., 2009), silkworm spermatocytes (Werner et al., 2007; Reymann et al., 2016; Green et al., 2012; Su et al., 2014; Khaliullin et al., 2018; Turlier et al., 2014; D'Avino et al., 2015; Chen et al., 2008; Dechant and Glotzer, 2003; Salbreux et al., 2009; D'Avino et al., 2006; DeBiasio et al., 1996; Uehara et al., 2016; Zhou and Wang, 2008; Nishimura and Yonemura, 2006; Argiros et al., 2012; Nguyen et al., 2014), and mammalian cultured cells (White and Glotzer, 2012; Sedzinski et al., 2011; Mishima, 2016; Wolpert, 1960; Murthy and Wadsworth, 2008; White and Borisy, 1983). Polar relaxation contributes to furrow formation often by promoting the flow of the actomyosin network within the cell cortex (cortical flow) (Rodrigues et al., 2015; Grill et al., 2001; Dan, 1954; Pecreaux et al., 2006; Werner et al., 2007; Saunders et al., 2007; Chen et al., 2008; Panbianco et al., 2008; Murthy and Wadsworth, 2008; Nahaboo et al., 2015; Khaliullin et al., 2018; Lee et al., 2015; Maton et al., 2015; Kiyomitsu and Cheeseman, 2013; Kotak et al., 2012; Schmidt et al., 2017; Fielmich et al., 2018), which contributes to the assembly of the actomyosin contractile ring (Dassow, 2009; Saunders et al., 2007; Mishima, 2016; Reymann et al., 2016; Lee et

al., 2015; Turlier et al., 2014; Nahaboo et al., 2015; Salbreux et al., 2009; Maton et al., 2015; DeBiasio et al., 1996; Zhou and Wang, 2008) and releases cytoplasmic hydrostatic pressure, which would otherwise destabilize the furrow (Su et al., 2014; Verbrugghe and White, 2007; D'Avino et al., 2006; Sedzinski et al., 2011; Uehara et al., 2016; Werner et al., 2007; Nishimura and Yonemura, 2006; Bringmann et al., 2007; Argiros et al., 2012; Dechant and Glotzer, 2003; Nguyen et al., 2014). However, the molecular mechanisms responsible have remained unclear.

An important action of astral microtubules is to generate the mechanical forces that pull the spindle poles towards the cell cortex through the interaction with the cortically anchored subpopulations of dynein, a multi-functional minus-end directed motor (Wolpert, 1960; Carvalho et al., 2009; White and Borisy, 1983; Grill et al., 2001; Jordan et al., 2016; Pecreaux et al., 2006; Khaliullin et al., 2018; Saunders et al., 2007; Panbianco et al., 2008; Nahaboo et al., 2015; Lee et al., 2015; Maton et al., 2015; Kiyomitsu and Cheeseman, 2013; Kotak et al., 2012; Schmidt et al., 2017; Fielmich et al., 2018). After anaphase onset, more astral microtubules start to grow towards the cell cortex and, in one-cell stage *C. elegans* embryos, interact first with the posterior cortex and then with the anterior cortex to generate cortical pulling forces. These dynein-mediated cortical pulling forces drive asymmetric spindle positioning and spindle elongation in coordination with microtubule-sliding and polymerization at the spindle midzone (White and Glotzer, 2012; Werner et al., 2007; Rappaport, 1996; Saunders et al., 2007; Mishima, 2016; Reymann et al., 2016; Lee et al., 2015; Nahaboo et al., 2015; Maton et al., 2015). Although defects in cortical force generation alone do not prevent cleavage furrow formation, combinations with central spindle defects severely do (Wolpert, 1960; Dickinson et al., 2013; Rodrigues

et al., 2015; Verbrugghe and White, 2007; Murthy and Wadsworth, 2008; Werner et al., 2007; Bringmann et al., 2007; White and Borisy, 1983; Dechant and Glotzer, 2003; Khaliullin et al., 2018). This indicates that a dynein-mediated mechanism contributes to cleavage furrow formation. However, it remains unclear whether this is due to an indirect effect of the altered spindle geometry or due to a more direct influence of dynein and astral microtubules on the activities of the cortical actomyosin network.

In this manuscript, in which the term polar is used to refer to the non-equatorial zone, we report internalization of myosin II from the polar cortexes by astral microtubules and dynein, and investigate the possibility that this activity triggers the polar relaxation and facilitates the contractile ring assembly via the induction of a bidirectional cortical flow towards the cell equator.

Results

Dynein drives unidirectional transport of myosin II along astral microtubules during anaphase

Nonmuscle myosin II is a major component of the cortical actomyosin network and the cytokinetic contractile ring, and is crucial for animal cell cytokinesis. Myosin dynamics in dividing *C. elegans* embryos has previously been studied, but data are limited to a low temporal resolution ($> \sim 10$ s) for a 3D volume (Dan, 1954; Singh et al., 2019; Dassow, 2009; Carvalho et al., 2009; Werner et al., 2007; Jordan et al., 2016; Foe and Dassow, 2008; Khaliullin et al., 2018; Chen et al., 2008; Mishima, 2016; Murthy and Wadsworth, 2008; Dassow et al., 2009) or, to higher temporal resolution, but only at the cell surface (Field and Alberts, 1995; Werner et al., 2007; Oegema et

al., 2000; Reymann et al., 2016; Maddox et al., 2007; Piekny and Glotzer, 2008; Piekny and Maddox, 2010; Tse et al., 2011). To examine the rapid dynamics of myosin II on the mitotic spindle and astral microtubules, we performed fast time-lapse recording (~every second) of GFP-tagged nonmuscle myosin II (NMY-2::GFP), expressed from the endogenous locus (*nmy-2(cp13)*) (Reck-Peterson et al., 2018; Dickinson et al., 2013), imaging a 2 μm thick z-section at the embryo midplane with 0.5 μm z-steps. Embryos within the eggshell were immobilized on the surface of a cover glass without any deformation to avoid the influence of mechanical stress on the cortical actomyosin network (Gönczy et al., 1999; Singh et al., 2019). During metaphase, in addition to accumulation at the cell cortex, NMY-2::GFP was detected on the spindle (**Fig. 1A** metaphase). NMY-2::GFP was also observed as cytoplasmic particulate signals, which showed diffusive random motion and gradually disappeared during early mitosis and were nearly undetectable at the metaphase to anaphase transition (**Fig. 1A** metaphase, **Videos 1 and 2**). By contrast, after anaphase onset, unidirectional movement of cytoplasmic myosin II particles towards the spindle poles was detected (**Fig. 1A** anaphase, **Videos 2 and 3**), with a mean velocity of $\sim 0.7 \mu\text{m/s}$ (**Fig. 1 B and C**). Anillin, a multifunctional cytoskeletal protein that interacts with myosin II and other contractile ring components (Radulescu and Cleveland, 2010; Field and Alberts, 1995; Lorson et al., 2000; Oegema et al., 2000; Maddox et al., 2007; Piekny and Glotzer, 2008; Piekny and Maddox, 2010; Tse et al., 2011) also showed a similar unidirectional movement towards the spindle poles (**Fig. 1D**) although, in this case, such movement was less prominent than that of the particulate signals that are immobile relative to the spindle poles. Overlaying the trajectories of myosin II from multiple embryos revealed a bipolar spatial pattern similar to that of the spindle asters (**Fig. 1E**).

The direction and velocity of the cytoplasmic movement of myosin II particles were consistent with motility by dynein, a microtubule minus-end-directed motor (Firestone et al., 2012; Reck-Peterson et al., 2018). To test the role of microtubules in the myosin II particle movement, embryos were treated with a microtubule depolymerizing drug, nocodazole (**Fig. 2A**). As expected, the cytoplasmic movement of myosin II particles was quickly suppressed after nocodazole treatment (5/5 embryos), while it was not affected at all in control embryos treated with the drug vehicle (4/4). To test the role of dynein in the directional transport of myosin II as its potential cargo, we used RNAi to deplete embryos of the dynein heavy chain DHC-1 (Gönczy et al., 1999) or to deplete LIN-5, the ortholog of NuMA, a dynein regulator (Srinivasan et al., 2003; Radulescu and Cleveland, 2010; Panbianco et al., 2008; Lorson et al., 2000; Galli et al., 2011; Lee et al., 2015) (**Fig. 2B**). We also tested ciliobrevin D (Colombo et al., 2003; Firestone et al., 2012; Srinivasan et al., 2003; Gotta et al., 2003), an inhibitor of vertebrate cytoplasmic dynein, but this did not cause any mitotic or developmental abnormalities (data not shown). No unidirectional cytoplasmic movement of myosin II was observed in the *dhc-1(RNAi)* embryos (7/7 embryos) (**Fig. 2B *dhc-1(RNAi)*, Video 4**), which underwent massively disorganized cell division reflecting the multiple mitotic functions of dynein in centrosome separation, metaphase chromosome congression, spindle positioning and elongation, and chromosome segregation (Redemann et al., 2010; Gönczy et al., 1999; Tse et al., 2011). In *lin-5(RNAi)* embryos, mitosis proceeded more normally except for expected defects in spindle positioning and elongation (Nishikawa et al., 2017; Srinivasan et al., 2003; Panbianco et al., 2008; Galli et al., 2011; Lee et al., 2015). As in the case of *dhc-1(RNAi)* embryos, no cytoplasmic unidirectional

movement of myosin II particles was detected in the embryos depleted of LIN-5 (33/33 embryos) (**Fig. 2B** *lin-5(RNAi)*, **Video 5**) or its binding partners GPR-1/2, the Pins/LGN orthologs (Audhya et al., 2005; Colombo et al., 2003; Srinivasan et al., 2003; Gotta et al., 2003) (movies not shown, quantitative analysis in Fig. 7C to be described in a later section). These results indicate that the movement of myosin II from the cortex towards the spindle poles depends on astral microtubules and is driven by dynein.

Interestingly, we observed a significant delay in the furrow formation in the embryos depleted of LIN-5 or GPR-1/2 and thus lacking the internalization of the myosin II particles (**Fig. 2C**). While the first sign of furrow ingression was detected by visual inspection at ~150 s post anaphase onset (p.a.o.) in control embryos, it was delayed by more than 40 s in the *lin-5(RNAi)* and *gpr-1/2(RNAi)* embryos (**Fig. 2C**). These data are consistent with a positive role of dynein-mediated myosin II transport in cleavage furrow formation.

Myosin II removal from the cortex by dynein and astral microtubules is associated with the local relaxation of the cortical actomyosin network

The aster-dependent unidirectional motion of myosin II particles towards the spindle poles suggests that they are derived from the myosin II population that has been accumulated at the cell cortex but then has been removed from it by dynein and the astral microtubules. It has been reported that dynein and the astral microtubules cause invagination of the plasma membrane during anaphase, which becomes exaggerated upon disruption of the cortical actomyosin network by depletion of myosin II or by treatment with cytochalasin D, an actin inhibitor (Nishikawa et al.,

2017; Redemann et al., 2010; Tse et al., 2011). To examine the relationship between the internalization of myosin II and the invagination of the plasma membrane, we simultaneously observed NMY-2::tagRFP-T (myosin II tagged with a red fluorescent protein) (Kozlowski et al., 2007; Nishikawa et al., 2017) and PH(PLC δ 1)::GFP (a plasma membrane marker) (Mayer et al., 2010; Audhya et al., 2005) in the midplane of an embryo in anaphase. Some membrane invaginations were led by a myosin II particle (**Fig. 3A** and **Video 6**, arrow) while no myosin II peak was detected at the leading tip of other invaginations (**Fig. 3A** and **Video 6**, arrowhead). Colocalization of myosin II and the PH domain was detected on some cytoplasmic particles (**Fig. 3B** and **Video 7**, white arrowhead) but not on other particles (**Fig. 3B** and **Video 7**, magenta and green arrowheads). We examined 8 PH/Myosin dual-labeled embryos and observed 20 myosin II particles negative for the PH domain signal, 31 membrane signals positive for the PH domain but negative for myosin II, and 8 double positive particles. According to these observations, it is likely that dynein and astral microtubules act on the plasma membrane and the myosin II particles through distinct mechanisms. Although it is possible that internalization of other factors might play a role in the regulation of cytokinesis, we focused on the removal of the myosin II particles from the cell cortex in this work.

To further examine the action of the astral microtubules on the cortical actomyosin network, we next observed embryos expressing both NMY-2::tagRFP-T (Dechant and Glotzer, 2003; Nishikawa et al., 2017) and tubulin::YFP (Motegi et al., 2006; Kozlowski et al., 2007; Verbrugghe and White, 2007; Bouvrais et al., 2018) (**Fig. 3 C-F, Video 8**). We detected the internalization of single myosin II particles from the cell cortex at the site of contact with single astral microtubule fibers (**Fig. 3 C and E**,

Videos 9 and **10**, posterior and anterior cortexes, respectively). A particle (red arrowheads) traveled along an astral fiber towards the associated spindle pole, although we usually failed to track the particle until it reached the pole due to the focal drift of the spindle pole or the disappearance of the particle. Following the internalization of a myosin II particle, gradual separation of flanking peak signals of the cortical myosin II was observed (**Fig. 3 D and F, Videos 9 and 10**, blue arrowheads). Although it was difficult to systematically and statistically assess the frequency of this phenomenon, in 20 embryos, we could detect 39 cases of internalization, 30 of which were followed by local cortical relaxation. This was reminiscent of the cortical relaxation that is observed after laser ablation (Harris and Gewalt, 1989; Mayer et al., 2010) (**Fig. S1**). These observations suggest that the removal of myosin II from the cortex leads to a local reduction of the cortical tension/contractility.

Spindle pole positioning during anaphase generates a local minimum of the aster-cortex interaction at the future cleavage site

Trajectories of the myosin II particles overlaid across embryos illustrate that the internalization of myosin occurs less frequently around the equatorial zone (**Fig 1E**). To test whether the geometry of the anaphase spindle accounts for this spatial pattern of myosin II internalization, we examined the global distribution of the astral microtubules that reach the cell cortex. A local minimum of the microtubule density at the equatorial future cleavage site has previously been demonstrated in fixed and immunostained *C. elegans* one-cell stage embryos (Dechant and Glotzer, 2003). However, variable results have been reported by live imaging of microtubules within a plane near the cell surface of embryos that were flattened by being pressed onto a

coverglass with an overlaid agarose pad (Bouvrais et al., 2018; Motegi et al., 2006; Verbrugghe and White, 2007). To avoid the possible influence of the deformation of the embryos on the microtubule densities (Dechant and Glotzer, 2003; Harris and Gewalt, 1989; Redemann et al., 2017), we observed the embryos immobilized on coverglass surfaces without overlaying agarose pads. Recording of the midplane instead of the flattened surface allowed us to simultaneously observe the spindle poles and the astral microtubules that reach the cell cortex (**Fig. 4A**).

To quantify the microtubule density near the cell cortex, we measured the line profile of the YFP-tubulin signal along a curve 1 μm inside the cell boundary (**Fig. S2**). The height of the peaks that correspond to astral microtubules is very low in comparison with the background signals derived from unpolymerized tubulin, which itself is noisy and variable depending on the location, prohibiting a simple background subtraction (**Fig. 4A**). Thus, we calculated the signals that were significantly higher than the local background level (**Fig. S2**). The average across 9 embryos is shown in **Fig. 4B**. Reflecting the posterior shift of the spindle after anaphase onset, the density of the microtubules first started to increase at the posterior cortex, creating a global minimum at the anterior tip of the embryo. Then, a local minimum at a site slightly posterior to the equator became clearer around 60 s after anaphase onset and was stably maintained thereafter (**Fig. 4B**, green arrows). This spatial pattern is very similar to what was reported by immunostaining (Jiang et al., 1998; Dechant and Glotzer, 2003; Mollinari et al., 2002; Verbrugghe and White, 2004; Subramanian and Kapoor, 2012), and the spatio-temporal dynamics is consistent with the recent data by live imaging of flattened cell surfaces (Verbrugghe and White, 2004; Bouvrais et al., 2018; Lee et al., 2015).

Astral microtubules in anaphase have been reported to show an exponential length distribution (Mishima et al., 2002; Dechant and Glotzer, 2003; Davies et al., 2015; Redemann et al., 2017). The appearance of the local minimum of the microtubule density at the equatorial cortex described above might be explained simply as a consequence of the distance to the nearby spindle pole. We detected the position of the spindle poles as faint signals of myosin II-GFP (**Fig. 4C, yellow arrowheads**). The posterior pole approaching the posterior cortex created a local minimum in the pole-to-cortex distance on that tip of the embryo (**Fig. 4D, 20~60s**), corresponding to the rise of microtubule density on the posterior cortex that precedes the one on the anterior cortex. This was followed by the appearance of a local maximum in the pole-to-cortex distance at the equatorial region ($\sim 3 \mu\text{m}$ posterior from the exact center of the A-P axis (**Fig. 4D, green arrows**)), which gradually grew as the anterior spindle pole started to move towards the anterior tip (40~70 s), and predicted the site of cleavage furrow ingression. These observations indicate that the positioning of the spindle poles relative to the cell surface is the primary factor that determines the density of astral microtubules that can influence the cortical activity.

Promotion of the aster-polar cortex interaction facilitates furrow formation

So far, we have shown that astral microtubules and dynein remove myosin II preferentially from the polar/non-equatorial cortexes and that removal of myosin II from the cortex is frequently associated with local reduction of the cortical tension/contractility. This might provide a mechanism for the polar relaxation, which facilitates the equatorial accumulation of actomyosin. To test this idea, we examined the influence of modifying the geometry of the anaphase spindle relative to the cell

surface on the behavior of the cortical actomyosin network.

First, to enhance the aster-cortex interaction, we disrupted the central spindle, which mechanically links the two spindle poles and thus opposes spindle pole separation. We did this by depleting SPD-1, the ortholog of vertebrate PRC1, a highly conserved microtubule-bundling protein crucial for the central spindle formation (Jantsch-Plunger et al., 2000; Jiang et al., 1998; Mollinari et al., 2002; Verbrugghe and White, 2004; Subramanian and Kapoor, 2012) (**Fig. 5**). In *spd-1(RNAi)* embryos, as expected, the anterior and posterior asters were separated from each other immediately after anaphase onset as a consequence of the rupture of the mitotic spindle (**Fig. 5 A and B**) (Grill et al., 2001; Verbrugghe and White, 2004; Srayko et al., 2005; Lee et al., 2015; Han et al., 2015; Rankin and Wordeman, 2010). This accelerated the approaching of both the poles to the cortex, positioning both of them closer to the cortex than the final levels in the control embryos already within 40 s after anaphase onset (**Fig. 5B**). This established a deeper and wider minimum in the pole-to-cortex distance pattern in the anterior cortex, which covered the anterior tip of the embryos, earlier than in control embryos (**Fig. 5A vs Fig. 4D**, anterior zones in orange). As anticipated, the cytoplasmic transport of the myosin II particles was elevated in the *spd-1(RNAi)* embryos (**Fig. 5 C and D**, for details see **Materials and Methods and Fig. S3**) and reached a plateau earlier than in the control embryos. Importantly, despite the disruption of the central spindle, an important source of a positive signal for the contractile ring assembly, the initiation of cleavage furrow formation was accelerated by 50 s in the *spd-1(RNAi)* embryos (**Fig. 2C**).

Slight acceleration of initial furrow formation was also observed when the central

spindle was disrupted by depletion of CYK-4, the GAP component of centralspindlin, which is crucial for the microtubule-bundling activity as well as for signaling (O'Rourke et al., 2010; Mishima et al., 2002; Davies et al., 2015), although in this case the shallow furrow failed to deepen and regressed (**Fig. 2C** and **Fig. 6A**) (Khaliullin et al., 2018; Jantsch-Plunger et al., 2000; Reymann et al., 2016; Sugioka and Bowerman, 2018). This indicates that the acceleration of the initial furrow formation by disruption of the central spindle does not depend on the centralspindlin-mediated equatorial stimulation. A similar but milder acceleration of furrow initiation was observed by milder acceleration of the pole-to-pole separation (spindle elongation) by depletion of KLP-7, a microtubule depolymerizer that targets astral microtubules (Munro et al., 2004; Grill et al., 2001; Werner et al., 2007; Srayko et al., 2005; Han et al., 2015; Rankin and Wordeman, 2010), or by depletion of EFA-6, a negative regulator of the dynein-dependent aster-cell cortex interaction (Grill et al., 2001; O'Rourke et al., 2010; Srayko et al., 2005; Lee et al., 2015) (**Fig. 2C**). These observations further support the role of dynein-mediated myosin II internalization by astral microtubules in the induction of the cleavage furrow.

Myosin II removal from the polar cortexes triggers a bidirectional cortical flow towards the cell equator

Next, we investigated how dynein-mediated internalization of myosin II controls cleavage furrow formation. In order to track cortical myosin II dynamics in relation to the geometry of the spindle, we measured the NMY-2::GFP signal along the cell periphery from the anterior tip to the posterior tip at each time point (for details see **Materials and Methods** and **Fig. S4**). The average across embryos was displayed as a kymograph representing the dynamics of the distribution of the cortical myosin II

(**Fig. 5E**). Using the same data, the cortical flow along the cell periphery was computed in individual embryos by one-dimensional particle image velocimetry (PIV), which gave comparable values to the anterior-posterior component of the flow velocity obtained by two-dimensional PIV (Rose et al., 1995; Khaliullin et al., 2018; Tse et al., 2012; Reyman et al., 2016; Zhang and Glotzer, 2015; Sugioka and Bowerman, 2018; Maniscalco et al., 2020). The average across embryos was displayed as a kymograph in which the green signal indicates flow from the anterior to the posterior, and the magenta indicates flow from the posterior to the anterior (**Fig. 5F**).

In normal embryos, during metaphase, myosin II showed an asymmetric cortical localization slightly enriched at the anterior cortex (**Fig. 5E**, control, 0 s), as a remnant of post-fertilization polarization (Tse et al., 2012; Munro et al., 2004; Werner et al., 2007). After anaphase onset, the signal in the posterior cortex gradually increased and subsequently the global signal reached a uniform distribution at around 20 s p.a.o. (**Fig. 5E**, control, 0~20 s). This state was maintained until 40 s p.a.o., when the posterior pole approached within $\sim 10 \mu\text{m}$ from the posterior cortex, forming a minimum at the posterior pole (**Fig. 4D and Fig. 5B**, P-pole). At this point, the signal in the posterior cortex started to decline (**Fig. 5E**, control, ~ 40 s) and, simultaneously, a flow from the posterior to the anterior appeared, which gradually became stronger and extended towards the anterior cortex until ~ 100 s p.a.o. (**Fig. 5F**, control, 40~100 s, the magenta signal on the right half of the panel). This resulted in a broad and moderate accumulation of myosin II at the equatorial zone (**Fig. 5E**, control, ~ 100 s p.a.o.). By contrast, the signal on the anterior cortex stayed largely constant and without a strong flow (**Fig. 5E and F**, control, 40~100 s, the left

half of the panels). Meanwhile, the anterior spindle pole started to move slowly towards the anterior tip of the embryo (**Figs. 4D** and **5B**, A-pole). Following this period, around 100 s p.a.o., a drop in the myosin II signal was observed in the anterior cortex (**Fig. 5E**), concomitantly with the appearance of a strong flow towards the posterior (**Fig. 5F**, control, >100 s, the green signal on the left half of the panel). This resulted in a period of bidirectional flow, in which the flow towards the anterior from the posterior cortex converged with the flow towards the posterior from the anterior cortex at the cell equator. The bidirectional flow gradually sharpened the equatorial zone of myosin II accumulation (**Fig. 5E**, control) and resulted in furrow formation around 150 s. p.a.o (**Fig. 2C**).

As described above, the depletion of SPD-1 accelerated the approach of the centrosomes to the polar cortexes and promoted the dynein-mediated internalization of myosin II. This caused a drastic change in the dynamic behavior of myosin II at the cortex. Following the earlier appearance of the anterior minimum in the pole to cortex distance (**Fig. 5A and B**), a drop of myosin II signal at the anterior cortex was observed at 60 s p.a.o. (**Fig. 5E**, *spd-1(RNAi)*, 60 s p.a.o.), accompanied by an earlier rise of a stronger posterior-directed flow in the anterior cortex (**Fig. 5F**, *spd-1(RNAi)*). This accelerated entry into the phase of bidirectional flow by ~50 s (**Fig. 5F**, dashed white lines) and resulted in earlier and stronger accumulation of myosin II at the equator (**Fig. 5E**) together with furrow formation ~60 s earlier than in control embryos (**Fig. 2C**). Consistent with this, in the *cyk-4(RNAi)*, *efa-6(RNAi)* and *klp-7(RNAi)* embryos, in which the approaching of the spindle poles to the polar cortex is mildly accelerated for different reasons (Rose et al., 1995; Grill et al., 2001; Tse et al., 2012; Srayko et al., 2005; O'Rourke et al., 2010; Lee et al., 2015), the bidirectional

flow (**Fig. 6**) and the furrow ingression (**Fig. 2C**) occurred 20~30 s earlier than in control embryos. These observations are consistent with our hypothesis that asters mediate cortical relaxation by removing myosin II from the non-equatorial cortex, thus prompting flow towards the equator.

Centralspindlin and NOP-1, two upstream regulators of Rho GTPase, regulate the cortical flow in different ways

A nematode specific protein, NOP-1, has been reported to be another upstream activator of the RhoA GTPase and to work independently of the centralspindlin-ECT2 pathway (Verbrugghe and White, 2007; Rose et al., 1995; Werner et al., 2007; Tse et al., 2012; Bringmann et al., 2007; Zhang and Glotzer, 2015; Maton et al., 2015; Maniscalco et al., 2020; Dechant and Glotzer, 2003). In contrast to centralspindlin, it shows a diffuse localization throughout the cell with slight enrichment at the entire cortex and strong accumulation to the interphase nuclei (Canman et al., 2008; Tse et al., 2012; Davies et al., 2014). Consistent with its function as a positive regulator of Rho, depletion of NOP-1 significantly delayed furrow initiation (**Fig. 2C**) although all the embryos tested could complete cytokinesis as previously reported (Canman et al., 2008; Rose et al., 1995; Zhuravlev et al., 2017; Tse et al., 2012). Cortical flow, especially in the posterior cortex, was decreased in the *nop-1(RNAi)* embryos (**Fig. 6B**), indicating that efficient cortical flow requires the global activity of Rho, which partially depends on NOP-1.

Polar relaxation by dynein-mediated removal of cortical myosin II works in parallel with the equatorial stimulation

It has been shown that combinations of the depletion of the dynein regulators LIN-

5/NuMA or GPR-1/2/Pins/LGN with defects in the central spindle factors cause synthetic failure of furrow formation (Zhang and Glotzer, 2015; Verbrugghe and White, 2007; Tse et al., 2012; Werner et al., 2007; Loria et al., 2012; Bringmann et al., 2007; Maton et al., 2015; Dechant and Glotzer, 2003). These observations indicate that dynein and dynein regulators work in the furrow induction in parallel with the central spindle/centralspindlin-dependent pathway. However, it has remained unclear whether dynein and its regulators positively control cortical contractility by equatorial stimulation or negatively by polar relaxation.

Here we have found that the depletion of the dynein regulators LIN-5/NuMA and GPR-1/2/Pins/LGN completely eliminates the dynein-driven internalization of myosin II from the cell cortex (**Fig. 2B and 7C**). Since our results suggest that the removal of myosin II from the cortex leads to local cortical relaxation, it is likely that these dynein regulators work in polar relaxation rather than in equatorial stimulation. To test this and to clarify the contributions of the dynein-dependent and centralspindlin-dependent pathways to the furrow induction, we assessed the effect of the depletion of GPR-1/2 on the dynamics of cortical myosin II and examined the synthetic effect with the inhibition of the centralspindlin-dependent equatorial stimulation observed in *cyk-4(or749)* mutant embryos (**Fig. 7**). The point mutation of this temperature-sensitive allele (designated as *cyk-4(GAP)* hereafter) lies in the GAP domain of CYK-4 (Canman et al., 2008; Davies et al., 2014) and prevents the equatorial stimulation because the GAP domain plays a role in equatorial stimulation by inactivation of Rac (Wolpert, 1960; Canman et al., 2008; Zhuravlev et al., 2017) or by activation of Rho via the RhoGEF ECT-2 (Reymann et al., 2016; Zhang and Glotzer, 2015; Turlier et al., 2014; Tse et al., 2012; Salbreux et al., 2009; Loria et al., 2012;

DeBiasio et al., 1996; Zhou and Wang, 2008; Werner et al., 2007). Importantly, unlike other methods used to perturb the centralspindlin/central spindle-dependent pathway, such as *cyk-4(RNAi)* (**Figs. 2C, 6A and 6B**), the *cyk-4(GAP)* mutation, at the restrictive temperature, achieves this without affecting the formation and maintenance of the central spindle or the localization of centralspindlin to the spindle midzone (Su et al., 2014; Canman et al., 2008; D'Avino et al., 2006; Uehara et al., 2016; Nishimura and Yonemura, 2006; Argiros et al., 2012; Nguyen et al., 2014).

In the embryos with normal CYK-4 but depleted of GPR-1/2, myosin II showed hyperaccumulation to the posterior cortex during early anaphase (**Fig. 7A iii**, *gpr-1/2(RNAi)*, “density”, and **Fig. 7B** *gpr-1/2(RNAi)*), confirming the role of dynein-driven cytoplasmic transport of myosin II, which is almost completely eliminated in *gpr-1/2(RNAi)* embryos (**Fig. 7C** *gpr-1/2(RNAi)*), in the removal of myosin II from the cell cortex (polar relaxation). This was accompanied by a reversal of the global flow that lasts until 120 s p.a.o. (**Fig. 7A iii**, *gpr-1/2(RNAi)*, “flow”) and by slightly delayed furrow ingression (**Fig. 7D**, *gpr-1/2*). Although the bidirectional flow appeared after 120 s p.a.o., the posterior-to-anterior flow towards the equator in the posterior cortex was much slower than in the wild type embryos (**Fig. 7A**, “flow”, iii *gpr-1/2(RNAi)* vs i wild type). By contrast, in the *cyk-4(GAP)* embryos with a normal level of GPR-1/2, no obvious difference in the cytoplasmic transport of myosin II particles (**Fig. 7C**) or in its distribution and flow at the cortex was observed (**Fig. 7A**, ii *cyk-4(GAP)* vs i wild type), indicating a limited role of the centralspindlin-dependent equatorial stimulation in initial formation of the bidirectional cortical flow.

In the embryos that were defective for both the centralspindlin-dependent and dynein-dependent pathways (**Fig. 7A, iv** *cyk-4(GAP) gpr-1/2(RNAi)*), myosin II showed hyper-accumulation at the posterior cortex in a similar manner to that observed in the *gpr-1/2(RNAi)* embryos, with a slight enhancement of the peak height and a delayed clearance (**Fig. 7A, iii vs iv, density, and 7B**). Initially, the entire cortex showed a flow towards the posterior side, where myosin II was enriched, indicating that the contractility per se was not eliminated. Interestingly, however, in these double mutant embryos, the bidirectional flow towards the equator was almost completely abolished (**Fig. 7A, iv, flow**), suggesting that the mechanisms for organizing the contractility into properly patterned flows were perturbed. These results indicate that the weak bidirectional cortical flow observed in the *gpr-1/2(RNAi)* embryos in the presence of normal centralspindlin was caused by the centralspindlin-dependent equatorial stimulation. While all the embryos defective only for the centralspindlin-dependent pathway or the dynein-dependent pathway could initiate furrowing, albeit with some delays, about half of the embryos defective for both pathways failed to form a furrow (**Fig. 7D**). The rest of the embryos could only form a very shallow furrow, which eventually regressed after a long delay. The depletion of LIN-5 also prevented furrow formation or deepening in the *cyk-4* mutant background (**Fig. S5**). Taken together, these data indicate that, in the *C. elegans* one-cell stage embryo, the polar relaxation triggered by dynein-driven myosin II transport contributes to the furrow induction in parallel with the centralspindlin-dependent equatorial stimulation, especially during the early establishment of the bidirectional cortical flow (**Fig. 8**) although our data do not exclude the possibility that dynein might also contribute to the equatorial stimulation by an unknown mechanism.

Discussion

Since the first proposal of the polar relaxation by astral microtubules (Hutterer et al., 2009; Wolpert, 1960; Douglas et al., 2010), its molecular mechanism has remained unclear. Here we present evidence for a fundamental mechanism in which centrosome-directed transport of myosin II along astral microtubules, driven by dynein, reduces the contractility of the polar/non-equatorial cortexes by removing myosin II from them (**Fig. 8**). The local reduction of cortical contractility triggers a global cortical flow, and the geometry of the mitotic apparatus plays a key role in switching the flow from a unidirectional mode to a bidirectional mode. The bidirectional cortical flow towards the cell equator contributes to the equatorial accumulation of the actomyosin network and to the cleavage furrow formation (Lekomtsev et al., 2012; Reymann et al., 2016; Basant et al., 2015; Turlier et al., 2014; Salbreux et al., 2009; DeBiasio et al., 1996; Zhou and Wang, 2008; Werner et al., 2007).

In addition to the central spindle, the molecules responsible for equatorial stimulation, such as centralspindlin, can be recruited to the anti-parallel overlaps of equatorial astral microtubules (Adriaans et al., 2019; Su et al., 2014; D'Avino et al., 2006; Uehara et al., 2016; Nishimura and Yonemura, 2006; Argiros et al., 2012; Nguyen et al., 2014). Association of the equatorial pools of centralspindlin with the plasma membrane, which is enhanced by its oligomerization, regulated by Aurora B kinase and 14-3-3 proteins (Hutterer et al., 2009; Douglas et al., 2010), has been shown to promote RhoA activation and the tethering of the plasma membrane at the midbody (Lekomtsev et al., 2012; Basant et al., 2015). Based on these, the accelerated furrowing that follows the rupture of the central spindle by SPD-1/PRC1 depletion

might be explained by an earlier release of centralspindlin from the central spindle onto the equatorial asters and the plasma membrane (Adriaans et al., 2019). However, the accelerated induction of a bidirectional cortical flow and furrow initiation by depletion of the CYK-4 GAP subunit (**Figs. 2C** and **6B**) is difficult to explain by this scenario. Our observations rather indicate that the promotion of the aster-cortex interaction can facilitate the furrow formation via the bidirectional cortical flow independently of the equatorial stimulation by centralspindlin; however, we cannot exclude the possibility that a positive signal is released from the central spindle upon its rupture (Baruni et al., 2008; Adriaans et al., 2019) and delivered to the cortex via astral microtubules by an unknown motor protein (Atilgan et al., 2012) earlier than normal.

The equatorial microtubule structures that recruit centralspindlin during natural cytokinesis are organized in an anti-parallel manner and placed lateral to the plasma membrane. In experimentally induced monopolar cytokinesis, the astral microtubules that asymmetrically recruit the midzone proteins to induce furrowing are also laterally associated with the furrow cortex although they are often bundled in a parallel configuration. Here we have shown that the astral microtubules that contact the cell cortex with their plus ends can induce local cortical relaxation by removing myosin II from the cortex. It has also been reported that precise microtubule polymerization/depolymerization dynamics does not play a crucial role in furrow induction (Strickland et al., 2005). Considering these observations, we propose a rule of thumb for the action of microtubules on the nearby cell cortex: microtubules laterally associated with the cortex, which are often in an anti-parallel configuration, promote contractility, whereas microtubules pointing towards the cortex relax

contractility. It remains unclear whether and how unbundled microtubules approaching the cortex at a shallow angle, which are observed at the cell equator in early anaphase, control cortical contractility.

How myosin II is linked to dynein is currently unclear. Whether and how regulators of the cortical pulling forces (Sugioka and Bowerman, 2018; Schmidt et al., 2017; Fielmich et al., 2018; Rodriguez-Garcia et al., 2018; Kiyomitsu and Cheeseman, 2013; Kotak et al., 2014) and cortical asymmetry (Chartier et al., 2011; Pacquelet et al., 2015; Sugioka et al., 2018), as well as membrane additions (Gudejko et al., 2012) or invaginations/endocytosis (Tse et al., 2011; Redemann et al., 2010), are involved in dynein-driven transport of myosin II and regulation of other cortical activities (and, vice versa) will be important future questions. Although myosin II showed hyper-accumulation to the posterior cortex in the absence of the dynein activities, it was gradually cleared off, albeit with a significant delay, indicating the presence of other mechanisms for clearing myosin II from the cortex outside of the cleavage furrow. Another interesting question will be how the dynein-driven removal of myosin II cooperates with or is regulated by mitotic kinases and phosphatases or GTPases that control the contractile actomyosin networks, such as Aurora A, which accumulates at the centrosomes and breaks the symmetry of the cell cortex during zygotic polarity establishment and cytokinesis (Zhao et al., 2019; Mangal et al., 2018; Klinkert et al., 2019).

Based on the model we propose, we can say that compared to a mechanism that depends on a diffusive signal, direct remodeling of the actomyosin network through a physical interaction with microtubules has the advantage of keeping the effect

precisely localized at a long distance. We speculate that the dynein-driven myosin II transport mechanism revealed in this work may have a broader biological role in controlling cell shape in other cell/tissue contexts, such as in cell migration and epithelial morphogenesis, since both dynein and myosin II are universally expressed in metazoan cells.

Acknowledgments

We thank Behrooz Esmaeili for help in *C. elegans* maintenance, Rob Cross for useful comments on the text, and CAMDU (Computing and Advanced Microscopy Unit, Warwick Medical School) for support on imaging. We thank Stephan Grill, Julie Canman, Tim Davies and the Caenorhabditis Genetics Center for providing worm strains. This work was supported by the Wellcome Trust Senior Investigator Award (WT101885MA) and European Research Council Advanced Grant (ERC-2014-ADG No. 671083) to M.K.B., Cancer Research UK program grant (C19769/A11985) to M.M., Wellcome-Warwick Quantitative Biomedicine Programme (WQBP, Institutional Strategic Support Fund: 105627/Z/14/Z), Seed Funding to M.M. and WQBP Interdisciplinary Summer School scholarship to A.W. The authors declare no competing financial interests.

Author contributions

B.C., M.K.B. and M.M. initiated the project. B.C., A.W. and M.M. generated the new *C. elegans* strains. B.C. and M.H. performed live imaging. M.M. performed image analysis. B.C., M.K.B. and M.M. prepared the manuscript. M.K.B. and M.M. supervised the project.

Figure Legends

Fig. 1 Centrosome-directed movement of myosin II particles along astral microtubules during anaphase.

(A) GFP-tagged non-muscle myosin II (NMY-2) in the mid-plane of *C. elegans* embryos was imaged by spinning disk confocal microscopy every 0.83 s and is presented with temporal color-coding. Unidirectional movement of the myosin II in the cytoplasm (arrows) was observed during anaphase. (B) Kymographs of the myosin II particles labeled I and II in the anaphase embryo in panel (A). (C) Velocity distributions of the unidirectional movement of the cytoplasmic myosin II particles found in anterior (A) and posterior (P) sides of anaphase embryos (n=19 and 10, respectively, from 4 embryos). (D) Anillin::GFP in the mid-plane was imaged every 0.72 s and is presented with temporal color-coding. Unidirectional transport of anillin similar to that of myosin II was detected (arrows) although particles immobile relative to the spindle poles were observed more frequently (e.g. those surrounded by a dotted curve). (E) The trajectories of the cytoplasmic transport of myosin II particles from 22 embryos were overlaid and temporally color-coded based on the time after anaphase onset.

Fig. 2 Cytoplasmic transport of myosin II particles and timely furrow formation depend on the astral microtubules and dynein.

(A) Embryos expressing NMY-2::GFP were treated with 15 μ M nocodazole, a microtubule poison, or the drug vehicle (control) during anaphase (at 0 s). Nocodazole strongly suppressed the cytoplasmic motility of myosin II. (B) Depletion of the cytoplasmic dynein heavy chain (*dhc-1(RNAi)*) or the dynein regulator LIN-5/NuMA (*lin-5(RNAi)*) eliminated the cytoplasmic unidirectional movement of myosin

II. (C) Timing of the initial sign of furrow formation determined by visual inspection of anonymized movies (mean \pm s.e.). Statistical significance was tested against the control by linear modeling with correction for multiple comparisons by Dunnett's method. *, ** and *** indicate $p < 0.02$, 0.01 and 0.001 , respectively.

Fig. 3 Local cortical relaxation is observed after removal of myosin II from the cortex.

(A, B) Myosin II-tagRFP-T (magenta) and PH-GFP (green) were simultaneously imaged every 0.77 s with a dual camera spinning disk confocal microscope. (A) The invagination of the plasma membrane marked with arrows was led by an internalizing myosin II particle. On the other hand, no myosin particle was found at the tip of another invagination (marked by arrowheads). See Video 5 for the temporal dynamics. (B) Some cytoplasmic particles (white arrowhead) showed the signals of both myosin II and the PH domain, while others showed exclusive accumulation of the myosin II (magenta arrowheads) or the PH domain (green arrowheads) signals. See Video 6 for the temporal dynamics. (C, E) Embryos expressing myosin II-tagRFP-T (magenta) and tubulin-YFP (green) were imaged by spinning disk confocal microscopy (every 1.7 and 1.8 s in C and E, respectively) and time series of the posterior (C) and the anterior (E) side of two representative embryos are presented. Red arrows indicate the myosin-II particles that were detached from the cortex and traveled along a microtubule fiber towards the centrosome. (D, F) Line profiles of myosin-II on the regions of the cell cortex indicated in (C) and (E), respectively, are presented as kymographs. The particles marked with a red arrowhead were removed from the cortex. Following this, the flanking myosin peaks (blue arrows) were gradually separated from each other, indicating the local relaxation of the

cortical actomyosin network.

Fig. 4 The position of the spindle poles during anaphase defines a local minimum in the density of astral microtubules that reach the cell cortex at the future cleavage site.

(A, B) Density of microtubules near the cell cortex after anaphase onset. (A) Stills from a time-lapse recording of embryos expressing tubulin-YFP. (B) The density of the microtubules along the cell boundary from the anterior tip (0 as the relative position) to the posterior tip (1 as the relative position) in the indicated time window (seconds after anaphase onset). The tubulin/microtubule signals were measured along a line of 1 μm width placed 1.5 μm inside the cell boundary. The intensity above the local background defined by a threshold that corresponds to the top 5% level within a local spatial window (0.2 relative length) was scored as the density of microtubules (see Materials and Methods and Fig. S2 for more details) and averaged across the embryos ($n=9$, mean \pm s.e.). (C, D) Distance from the cell cortex to the nearest spindle pole. (C) The positions of the spindle poles (yellow arrowheads) and the chromosomes (magenta arrows) can be determined in the myosin II-GFP movies. (D) The distance from each point on the cell periphery to the nearest spindle pole was measured and presented with color-coding (left) and by line plotting (right). The equatorial maximum became prominent after ~ 40 s and went above a virtual level that linearly links the anterior and posterior tips (gray dotted line). Green arrows indicate the positions of the local minima in the microtubule density (A and B) or the local maxima in the distance to the nearby pole (A and D).

Fig. 5 The spindle pole-cortex geometry regulates the distribution and flow of

cortical myosin II.

(A and B) Distance between the spindle poles and the cell cortex. (A) The distance from each point on the cell periphery to the nearest spindle pole in embryos depleted of the key microtubule-bundler SPD-1/PRC1 (*spd-1(RNAi)*) was measured and is presented in the same way as the control embryos shown in Fig. 4D. (B) Distance from the anterior pole (A-pole) or the posterior pole (P-pole) to the nearest cortex. Rupture of the central spindle by *spd-1(RNAi)* accelerated the spindle's pole-to-pole elongation, and thus, the decrease in pole-to-cortex distance. (C and D) Effect of depletion of SPD-1 on the cytoplasmic myosin II particles. (C) Temporal color-coded trajectories of NMY-2::GFP from control (n=22) or *spd-1(RNAi)* (n=23) embryos were overlaid by maximum projection. (D) Cytoplasmic myosin II particles were tracked by automation and the activity of the cytoplasmic transport towards the centrosomes is presented as cumulative sums after anaphase onset (see Materials and Methods and Fig. S3 for more details) (mean \pm s.e., n=22 and 23, respectively). (E and F) Density (E) and flow (F) of myosin II in the cell cortex. (E) Cortical myosin II (NMY-2::GFP) was quantified along the cell periphery from the anterior tip to the posterior tip and normalized with the local background and cytoplasmic levels. Average across the embryos from two sets of recordings (22 control and 23 *spd-1(RNAi)* embryos for 0 to 140 s p.a.o. and 56 control and 19 *spd-1(RNAi)* embryos for 70 to 210 s p.a.o.) were merged and are presented as a kymograph for each condition (see Materials and Methods and Fig. S4 for more details). (F) The flow of myosin II along the cell periphery was computed by one-dimensional particle image velocimetry and is presented as a kymograph. The green signal indicates flow towards the posterior and the magenta towards the anterior. The timing of the transition of the cortical flow into the bidirectional mode (white dashed line) is accelerated in the embryos

depleted of SPD-1, in which the spindle asters are positioned closer to the cortex.

Fig. 6 Promotion of the aster-cortex interaction facilitates the formation of the bidirectional cortical flow.

Further evidence of the acceleration of the formation of the bidirectional cortical flow and the initial furrowing. (A) Stills from a video sequence of an embryo expressing myosin II-GFP depleted of CYK-4, the GAP subunit of centralspindlin, in which the central spindle was ruptured (yellow arrowheads) due to defective microtubule-bundling, allowing earlier approach of the spindle poles to the polar cortexes. The initial sign of furrowing (green arrows) was detected earlier than in the control embryos (see Fig. 2C for statistics) but the furrow failed to deepen. (B, C) Density (top) and flow (bottom) of the cortical myosin II. (B) In addition to the microtubule-bundling activity, centralspindlin has been shown to interact with a RhoGEF, ECT2, and to promote the activity of the actomyosin network. This activity is partially redundant with another upstream activator of ECT2, NOP-1. Consistent with the earlier formation of the furrow (A, Fig. 2C), depletion of CYK-4 accelerated the timing of the formation of the bidirectional cortical flow (white dashed line). In contrast, in the embryos depleted of NOP-1, the timing of the formation of the bidirectional flow was not affected although the flow from the posterior to anterior was weaker and the timing of initial furrowing was delayed (Fig. 2C). (C) The aster-cortex interaction are known to be regulated by a microtubule depolymerizer, KLP-7/MCAK, and by a negative regulator of the cortical dynein, EFA-6. In the embryos depleted of either of them, the bidirectional flow formed earlier than in the control embryos, consistent with the earlier initial sign of the furrow formation (Fig. 2C).

Fig. 7 Polar relaxation by dynein-mediated removal of cortical myosin II induces furrow formation in parallel with equatorial stimulation.

Synthetic effect of the simultaneous inhibition of dynein activity (*gpr-1/2(RNAi)*) and centralspindlin-mediated signaling (*cyk-4(GAP)*) was examined by observing embryos with these mutations in the following combinations: *cyk-4(+)* *control(RNAi)* (n=12), *cyk-4(+)* *gpr-1/2(RNAi)* (n=12), *cyk-4(GAP)* *control(RNAi)* (n=13), and *cyk-4(GAP)* *gpr-1/2(RNAi)* (n=13). In contrast to *cyk-4(RNAi)*, a point mutation in the GAP domain of CYK-4 (*cyk-4(GAP)*) specifically inactivates the equatorial stimulation via the centralspindlin-ECT2 pathway without affecting the spindle geometry. Since the *cyk-4(GAP)* allele is temperature-sensitive, embryos were imaged at 23°C. (A) Density and flow of myosin II in the cell cortex for the indicated strains and conditions was obtained in a similar manner to Fig. 5E and 5F. (B) Temporal change of the density of myosin II in the posterior cortex within 10 µm from the posterior tip was plotted (mean across embryos ± s.e.). (C) Cumulative activities of the centrosome-directed cytoplasmic transport of myosin II particles were plotted (mean ± s.e.). Inhibition of dynein activity almost completely abolished the myosin transport irrespective of the genotype of *cyk-4* (*gpr-1/2(RNAi)* or *cyk-4(GAP)* *gpr-1/2(RNAi)*). (D) Timing of the initial sign of furrow formation. The numbers on the top indicate the number of embryos that formed the cleavage furrow out of the total number of observed embryos. The defect in the centralspindlin-dependent equatorial stimulation did not severely affect the patterns of the myosin II distribution and the cortical flow (A ii, *cyk-4(GAP)* vs A i, wild type). In contrast, inactivation of the cortical dynein activity, which nearly completely eliminated the cytoplasmic transport of myosin II (shown in C), drastically changed the cortical distribution and flow of myosin II (A iii, *gpr-1/2(RNAi)* vs A i, wild type, and B *gpr-1/2(RNAi)*) although it still

allowed a weaker bidirectional flow (A iii) and furrow formation in the presence of wild type centralspindlin (D, *gpr-1/2*). Inhibition of both the centralspindlin-dependent pathway and the dynein-dependent pathway almost completely abolished the bidirectional flow (A iv, *cyk-4(GAP) gpr-1/2(RNAi)*, and B, *cyk-4 gpr-1/2*) and severely affected the furrow formation (D, *cyk-4 gpr-1/2*, see also Fig. S5).

Fig. 8 A model of furrow induction by polar relaxation

Schematic showing how bidirectional cortical flow triggered by dynein-dependent removal of myosin II from the polar cortexes leads to cleavage furrow formation. In the one-cell stage *C. elegans* embryo, this pathway provides a mechanism crucial for the initial induction of the cleavage furrow before the equatorial stimulation from the central spindle becomes effective.

Materials and Methods

***C. elegans* strains and culture conditions.** The *C. elegans* strains used in this study are listed in **Supplementary Table 1** and were maintained at 20°C, except for those containing the temperature-sensitive *cyk-4(or749)* allele, which were maintained at 15°C. LP162 strain (*nmy-2(cp13)*) (Dickinson et al., 2013) was crossed with OD56 (*ItIs37*) (Essex et al., 2009) and EU1404 (*cyk-4(or749)*) (Canman et al., 2008) strains to create QM160 (*nmy-2(cp13[nmy-2::gfp + LoxP]) I; ItIs37[(pAA64) pie-1p::mCherry::his-58] IV*) and QM196 (*nmy-2(cp13[nmy-2::gfp + LoxP]) I; cyk-4(or749) III*) strains, respectively. SWG008 strain (*nmy-2(ges6[nmy-2::tagRFP-T + unc-119(+)] I*) was crossed with TH65 (*ddl15[pie-1p::tba-2::YFP]*) and OD58 (*ItIs38[pie-1p::GFP::PH (PLC1delta1)]*) to create QM168 (*nmy-2(ges6[nmy-2::tagRFP-T]) I; ddl15[pie-1p::tba-2::YFP]*) and QM169 (*nmy-*

2(ges6[nmy-2::tagRFP-T]) ; ItIs38[pie-1p::GFP::PH (PLC1delta1)]) strains, respectively. The *gpr-1/2* RNAi clone was from the Vidal Library (Rual et al., 2004) (Source BioScience). The *spd-1* RNAi clone was previously described (Carvalho et al., 2011; Lee et al., 2015). The other RNAi clones were from the Ahringer Library (Edelstein et al., 2014; Kamath et al., 2003) (Source BioScience). For RNAi in the non-temperature sensitive strains, L4 worms were plated on the RNAi plates and incubated at 20 °C for 48 h before dissection to obtain embryos. For RNAi in the temperature sensitive strains, L4 worms were plated and incubated at 16°C for 72 h.

Live microscopy. For fluorescence live imaging, *C. elegans* embryos were immobilized on the cover glass surface of a 35 mm glass-bottom dish (Fluorodish, FD35, World Precision) and cultured in a drop of an osmolality controlled medium. Before experiments, the center of the cover glass was coated with 20 µl of 0.1 mg/ml poly-L-lysine (MW>300,000, Sigma P1524) for 1 h. After a quick wash with water, a circle was drawn with a PAP pen (Sigma, Z377821) around the poly-lysine-coated area, leaving a hydrophilic surface in the center (~10 mm diameter). A piece of filter paper cut into a donut shape of 32 mm diameter with an 18 mm diameter hole was placed onto the bottom of the culture dish and made wet with water. Gravid hermaphrodite animals were dissected in 1 µl of dissection buffer (90 mM sucrose, 50 mM EGTA, 5 mM MgCl₂, 50 mM potassium acetate, 50 mM PIPES-NaOH pH 7) dropped at the center of the poly-L-lysine coated area of the cover glass, immediately followed by addition of 100 µl of an isotonic medium based on Leibovitz's L-15 medium (Gibco 21083-027) supplemented with 10% (v/v) fetal bovine serum, 35 mM sucrose, 100 units/mL penicillin and 100 µg/mL streptomycin (Schindelin et al., 2012; Edgar, 1995; Christensen et al., 2002). The sucrose

concentration was optimized for the viability and normal embryonic divisions upon eggshell permeabilization by *perm-1* RNAi (Sage et al., 2017; Carvalho et al., 2011). To remove unattached embryos and the corpses of the mothers, the medium was replaced with another 100 μ l of the same medium. The culture dish was then covered with a 50 mm diameter cover glass and mounted on the stage of an Andor Revolution XD spinning disk confocal microscopy system based on a Nikon Eclipse Ti inverted microscope equipped with a Nikon CFI Apochromat Lambda S 60 \times /1.40 NA oil-immersion objective lens, a spinning-disk unit (Yokogawa CSU-X1) and an Andor iXon Ultra EM-CCD camera. Images were acquired using Andor IQ3 software. Fluorophores were excited by laser lines at wavelengths of 488 nm for GFP or 561 nm for mCherry. The temperature of the sample, which was monitored by a FLIR One thermal imaging camera (FLIR Systems, Inc.), was maintained at 21 $^{\circ}$ C (**Figs. 1 to 6**) or at 23 $^{\circ}$ C (**Fig. 7**) by circulating cold air within an environmental chamber with a side panel removed. Embryos were staged by differential interference contrast (DIC) observation and the progress of mitosis was monitored every 5 s by observing the state of chromosomes with histone-mCherry (QM160 strain in **Figs. 5 and 6**) or as dark zones in NMY-2::GFP (QM168 and QM169 strains in **Figs. 3 and 4** and QM196 strain in **Fig. 7**). For fast recording of NMY-2::GFP, image acquisition was started immediately after anaphase onset (**Figs. 4 , 5, 6A, 6B and 7**) or 70 s later (**Figs. 5 and 6C**) to capture a set of 5 z-slice images (100 ms exposure, 133 nm/pixel) with 0.5 μ m z-steps every 0.83 s (**Figs. 1 and 2**), 1.7 s (**Fig. 3C**), 1.8 s (**Fig. 3E**), 1.6 s (**Fig. 4**), 0.72 s (**Fig. 5 and 6C**) or 1.25 s (**Figs. 5A, 5B and 7**).

Live microscopy of strain QM169 (**Fig. 3 A and B**) was performed in a similar manner to that indicated above (100 ms exposure, 0.5 μ m z-steps every 0.77 s) but

using an Andor TuCam system equipped with a Nikon ECLIPSE Ti inverted microscope, a Nikon Plan Apo Lambda 100×/1.45-NA oil-immersion objective lens, a spinning-disk confocal system (CSU-X1; Yokogawa Electric Corporation), two Andor iXon Ultra EMCCD cameras, and a wavelength filter set consisting of a 561 nm single-edge laser-flat dichroic beamsplitter, a 514/30 nm single-band bandpass filter and a 568 nm ultrasteep long-pass edge filter (Semrock). Andor IQ3 software was used for simultaneous image acquisition (69 nm/pixel) of specimens with fluorophores excited by laser lines at wavelengths of 488 nm and 561 nm.

For laser ablation of the anterior cortex in **Fig. S1**, embryos prepared as above were subsequently mounted on the stage of a 3i Marianas SDC inverted microscope equipped with a Zeiss α Plan Apo 100x/1.46 NA oil-immersion objective lens, a Yokogawa CSU X-10 spinning disk unit, and a 3i Ablate! laser ablation system (532 nm, 1.3 ns pulsed laser). Images were acquired and ablations were performed using 3i SlideBook 6 software with the following parameters for ablation: laser intensity = 100%, duration = 10 ms, repetitions = 1, raster block size = 3, and rectangle size = 3. GFP was excited with a laser line at a wavelength of 488 nm.

For DIC microscopy in **Fig. S5**, the embryos were mounted between a cover glass and a 2% (w/v) agarose pad in 0.7× egg salt on an FCS2 cooling device (Bioptechs) set at 25 °C, and then filmed with an Olympus BX-51 upright microscope equipped with a UPlanSApo 100×/1.4 NA objective, DIC optics, and a CoolSNAP HQ2 CCD camera (Photometrics) controlled by MicroManager (<https://www.micro-manager.org/>) (Kirshner et al., 2013; Edelstein et al., 2014).

Image analysis. The microscope images were processed and analyzed by custom scripts written with the macro language of Fiji/ImageJ (<https://fiji.sc>) (Arganda-Carreras et al., 2017; Schindelin et al., 2012) and R language (<https://www.r-project.org/>). The 4D images of NMY-2::GFP were deconvolved time frame by time frame with the DeconvolutionLab2 plugin (<http://bigwww.epfl.ch/deconvolution/>) (Oegema et al., 2001; Sage et al., 2017) using the Richardson-Lucy algorithm with total-variation regularization and a point spread function calculated by the PSF generator plugin (Kirshner et al., 2013) with the Born and Wolf model. The coordinates of the cell periphery were determined frame by frame in the bleach-corrected, average z-projections of non-deconvolved images using the Trainable Weka Segmentation plugin ([https://imagej.net/Trainable Weka Segmentation](https://imagej.net/Trainable_Weka_Segmentation)) (Arganda-Carreras et al., 2017). The positions of the spindle poles were determined by human visual detection of the weak NMY-2 localization to the spindle and the spindle poles in anonymized movies.

The microtubule density near the cell cortex was quantified according to a procedure summarized in **Fig. S2**. The background outside of the cell was subtracted from the line profile along a curve of 1 μm width, placed 1.5 μm inside the cell boundary (**Fig. S2A**) and standardized with the mean intensity (**Fig. S2B**). The position along the curve from the anterior pole to the posterior pole was rescaled from 0 to 1. Data from the top and bottom sides of an embryo were pooled for 10 time points (16 s) (**Fig. S2C**). The intensity data that were ranked within the top 5% of the local area of 0.2 width window were treated as the signal from a microtubule (red dots in **Fig. S2D**).

After subtracting the threshold levels, the data from 9 embryos were averaged (**Fig. S2E**).

The velocity of the cytoplasmic NMY-2::GFP particles (**Fig. 1C**) was determined by making kymographs of all the trajectories in four anaphase embryos and measuring their gradients. For automated scoring in **Figs. 1E, 5D, and 7C**, cytoplasmic NMY-2::GFP particles were detected in each time frame of the deconvolved and average z-projected movies by finding maxima within the boundary of each embryo using MaximumFinder in Fiji/ImageJ. The particles detected in two consecutive time frames less than 7 pixels apart, which correspond to movement at 1.295 $\mu\text{m/s}$, were stitched into a trajectory by using a custom R script and overlaid on the original movies to be checked by visual inspection (not shown). The particles that had been trapped on the spindle from metaphase and the false signals derived from the ingressing cleavage furrow were manually omitted. The distance to the closest cortical point was measured for each time point of a trajectory. The direction of the movement of a trajectory relative to the nearby cortex was assessed by the average rate of the increase of the distance to the closest point on the cortex (V_{cortex}). For **Fig. 1E**, the trajectories from 22 *control(RNAi)* embryos that appeared in 6 or more time frames (4.32 s) moving away from the cortex ($V_{\text{cortex}} > 0$) were overlaid with temporal color coding.

The activity of cytoplasmic transport of myosin II towards the centrosome was scored according to the procedure shown in Fig. S3. The force to move a particle of the same size and shape at velocity v in a viscous medium is proportional to v and thus the power for the movement is proportional to v^2 . Although we do not know the

exact size and shape of individual particles, as an estimator of the transport activity of a trajectory per frame, we scored $I \times v^2$, where I is the intensity of the fluorescent signal of each particle. To quantify the transport activity towards the spindle pole, the movement of the spindle pole needs to be considered. Although the position of the spindle pole was difficult to determine by automation, that of a chromosome linked to the pole, with the kinetochore microtubules of a constant length (anaphase A is absent in *C. elegans* embryos (Oegema et al., 2001)), could be tracked by semi-automation, i.e. by manually specifying the dark spots corresponding to the chromosomes in the first time frame and by repeating identification of a nearby minimum in the next time frame. Thus, the movement of the particles was compensated for the movement of the spindle pole in the A-P direction using the A-P component of the chromosome movement (**Fig. S3A**). The (x, y) coordinates of a trajectory (**Fig. S3B**) were converted to the centrosome-directed components of the movement, z , considering the main axis of the movement and the centrosome/chromosome movement (**Fig. S3C**). After smoothing z to z' , the $I \times dz'^2$ was calculated for each time point for which $dz' > 0$ (**Fig. S3D**) and summed up for all the particles moving away from the cortex (**Fig. S3E**, thick lines). For **Fig. 5D** and **Fig. 7C**, the cumulative sum was calculated (**Fig. S3F**). This is the total transport activity from anaphase onset (time 0) and is supposed to represent the overall work executed by dynein for the cytoplasmic transport of the myosin II particles.

For analysis of the temporal change of the cortical density and flow of NMY-2::GFP, first, the region 40 pixels (5.33 μm) inside and outside of the edge of the cell in the average z-projected image was straightened so that the periphery of the embryo that was traced counter-clockwise starting from the anterior tip was placed from left to

right and thus the posterior tip ended at the center (**Fig. S4**). The intensity of the cortical NMY-2::GFP signal, which appeared now as a horizontal line in the middle of the straightened “edge” image, was normalized so that the local background (outside of the cell) is 0 and the local cytoplasmic level is 1. The kymograph of the normalized density at the cell cortex was then generated by reslicing the stack of the time series of the straightened edge images with horizontal lines at intervals of one pixel and averaging the five consecutive best focal planes, which corresponds to the peak of the NMY-2::GFP signal at the cell periphery (cortex) of 0.67 μm width. After averaging across multiple embryos, the kymograph was folded back at the center (= at the posterior tip) so that the anterior and posterior tips were placed on the left and right ends, respectively. For **Fig. 5**, the average kymograph from the dataset obtained from 0 to 140 s p.a.o. and that from 70 to 215 s p.a.o. were merged by linearly changing the blending ratio for the overlapping period (70 s to 140 s p.a.o.) after correction for photobleaching, which is more profound at the cortex than in the cytoplasm due to slower exchange with unbleached molecules.

One-dimensional particle image velocimetry (1D-PIV) was performed by a custom R script that compares the one-dimensional distribution pattern of the cortical NMY-2 signal along the cell periphery in a time frame with that of the next time frame. The spatial resolution of the kymograph of the NMY-2 signal was increased 5-fold (**Figs. 5 and 6**) or 10-fold (**Fig. 7**) by interpolation. The local velocity was determined as the spatial shift needed to maximize the cross-correlation between the windows of the size of 256 pixels (6.83 μm , **Figs. 5 and 6**) or 384 pixels (5.11 μm , **Fig. 7**) from the two consecutive time points. This was scanned along the cell periphery and repeated through the temporal dimension to make a kymograph of the flow of NMY-2 and the

spatial resolution was set back to the original one by averaging. After averaging across multiple embryos, the kymograph of the flow was folded back so that the anterior and posterior tips were placed on the left and right ends, respectively. Positive (anterior to posterior) and negative (posterior to anterior) flow velocities were presented by pseudocoloring with green and magenta, respectively.

Online supplemental material

Fig. S1 shows the relaxation of the cell cortex induced by laser ablation. Figs. S2, S3 and S4 illustrate the methods to measure the microtubule density near the cell cortex, the activity of cytoplasmic transport of myosin II, and the density and flow of cortical myosin II, respectively. Fig. S5 shows the synthetic effect of the CYK-4 GAP mutation and the depletion of LIN-5/NuMA or GPR-1/2/Pins/LGN. Videos 1, 2 and 3 show myosin II in wild type embryos in metaphase, from prometaphase to anaphase and in anaphase, respectively. Videos 4 and 5 show myosin II during anaphase in embryos depleted of DHC-1/dynein heavy chain and LIN-5/NuMA, respectively. Videos 6 and 7 show the dual-color imaging of myosin II and a membrane marker (PH domain). Videos 8, 9 and 10 show the dual-color imaging of myosin II and tubulin/microtubules. Supplementary Table 1 contains the list of the *C. elegans* strains used in this study.

References

- Adriaans, I.E., A. Basant, B. Ponsioen, M. Glotzer, and S.M.A. Lens. 2019. PLK1 plays dual roles in centralspindlin regulation during cytokinesis. *J Cell Biol.* doi:10.1083/jcb.201805036.
- Arganda-Carreras, I., V. Kaynig, C. Rueden, K.W. Eliceiri, J. Schindelin, A. Cardona, and H. Sebastian Seung. 2017. Trainable Weka Segmentation: a machine learning tool for microscopy pixel classification. *Bioinformatics.* 33:2424–2426. doi:10.1093/bioinformatics/btx180.
- Argiros, H., L. Henson, C. Holguin, V. Foe, and C.B. Shuster. 2012. Centralspindlin and chromosomal passenger complex behavior during normal and Rappaport furrow specification in echinoderm embryos. *Cytoskeleton.* 69:840–853. doi:10.1002/cm.21061.
- Atilgan, E., D. Burgess, and F. Chang. 2012. Localization of cytokinesis factors to the future cell division site by microtubule-dependent transport. *Cytoskeleton (Hoboken).* 69:973–982. doi:10.1002/cm.21068.
- Audhya, A., F. Hyndman, I.X. McLeod, A.S. Maddox, J.R. Yates III, A. Desai, and K. Oegema. 2005. A complex containing the Sm protein CAR-1 and the RNA helicase CGH-1 is required for embryonic cytokinesis in *Caenorhabditis elegans*. *J Cell Biol.* 171:267–279. doi:10.1083/jcb.200506124.
- Baruni, J.K., E.M. Munro, and G. von Dassow. 2008. Cytokinetic furrowing in toroidal, binucleate and anucleate cells in *C. elegans* embryos. *J Cell Sci.* 121:306–316. doi:10.1242/jcs.022897.
- Basant, A., S. Lekomtsev, Y.C. Tse, D. Zhang, K.M. Longhini, M. Petronczki, and M. Glotzer. 2015. Aurora B kinase promotes cytokinesis by inducing centralspindlin oligomers that associate with the plasma membrane. *Dev. Cell.* 33:204–215. doi:10.1016/j.devcel.2015.03.015.
- Bouvrais, H., L. Chesneau, S. Pastezeur, D. Fairbrass, M. Delattre, and J. Pecreaux. 2018. Microtubule Feedback and LET-99-Dependent Control of Pulling Forces Ensure Robust Spindle Position. *Biophys J.* 115:2189–2205. doi:10.1016/j.bpj.2018.10.010.
- Bringmann, H., C.R. Cowan, J. Kong, and A.A. Hyman. 2007. LET-99, GOA-1/GPA-16, and GPR-1/2 are required for aster-positioned cytokinesis. *Curr. Biol.* 17:185–191. doi:10.1016/j.cub.2006.11.070.
- Canman, J.C., L. Lewellyn, K. Laband, S.J. Smerdon, A. Desai, B. Bowerman, and K. Oegema. 2008. Inhibition of Rac by the GAP activity of centralspindlin is essential for cytokinesis. *Science.* 322:1543–1546. doi:10.1126/science.1163086.
- Canman, J.C., L.A. Cameron, P.S. Maddox, A. Straight, J.S. Tirnauer, T.J. Mitchison, G. Fang, T.M. Kapoor, and E.D. Salmon. 2003. Determining the position of the cell division plane. *Nature.* 424:1074–1078. doi:10.1038/nature01860.
- Carvalho, A., A. Desai, and K. Oegema. 2009. Structural memory in the contractile

- ring makes the duration of cytokinesis independent of cell size. *Cell*. 137:926–937. doi:10.1016/j.cell.2009.03.021.
- Carvalho, A., S.K. Olson, E. Gutierrez, K. Zhang, L.B. Noble, E. Zanin, A. Desai, A. Groisman, and K. Oegema. 2011. Acute drug treatment in the early *C. elegans* embryo. *PLoS ONE*. 6:e24656. doi:10.1371/journal.pone.0024656.
- Chartier, N.T., D.P. Salazar Ospina, L. Benkemoun, M. Mayer, S.W. Grill, A.S. Maddox, and J.C. Labbé. 2011. PAR-4/LKB1 mobilizes nonmuscle myosin through anillin to regulate *C. elegans* embryonic polarization and cytokinesis. *Curr. Biol*. 21:259–269. doi:10.1016/j.cub.2011.01.010.
- Chen, W., M. Foss, K.-F. Tseng, and D. Zhang. 2008. Redundant mechanisms recruit actin into the contractile ring in silkworm spermatocytes. *PLoS Biol*. 6:e209. doi:10.1371/journal.pbio.0060209.
- Christensen, M., A. Estevez, X. Yin, R. Fox, R. Morrison, M. McDonnell, C. Gleason, D.M. Miller, and K. Strange. 2002. A primary culture system for functional analysis of *C. elegans* neurons and muscle cells. *Neuron*. 33:503–514.
- Colombo, K., S.W. Grill, R.J. Kimple, F.S. Willard, D.P. Siderovski, and P. Gönczy. 2003. Translation of polarity cues into asymmetric spindle positioning in *Caenorhabditis elegans* embryos. *Science*. 300:1957–1961. doi:10.1126/science.1084146.
- D'Avino, P.P., M.G. Giansanti, and M. Petronczki. 2015. Cytokinesis in animal cells. *Cold Spring Harb Perspect Biol*. 7:a015834. doi:10.1101/cshperspect.a015834.
- D'Avino, P.P., M.S. Savoian, L. Capalbo, and D.M. Glover. 2006. RacGAP50C is sufficient to signal cleavage furrow formation during cytokinesis. *J Cell Sci*. 119:4402–4408. doi:10.1242/jcs.03210.
- Dan, K. 1954. The cortical movement in *Arbacia punctulata* eggs through cleavage cycles. *Embryologia*. 2:115–122.
- Davies, T., N. Kodera, G.S. Kaminski Schierle, E. Rees, M. Erdelyi, C.F. Kaminski, T. Ando, and M. Mishima. 2015. CYK4 promotes antiparallel microtubule bundling by optimizing MKLP1 neck conformation. *PLoS Biol*. 13:e1002121. doi:10.1371/journal.pbio.1002121.
- Davies, T., S.N. Jordan, V. Chand, J.A. Sees, K. Laband, A.X. Carvalho, M. Shirasu-Hiza, D.R. Kovar, J. Dumont, and J.C. Canman. 2014. High-resolution temporal analysis reveals a functional timeline for the molecular regulation of cytokinesis. *Dev. Cell*. 30:209–223. doi:10.1016/j.devcel.2014.05.009.
- DeBiasio, R.L., G.M. LaRocca, P.L. Post, and D.L. Taylor. 1996. Myosin II transport, organization, and phosphorylation: evidence for cortical flow/solution-contraction coupling during cytokinesis and cell locomotion. *Mol. Biol. Cell*. 7:1259–1282.
- Dechant, R., and M. Glotzer. 2003. Centrosome separation and central spindle assembly act in redundant pathways that regulate microtubule density and trigger cleavage furrow formation. *Dev. Cell*. 4:333–344.

- Dickinson, D.J., J.D. Ward, D.J. Reiner, and B. Goldstein. 2013. Engineering the *Caenorhabditis elegans* genome using Cas9-triggered homologous recombination. *Nat Methods*. 10:1028–1034. doi:10.1038/nmeth.2641.
- Douglas, M.E., T. Davies, N. Joseph, and M. Mishima. 2010. Aurora B and 14-3-3 coordinately regulate clustering of centralspindlin during cytokinesis. *Curr. Biol*. 20:927–933. doi:10.1016/j.cub.2010.03.055.
- Edelstein, A.D., M.A. Tsuchida, N. Amodaj, H. Pinkard, R.D. Vale, and N. Stuurman. 2014. Advanced methods of microscope control using μ Manager software. *J Biol Methods*. 1:e10. doi:10.14440/jbm.2014.36.
- Edgar, L.G. 1995. Blastomere culture and analysis. *Methods Cell Biol*. 48:303–321.
- Essex, A., A. Dammermann, L. Lewellyn, K. Oegema, and A. Desai. 2009. Systematic analysis in *Caenorhabditis elegans* reveals that the spindle checkpoint is composed of two largely independent branches. *Mol. Biol. Cell*. 20:1252–1267. doi:10.1091/mbc.E08-10-1047.
- Field, C.M., and B.M. Alberts. 1995. Anillin, a contractile ring protein that cycles from the nucleus to the cell cortex. *J Cell Biol*. 131:165–178.
- Fielmich, L.-E., R. Schmidt, D.J. Dickinson, B. Goldstein, A. Akhmanova, and S. van den Heuvel. 2018. Optogenetic dissection of mitotic spindle positioning in vivo. *Elife*. 7:e38198. doi:10.7554/eLife.38198.
- Firestone, A.J., J.S. Weinger, M. Maldonado, K. Barlan, L.D. Langston, M. O'Donnell, V.I. Gelfand, T.M. Kapoor, and J.K. Chen. 2012. Small-molecule inhibitors of the AAA+ ATPase motor cytoplasmic dynein. *Nature*. 484:125–129. doi:10.1038/nature10936.
- Foe, V.E., and G. von Dassow. 2008. Stable and dynamic microtubules coordinately shape the myosin activation zone during cytokinetic furrow formation. *J Cell Biol*. 183:457–470. doi:10.1083/jcb.200807128.
- Galli, M., J. Muñoz, V. Portegijs, M. Boxem, S.W. Grill, A.J.R. Heck, and S. van den Heuvel. 2011. aPKC phosphorylates NuMA-related LIN-5 to position the mitotic spindle during asymmetric division. *Nat Cell Biol*. 13:1132–1138. doi:10.1038/ncb2315.
- Gotta, M., Y. Dong, Y.K. Peterson, S.M. Lanier, and J. Ahringer. 2003. Asymmetrically distributed *C. elegans* homologs of AGS3/PINS control spindle position in the early embryo. *Curr. Biol*. 13:1029–1037.
- Gönczy, P., S. Pichler, M. Kirkham, and A.A. Hyman. 1999. Cytoplasmic dynein is required for distinct aspects of MTOC positioning, including centrosome separation, in the one cell stage *Caenorhabditis elegans* embryo. *J Cell Biol*. 147:135–150.
- Green, R.A., E. Paluch, and K. Oegema. 2012. Cytokinesis in animal cells. *Annu Rev Cell Dev Biol*. 28:29–58. doi:10.1146/annurev-cellbio-101011-155718.

- Grill, S.W., P. Gönczy, E.H. Stelzer, and A.A. Hyman. 2001. Polarity controls forces governing asymmetric spindle positioning in the *Caenorhabditis elegans* embryo. *Nature*. 409:630–633. doi:10.1038/35054572.
- Gudejko, H.F.M., L.M. Alford, and D.R. Burgess. 2012. Polar expansion during cytokinesis. *Cytoskeleton (Hoboken)*. 69:1000–1009. doi:10.1002/cm.21078.
- Han, X., K. Adames, E.M.E. Sykes, and M. Srayko. 2015. The KLP-7 residue S546 is a putative aurora kinase site required for microtubule regulation at the centrosome in *C. elegans*. *PLoS ONE*. 10:e0132593. doi:10.1371/journal.pone.0132593.
- Harris, A.K., and S.L. Gewalt. 1989. Simulation testing of mechanisms for inducing the formation of the contractile ring in cytokinesis. *J Cell Biol*. 109:2215–2223.
- Hu, C.K., M. Coughlin, C.M. Field, and T.J. Mitchison. 2008. Cell polarization during monopolar cytokinesis. *J Cell Biol*. 181:195–202. doi:10.1083/jcb.200711105.
- Hutterer, A., M. Glotzer, and M. Mishima. 2009. Clustering of centralspindlin is essential for its accumulation to the central spindle and the midbody. *Curr. Biol*. 19:2043–2049. doi:10.1016/j.cub.2009.10.050.
- Jantsch-Plunger, V., P. Gönczy, A. Romano, H. Schnabel, D.R. Hamill, R. Schnabel, A.A. Hyman, and M. Glotzer. 2000. CYK-4: a Rho family GTPase activating protein (GAP) required for central spindle formation and cytokinesis. *J Cell Biol*. 149:1391–1404.
- Jiang, W., G. Jimenez, N.J. Wells, T.J. Hope, G.M. Wahl, T. Hunter, and R. Fukunaga. 1998. PRC1: a human mitotic spindle-associated CDK substrate protein required for cytokinesis. *Mol Cell*. 2:877–885.
- Jordan, S.N., T. Davies, Y. Zhuravlev, J. Dumont, M. Shirasu-Hiza, and J.C. Canman. 2016. Cortical PAR polarity proteins promote robust cytokinesis during asymmetric cell division. *J Cell Biol*. 212:39–49. doi:10.1083/jcb.201510063.
- Kamath, R.S., A.G. Fraser, Y. Dong, G. Poulin, R. Durbin, M. Gotta, A. Kanapin, N. Le Bot, S. Moreno, M. Sohrmann, D.P. Welchman, P. Zipperlen, and J. Ahringer. 2003. Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature*. 421:231–237. doi:10.1038/nature01278.
- Khaliullin, R.N., R.A. Green, L.Z. Shi, J.S. Gomez-Cavazos, M.W. Berns, A. Desai, and K. Oegema. 2018. A positive-feedback-based mechanism for constriction rate acceleration during cytokinesis in *Caenorhabditis elegans*. *Elife*. 7:e36073. doi:10.7554/eLife.36073.
- Kirshner, H., F. Aguet, D. Sage, and M. Unser. 2013. 3-D PSF fitting for fluorescence microscopy: implementation and localization application. *J Microsc*. 249:13–25. doi:10.1111/j.1365-2818.2012.03675.x.
- Kitagawa, M., S.Y.S. Fung, N. Onishi, H. Saya, and S.H. Lee. 2013. Targeting Aurora B to the equatorial cortex by MKlp2 is required for cytokinesis. *PLoS ONE*. 8:e64826. doi:10.1371/journal.pone.0064826.

- Kiyomitsu, T., and I.M. Cheeseman. 2013. Cortical dynein and asymmetric membrane elongation coordinately position the spindle in anaphase. *Cell*. 154:391–402. doi:10.1016/j.cell.2013.06.010.
- Klinkert, K., N. Levernier, P. Gross, C. Gentili, L. von Tobel, M. Pierron, C. Busso, S. Herrman, S.W. Grill, K. Kruse, and P. Gönczy. 2019. Aurora A depletion reveals centrosome-independent polarization mechanism in *Caenorhabditis elegans*. *Elife*. 8. doi:10.7554/eLife.44552.
- Kotak, S., C. Busso, and P. Gönczy. 2012. Cortical dynein is critical for proper spindle positioning in human cells. *J Cell Biol*. 199:97–110. doi:10.1083/jcb.201203166.
- Kotak, S., C. Busso, and P. Gönczy. 2014. NuMA interacts with phosphoinositides and links the mitotic spindle with the plasma membrane. *EMBO J*. 33:1815–1830. doi:10.15252/embj.201488147.
- Kozlowski, C., M. Srayko, and F. Nédélec. 2007. Cortical microtubule contacts position the spindle in *C. elegans* embryos. *Cell*. 129:499–510. doi:10.1016/j.cell.2007.03.027.
- Lee, K.-Y., B. Esmaili, B. Zealley, and M. Mishima. 2015. Direct interaction between centralspindlin and PRC1 reinforces mechanical resilience of the central spindle. *Nat Commun*. 6:7290. doi:10.1038/ncomms8290.
- Lekomtsev, S., K.-C. Su, V.E. Pye, K. Blight, S. Sundaramoorthy, T. Takaki, L.M. Collinson, P. Cherepanov, N. Divecha, and M. Petronczki. 2012. Centralspindlin links the mitotic spindle to the plasma membrane during cytokinesis. *Nature*. 492:276–279. doi:10.1038/nature11773.
- Loria, A., K.M. Longhini, and M. Glotzer. 2012. The RhoGAP domain of CYK-4 has an essential role in RhoA activation. *Curr. Biol*. 22:213–219. doi:10.1016/j.cub.2011.12.019.
- Lorson, M.A., H.R. Horvitz, and S. van den Heuvel. 2000. LIN-5 is a novel component of the spindle apparatus required for chromosome segregation and cleavage plane specification in *Caenorhabditis elegans*. *J Cell Biol*. 148:73–86.
- Maddox, A.S., L. Lewellyn, A. Desai, and K. Oegema. 2007. Anillin and the septins promote asymmetric ingression of the cytokinetic furrow. *Dev. Cell*. 12:827–835. doi:10.1016/j.devcel.2007.02.018.
- Mangal, S., J. Sacher, T. Kim, D.S. Osório, F. Motegi, A.X. Carvalho, K. Oegema, and E. Zanin. 2018. TPXL-1 activates Aurora A to clear contractile ring components from the polar cortex during cytokinesis. *J Cell Biol*. doi:10.1083/jcb.201706021.
- Maniscalco, C., A.E. Hall, and J. Nance. 2020. An interphase contractile ring reshapes primordial germ cells to allow bulk cytoplasmic remodeling. *J Cell Biol*. 219. doi:10.1083/jcb.201906185.
- Maton, G., F. Edwards, B. Lacroix, M. Stefanutti, K. Laband, T. Lieury, T. Kim, J.

- Espeut, J.C. Canman, and J. Dumont. 2015. Kinetochore components are required for central spindle assembly. *Nat Cell Biol.* 17:697–705. doi:10.1038/ncb3150.
- Mayer, M., M. Depken, J.S. Bois, F. Jülicher, and S.W. Grill. 2010. Anisotropies in cortical tension reveal the physical basis of polarizing cortical flows. *Nature.* 467:617–621. doi:10.1038/nature09376.
- Mishima, M. 2016. Centralspindlin in Rappaport's cleavage signaling. *Semin. Cell Dev. Biol.* 53:45–56. doi:10.1016/j.semcdb.2016.03.006.
- Mishima, M., S. Kaitna, and M. Glotzer. 2002. Central spindle assembly and cytokinesis require a kinesin-like protein/RhoGAP complex with microtubule bundling activity. *Dev. Cell.* 2:41–54.
- Mollinari, C., J.-P. Kleman, W. Jiang, G. Schoehn, T. Hunter, and R.L. Margolis. 2002. PRC1 is a microtubule binding and bundling protein essential to maintain the mitotic spindle midzone. *J Cell Biol.* 157:1175–1186. doi:10.1083/jcb.200111052.
- Motegi, F., N.V. Velarde, F. Piano, and A. Sugimoto. 2006. Two phases of astral microtubule activity during cytokinesis in *C. elegans* embryos. *Dev. Cell.* 10:509–520. doi:10.1016/j.devcel.2006.03.001.
- Munro, E., J. Nance, and J.R. Priess. 2004. Cortical flows powered by asymmetrical contraction transport PAR proteins to establish and maintain anterior-posterior polarity in the early *C. elegans* embryo. *Dev. Cell.* 7:413–424. doi:10.1016/j.devcel.2004.08.001.
- Murthy, K., and P. Wadsworth. 2008. Dual role for microtubules in regulating cortical contractility during cytokinesis. *J Cell Sci.* 121:2350–2359. doi:10.1242/jcs.027052.
- Nahaboo, W., M. Zouak, P. Askjaer, and M. Delattre. 2015. Chromatids segregate without centrosomes during *Caenorhabditis elegans* mitosis in a Ran- and CLASP-dependent manner. *Mol. Biol. Cell.* 26:2020–2029. doi:10.1091/mbc.E14-12-1577.
- Nguyen, P.A., A.C. Groen, M. Loose, K. Ishihara, M. Wühr, C.M. Field, and T.J. Mitchison. 2014. Spatial organization of cytokinesis signaling reconstituted in a cell-free system. *Science.* 346:244–247. doi:10.1126/science.1256773.
- Nishikawa, M., S.R. Naganathan, F. Jülicher, and S.W. Grill. 2017. Controlling contractile instabilities in the actomyosin cortex. *Elife.* 6. doi:10.7554/eLife.19595.
- Nishimura, Y., and S. Yonemura. 2006. Centralspindlin regulates ECT2 and RhoA accumulation at the equatorial cortex during cytokinesis. *J Cell Sci.* 119:104–114. doi:10.1242/jcs.02737.
- O'Rourke, S.M., S.N. Christensen, and B. Bowerman. 2010. *Caenorhabditis elegans* EFA-6 limits microtubule growth at the cell cortex. *Nat Cell Biol.* 12:1235–1241. doi:10.1038/ncb2128.

- Oegema, K., A. Desai, S. Rybina, M. Kirkham, and A.A. Hyman. 2001. Functional analysis of kinetochore assembly in *Caenorhabditis elegans*. *J Cell Biol.* 153:1209–1226.
- Oegema, K., M.S. Savoian, T.J. Mitchison, and C.M. Field. 2000. Functional analysis of a human homologue of the *Drosophila* actin binding protein anillin suggests a role in cytokinesis. *J Cell Biol.* 150:539–552.
- Pacquelet, A., P. Uhart, J.-P. Tassan, and G. Michaux. 2015. PAR-4 and anillin regulate myosin to coordinate spindle and furrow position during asymmetric division. *J Cell Biol.* 210:1085–1099. doi:10.1083/jcb.201503006.
- Panbianco, C., D. Weinkove, E. Zanin, D. Jones, N. Divecha, M. Gotta, and J. Ahringer. 2008. A casein kinase 1 and PAR proteins regulate asymmetry of a PIP(2) synthesis enzyme for asymmetric spindle positioning. *Dev. Cell.* 15:198–208. doi:10.1016/j.devcel.2008.06.002.
- Pecreaux, J., J.-C. Röper, K. Kruse, F. Jülicher, A.A. Hyman, S.W. Grill, and J. Howard. 2006. Spindle oscillations during asymmetric cell division require a threshold number of active cortical force generators. *Curr. Biol.* 16:2111–2122. doi:10.1016/j.cub.2006.09.030.
- Piekny, A.J., and A.S. Maddox. 2010. The myriad roles of Anillin during cytokinesis. *Semin. Cell Dev. Biol.* 21:881–891. doi:10.1016/j.semcdb.2010.08.002.
- Piekny, A.J., and M. Glotzer. 2008. Anillin is a scaffold protein that links RhoA, actin, and myosin during cytokinesis. *Curr. Biol.* 18:30–36. doi:10.1016/j.cub.2007.11.068.
- Radulescu, A.E., and D.W. Cleveland. 2010. NuMA after 30 years: the matrix revisited. *Trends Cell Biol.* 20:214–222. doi:10.1016/j.tcb.2010.01.003.
- Rankin, K.E., and L. Wordeman. 2010. Long astral microtubules uncouple mitotic spindles from the cytokinetic furrow. *J Cell Biol.* 190:35–43. doi:10.1083/jcb.201004017.
- Rappaport, R. 1996. Cytokinesis in animal cells. Cambridge University Press. 1 pp.
- Reck-Peterson, S.L., W.B. Redwine, R.D. Vale, and A.P. Carter. 2018. The cytoplasmic dynein transport machinery and its many cargoes. *Nat Rev Mol Cell Biol.* 19:382–398. doi:10.1038/s41580-018-0004-3.
- Redemann, S., J. Baumgart, N. Lindow, M. Shelley, E. Nazockdast, A. Kratz, S. Prohaska, J. Brugués, S. Fürthauer, and T. Müller-Reichert. 2017. *C. elegans* chromosomes connect to centrosomes by anchoring into the spindle network. *Nat Commun.* 8:15288. doi:10.1038/ncomms15288.
- Redemann, S., J. Pecreaux, N.W. Goehring, K. Khairy, E.H.K. Stelzer, A.A. Hyman, and J. Howard. 2010. Membrane invaginations reveal cortical sites that pull on mitotic spindles in one-cell *C. elegans* embryos. *PLoS ONE.* 5:e12301. doi:10.1371/journal.pone.0012301.

- Reymann, A.-C., F. Staniscia, A. Erzberger, G. Salbreux, and S.W. Grill. 2016. Cortical flow aligns actin filaments to form a furrow. *Elife*. 5:e17807. doi:10.7554/eLife.17807.
- Rodrigues, N.T.L., S. Lekomtsev, S. Jananji, J. Kriston-Vizi, G.R.X. Hickson, and B. Baum. 2015. Kinetochore-localized PP1-Sds22 couples chromosome segregation to polar relaxation. *Nature*. 524:489–492. doi:10.1038/nature14496.
- Rodriguez-Garcia, R., L. Chesneau, S. Pastezeur, J. Roul, M. Tramier, and J. Pecreaux. 2018. The polarity-induced force imbalance in *Caenorhabditis elegans* embryos is caused by asymmetric binding rates of dynein to the cortex. *Mol. Biol. Cell*. mbcE17110653. doi:10.1091/mbc.E17-11-0653.
- Rose, L.S., M.L. Lamb, S.N. Hird, and K.J. Kemphues. 1995. Pseudocleavage is dispensable for polarity and development in *C. elegans* embryos. *Dev Biol*. 168:479–489. doi:10.1006/dbio.1995.1096.
- Rual, J.-F., J. Ceron, J. Koreth, T. Hao, A.-S. Nicot, T. Hirozane-Kishikawa, J. Vandenhoute, S.H. Orkin, D.E. Hill, S. van den Heuvel, and M. Vidal. 2004. Toward improving *Caenorhabditis elegans* phenome mapping with an ORFeome-based RNAi library. *Genome Res*. 14:2162–2168. doi:10.1101/gr.2505604.
- Sage, D., L. Donati, F. Soulez, D. Fortun, G. Schmit, A. Seitz, R. Guet, C. Vonesch, and M. Unser. 2017. DeconvolutionLab2: An open-source software for deconvolution microscopy. *Methods*. 115:28–41. doi:10.1016/j.ymeth.2016.12.015.
- Salbreux, G., J. Prost, and J.F. Joanny. 2009. Hydrodynamics of cellular cortical flows and the formation of contractile rings. *Phys Rev Lett*. 103:058102. doi:10.1103/PhysRevLett.103.058102.
- Saunders, A.M., J. Powers, S. Strome, and W.M. Saxton. 2007. Kinesin-5 acts as a brake in anaphase spindle elongation. *Curr. Biol*. 17:R453–R454. doi:10.1016/j.cub.2007.05.001.
- Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods*. 9:676–682. doi:10.1038/nmeth.2019.
- Schmidt, R., L.-E. Fielmich, I. Grigoriev, E.A. Katrukha, A. Akhmanova, and S. van den Heuvel. 2017. Two populations of cytoplasmic dynein contribute to spindle positioning in *C. elegans* embryos. *J Cell Biol*. 216:2777–2793. doi:10.1083/jcb.201607038.
- Sedzinski, J., M. Biro, A. Oswald, J.-Y. Tinevez, G. Salbreux, and E. Paluch. 2011. Polar actomyosin contractility destabilizes the position of the cytokinetic furrow. *Nature*. 476:462–466. doi:10.1038/nature10286.
- Shrestha, S., L.J. Wilmeth, J. Eyer, and C.B. Shuster. 2012. PRC1 controls spindle

- polarization and recruitment of cytokinetic factors during monopolar cytokinesis. *Mol. Biol. Cell.* 23:1196–1207. doi:10.1091/mbc.E11-12-1008.
- Singh, D., D. Odedra, P. Dutta, and C. Pohl. 2019. Mechanical stress induces a scalable switch in cortical flow polarization during cytokinesis. *J Cell Sci.* 132. doi:10.1242/jcs.231357.
- Srayko, M., A. Kaya, J. Stamford, and A.A. Hyman. 2005. Identification and characterization of factors required for microtubule growth and nucleation in the early *C. elegans* embryo. *Dev. Cell.* 9:223–236. doi:10.1016/j.devcel.2005.07.003.
- Srinivasan, D.G., R.M. Fisk, H. Xu, and S. van den Heuvel. 2003. A complex of LIN-5 and GPR proteins regulates G protein signaling and spindle function in *C. elegans*. *Genes Dev.* 17:1225–1239. doi:10.1101/gad.1081203.
- Strickland, L.I., E.J. Donnelly, and D.R. Burgess. 2005. Induction of cytokinesis is independent of precisely regulated microtubule dynamics. *Mol. Biol. Cell.* 16:4485–4494. doi:10.1091/mbc.E05-04-0305.
- Su, K.-C., W.M. Bement, M. Petronczki, and G. von Dassow. 2014. An astral simulacrum of the central spindle accounts for normal, spindle-less, and anucleate cytokinesis in echinoderm embryos. *Mol. Biol. Cell.* 25:4049–4062. doi:10.1091/mbc.E14-04-0859.
- Subramanian, R., and T.M. Kapoor. 2012. Building complexity: insights into self-organized assembly of microtubule-based architectures. *Dev. Cell.* 23:874–885. doi:10.1016/j.devcel.2012.10.011.
- Sugioka, K., and B. Bowerman. 2018. Combinatorial contact cues specify cell division orientation by directing cortical myosin flows. *Dev. Cell.* 46:257–270. doi:10.1016/j.devcel.2018.06.020.
- Sugioka, K., L.-E. Fielmich, K. Mizumoto, B. Bowerman, S. van den Heuvel, A. Kimura, and H. Sawa. 2018. Tumor suppressor APC is an attenuator of spindle-pulling forces during *C. elegans* asymmetric cell division. *Proc Natl Acad Sci USA.* 115:E954–E963. doi:10.1073/pnas.1712052115.
- Tse, Y.C., A. Piekny, and M. Glotzer. 2011. Anillin promotes astral microtubule-directed cortical myosin polarization. *Mol. Biol. Cell.* 22:3165–3175. doi:10.1091/mbc.E11-05-0399.
- Tse, Y.C., M. Werner, K.M. Longhini, J.C. Labbé, B. Goldstein, and M. Glotzer. 2012. RhoA activation during polarization and cytokinesis of the early *Caenorhabditis elegans* embryo is differentially dependent on NOP-1 and CYK-4. *Mol. Biol. Cell.* 23:4020–4031. doi:10.1091/mbc.E12-04-0268.
- Turlier, H., B. Audoly, J. Prost, and J.-F. Joanny. 2014. Furrow constriction in animal cell cytokinesis. *Biophys J.* 106:114–123. doi:10.1016/j.bpj.2013.11.014.
- Uehara, R., T. Kamasaki, S. Hiruma, I. Poser, K. Yoda, J. Yajima, D.W. Gerlich, and G. Goshima. 2016. Augmin shapes the anaphase spindle for efficient cytokinetic

- furrow ingression and abscission. *Mol. Biol. Cell.* 27:812–827. doi:10.1091/mbc.E15-02-0101.
- Verbrugghe, K.J.C., and J.G. White. 2004. SPD-1 is required for the formation of the spindle midzone but is not essential for the completion of cytokinesis in *C. elegans* embryos. *Curr. Biol.* 14:1755–1760. doi:10.1016/j.cub.2004.09.055.
- Verbrugghe, K.J.C., and J.G. White. 2007. Cortical centralspindlin and Gα have parallel roles in furrow initiation in early *C. elegans* embryos. *J Cell Sci.* 120:1772–1778. doi:10.1242/jcs.03447.
- von Dassow, G. 2009. Concurrent cues for cytokinetic furrow induction in animal cells. *Trends Cell Biol.* 19:165–173. doi:10.1016/j.tcb.2009.01.008.
- von Dassow, G., K.J.C. Verbrugghe, A.L. Miller, J.R. Sider, and W.M. Bement. 2009. Action at a distance during cytokinesis. *J Cell Biol.* 187:831–845. doi:10.1083/jcb.200907090.
- Werner, M., E. Munro, and M. Glotzer. 2007. Astral signals spatially bias cortical myosin recruitment to break symmetry and promote cytokinesis. *Curr. Biol.* 17:1286–1297. doi:10.1016/j.cub.2007.06.070.
- White, E.A., and M. Glotzer. 2012. Centralspindlin: At the heart of cytokinesis. *Cytoskeleton (Hoboken)*. 69:882–892. doi:10.1002/cm.21065.
- White, J.G., and G.G. Borisy. 1983. On the mechanisms of cytokinesis in animal cells. *J Theor Biol.* 101:289–316.
- Wolpert, L. 1960. The mechanics and mechanism of cleavage. *Int Rev Cytol.* 10:163–216.
- Zhang, D., and M. Glotzer. 2015. The RhoGAP activity of CYK-4/MgcRacGAP functions non-canonically by promoting RhoA activation during cytokinesis. *Elife.* 4:e08898. doi:10.7554/eLife.08898.
- Zhao, P., X. Teng, S.N. Tantirimudalige, M. Nishikawa, T. Wohland, Y. Toyama, and F. Motegi. 2019. Aurora-A Breaks Symmetry in Contractile Actomyosin Networks Independently of Its Role in Centrosome Maturation. *Dev. Cell.* 48:631–645.e6. doi:10.1016/j.devcel.2019.02.012.
- Zhou, M., and Y.-L. Wang. 2008. Distinct pathways for the early recruitment of myosin II and actin to the cytokinetic furrow. *Mol. Biol. Cell.* 19:318–326. doi:10.1091/mbc.E07-08-0783.
- Zhuravlev, Y., S.M. Hirsch, S.N. Jordan, J. Dumont, M. Shirasu-Hiza, and J.C. Canman. 2017. CYK-4 regulates Rac, but not Rho, during cytokinesis. *Mol. Biol. Cell.* 28:1258–1270. doi:10.1091/mbc.E17-01-0020.