

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/142821>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Title: Preterm children’s long-term academic performance after adaptive computerized training: an efficacy and process analysis of a randomized controlled trial

Authors: Julia Jaekel *^{1 2}, Katharina M. Heuser^{3 4}, Antonia Zapf⁵, Claudia Roll⁶, Francisco Brevis Nuñez⁷, Peter Bartmann⁸, Dieter Wolke², Ursula Felderhoff-Mueser³, & Britta Huening³

Institutional affiliations:

¹ Child and Family Studies | Psychology; University of Tennessee Knoxville; Knoxville, TN

² Psychology; University of Warwick; Coventry; UK

³ Pediatrics I, Neonatology, Pediatric Intensive Care, Pediatric Neurology; University Hospital Essen; Essen; Germany

⁴ Psychology; Ruhr-University Bochum; Bochum; Germany

⁵ Medical Biometry and Epidemiology; Universitätsklinikum Hamburg-Eppendorf; Hamburg; Germany

⁶ Vest Children's Hospital Datteln, University Witten-Herdecke; Datteln, Germany

⁷ Pädiatrische Intensivmedizin; Sana-Kliniken Duisburg; Duisburg; Germany

⁸ Children's Hospital; University Hospital Bonn; Bonn; Germany

Corresponding author: Julia Jaekel, Department of Child and Family Studies; University of Tennessee Knoxville, 1215 W. Cumberland Ave., Knoxville, TN 37996, Phone: 865-249-4479, jjjaekel@utk.edu

Authorship contributions:

- Substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data: all authors
- Drafting the article or revising it critically for important intellectual content: Jaekel, Heuser, Huening, Zapf, Bartmann, Wolke
- Final approval of the version to be published: all authors

Statement of financial support: This study was funded by Stiftung Mercator (grant number PR-2014-0039).

Disclosure statement: The authors have nothing to disclose.

Category of study: Clinical Research Article, RCT; registered online with the German Clinical Trials Register (DRKS; ID DRKS00007685).

Statement of consent: Participants provided written informed consent and assent.

Impact:

- Adaptive computerized math training may help improve preterm children's short-term school performance
- Computerized math training provides a novel avenue towards intervention after preterm birth
- Well-powered randomized-controlled studies of math intervention effectiveness for preterm school-aged children are warranted

ABSTRACT

Background: Adaptive computerized interventions may help improve preterm children's academic success, but randomized trials are rare. We tested whether a math training (XtraMath®) versus an active control condition (Cogmed®; working memory) improved school performance. Training feasibility was also evaluated.

Methods: Preterm born first graders, $N=65$ (28 - 35+6 weeks gestation) were recruited into a prospective randomized-controlled multi-centre trial and received one of two computerized trainings at home for five weeks. Teachers rated academic performance in math, reading/writing, and attention compared to classmates before (baseline), directly after (post), and 12 months after the intervention (follow-up). Total academic performance growth was calculated as change from baseline (hierarchically ordered - post first, follow-up second).

Results: Bootstrapped linear regressions showed that academic growth to post-test was significantly higher in the math intervention group ($B= .25$ [95% CI: .04 to .50], $p=.039$), but this difference was not sustained at the 12-months follow-up ($B=.00$ [-.31 to .34], $p=.996$). Parents in the XtraMath group reported higher acceptance compared with the Cogmed group (*mean difference*: -0.49, [-0.90 to -0.08], $p=.037$).

Conclusions: Our findings do not show a sustained difference in efficacy between both trainings. Studies of math intervention effectiveness for preterm school-aged children are warranted.

INTRODUCTION

About 15 million babies worldwide (10.6% of all births) are born preterm (< 37 weeks gestational age (GA)) every year and preterm birth represents a significant cause of lifelong morbidity (1). Delivery at any gestation other than full term (39-41 weeks GA) may result in altered brain development and risk for adverse neurocognitive outcomes and academic underachievement (2, 3). While problems of moderately and late preterm (MLP) children are subtle they represent about 80-85% of all preterm born children, and even small increases in cognitive abilities may have large effects on academic performance on a population level (4). MLP children have been neglected in follow-up services although they may benefit most from intervention (5).

With regard to specific difficulties, preterm birth has been consistently associated with low attention, working memory and mathematic scores (6, 7). These areas are closely associated with academic achievement. To reduce adverse outcomes it is timely and essential to invest into the development of interventions that can increase academic performance of children born preterm.

There is increasing evidence that the start of schooling may be a critical window of opportunity for intervention in preterm populations (8). Some have reported that an adaptive working memory training, Cogmed® (Pearson Education Inc.) may improve working memory up to seven months post-training in very low birth weight (< 1,500 g) preschool children (9) and extremely low birth weight (< 1,000 g) adolescents (10). However, effects were rebutted by a recent study reporting no long-term benefits of Cogmed in extremely preterm school-aged children (11). Moreover, a population-based randomized controlled clinical trial found no

evidence for short- and long-term effects of Cogmed on academic performance among children with working memory difficulties (12). Thus, interventions targeting general working memory skills may not result in beneficial effects on preterm children's school performance, in particular in mathematics, and more targeted specific intervention may be needed.

Individual mathematic differences are determined early in life (13, 14). In elementary school, the math curriculum comprises different domains such as numeration, arithmetic, problem solving, geometry, and algorithmic computation; however, we still know little about the domain-specific and domain-general cognitive processes involved in each domain (15). There is agreement that complex mathematical problem solving requires both short-term/working memory (e.g., processing of new information) and long-term memory (e.g., arithmetic fact recall). Development of fast arithmetic fact recall (i.e., computational fluency) comprises automaticity and efficient use of limited cognitive resources, and is thus essential for successful math learning and progress. The assumption underlying this argument is that fluid abilities such as attention, working memory, and processing speed are essential for acquiring knowledge (16). To master higher-level math skills, children thus need to transition from finger counting or slowly calculating basic math facts to recalling all four operations (addition, subtraction, multiplication, and division) accurately, quickly, and effortlessly. Such progress to automatized recall is associated with freeing up working memory resources during problem solving (3, 17). While fact fluency is requisite for later mathematic success (17, 18), inattentive behaviour and poor processing speed may inhibit the age-appropriate development of such computational skills (15, 19). Preterm children's processing speed (20) and cognitive workload deficits (3) may make them vulnerable for delays with basic math fact fluency. Thus,

interventions targeting fact fluency may be particularly effective for increasing preterm children's school success. XtraMath® is such a program, but has never been systematically evaluated. In addition, while cognitive training in general may have beneficial effects (21), there is little research on feasibility and acceptability of computerized interventions at school age (22), especially among preterm children and their parents (11, 23, 24).

In this randomized controlled trial, we investigated whether an adaptive online math training (XtraMath) would result in short- and long-term (i.e., directly after and 12 months after the intervention) changes in academic performance in preterm elementary school children, compared with an active control condition (Cogmed working memory training). Second, we evaluated children's and parents' training satisfaction, motivation, and general feasibility of both trainings.

METHODS

Procedure

Participants of this prospective, multi-centre randomized controlled trial (RCT) were recruited from birth registries of seven neonatal intensive care (NICU) units, level 3, in the German State of North-Rhine Westphalia. The study was conducted according to CONSORT guidelines and the Declaration of Helsinki. The study protocol was approved by the Ethics Committees of the Faculty of Psychology at the Ruhr-University Bochum (Votum 134) and the Medical Faculty of the University of Essen (14-6163-BO). The trial was registered online with the German Clinical Trials Register (DRKS; ID DRKS00007685).

First, participants were screened for eligibility based on the information available in their birth records. Overall, $N=1,026$ infants were born preterm between October 1, 2008 and September 30, 2009 in the participating centres. Of these, $n=494$ did not meet the predefined inclusion criteria (see Figure 1 for details). Accordingly, the parents of $n=532$ children were invited via mail to participate in the RCT. In line with German medical records data protection regulations, only those who actively responded and agreed to a screening interview could be interviewed via telephone ($n=88$). Of these, $n=7$ did not meet inclusion criteria, $n=16$ parents declined to participate in the trial, and $n=65$ agreed. Participants (parents and children) provided written informed consent and assent before being randomized.

- Figure 1 about here -

Inclusion / Exclusion Criteria. Preterm children (GA < 37 weeks) who started elementary school in August 2015 were included. Only children and parents with sufficient German language abilities to participate in standardized assessments were included; their place of residence was < 100 km distance from at least one of the study centres to increase feasibility of participation. Preterm children with non-correctable handicaps (London Handicap Scale, function scale > 2), moderate/severe cerebral palsy (CP), hemiparesis, or intraventricular haemorrhage (IVH) > grade 2 combined with significant leucomalacia/hydrocephalus were excluded to allow participants to successfully complete the training. Families with twins or higher multiples were excluded as parental supervision of ≥ 2 children's completion of daily training after school was not considered feasible. Children with a diagnosis of Oppositional defiant disorder (ODD) or conduct problems were excluded as they were expected to resist

daily training. Children whose school entry was delayed were excluded as their academic progress (primary outcome) could not be assessed.

Intervention

Children were stratified according to gender and gestational age, and then randomly assigned to either the adaptive math or working memory training group using central blockwise randomization. AZ generated the random allocation sequence, research nurses enrolled participants and assigned them to intervention groups. In addition, children were grouped into four blocks and started assessments and trainings successively during their first year of formal schooling.

Intervention group: Children in the intervention group played an online, computerized math program (XtraMath®) that continuously adapts to children's abilities. XtraMath aims to help students transition from counting or calculating the basic math facts to automatically recalling them (25). The program uses timed activities with three-second thresholds to encourage students to answer questions as quickly as possible combined with spaced repetition. XtraMath has four operation components: addition, subtraction, multiplication, and division. Children usually progress through these operations sequentially, based on their mastery (e.g., once children achieve 100% mastery in addition, they will start with subtraction, and so on). Each of the four components starts with a placement quiz that determines the individual initial mastery score and the subsequent math problems the child is presented with. Parents receive weekly progress report emails and they can check training progress online. The XtraMath training itself is not language dependent while the instructions are available in nine languages, including German. Children randomized into the intervention group used the

XtraMath standard protocol (at least 5x/weekly practice for 10-15 minutes) for five weeks. In this study, only one-digit addition tasks were used (0+0 to 9+9) in order to standardize training content in line with child age and training duration.

Active control group: For the purpose of rigorous trial evaluation, an adaptive computerized working memory training, Cogmed (version JM for children aged 4-6 years), that has recently been shown to not facilitate school success (12) was administered to the preterm children randomized into the active control group. The control training comprised several games designed as fairground rides (e.g., Ferris wheel, rollercoaster). Children were instructed to recall visual-spatially distributed sequences. The training started at a low level of complexity and was administered according to its standard protocol, 5x/weekly for 15-20 minutes per day for five weeks.

In both groups, a trained coach provided weekly in-person and phone-based technical and motivational support for parents during the 5-week intervention period.

Outcome Measures

Children's academic performance was assessed via teacher ratings and standardized tests, administered by trained psychologists before (baseline), directly after (post), and 12 months after the intervention (follow-up). All teachers were blind to children's intervention group memberships. Blind assessment of secondary outcomes was not possible because post-intervention questionnaires on training motivation and general feasibility differed by training program.

Primary outcome: Participants' teachers were asked to compare the individual child's performance to the class average expected performance levels, using the Teacher Academic

Attainment Scale (TAAS) (26). The TAAS is a brief and psychometrically sound teacher report of achievement in the following three core dimensions: mathematics, reading/writing, as well as attention/concentration in class. Ratings are completed on 5-point scales (1 to 5) with 3 representing average class performance. The TAAS has been validated in general and preterm populations, shows high concurrent correlations with achievement tests (range: $r=.69-.82$), high test-retest reliability ($r=.77$), and has high sensitivity and specificity in identifying learning difficulties (26). The TAAS was administered at baseline, post-test, and at 12 months follow-up to assess training-induced changes in individual trajectories of academic achievement. Students in North Rhine-Westphalia usually keep their main subjects' classroom teachers throughout the four years of elementary school. Thus, the three TAAS ratings were completed by the same teacher for each child, while all participating children were at different schools. The two primary hypotheses (post-test versus baseline and 12 months versus baseline) were ordered hierarchically: post-test first, 12 months second. Because of the hierarchical testing, no adjustment for multiplicity was necessary.

Secondary outcomes: In addition to teacher ratings, children performed standardized tests of school achievement in mathematics (DEMAT 1+, 2+) (27, 28). At baseline and post-test, the DEMAT 1+ (Deutscher Mathematiktest für erste Klassen) was used and at 12 months follow up the DEMAT 2+ (Deutscher Mathematiktest für zweite Klassen) was administered. Both tests comprise different tasks (e.g., number range, addition and subtraction, geometry) based on the respective math curricula.

For the process analysis, $n=60$ children answered 16 questions on their experiences with the online training at the post-intervention assessment; 14 of these were adapted from the

Intrinsic Motivation Inventory (IMI) (29, 30) and translated into German, two were self-created. The IMI assesses participants' experiences related to a target activity. Items used in this study were chosen from the four IMI subscales interest/enjoyment, perceived competence, effort/importance, and value/usefulness. Children responded on an adapted three-point Likert-type scale (1=no, 2=sometimes, 3=yes). Factor structure (PCA) and reliability analyses revealed that eleven of the 16 items loaded on one single factor that explained 36.9% of the overall variance (Cronbach's $\alpha=.81$), and these were included in final analyses. Supplemental Table S1 (online) displays these items ordered by IMI subscale. Individual child responses were z-standardized and averaged into an index scale of *Training Motivation*.

In addition, $n=59$ parents completed an in-house questionnaire containing 16 items assessing training satisfaction, motivation, and general feasibility. Z-standardized items were averaged and combined into index scales of *Training Satisfaction* (Cronbach's $\alpha=.74$), *Child's Motivation* ($\alpha=.81$), and *General Feasibility* ($\alpha=.76$; for details see Supplemental Table S2 (online)).

Perinatal variables: Information on child sex, gestational age, birth weight, infections, and brain injury was drawn from birth records.

Social variables: Parental education was assessed as part of a screening interview performed with the parents during the recruitment phase. According to International Standard Classification of Education (ISCED) categories, all parents had medium to high education. Mothers' and fathers' scores were thus dichotomized into medium vs. high education. In addition, parents completed questionnaires on their child's behaviour and family background information at baseline, post-intervention and 12 months follow up.

Data Analysis

This is a randomized, multi-centre trial with two arms. The primary analysis is conducted on the intention-to-treat (ITT) population, control of compliance was conducted via log-files. 65 participants were included, allowing to detect a group difference with an effect size of $d=0.74$. Values of participants who were lost to follow-up were not imputed, resulting in a final sample size of XtraMath $n=29$ versus controls $n=27$ (TAAS) / $n=25$ (DEMAT). The primary hypothesis was that children who were randomized into the adaptive XtraMath versus the working memory training would have more academic growth, assessed as TAAS change from baseline (calculated as a difference score of (1) TAAS at post-test, directly after the intervention, minus TAAS at baseline; and (2) TAAS at 12 months follow-up minus TAAS at baseline). Two bootstrapped linear regression analyses were carried out with the dependent variables of academic growth, intervention as main factor, and baseline TAAS as covariate. The Type-I-error was set to 5% two-sided (no adjustment because of hierarchical testing), yielding >80% statistical power to detect effects of $d=0.78$ and $R^2=.16$, respectively. Analyses were repeated controlling for child sex, GA, and intervention block. Results remained stable and there were no effects of confounding variables, due to stratification, thus unadjusted results are reported to preserve statistical power. The secondary math test outcomes were assessed with corresponding regression analyses. A process analysis was carried out using bootstrapped independent-samples *t*-tests to examine differences between training programs on process analysis index scales.

RESULTS

Descriptive sample characteristics

Table 1 shows descriptive characteristics of the two intervention groups at baseline. There were no differences with regard to child sex, age at assessments, or GA due to stratification. Baseline z-standardized TAAS scores were equally distributed across the two groups (TAAS T1: *Mean (SD)* XtraMath: -0.03 (1.15) versus Cogmed: 0.03 (0.84), *Mean Difference*: 0.06 [95% *CI*: -0.44 to 0.56], $p=.811$). The same was true for the math test scores (DEMAT T1: *Mean (SD)* XtraMath: 0.19 (0.97) versus Cogmed: 0.09 (0.98), *Mean Difference*: -0.10 [95% *CI*: -0.66 to 0.44], $p=.721$). In addition, we confirmed that children's working memory (WISC digit span backwards score: *Mean (SD)* XtraMath: 5.29 (1.24) versus Cogmed: 5.15 (1.10), *Mean Difference*: -0.14 [95% *CI*: -0.77 to 0.50], $p=.666$) and arithmetic (K-ABC calculation standard score: *Mean (SD)* XtraMath: 107.71 (10.14) versus Cogmed: 108.11 (8.99), *Mean Difference*: 0.40 [95% *CI*: -4.79 to 5.59], $p=.879$) abilities at baseline did not differ between the two groups.

- Table 1 about here -

With regard to training compliance across the 5-week intervention period, log-files showed that, on average, children in the XtraMath group completed fewer sessions per week (4.41 (0.98)) than children in the Cogmed group (5.11 (0.47) *Mean Difference*: 0.70 [95% *CI*: 0.28 to 1.12], $p=.002$). However, preliminary analyses showed that there was no association between training compliance and primary or secondary outcomes, and main analyses were conducted on the intention-to-treat (ITT) population as planned.

Training effects on academic performance growth

Bootstrapped linear regressions with intervention as main factor and baseline TAAS (T1) as covariate showed that total academic growth to post-test was significantly higher in the XtraMath group ($B = .25$ [95% CI: .04 to .50], $p = .039$; $R^2 = .14$), but this difference was not sustained at the 12-months follow-up ($B = .00$ [-.31 to .34], $p = .996$; $R^2 = .14$). Figure 2 suggests that, directly after the intervention (post-test) children in the XtraMath group showed more academic growth than children in the active control group, however, these short-term gains were not sustained.

There were no relevant group differences in math test score growth according to training condition (baseline DEMAT entered as covariate) post-test: $B = .05$ [-.20 to .29], $p = .697$; 12-months follow-up: $B = .16$ [-.13 to .49], $p = .346$), however, inspection of Figure 3 suggests that children in the XtraMath group achieved stable performance while children in the active control group (Cogmed), on average, showed a tendency for decreasing math test performance after participating in the training.

- Figures 2 and 3 about here -

Process analysis

Overall, acceptance of both trainings was high. For example, most children reported the training was fun (*Mean (SD)* Cogmed: 2.55 (0.69), XtraMath: 2.58 (0.72), range 1 (low) – 3 (high)), and parents rated integration into everyday life as easy (Cogmed: 2.79 (0.69), XtraMath: 2.97 (0.61), range 1 – 4). Bootstrapped independent-samples *t*-tests showed that, on average, parents in the XtraMath group reported higher general feasibility of the training compared with the Cogmed group (see Table 2), representing a medium-sized effect, $d = 0.56$.

- Table 2 about here -

DISCUSSION

This multi-centre RCT examined the efficacy and feasibility of an adaptive online math training (XtraMath) to improve short- and long-term academic performance in preterm elementary school children, compared with an active control condition (Cogmed adaptive working memory training). Results revealed that the computerized math training promoted higher short-term academic performance growth than the active control condition, but group differences in academic growth were not sustained to 12 months follow-up. Overall, parents in the math training group rated the training as more feasible than in the control group.

First, although differences between training groups were not significant 12 months after the completion of the intervention, the findings of this study provide tentative novel evidence that computerized trainings targeting specific math skills may help support preterm children's success in school short term. XtraMath has been specifically designed to improve automatic recall of arithmetic facts, however, the short-term training effect found here applied to academic performance across all main subjects, suggesting potential broader transfer effects to other domains.

Complex mathematical problem solving requires both working memory (e.g., processing of new information) and long-term memory (e.g., involving arithmetic fact retrieval). XtraMath involves a simultaneous combination of multiple cognitive processes required in mental arithmetic, such as processing speed, working memory, visual-spatial skills, and fact retrieval (15, 19, 31). Developing fast arithmetic fact recall comprises automaticity and efficient use of

limited cognitive resources, and is thus essential for successful math learning. The assumption underlying this argument is that fluid abilities such as working memory and processing speed are essential for acquiring knowledge (16), but these are not the only requirements. To master higher-level math skills, students need to obtain and understand numeric symbols and rules, and transition from finger counting or calculating basic math facts to recalling operations accurately, quickly, and effortlessly. Such progress to automatized retrieval (i.e., recall) has been associated with freeing up working memory resources during problem solving (17) and with functional changes in the left inferior parietal cortex (32). Training arithmetic fact recall may help preterm children who are struggling with mathematics become more fluent and provide a foundation for successfully mastering more complex math problems later in life.

There is mixed evidence in intervention research regarding cost-effectiveness of training intensity, scope, and duration. The data presented here are from an efficacy study, as the intervention was implemented under ideal conditions in highly motivated families and with the support of a specifically trained coach (33). Training effectiveness under real-life conditions has yet to be evaluated.

Our findings suggest that computerized trainings represent a motivating and feasible avenue towards intervention for school-aged preterm children. Results revealed small differences in training acceptance. XtraMath training tasks may be more similar to children's everyday activities at school than the more playful approach of Cogmed, which may have contributed to differences in parent-rated feasibility of both trainings. Overall, the adaptive, open-access XtraMath training may be more cost-effective and easier to implement into everyday life. Nevertheless, the experiences with recruiting families of school-aged preterm

children into this randomized trial suggest that a majority of German parents may be reluctant to participate in computerized interventions, a potentially culturally-specific attitude. As Figure 1 shows, only 88 out of 532 parents of preterm children who fulfilled the inclusion criteria showed interest to participate in a computerized training. Of these 88, however, 16 voiced critical concerns about increased daily screen time as part of the intervention and refused to participate.

With regard to teachers' roles in supporting preterm children at school, previous research has shown that teachers have poor knowledge about preterm children's specific needs and difficulties, and it is crucial to provide strategies for supporting their specific needs in the classroom (34). Applying and rigorously evaluating different approaches to improve preterm children's academic performance provides the best avenue towards improving education and health services for this population. Approaches may include novel e-learning resources to increase teachers' knowledge of preterm birth and how to support preterm children (35), recommendations for specific changes in classroom teaching methods, as well as adaptive computerized interventions, as presented here.

In general, and even more in light of recent global developments related to the COVID-19 epidemic, teachers are increasingly challenged to integrate technologies and tools that support distance learning in education. Specialised training programs may help children obtain crucial abilities, however, teachers and parents often struggle to identify innovative high-quality resources, integrate the new materials in regular routines (36), and worry about potential negative consequences of extended screen exposure (37). Computerized trainings can complement but not substitute classroom teaching. With distance and classic in-school learning

alike, the aim is to design a well-rounded curriculum that incorporates educational technology elements but does not solely rely on them (37). Accordingly, the strength of a computational fluency program such as XtraMath may be that it trains foundational skills while leaving strategy instruction to teachers.

Strengths and limitations. This study has several methodological strengths, including a prospective design investigating short- and long-term intervention-induced effects and stratified random assignment of participants to training conditions. The intervention was compared with an active control condition that also adapted to children's learning progress, which helped control effects of maturation and practice as well as irrelevant training aspects such as expectancy effects (38). Despite the relatively low recruitment rate discussed above, a high retention rate was achieved, with 86% of children participating to 12-months follow up. The primary outcome were teacher ratings, while previous studies on the effectiveness of computerized trainings on academic functioning used standardized tests of school performance (11, 12). In this study, teachers who were blind to intervention group membership rated participants' academic attainment to evaluate the feasibility of computerized home-based interventions with an ecologically valid measure. Teachers are highly experienced in comparing an individual child's performance to expected grade levels in school. Moreover, parents are particularly interested in outcomes that make a difference in real life, such as education and later life chances, rather than changes in standardized test scores. Participants were born in multiple centres across a densely populated region of Germany, increasing the generalizability of findings. Children were born very preterm as well as moderately and late preterm and stratified across groups, thereby extending the range of gestational age for evaluating

intervention efficacy in comparison to previous studies (9-11, 39). Finally, data was analysed with an intention-to-treat approach, including all participants as randomized irrespective of their training compliance.

This study also has limitations. Blind assessment of secondary outcomes was not possible because post-intervention questionnaires were different for the two training groups. Second, the sample size was small and statistical power thus only sufficient to detect a large effect size, however, bootstrapping was employed in order to help alleviate some statistical power limitations. Finally, some child characteristics, such as fine motor skills and computer affinity, which may have influenced training performance, were not assessed and accounted for in analyses.

In conclusion, findings of this RCT do not provide evidence that adaptive computerized math training better supports preterm children's long-term academic performance than working memory training. However, children in the XtraMath showed significantly more short-term academic growth than children in the active control group. Overall, results of this study suggest that home-based delivery of computerized training as intervention for school-aged preterm children is feasible. More research on individualized interventions and classroom teaching strategies that cater to preterm children's specific educational needs is warranted.

ACKNOWLEDGEMENTS

This study was funded by Stiftung Mercator (grant number PR-2014-0039). We thank all cooperating hospitals, Katholisches Klinikum Bochum, Klinikum Duisburg, Elisabeth Hospital Essen, Vest Children's Hospital Datteln, University Hospital Duesseldorf, University Hospital

Essen, and Helios Klinikum Niederberg in Velbert, for supporting this project. Special thanks to Frauke Dransfeld, Eva Weishaupt, and Gisela Axmacher-Bopp for assistance with recruitment, intervention implementation, data collection and management, and to the study participants, their families, and teachers.

REFERENCES

1. March of Dimes, PMNCH, Save the Children, WHO. Born Too Soon: The Global Action Report on Preterm Birth. Howson C, Kinney M, Lawn J, editors. Geneva: World Health Organization; 2012.
2. Davis EP et al. Children's brain development benefits from longer gestation. *Frontiers in Psychology*. 2011;2.
3. Jaekel J, Baumann N, Wolke D. Effects of gestational age at birth on cognitive performance: A function of cognitive workload demands. *PLoS ONE*. 2013;8(5):e65219.
4. MacKay DF, Smith GCS, Dobbie R, Pell JP. Gestational age at delivery and special educational need: retrospective cohort study of 407,503 schoolchildren. *PLoS Med*. 2010;7(6):e1000289.
5. McCormick MC et al. Early intervention in low birth weight premature infants: Results at 18 years of age for the Infant Health and Development Program. *Pediatrics*. 2006;117(3):771-80.
6. Jaekel J, Bartmann P, Schneider W, Wolke D. Neurodevelopmental pathways to preterm children's specific and general mathematic abilities. *Early Human Development*. 2014;90(10):639-44.
7. Wolke D et al. Universal gestational age effects on cognitive and basic mathematic processing: 2 cohorts in 2 countries. *The Journal of Pediatrics*. 2015.
8. van Veen S et al. Very preterm born children at early school age: Healthcare therapies and educational provisions. *Early Hum Dev*. 2018;117:39-43.

9. Grunewaldt KH, Skranes J, Brubakk AM, Låhaugen GC. Computerized working memory training has positive long-term effect in very low birthweight preschool children. *Dev Med Child Neurol.* 2016;58(2):195-201.
10. Lohaugen GCC et al. Computerized working memory training improves function in adolescents born at extremely low birth weight. *J Pediatr.* 2011;158(4):555-U56.
11. Anderson PJ et al. Long-Term Academic Functioning Following Cogmed Working Memory Training for Children Born Extremely Preterm: A Randomized Controlled Trial. *J Pediatr.* 2018;202:92-7.e4.
12. Roberts G et al. Academic outcomes 2 years after working memory training for children with low working memory: a randomized clinical trial. *JAMA Pediatr.* 2016;170(5):e154568.
13. Stern E. The development of mathematical competencies: Sources of individual differences and their developmental trajectories. In: W. Schneider, Bullock M, editors. *Human development from early childhood to early adulthood: Evidence from the Munich Longitudinal Study on the Genesis of Individual Competencies (LOGIC)*. Mahwah, NJ: Erlbaum; 2009. p. 221-37.
14. Gilmore CK, Papadatou-Pastou M. Patterns of individual differences in conceptual understanding and arithmetical skill: A meta-analysis. *Mathematical Thinking and Learning.* 2009;11(1-2):25-40.
15. Fuchs LS et al. Problem solving and computational skill: Are they shared or distinct aspects of mathematical cognition? *Journal of Educational Psychology.* 2008;100(1):30-47.
16. Blair C, Raver CC. School readiness and self-regulation: a developmental psychobiological approach. *Annu Rev Psychol.* 2015;66:711-31.

17. Vasilyeva M, Laski EV, Shen C. Computational fluency and strategy choice predict individual and cross-national differences in complex arithmetic. *Dev Psychol.* 2015;51(10):1489-500.
18. Carr M, Alexeev N. Fluency, accuracy, and gender predict developmental trajectories of arithmetic strategies. *Journal of Educational Psychology.* 2011;103(3):617-31.
19. Cirino PT, Fuchs LS, Elias JT, Powell SR, Schumacher RF. Cognitive and mathematical profiles for different forms of learning difficulties. *J Learn Disabil.* 2015;48(2):156-75.
20. Mulder H, Pitchford NJ, Marlow N. Inattentive behaviour is associated with poor working memory and slow processing speed in very pre-term children in middle childhood. *British Journal of Educational Psychology.* 2011;81:147-60.
21. Jolles D, Crone EA. Training the developing brain: a neurocognitive perspective. *Frontiers in Human Neuroscience.* 2012;6.
22. Roberts G et al. Can improving working memory prevent academic difficulties? A school based randomised controlled trial. *BMC Pediatr.* 2011;11:57.
23. Pascoe L et al. Preventing academic difficulties in preterm children: a randomised controlled trial of an adaptive working memory training intervention - IMPRINT study. *BMC Pediatr.* 2013;13:144.
24. Aarnoudse-Moens CSH et al. Executive Function Computerized Training in Very Preterm-Born Children: A Pilot Study. *Games Health J.* 2018;7(3):175-81.
25. XtraMath. <https://xtramath.org/#/home/faq> 2016 [

26. Johnson S, Marlow N, Wolke D. Assessing educational outcomes in middle childhood: validation of the Teacher Academic Attainment Scale. *Developmental Medicine & Child Neurology*. 2012:no-no.
27. Krajewski K, Küspert P, Schneider W. Deutscher Mathematiktest für erste Klassen (DEMAT 1+). Göttingen: Hogrefe; 2002.
28. Krajewski K, Liehm S, Schneider W. Deutscher Mathematiktest für zweite Klassen (DEMAT 2+). Göttingen: Hogrefe; 2004.
29. McAuley E, Duncan T, Tammen VV. Psychometric properties of the Intrinsic Motivation Inventory in a competitive sport setting: a confirmatory factor analysis. *Res Q Exerc Sport*. 1989;60(1):48-58.
30. Ryan RM, Deci EL. Intrinsic and extrinsic motivations: Classic definitions and new directions. *Contemporary Educational Psychology*. 2000;25(1):54-67.
31. Simms V et al. Nature and origins of mathematics difficulties in very preterm children: a different etiology than developmental dyscalculia. *Pediatr Res*. 2015;77(2):389-95.
32. Rivera SM, Reiss AL, Eckert MA, Menon V. Developmental changes in mental arithmetic: evidence for increased functional specialization in the left inferior parietal cortex. *Cereb Cortex*. 2005;15(11):1779-90.
33. Singal AG, Higgins PDR, Waljee AK. A primer on effectiveness and efficacy trials. *Clinical and translational gastroenterology*. 2014;5(1):e45-e.
34. Johnson S, Gilmore C, Gallimore I, Jaekel J, Wolke D. The long-term consequences of preterm birth: what do teachers know? *Dev Med Child Neurol*. 2015;57(6):571-7.

35. Johnson S et al. Improving developmental and educational support for children born preterm: evaluation of an e-learning resource for education professionals. *BMJ open*. 2019;9(6):e029720.
36. Cavanagh S. Districts put open ed. resources to work. *Education Week*. 2016;Spotlight On Digital Math Instruction:2.
37. Herold B. Doing ed-tech right in the early years. *Education week*. 2016;Spotlight On Digital Math Instruction:12-3.
38. Melby-Lervåg M, Hulme C. Is working memory training effective? A meta-analytic review. *Dev Psychol*. 2013;49(2):270-91.
39. Grunewaldt KH, Løhaugen GC, Austeng D, Brubakk AM, Skranes J. Working memory training improves cognitive function in VLBW preschoolers. *Pediatrics*. 2013;131(3):e747-54.

Table 1. Descriptive characteristics of preterm children randomized into intervention groups

	Active control group (Cogmed; <i>n</i> = 32)	Intervention group (XtraMath; <i>n</i> = 33)	<i>p</i>
Demographic characteristics			
Age at baseline	6.91 (0.44)	6.99 (0.31)	.427
Sex (% male)	47	49	.897
Maternal education (% high)	56	48	.835
Paternal education (% high)	50	42	.575
Monthly net income (€, median)	3,000-4,000	3,000-4,000	1.000
Perinatal characteristics			
Gestational age at birth (weeks)	32.81 (2.10)	32.73 (2.14)	.870
Birth weight (grams)	1937 (511)	2058 (599)	.385
Hypoxia (% no/yes/unknown)	100/0/0	97/0/3	.484
Amnion infection syndrome (% no/yes/unknown)	91/9/0	88/9/3	1.000
Cerebral ultrasonography (postnatal period)			
Intraventricular haemorrhage (% no/yes/unknown)	100/0/0	91/6/3	.107
Periventricular leukomalacia at term- equivalent age (% no/yes/unknown)	100/0/0	97/0/3	.484

Data are reported as mean (SD) if not indicated otherwise.

Table 2. Comparison of training programs on process analysis index scales

Index Scales	Active control group		Intervention group		<i>Mean difference</i>	<i>95% CI</i>
	(Cogmed)		(XtraMath)			
	<i>n</i> = 29		<i>n</i> = 31			
Child report: Training Motivation ^a	-0.04	(0.56)	0.04	(0.63)	-0.08	(-0.35 – 0.20)
Parent report: Training Satisfaction ^{a, b}	-0.06	(0.64)	0.04	(0.59)	-0.10	(-0.44 – 0.23)
Parent report: Child’s Motivation ^{a, b}	0.03	(0.61)	-0.03	(0.73)	0.06	(-0.24 – 0.36)
Parent report: General Feasibility ^{a, b, c}	-0.26	(0.91)	0.23	(0.83)	-0.49	(-0.90 – -0.08)

Note. Data are presented as *mean (SD)*. 95% confidence intervals (CI) are bias-corrected and accelerated, bootstrapping based on 2,000 samples.

^a Index scales are based on z-standardized items.

^b For index scales based on parent reports statistics are based on *n* = 28 cases for Cogmed© training.

^c For index scale *General Feasibility* statistics are based on *n* = 29 cases for XtraMath© training.




Figure 1. Randomization Flow Chart

Note. HIE: Hypoxic Ischemic Encephalopathy; PVL: Periventricular Leukomalacia; IVH: Intraventricular Hemorrhage; ODD: Oppositional Defiant Disorder




Figure 2. Teacher-rated Academic Attainment Scores (TAAS) by Intervention Group (XtraMath ($n=29$, bolded) versus Cogmed ($n=27$)) at Baseline, Post-test, and 12 Months Follow-up

Note: Bolded bars represent the XtraMath Intervention group




Figure 3. Math Test Scores (DEMAT) by Intervention Group (XtraMath ($n=28$, bolded) versus Cogmed ($n=25$)) at Baseline, Post-test, and 12 Months Follow-up

Note: Bolded bars represent the XtraMath Intervention group.

Supplemental Table S1. Adapted items of the Intrinsic Motivation Inventory (IMI) from the post-intervention questionnaire

answered by children, ordered by IMI subscale.

Subscale of IMI	Question included in index scale <i>Training Motivation</i>
Interest / Enjoyment	I thought this was a boring training (recoded).
Interest / Enjoyment	I found it easy to concentrate on this training.
Interest / Enjoyment	This training was fun to do.
Interest / Enjoyment	I would describe the training as interesting.
Interest / Enjoyment	I liked this training very much.
Value / Usefulness	I think doing this training could help me to do better at school.
Value / Usefulness	I would be willing to do the training again because it has value to me.
Value / Usefulness	I think this is an important training.
Value / Usefulness	I think this is an important training because it can help me at school.
Perceived competence	I think I was pretty good at this training.
Effort / Importance	It was important to me to do well at this training.

Note. Items were answered on a three-point Likert-type scale (1=no, 2=sometimes, 3=yes).

Supplemental Table S2. Items from the in-house post-intervention questionnaire answered by parents ordered by index scale.

Index scale	Question	Response format
Training Satisfaction	Overall, I think the user-friendliness (i.e., simple operation, etc.) of the training program was ...	1 = very bad, 2 = bad, 3 = good, 4 = very good
Training Satisfaction	I think the training program was understandable (all information required was available to ensure that everything ran smoothly).	1 = Not at all true, 2 = Rather not true, 3 = Rather true, 4 = Absolutely true
Training Satisfaction	Overall, I think the difficulty level of the individual training task was ...	1 = Too low, 2 = Appropriate, 3 = Too high
Training Satisfaction	Overall, I think the individual training tasks were too monotonous.	1 = Absolutely true, 2 = Rather true, 3 = Rather not true, 4 = Not at all true
Training Satisfaction	How would you judge the amount of time spent on the training (frequency and duration of the individual sessions)?	1 = Very high, 2 = Appropriate, 3 = Very low
Training Satisfaction	Were your individual expectations for the training met?	1 = Not at all, 2 = A little, 3 = A lot
Training Satisfaction	Would you recommend this training to your friends?	1 = Not at all, 2 = A little, 3 = A lot
Child's Motivation	Did your child feel overburdened with performing the training?	1 = Always, 2 = Often, 3 = Rarely, 4 = Never
Child's Motivation	Has your child had any problems performing the training or with individual training tasks?	1 = Always, 2 = Often, 3 = Rarely, 4 = Never
Child's Motivation	Was your child frustrated after performing the training?	1 = Always, 2 = Often, 3 = Rarely, 4 = Never
Child's Motivation	Did it happen that you had to additionally motivate your child to performing the training (e.g., additional rewards)?	1 = Always, 2 = Often, 3 = Rarely, 4 = Never
Child's Motivation	Did your child have fun performing the training?	1 = Never, 2 = Rarely, 3 = Often, 4 = Always
Child's Motivation	Did your child complain about having to perform the training?	1 = Always, 2 = Often, 3 = Rarely, 4 = Never

Child's Motivation	My child felt the training was motivating.	1 = Not at all true, 2 = Rather not true, 3 = Rather true, 4 = Absolutely true
General Feasibility	How easy was the integration of the training program into your everyday life in general?	1 = Very difficult, 2 = Difficult, 3 = Easy, 4 = Very easy
General Feasibility	Did you find the implementation of the training program was an additional burden in your everyday life?	1 = A lot, 2 = A little, 3 = Not at all