Environmental Microbiology (2021) 23(9), 5069-5086



Proteomics insights into the *Burkholderia cenocepacia* phosphorus stress response

Holly Shropshire,^{1,2} Rebekah A. Jones,² María M. Aguilo-Ferretjans,² David J. Scanlan ¹⁰² and Yin Chen ^{102*}

¹BBSRC Midlands Integrative Biosciences Training Partnership, University of Warwick, Coventry, CV4 7AL. UK.

²School of Life Sciences, University of Warwick, Coventry, CV4 7AL, UK.

Summary

The Burkholderia cepacia complex is a group of Burkholderia species that are opportunistic pathogens causing high mortality rates in patients with cystic fibrosis. An environmental stress often encountered by these soil-dwelling and pathogenic bacteria is phosphorus limitation, an essential element for cellular processes. Here, we describe cellular and extracellular proteins differentially regulated between phosphate-deplete (0 mM, no added phosphate) and phosphate-replete (1 mM) growth conditions using a comparative proteomics (LC-MS/MS) approach. We observed a total of 128 and 65 unique proteins were downregulated and upregulated respectively, in the B. cenocepacia proteome. Of those downregulated proteins, many have functions in amino acid transport/ metabolism. We have identified 24 upregulated proteins that are directly/indirectly involved in inorganic phosphate or organic phosphorus acquisition. Also, proteins involved in virulence and antimicrobial resistance were differentially regulated, suggesting B. cenocepacia experiences a dramatic shift in metabolism under these stress conditions. Overall, this study provides a baseline for further research into the biology of Burkholderia in response to phosphorus stress.

Introduction

Burkholderia cenocepacia is a member of the Burkholderia cepacia complex (Bcc), a group of closely

related Burkholderia species (Mahenthiralingam et al., 2005; Vial et al., 2011). The genus Burkholderia was separated from Pseudomonas species and validated in 1992 (Burkholder, 1950; Yabuuchi et al., 1992). Although first described as a plant pathogen causing onion rot, Bcc species are also opportunistic human pathogens that cause infections in patients with cystic fibrosis (CF) and chronic granulomatous disease (Mahenthiralingam et al., 2005; Vial et al., 2011). Bcc species cause under 5% of infections in CF patients, but these types of infections are feared due to easy transmission, high intrinsic resistance to antibiotics and the unpredictability of patient outcome, i.e. an asymptomatic infection or the development of fatal cepacia syndrome (Leitão et al., 2017).

One of the most well-studied bacterial stress responses is phosphate limitation, which has not been extensively studied in Bcc species. Phosphorus is an essential element required for many cellular processes and is a major component of membranes and genetic material. Bacteria largely acquire phosphorus as inorganic phosphate (Pi) (Wanner, 1996; Lamarche et al., 2008b; Crépin et al., 2011; Santos-Beneit, 2015). In Gram-positive bacteria, the Pho regulon is well studied in the genus Bacillus. The two-component system (TCS) PhoPR is known to mediate a specific response to phosphate stress, many of which are involved in teichoic acid turnover in the cell wall, a feature specific to Grampositive bacteria (Antelmann et al., 2000; Prunty et al., 2018). In Gram-negative bacteria, however, the response to Pi stress is orchestrated by the PhoBR TCS. At high external Pi concentrations, PhoR is hypothesized to form a repression complex with the high-affinity Pi transporter, PstSCAB. However, when the environmental Pi concentration falls below a few micromolar. PhoR is liberated from this repression and can activate its transcriptional regulator, PhoB (Wanner, 1996; Lamarche et al., 2008b; Santos-Beneit, 2015). Genes directly controlled by PhoB are collectively termed the Pho regulon and contain a conserved Pho box in their promoter region which PhoB can bind to. In Escherichia coli, the Pho regulon originally comprised eight transcriptional units containing 31 genes (Wanner, 1990, 1993, 1996). However, over 47 genes have now been assigned to the

© 2021 The Authors. *Environmental Microbiology* published by Society for Applied Microbiology and John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Received 21 July, 2020; accepted 2 March, 2021. *For correspondence. E-mail y.chen.25@warwick.ac.uk; Tel. +44 24 76528976; Fax +44 (0) 24765 23701.

E. coli Pho regulon, some of the newer genes include an AMP nucleosidase (amn) and a metal ion stress response gene (vibD) (Baek and Lee, 2006; Lamarche et al., 2008b). ChIP-seg analysis of PhoB has identified 19 directly regulated genes that have varying roles in metabolism, transport and transcriptional regulation (Yang et al., 2012). However, proteomic studies have suggested the response to Pi limitation could involve over 400 proteins (Van Bogelen et al., 1996). Although the soil and plant environments are known to have limited Pi availability, whether human pathogens could experience Pi stress during host invasion is not completely understood. One hypothesis is that intracellular pathogens such as Salmonella experience Pi limitation in the macrophage vacuole during the host immune response. This theory is supported by the expression of PhoBR and other genes involved in Pi acquisition (Eriksson et al., 2003; Srikumar et al., 2015; Choi et al., 2019). Such a scenario is also possible for Burkholderia species as they are intracellular pathogens able to survive in immune cells like macrophages where they could encounter Pi stress (Valvano, 2015). The human bronchial epithelium is also a low Pi environment and the induction of Pi acquisition genes has already been reported during Pseudomonas aeruginosa infections in this location (Frisk et al., 2004; Chugani and Greenberg, 2007).

Not only is the response to Pi stress critical for cell survival, but it is often linked to virulence and antimicrobial resistance. This has been extensively studied in pathogenic species of E. coli (Lamarche et al., 2008b; Bertrand et al., 2010; Yoshida et al., 2010; Crépin et al., 2011) and Vibrio cholerae, the human intestinal pathogen causing cholera (Von Krüger et al., 1999). Similarly, in the common CF pathogen, P. aeruginosa, Pi limitation increases swarming motility and cytotoxicity to human bronchial epithelia (Bains et al., 2012) whilst Pi depletion was also a trigger for the lethal red death phenotype in a Caenorhabditis elegans infection model (Zaborin et al., 2009). Proteomic studies have also shown Pi limitation increases the expression of virulence factors in many pathogens, including Salmonella typhimurium and V. cholerae (Von Krüger et al., 2006; Jiang et al., 2018). Many cell wall modifications can occur under Pi depletion, such as the remodelling of the cell membrane with non-phosphorus-containing lipids (Taylor et al., 1998; Zavaleta-Pastor et al., 2010). Changes in the cell wall due to Pi depletion can also lead to increased antimicrobial resistance. For example, the modification of the lipid A moiety of lipopolysaccharide causes increased resistance to polymyxins (Lamarche et al., 2008a; Crépin et al., 2011). Furthermore, the PstSCAB Pi transporter is known to be involved in the efflux of fluoroquinolone antibiotics (Baneriee et al., 2000; Li and Nikaido, 2004) and

the TolC efflux pump, which is implicated in drug resistance, is upregulated in *V. cholerae* under Pi stress (Von Krüger *et al.*, 2006).

In this work, we set out to identify the Pi limitation stress response in B. cenocepacia using comparative proteomics of the cellular and extracellular proteomes. In the last 10 years, advances in proteomic methods have facilitated studies of the exoproteome. This extracellular 'compartment' is a key area for secretion of toxins, virulence factors and degradative enzymes (Brown et al., 2012; Gagic et al., 2016). Kvoto Encvclopedia of Genes and Genomes (KEGG) analysis of our data showed differential regulation of a wide range of cellular processes, particularly proteins related to phosphate regulation, virulence and antimicrobial resistance. These results not only identify the proteins that are regulated by phosphorus limitation in Burkholderia species, but also highlight the complexity of the Pi stress response and begin to unravel the consequences of Pi limitation on pathogenesis and antimicrobial resistance in this key human pathogen.

Results and discussion

The effects of Pi stress on growth and alkaline phosphatase activity in B. cenocepacia

To examine the Burkholderia Pi stress response, B. cenocepacia K56-2 was cultivated under both Pi-deplete (0 mM) and Pi-replete (1 mM) conditions. Burkholderia cenocepacia had a significantly slower growth rate under Pi depletion (P < 0.01, Welch *t*-test) (Fig. 1A). To ensure these cultures were experiencing Pi stress, we used alkaline phosphatase activity as an indicator of Pi limitation since these secreted proteins scavenge Pi from their environment and are known to be regulated by the PhoBR TCS (Hong et al., 2007; Richards and Vanderpool, 2012). Burkholderia cenocepacia showed a significant increase in alkaline phosphatase activity 4 h after exposure to Pi-deplete conditions (P < 0.01, Welch t-test) (Fig. 1B). These results confirmed the experimental setup was sufficient for inducing Pi limitation in B. cenocepacia.

Environmental Pi limitation causes significant shifts in cellular and extracellular proteomes in B. cenocepacia

To begin to unravel the proteins regulated by Pi availability in *Burkholderia* species, we used comparative proteomics to study the response of *B. cenocepacia* K56-2 to Pi stress. Samples for LC–MS/MS were taken 8 h after exposure to Pi-deplete (0 mM) and Pi-replete (1 mM) conditions (Fig. 1A). The cellular proteome was analysed from pelleted cells and the extracellular proteome from



Fig. 1. A. Growth of *B. cenocepacia* K56-2 under Pi-deplete (0 mM, red) and Pi-replete (1 mM, black) conditions. Samples for proteomics were taken at 8 h

B. At each growth time point, alkaline phosphatase activity was measured using the *p*NPP assay to assess the activation of the Pho regulon. Alkaline phosphate activity was measured by calculating the *p*NP production in $\mu g^{-1} h^{-1}$ normalized to cell density of the sample (OD₆₀₀). Error bars represent standard error of the mean of triplicate cultures.

filtered supernatant after removing cells by centrifugation. A total of 1728 and 332 proteins were detected in the cellular and extracellular proteome respectively (Suppl. Table 1). Of these proteins, 182 and 23 were significantly differentially regulated ($P \le 0.05$) in Pi-deplete compared with Pi-replete conditions in the cellular and extracellular proteome respectively (Fig. 2). Twelve of these proteins were significantly differentially regulated in both the cellular and exoproteome fractions. In the cellular proteome, 63 proteins were significantly downregulated ($P \le 0.05$, Fig. 2A). In the exoproteome, 11 proteins were significantly upregulated and 12 proteins were significantly downregulated ($P \le 0.05$, Fig. 2B; Suppl. Table 1).

In terms of abundance, 13 proteins made up at least 1% of the total cellular proteome in either Pi-deplete or Pi-replete conditions, two of which were significantly differentially regulated (Table 1). Noticeably, PstS showed the largest difference in relative abundance (7.02%) in Pi-deplete compared with Pi-replete conditions. PstS is the periplasmic substrate-binding protein of the highaffinity Pi transporter, PstSCAB (Rosenberg et al., 1977; Torriani, 1990; Yuan et al., 2006a; Yuan et al., 2006b). PstSCAB expression has been shown to be controlled by PhoB in many bacterial species and is a wellcharacterized member of the Pho regulon (Rosenberg et al., 1977; Yamada et al., 1989; Yuan et al., 2006a; Yuan et al., 2006b; Lubin et al., 2016). Therefore, this sharp increase in PstS abundance in our results confirms the activation of the Pi stress response. In the extracellular proteome, 17 proteins made up at least 1% of the total exoproteome in either Pi-deplete or Pi-replete conditions, none of which, however, were significantly differentially regulated (Table 1). Noticeably, FliC, a flagellar filament structural protein accounted for >20% of the total exoproteome.

Proteins classified by clusters of orthologous groups and KEGG analysis further demonstrate the shift occurring in metabolism under Pi stress

Pi stress elicited a major change in the cellular and extracellular proteome with a total of ~200 proteins being significantly affected (Fig. 2). To gain a better understanding of the metabolic pathways affected by Pi limitation, significantly differentially regulated proteins (Fig. 2) were subjected to KEGG and Clusters of Orthologous Groups (COG) analysis (Table 2 and Suppl. Table 2 respectively). KEGG analysis highlights protein functions, whereas COG analysis gives a general overview of the cellular processes each protein contributes to. In the KEGG analysis (Table 2), there were a large number of enzymes upregulated and downregulated in both fractions of the Pi-deplete proteome compared with the Pireplete proteome. The most striking result was the high number of downregulated transporters in the Pi-deplete cellular proteome. A deeper analysis into the function of transporters using the KEGG identifier ko02000 (Table 2) revealed four of the upregulated proteins comprised the PstSCAB Pi transporter, whereas the downregulated transporters were involved in amino acid, sulfur, carbon, nitrogen and iron transport (Fig. 3).

Other observations in the KEGG analysis showed three proteins related to TCSs were upregulated (Table 2, Suppl. Table 2), including PhoB and PhoR, the TCS for the Pi stress response and BCAM1199, the homologue of which is involved in chemotaxis in *P. aeruginosa* (Joshi *et al.*, 2019). The only TCS down-regulated in response to Pi stress was the response regulator CheY that is involved in chemotaxis in *B. cenocepacia* (Bazzini *et al.*, 2011). Proteins related to genetic information processing were mostly down-regulated with a few exceptions, such as mRNA

Table 1. Proteins detected in the cellular and exoproteomes comprising at least 1% of the total cellular or exoproteome in either Pi-deplete (0 mM) or Pi-replete (1 mM) conditions.

P < 0.05	Log2-fold	Locus tag (gene name)	Annotation	Sequence	Relative abundance (%)		
7 ≪ 0.05	change)	Locus lag (gene hame)	Annotation	lengui	+Pi	-Pi	Difference
Cellular pr	roteome						
+	+3.054	BURCENK562V_C1990 (pstS)	Phosphate ABC transporter substrate- binding protein, PhoT family	344	0.92	7.94	+7.02
	+0.946	BURCENK562V_C0963 (phaP)	Phasin family protein	188	2.33	4.67	+2.34
	+0.343	BURCENK562V_C5730 (aroL1)	Chaperonin GroEL	546	1.92	2.53	+0.61
	+0.103	BURCENK562V_C1183 (BPSL2096)	Peroxiredoxin, alkyl hydroperoxide reductase subunit C	182	3.11	3.48	+0.37
	+0.370	BURCENK562V_C0143	Superoxide dismutase, Fe-Mn family	192	0.75	1.01	+0.26
	+0.284	BURCENK562V_C2988	Malate dehydrogenase, NAD	328	0.96	1.21	+0.25
	+0.024	BURCENK562V_C6532	Translation elongation factor 1A, EF-1A/	396	3.01	3.19	+0.18
	-0.156	BURCENK562V_C5942	OmpA-OmpF porin, OOP family	222	1.90	1.78	-0.12
	-0.104	BURCENK562V_C2161	HU family DNA-binding protein	92	4.24	4.10	-0.14
	-0.323	BURCENK562V_C3041	DNA-binding protein	149	2.24	1.86	-0.38
	-0.803	BURCENK562V_C5678	1-Cys peroxiredoxin	212	1.36	0.81	-0.55
	-1.026	BURCENK562V_C7254	Inclusion body protein	167	1.98	1.01	-0.97
+	-3.292	BURCENK562V_C2529	Chaperonin GroEL	546	2.28	0.24	-2.04
Extracellul	ar proteome						
	+0.849	BURCENK562V_C6831	Flagellin	384	21.76	43.19	+21.43
	+2.330	BURCENK562V_C7127 (<i>zmpA</i>)	ZmpA-like peptidase, Metallopeptidase, MEROPS family M04	565	0.24	1.32	+1.08
	-0.018	BURCENK562V_C6346	Protein of unknown function (DUF4399)	178	1.78	1.94	+0.16
	-0.064	BURCENK562V_C4415 (<i>livK1</i>)	L-leucine-binding protein/L-isoleucine- binding protein/L-valine-binding protein	372	0.96	1.01	+0.05
	-0.514	BURCENK562V_C1183 (BPSL2096)	Peroxiredoxin (alkyl hydroperoxide reductase subunit C)	182	1.08	0.83	-0.25
	-0.476	BURCENK562V_C4051 (BCAM1931)	Outer membrane protein, porin	359	1.23	0.97	-0.26
	-0.254	BURCENK562V_C1023	Peptidyl-prolyl cis-trans isomerase A, cyclophilin A	191	3.41	3.15	-0.26
	-0.746	BURCENK562V_C6611 (BCAL 0305)	Phospholipid transport system substrate- binding protein	210	1.23	0.81	-0.42
	-0.609	BURCENK562V_C1663 (BCAL 1577)	Hypothetical protein	315	1.57	1.14	-0.43
	-0.564	BURCENK562V_C6135	Iron(III) transport system substrate- binding protein	347	1.89	1.41	-0.48
	-0.183	BURCENK562V_C1990	Phosphate ABC transporter substrate-	344	19.70	19.13	-0.57
	-1.016	BURCENK562V_C0461	D-methionine transport system substrate-	271	1.48	0.81	-0.67
	-0.904	BURCENK562V_C0550	Thiol:disulfide interchange protein DsbA	212	1.75	1.03	-0.72
	-1.772	BURCENK562V_C4473	Fungalysin metallopeptidase, M36	570	1.48	0.48	-1.00
	-1.682	BURCENK562V_C6106	Amino acid ABC transporter substrate-	257	1.67	0.57	-1.10
	-1.501	BURCENK562V_C1576	Monosaccharide ABC transporter	316	1.32	0.51	-1.31
	-0.652	BURCENK562V_C2355 (gltl)	L-glutamate-binding protein/L-aspartate- binding protein	297	8.82	6.18	-2.64

Changes in the phosphate-deplete condition were compared with the phosphate-replete condition. Significance ($P \leq 0.05$) is related to the log2-fold change of their relative abundance in proteomes.



Fig. 2. Protein expression fold changes when comparing *B. cenocepacia* K56-2 grown in Pi-deplete (0 mM) to Pi-replete (1 mM) conditions.

A. Cellular proteome expression changes.

B. Exoproteome expression changes. Upregulated and downregulated proteins are shown in green/blue and red/pink respectively. Pink/green and blue/red represent $P \leq 0.01$ and $P \leq 0.05$ respectively. Black proteins are not significantly different.

synthesis and ribosome biogenesis. Results from the KEGG analysis give the overall impression of metabolic adaptation occurring in response to Pi stress in *B. cenocepacia*. A similar adaptation has been described in other bacterial species such as *Salmonella*, *Corynebacterium* and *Streptomyces* whereby primary metabolic processes including transcription, amino acid metabolism and protein synthesis have previously been shown to decrease in Pi-limited environments (Ishige *et al.*, 2003; Novotna *et al.*, 2003; Rodríguez-García *et al.*, 2007; Jiang *et al.*, 2018). As seen in both analyses (Table 2, Suppl. Table 2), a large fraction of the differentially regulated proteins were unclassified, i.e. encoding hypothetical proteins with unknown function.

Proteins related to Pi regulation are significantly upregulated under Pi depletion

Out of 63 and 11 significantly upregulated cellular and extracellular proteins respectively, 22 and 6 were identified as being involved in Pi regulation (Table 3). In B. cenocepacia, the PhoBR TCS forms a putative operon with the high-affinity Pi transporter PstSCAB and the regulator PhoU, which were all upregulated under Pi stress in the cellular proteome (Fig. 4A; Table 3). Other classic members of the E. coli Pho regulon include PsiA (a.k.a. PhoA, an alkaline phosphatase) and PsiF (a protein of unknown function) which form an operon in E. coli (Wanner, 1996; Luu et al., 2018). PhoA homologues (BCAM0389 and BCAM0390, 31% and 32% identity respectively) were upregulated in both fractions of the proteome (Table 3), whereas the PsiF homologue (BURCENK562V_C1248) was not significantly differentially expressed and does not form an operon with the PhoA homologues. Burkholderia cenocepacia also has two homologues of the PhoH pro-(BURCENK562V_C0301 BURtein and CENK562V_C1266, 48% and 36% identity respectively), a protein of unknown function (Wanner, 1996; Goldsmith et al., 2011). However, neither protein was significantly differentially regulated under Pi stress. Burkholderia cenocepacia has homologues for the UgpBAECQ operon involved in sn-glycerol-3-phosphate (G3P) uptake (BURCENK562V C6584-C6588), however, they were not significantly regulated (Suppl. Table 1). Burkholderia cenocepacia does not have homologues for the other three members of the E. coli Pho regulon, which are the phosphonate uptake/degradation operon, PhnC-PhnP, the porin. PhoE and PsiE. a protein of unknown function (Wanner, 1996). Therefore, only nine of the cellular upregulated proteins (Table 3) can be linked to the classic E. coli Pho regulon.

Other significantly differentially expressed proteins in В. cenocepacia include Ppk/Ppk2 encodina polyphosphate kinases involved in the storage of Pi, PhnA encoding a phosphonoacetate hydrolase involved in releasing Pi and acetate from phosphonoacetate, and GlpQ, a glycerophosphoryl diester phosphodiesterase involved in releasing Pi from deacetvlated phospholipids (Nocek et al., 2008; Gilbert et al., 2009; Racki et al., 2017; Jorge et al., 2018). Homologues of the following upregulated proteins (Table 3) are induced under Pi starvation in other bacteria: PlcN, BPSL1375 (OlsB), BCAM1663, BCAM0451, BCAL1116, BPSL1190 and BCAL2364. The majority of these proteins are involved in increasing the availability of Pi by degrading phospholipids or extracellular DNA (eDNA) (Korbsrisate et al., 2007; Gangaiah et al., 2009; Mulcahy et al., 2010; Zavaleta-Pastor et al., 2010; Diercks et al., 2015; Ball et al., 2016; Wilton et al., 2018). BCAL1083, encoding a putative alkaline phosphatase, is significantly overexpressed during Pi starvation in B. cenocepacia K56-2. Interestingly, this gene was shown to be highly induced in a polymyxin B-sensitive mutant of B. cenocepacia K56-2 (Loutet et al., 2011). Similarly, three putative phospholipases (BCAM0164, BCAM2729 and BCAM0408) are also highly overrepresented in B. cenocepacia K56-2

Table 2. Significantly, differentially regulated proteins in the Pi-deplete compared with the Pi-replete condition ($P \le 0.05$, identified in Fig. 2) were assigned KEGG BRITE identifiers using BlastKOALA (version 2.2).

KEGG BRITE reference hierarchy identifier	Number of proteins identified						
	Cellula	lar only Extracell		ellular only	Both	fractions	
	Upregulated	Downregulated	Upregulated	Downregulated	Upregulated	Downregulated	
Protein families: Metabolism	05	20	2	2	Λ	1	
koutouu Enzymes	20	30	2	3	4	I	
koulour Fluten kinases	י ס	-	-	-	-	-	
ko01002 Feblidases and minibilions	۲ ۱	1	-	I	-	-	
ko01004 Lipid biosynthesis proteins	1	-	-	-	-	-	
biogynthesis protoins	1	-	-	-	-	-	
ko01007 Amino acid_related enzymes	2	1					
ko01011 Pentidoglycan biosynthesis	_	1	-	-	-	-	
and degradation proteins	_		-	-	-	-	
ko00194 Photosynthesis proteins	1	_	_	_	_	_	
Protein families: Signalling and cellular							
processes			-				
ko02000 Transporters	8	25	-	-	-	2	
ko02022 Two-component system	3	1	-	_	-	-	
ko02035 Bacterial motility proteins	1	2	-	2	_	-	
ko02042 Bacterial toxins	1	-	2	-	-	-	
ko02048 Prokaryotic defence system	-	1	-	-	-	-	
ko04147 Exosome	_	1	-	-	2	-	
ko04812 Cytoskeleton proteins	1	1	-	-	-	-	
ko01504 Antimicrobial resistance	_	1	-	-	-	-	
genes ko00527 Clycocylphocphatidylinacital					0		
(CPI) anabarad protaina	_	-	-	-	2	-	
(GFI)-anchored proteins							
processing							
ko03000 Transcription factors	_	2	_	_	_	_	
ko03009 Ribosome biogenesis	1	_	_	_	_	_	
ko03011 Ribosome	_	1	_	_	_	_	
Ko03016 Transfer RNA biogenesis	_	_	_	1	_	_	
ko03019 Messenger RNA	2	1	_	_	_	_	
biosynthesis							
ko03021 Transcription machinery	1	_	_	-	_	_	
ko03029 Mitochondrial biogenesis	_	2	_	-	_	_	
ko03036 Chromosome and associated	1	2	-	_	_	_	
proteins							
ko03110 Chaperones and folding	-	4	-	-	-	-	
catalysts	10	10			-		
Unclassified	18	48	-	4	5	-	

The number of proteins associated with each KEGG BRITE identifier was calculated for each fraction of the proteome.

during Pi starvation, one of which (BCAM0408) has been shown previously to be highly expressed in a clinical *B. cenocepacia* isolate (IST439) obtained from a CF patient who died from cepacia syndrome (Mira *et al.*, 2011; Stone *et al.*, 2014).

Using the upstream DNA sequences of the upregulated extracellular proteins identified in *B. cenocepacia* K56-2, Multiple Em for Motif Elicitation analysis was undertaken to detect a putative consensus sequence for a *Burkholderia* Pho box (Suppl. Fig. 1). The promoter region of *pstS* from *Burkholderia thailandensis* was also included, which has been shown previously to bind to the PhoB protein in a gel shift assay (Nowak-Lovato *et al.*, 2012). The consensus *Burkholderia* Pho box sequence was identified as follows: 5'-C(TC)

GTCACNNNN(GT)(TC)TTC(AC)(TG)-3'. Interestingly, this Pho box differs from the Pho box consensus sequences described in other bacterial species, e.g. *E. coli* Pho box: 5'-CTGTCATNNNNCTGT(CA)A(CT)-3' (Wanner, 1993; Yuan *et al.*, 2006a; Yuan *et al.*, 2006b; Diniz *et al.*, 2011).

Differential expression of putative virulence factors occurs in response to Pi limitation

Apart from PhoB and PstSCAB, a total of 11 and 18 unique proteins involved in virulence were upregulated and downregulated respectively (Table 4). The following proteins, previously identified as playing a



Fig. 3. Cellular processes related to each transporter identified in the KEGG analysis (Table 2). Transporters highlighted in red are downregulated and those in green are upregulated in Pi-deplete compared with Pi-replete conditions. Some proteins have putative roles in more than one cellular process. Specific substrates related to each protein are shown in brackets.

role in Pi regulation (see section above), also have a role in virulence: GlpQ, BPSL1375, Ppk, BCAM0451, BCAM2720, PlcN and BCAM0408. For respiratory pathogens like B. cenocepacia, pulmonary surfactant can act as a source of Pi as it is comprised of a mixture of lipids and proteins. Between ~70% and 80% of the phospholipid portion of pulmonary surfactant is phosphatidylcholine (PC) which can be degraded by phospholipases produced by the lung tissues, converting PC into glycerophosphocholine (GPC) (Batenburg, 1992; Hurley and McCormick, 2008). In the respiratory pathogen Mycoplasma pneumoniae, GPC is imported into the cell and degraded by GlpQ to release glycerol-3-phosphate which can then be used for many cellular processes, such as phospholipid synthesis. GlpQ is required for pathogenicity against HeLa cells due to its indirect role in hydrogen peroxide formation (Schmidl et al., 2011). A similar observation was found in Streptococcus pneumoniae whereby cytotoxicity was compromised in GlpQ mutants infecting human epithelial cells (Chuang et al., 2015). Ppk was shown in Burkholderia pseudomallei to be important for swarming and swimming motility, as well as biofilm formation (Tunpiboonsak et al., 2010). PlcN and BCAM0408 homologues were shown to cause cytotoxicity in HeLa cells and the BCAM2720 homologue was significantly upregulated in a hamster model for melioidosis (Tuanyok et al., 2006; Korbsrisate et al., 2007; Mira et al., 2011; Stone et al., 2014). BCAM0451 is a homologue of PA3909 (EddB) in P. aeruginosa encoding a DNase which is secreted to degrade eDNA. It has been shown that EddB can be used as a defence mechanism to evade the host immune system, specifically neutrophil extracellular traps. These are DNA lattices containing granular proteins with antimicrobial activity (Wilton et al., 2018). OIsB (BPSL1375) is an enzyme in the first step of creating Pifree ornithine lipids (OLs), which has been well studied in B. cenocepacia (Vences-Guzmán et al., 2012). In P. aeruginosa, OIsB overexpression leads to an increase in biofilm production and better evasion of host immune responses (Kim et al., 2018). These results suggest upregulated putative Pho regulon members are not only

Table 3. Significantly differentially expressed proteins in the Pi-deplete compared with the Pi-replete condition ($P \le 0.05$, identified in Fig. 2) associated with phosphate metabolism.

<i>P</i> ≤ 0.05	Log2 (fold change)	Locus tag (gene name)	Annotation	Sequence lenath	Relat	Relative abundance (%)		
	onango)	20000 kag (gono namo)		longui	+Pi	-Pi	Difference	
Cellular pro	oteome							
+	+2.150	BURCENK562V_C1978	Ornithine-acyl [acyl carrier protein] N-	268	0.0016	0.0072	+0.0056	
++	+7.112	(0/8B) BURCENK562V_C6451 (BPSI 1190)	Glycosyltransferase involved in cell wall biosynthesis	344	0.0006	0.0851	+0.0845	
++	+6.949	BURCENK562V_C6126 (BCAL1083)	Hypothetical protein	654	0.0021	0.2756	+0.2735	
++	+5.228	BURCENK562V_C5565 (BCAM0451)	Hypothetical protein	604	0.0004	0.0158	+0.0154	
++	+4.915	BURCENK562V_C5497	Alkaline phosphatase	472	0.0006	0.0184	+0.0178	
++	+4.863	BURCENK562V_C5498	Alkaline phosphatase	463	0.0004	0.0123	+0.0119	
++	+4.824	BURCENK562V_C3463	Glycerophosphoryl diester	381	0.0010	0.0286	+0.0276	
++	+4.758	BURCENK562V_C1985	Two-component transcriptional	235	0.0023	0.0639	+0.0616	
++	+4.739	BURCENK562V_C1987	Phosphate ABC transporter ATP-	282	0.0072	0.2011	+0.1939	
++	+4.563	BURCENK562V_C6086	Phospholipase C	714	0.0019	0.0463	+0.0444	
++	+4.116	BURCENK562V_C1983	Polyphosphate kinase	687	0.0007	0.0130	+0.0123	
++	+3.455	BURCENK562V_C1984	PAS/PAC sensor signal transduction	439	0.0036	0.0411	+0.0375	
++	+3.054	BURCENK562V_C1990	Phosphate ABC transporter substrate- binding protein. PhoT family	344	0.9185	7.9420	+7.0235	
++	+2.987	BURCENK562V_C1986	Phosphate uptake regulator	234	0.0576	0.4757	+0.4181	
++	+2.969	BURCENK562V_C6165 (BCAL1116)	Putative spermidine/putrescine transport system substrate-binding	342	0.0009	0.0072	+0.0063	
++	+2.903	BURCENK562V_C1988	Phosphate ABC transporter membrane	292	0.0009	0.0070	+0.0061	
++	+2.737	BURCENK562V_C3379	Polyphosphate kinase 2, PA0141	316	0.0009	0.0065	+0.0056	
++	+2.732	BURCENK562V_C1989	Phosphate ABC transporter membrane	328	0.0009	0.0060	+0.0051	
+	+2.157	BURCENK562V_C5264 (BCAM0164)	Histidine phosphatase superfamily (branch 2)	542	0.0005	0.0024	+0.0019	
+	+2.124	BURCENK562V_C4699 (phnA)	Phosphonoacetate hydrolase	436	0.0006	0.0028	+0.0022	
+	+1.656	BURCENK562V_C6452 (BCAL2364)	UDP-2,3-diacylglucosamine	312	0.0009	0.0029	+0.0020	
+	+1.542	BURCENK562V_C3774 (BCAM1663)	Calcineurin-like phosphoesterase	561	0.0007	0.0022	+0.0015	
Extracellula	ar proteome	, , , , , , , , , , , , , , , , , , ,						
+	+8.719	BURCENK562V_C3774 (BCAM1663)	Calcineurin-like phosphoesterase	561	0.0001	0.0356	+0.0355	
++	+7.842	BURCENK562V_C5497 (BCAM0389)	Alkaline phosphatase	472	0.0001	0.0274	+0.0273	
+	+6.944	BURCENK562V_C5498 (BCAM0390)	Alkaline phosphatase	463	0.0001	0.0194	+0.0193	
++	+4.676	BURCENK562V_C5264 (BCAM0164)	Histidine phosphatase superfamily (branch 2)	542	0.0020	0.0577	+0.0557	
++	+3.631	BURCENK562V_C5565 (BCAM0451)	Hypothetical protein	604	0.0727	0.9928	+0.9201	
++	+3.334	BURCENK562V_C4951 (BCAM2720)	Phospholipase C	779	0.0014	0.0159	+0.0145	
+	+2.611	BURCENK562V_C6126 (BCAL1083)	Hypothetical protein	654	0.0307	0.2068	+0.1761	

(Continues)

<i>P</i> ≤ 0.05	Log2 (fold	Locus tag (gene name)	Annotation	Sequence	Relat	tive abund	ance (%)
	onange)	Loods tag (gene hame)	Annotation	longui	+Pi	Pi [0.0061	Difference
+	+2.513	BURCENK562V_C3463 (BCAM1396)	Glycerophosphoryl diester	381	0.0010	0.0061	+0.0051
+	+2.082	BURCENK562V_C5518 (BCAM0408)	Phospholipase C	815	0.0010	0.0045	+0.0035

Significance is related to the log2-fold change of their relative abundance in proteomes.

 $+P \leq 0.05; ++P \leq 0.01.$

involved in Pi regulation but have other direct or indirect roles in virulence.

There are at least six other proteins upregulated under Pi limitation that are not bona fide members of the classic Pho regulon (Table 4). CpdB is a well-characterized phosphodiesterase in E. coli involved in degrading nontransportable organophosphates, which is regulated by carbon source availability (Wanner, 1996). In avian pathogenic E. coli and Salmonella enterica, CpdB is thought to be important for long-term colonization of avian tissues (Liu et al., 2015, 2017). However, CpdB is not involved in virulence in a mouse model for the pathogen Yersinia enterocolitica (Trülzsch et al., 2001). This suggests it may only be a specific virulence factor for avian infections through the gut. The homologue of BCAM1283 in Burkholderia mallei, BMAA0553, was discovered to be a novel virulence factor that interferes with the host cytoskeleton arrangement and has other roles in signalling and adhesion (Memišević et al., 2013, 2015; Bozue et al., 2016). Proteases BPSL0627 and BCAL0849 are likely to be involved in host tissue degradation, though BPSL0627 has yet to be characterized in any species of bacteria (Drevinek et al., 2008). The putative citrate synthase, BCAS0207, is required for optimal virulence and biofilm formation in B. cenocepacia and is less virulent in an alfalfa infection model (Subramoni et al., 2011). Finally, BCAM1453 is a putative LysR family transcriptional regulator which typically regulates genes involved in quorum sensing and virulence (Maddocks and Oyston, 2008). However, its role in B. cenocepacia is unknown.

Although there are a large number of upregulated virulence factors, our results also show many virulence factors are downregulated (Table 4). This includes many members of the siderophore (ornibactin) gene cluster (BURCENK562V_C1523 to BURCENK562V_C1535, Table 5) suggesting a reduction in the amount of siderophore produced under Pi limitation (Agnoli *et al.*, 2006). Another LysR transcriptional regulator, BCAL3178, is downregulated, which is involved in the regulation of the Hmq operon, a quorum-sensing system in many

Burkholderia species (Coulon et al., 2019). Some of the proteins downregulated are involved in defence against macrophages. For example, BPSS1342 is homologous to MAV2054 in Mycobacterium avium, which induces macrophage apoptosis (Lee et al., 2016). BCAM0276, a putative universal stress protein, is highly expressed during intracellular infection. However, its role in protection from macrophage attack is not fully understood (Tolman and Valvano, 2012). Another downregulated protein is an acid phosphatase (BCAM1576) whose homologue in B. thailandensis (BTH_I320) shows similarity to a phospholipase C (PIcH) from P. aeruginosa which is a hemolytic exotoxin (Franco et al., 2018). The final cellular downregulated protein, BCAM0580, has been characterized in B. cenocepacia as RfpR, a key regulator of cyclic-di-GMP signalling. This type of signalling upregulates a range of virulence factors involved in biofilm formation, swarming motility and protease activity (Kumar et al., 2018; Richter et al., 2019). In the extracellular fraction, five proteins are downregulated, three of which are part of the flagellum (FlgL, FliD1 and FliK). Flagella are important for motility as well as adhesion to host cells, both of which are important for virulence (Williams et al., 1996; Ryan et al., 2005; Hanuszkiewicz et al., 2014). Indeed, when swarming assays were performed, B. cenocepacia K56-2 had decreased motility in Pi-deplete conditions (Fig. 4B), likely due to the decrease in flagellar proteins. Another exoprotein involved in adhesion/motility is BCAL1681, part of an operon involved in the assembly of the pilus machinery (Inhülsen et al., 2012). Lastly, BCAL1722 is an exported chitinase that most likely plays a role in plant virulence (O'Grady et al., 2009; Burtnick et al., 2014).

Overall, these results suggest Pi limitation stimulates a shift in virulence factor expression when *B. cenocepacia* experiences Pi stress. In previous studies, although some virulence factors such as swarming motility, exopolysaccharide production and host colonization are known to be increased during Pi stress, many suggest virulence is overall decreased under Pi limitation (Rüberg *et al.*, 1999; Von Krüger *et al.*, 1999; Lamarche

Table 4. Significantly differentially expressed proteins in the Pi-deplete compared with the Pi-replete condition ($P \leq 0.05$, identified in Fig. 2) associated with virulence.

<i>P</i> ≤ 0.05	Log2 (fold change)	Locus tag (gene name)	Annotation	Sequence length	Relative	abundance (%)	
					+Pi	-Pi	Difference
Cellular pro	oteome						
++	+5.228	BURCENK562V_C5565 (BCAM0451)	Hypothetical protein	604	0.0004	0.0158	+0.0154
++	+4.824	BURCENK562V_C3463	Glycerophosphoryl diester phosphodiesterase	381	0.0010	0.0286	+0.0276
++	+4.563	BURCENK562V_C6086 (plcN)	Phospholipase C	714	0.0019	0.0463	+0.0444
++	+4.116	BURCENK562V_C1983	Polyphosphate kinase	687	0.0007	0.0130	+0.0123
++	+4.079	BURCENK562V_C0530	2,3-Cyclic-nucleotide 2-phosphodiesterase/3-nucleotidase	689	0.0003	0.0047	+0.0044
++	+4.010	BURCENK562V_C0366	Putative metalloprotease	253	0.0010	0.0167	+0.0157
++	+3.513	BURCENK562V_C3335	Calcineurin-like phosphoesterase	487	0.0007	0.0080	+0.0073
++	+2.954	BURCENK562V_C3526	Transcriptional regulator, LysR family	319	0.0006	0.0052	+0.0046
+	+2.150	BURCENK562V_C1978	Ornithine-acyl [acyl carrier protein] N-	268	0.0016	0.0072	+0.0056
+	+1.429	BURCENK562V_C7355	Citrate synthase	414	0.0058	0.0164	+0.0106
+	+1.366	(BCAS0207) BURCENK562V_C0351	Protease I	193	0.0147	0.0395	+0.0248
+	-1.612	BURCENK562V_C5692	Transcriptional regulator, LysR family	327	0.1359	0.0463	-0.0896
+	-1.751	BURCENK562V_C2564	Diguanylate cyclase/	667	0.0016	0.0005	-0.0011
+	-1.764	BURCENK562V_C3669	Acid phosphatase	554	0.0213	0.0065	-0.0148
+	-1.817	BURCENK562V_C1530	Putative ATP-binding cassette	581	0.0052	0.0015	-0.0037
++	-2.331	(orbE) BURCENK562V_C1524	Formyl transferase	279	0.0083	0.0017	-0.0066
++	-2.417	(orbF) BURCENK562V_C1528	Amino acid adenylation domain-	1669	0.0009	0.0002	-0.0007
++	-2.593	BURCENK562V_C1523	Protein N-acetyltransferase, RimJ/	338	0.0064	0.0011	0.0053
++	-3.451	(orbL) BURCENK562V_C1535	RimL family Taurine dioxygenase, alpha-	335	0.0101	0.0010	-0.0091
++	-3.476	BURCENK562V_C5387	Nucleotide-binding universal stress	156	0.0211	0.0020	-0.0191
++	-3.948	(BCAM0276) BURCENK562V_C1529 (orbl)	protein, UspA family Non-ribosomal peptide synthase domain TIGR01720/amino acid adenylation domain-containing	3234	0.0024	0.0002	-0.0022
++	-4.482	BURCENK562V_C2521	protein Major membrane protein 1 family	310	0.4929	0.0230	-0.4699
++	-5.254	(BPSS1342) BURCENK562V_C1526	protein ∟-ornithine N5-oxygenase	458	0.0220	0.0006	-0.0214
Extracellula	ar proteome	(pvdA)					
++	+6.330	BURCENK562V_C0530	2,3-Cyclic-nucleotide	689	0.0001	0.0067	+0.0066
++	+3.631	(cpdB) BURCENK562V_C5565	2-phosphodiesterase/3-nucleotidase Hypothetical protein	604	0.0727	0.9928	+0.9201
++	+3.334	(BCAM0451) BURCENK562V_C4951	Phospholipase C	779	0.0014	0.0159	+0.0145
+	+2.513	(BCAM2720) BURCENK562V_C3463	Glycerophosphoryl diester	381	0.0010	0.0061	+0.0051
+	+2.082	(BCAM1396) BURCENK562V_C5518	pnosphodiesterase Phospholipase C	815	0.0010	0.0045	+0.0035
+	-1.740	(BCAM0408) BURCENK562V_C1500 (BCAL1722)	Chitinase family 18	451	0.9304	0.3070	-0.6234

(Continues)

Table 4. Continued

<i>P</i> ≤ 0.05	Log2 (fold change)	l ocus tag (gene name)	Annotation	Sequence	Relative	Relative abundance (
	onango)	Loodo kag (gono hamo)		longui	+Pi	–Pi	Difference
+	-1.835	BURCENK562V_C0662 (flqL)	Flagellar hook-associated protein 3, FlgL	411	0.2751	0.0850	-0.1901
+	-1.977	BURCENK562V_C6832 (fliD1)	Flagellar hook-associated protein 2	502	0.0197	0.0055	-0.0142
+	-3.431	BURCENK562V_C1546 (BCAL1681)	Hypothetical protein	473	0.0537	0.0055	-0.0482
+	-6.226	BURCENK562V_C0711 (BCAL0529)	Hook-length control protein, FliK	450	0.0077	0.0001	-0.0076

Significance is related to the log2-fold change of their relative abundance in proteomes. + $P \leq 0.05$; ++ $P \leq 0.01$.

Table 5. Significantly differentially expressed proteins in the Pi-deplete compared with the Pi-replete condition ($P \le 0.05$, identified in Fig. 2) associated with antimicrobial resistance.

<i>P</i> ≤ 0.05	Log2 (fold	Locus tag (gene name)	Annotation	Sequence	Relat	tive abunda	ance (%)
	onunge)	Loods lag (gene hame)	Amolaton	lengui	+Pi	–Pi	Difference
Cellular pro	oteome			,			
++ '	+4.116	BURCENK562V_C1983 (ppk)	Polyphosphate kinase	687	0.0007	0.0130	+0.0123
++	+3.931	BURCENK562V_C3621 (BCAM1540)	Isoquinoline 1-oxidoreductase, beta subunit	739	0.0004	0.0065	+0.0061
+	+2.150	BURCENK562V_C1978 (olsB)	Ornithine-acyl [acyl carrier protein] N- acyltransferase	268	0.0016	0.0072	+0.0056
+	+1.857	BURCENK562V_C3815 (BCAM1700)	Protein N-acetyltransferase, RimJ/ RimL family	400	0.0009	0.0035	+0.0026
+	+1.578	BURCENK562V_C6450 (dagK)	Diacylglycerol kinase	177	0.0013	0.0041	+0.0028
+	-1.796	BURCENK562V_C1400 (BCAL1812)	Multidrug efflux pump subunit, AcrB	1048	0.0011	0.0003	-0.0008
++	-2.868	BURCENK562V_C4836 (BCAM2620)	D-methionine transport system substrate-binding protein	268	0.2400	0.0342	-0.2058
++	-3.766	BURCENK562V_C3648 (BCAM1559)	Acetyltransferase (GNAT) family protein	186	0.0249	0.0019	-0.0230
++	-5.253	BURCENK562V_C5411 (BCAM0300)	Metallo-beta-lactamase family protein	467	0.0164	0.0004	-0.0160
Extracellula	ar proteome		•				
+	-2.821	BURCENK562V_C4836 (BCAM2620)	D-methionine transport system substrate-binding protein	268	0.5332	0.0832	-0.4500

Significance is related to the log2-fold change of their relative abundance in proteomes. +P \leqslant 0.05; ++P \leqslant 0.01.

et al., 2008b; Bertrand *et al.*, 2010; Crépin *et al.*, 2011; Bains *et al.*, 2012). For example, PhoB has been shown to regulate operons involved in cyclic di-GMP production, a key signalling factor for quorum sensing and biofilm formation. Pi stress stimulates a decrease in c-di-GMP leading to reduced biofilm formation (Haddad *et al.*, 2009; Pratt *et al.*, 2009). Interestingly, here we show that RpfR (BCAM0580) is downregulated under Pi stress, RfpR being a key regulator of c-di-GMP in *Burkholderia* (Richter *et al.*, 2019). However, further experimentation is required to confirm whether this protein would lead to a decreased amount of c-di-GMP and a decrease in virulence under Pi limitation. Despite the suggestion PhoB activation results in decreased virulence, the deletion of PhoB in *Salmonella* causes hypervirulence which in turn decreases *Salmonella* survival in macrophages (Choi *et al.*, 2019). Pyocyanin production (a toxin) in *P. aeruginosa* is stimulated under Pi limitation and is PhoB-dependent, specifically when iron is also present in the media (Jensen *et al.*, 2006). From these examples, it appears that the Pho regulon can stimulate some aspects of virulence whilst indirectly inactivating others and overall has a very complex role to play in bacterial virulence, which could be specific to the pathogen.



Fig. 4. A. Fold change (log2) in protein expression between Pideplete and Pi-replete conditions for the phosphate transporter (PstSCAB) and two-component system (PhoBR) and their neighbouring genes. Proteins are colour-coded for significantly upregulated (green) and non-significantly regulated (yellow). B. Diameter of swarming colonies of *Burkholderia cenocepacia* under Pi-replete and Pi-deplete conditions. N = 5; ***P < 0.001.

Putative antimicrobial resistance proteins are differentially regulated in response to Pi limitation in B. cenocepacia

Burkholderia species are well-studied for their high tolerance to many different classes of antibiotics and antimicrobial peptides, as well as their ability to produce antifungals (Vial et al., 2007; Podnecky et al., 2015; Regan and Bhatt, 2019). Following our results indicating a shift in expression of putative virulence factors as a response to Pi deprivation, we similarly investigated whether the significantly differentially regulated proteins may be also involved in bacterial resistance to antimicrobials (Table 5). OlsB (BPSL1375) and Ppk have been mentioned previously for their roles in virulence and Pi storage. However, they also have roles in resistance. OIsB plays a role in producing OLs under Pi-stress conditions and the resulting change in lipid membrane composition confers resistance to antimicrobial peptides in P. aeruginosa (Kim et al., 2018). A ppk mutant in Campylobacter jejuni was more susceptible to a wide range of antibiotics including erythromycin, ciprofloxacin and cefotaxime (Gangaiah et al., 2009). Overexpression of diacylglycerol kinase (DagK) in Acinetobacter baumannii resulted in increased resistance to colistin through an unknown mechanism (Cafiso et al., 2019). The homoloaue of BCAM1700 in Klebsiella pneumoniae (KPN_00437) is thought to be associated with the functionality of SHV-11, а beta-lactamase (Mirvala

et al., 2020). Finally, the homologue of BCAM1540 in *Pseudomonas putida* (PP_2478) may be involved in tolerance to the toxic compound, 2,4,6-Trinitrotoluene (Fernández *et al.*, 2009).

There were also five proteins downregulated under Pi limitation related to antimicrobial resistance (Table 5). One was detected in both the cellular and extracellular proteome, BCAM2620, a putative lipoprotein that may confer resistance of B. cenocepacia to attack by P. aeruginosa (Bernier et al., 2018). BCAL1812 is the homologue of MexK in P. aeruginosa, one part of an efflux pump involved in expelling triclosan and tetracycline from the cell (Chuanchuen et al., 2002: Godov et al., 2010). The homologue of BCAM1559, AK961_03495, in Serratia marcescens is an anti-toxin protein thought to be involved in protection against nonribosomal peptides (Matteoli et al., 2018). The B. thailandensis homologue for BCAM0300 (BTH_II0462) is a putative metallo-beta-lactamase (Wagley et al., 2014). These results indicate a shift in AMR in response to Pi limitation and suggest that the consequences of this AMR shift could be as complex as the virulence changes we also observed under these conditions.

Conclusions

Inorganic phosphate (Pi) is an essential nutrient for life with many environments containing low Pi concentrations, including soil, oceans and some human host environments. As such, all bacteria require a stress response mechanism for when they experience Pi stress. In the Bcc species, B. cenocepacia K56-2, we identified a total of 182 and 23 proteins that are significantly differentially expressed in the cellular and exoproteome respectively. KEGG analysis highlighted the drastic upregulation and downregulation of different protein classes under Pi limitation, suggesting there is a major shift in cellular metabolism under this condition. Our comparisons have highlighted proteins of the Pho regulon which play a complex role in virulence, as well as other proteins involved in nutrient transport and antimicrobial resistance. Our results begin to unravel the phosphate stress response in Bcc species and serve as a baseline for further research into the biology of phosphorus metabolism in Burkholderia.

Experimental procedures

Cultivation of B. cenocepacia

Burkholderia cenocepacia K56-2 was obtained from the DSMZ culture collection (Germany) and routinely cultured in lysogeny broth (LB) and LB agar (1.5% wt./vol.) at 37°C. A modified minimal A medium, which has been

used in similar experiments with Pseudomonas species. was used to induce phosphate stress (Lidbury et al., 2016). This comprised Na-Succinate 5.4 g L^{-1} . NaCl 200 mg L⁻¹, NH₄Cl 450 mg L⁻¹, CaCl₂ 200 mg L⁻¹ KCl 200 mg L⁻¹, MgCl₂ 450 mg L⁻¹, FeCl₂ 10 mg L⁻¹, $MnCl_2$ 10 mg L⁻¹, 10 mM Hepes buffer at pH 7 and 0.01% (wt./vol.) yeast extract. Na₂HPO₄ was used as the phosphate source at a final concentration of 1 mM (phosphate-replete condition) and 0 mM (phosphate-deplete condition). All components were dissolved in deionized water and filter sterilized (through a 0.22 um pore size filter) before use. For proteomic analysis, a single colony of B. cenocepacia was grown in LB overnight at 37°C in triplicate. Cells were pelleted at 4000 rpm for 10 min at 4°C and gently re-suspended in modified minimal A media with the appropriate phosphate source. The optical density of B. cenocepacia cultures was measured at 600 nm (OD₆₀₀) across a 24-h growth period.

Alkaline phosphatase assay

Alkaline phosphatase activity was measured to determine whether cultures were experiencing phosphate stress. At each time point, samples were incubated in the dark with *p*-nitrophenol phosphate (*p*NPP) at a final concentration of 1 mM or with Tris buffer (pH 8) at a final concentration of 10 mM as a control. After 1 h, cells were pelleted at 13 000 rpm for 10 min. Yellow *p*-nitrophenol (*p*NP) supernatant was measured at 405 nm in triplicate (FLUOstar Omega plate reader, BMG Labtech). These measurements were compared with a *p*NP standard curve to calculate the amount of *p*NP produced in 1 h, normalized to cell density (OD₆₀₀).

Peptide isolation of cellular fraction

Isolation of peptides from the cellular fraction was carried out as previously described (Zadjelovic et al., 2020). After 8 h of growth in phosphate-replete/deplete conditions, biological triplicates of cells were collected by centrifugation at 4000 rpm at 4°C for 10 min. The cell pellet was collected and re-suspended in 3:1 water:LDS sample buffer (LDS with 1% beta-mercaptoethanol (vol./vol.)). Cells were lysed at 95°C with repeated vortexing. 20 µl of each sample was run on a NuPAGE 10% (wt./vol.) Bis-Tris protein gel for 5 min before Coomassie staining. Each protein gel band was excised and cut into 1 mm pieces before being de-stained with 50% (vol./vol.) ethanol, 50 mM ammonium bicarbonate (ABC). Gel pieces were dehydrated with 100% (vol./vol.) ethanol, followed by alkylation with 10 mM Tris(2-carboxyethyl)phosphine, 40 mM 2-chloroacetamide and 50 mM ABC at 70°C for 5 min with shaking. Gel pieces were washed with 50% (vol./vol.) ethanol 50 mM ABC and dehydrated with 100% (vol./vol.) ethanol. Proteins were digested with trypsin overnight and peptides were extracted using repeated 10 min sonication. Peptides were precipitated using a speed vacuum concentrator for 3 h at 45°C and re-suspended in 2% acetonitrile with 0.1% trifluoroacetic acid.

Isolation of peptides from the extracellular fraction

Isolation of peptides from the exoproteome was based on a previously published protocol (Christie-Oleza and Armengaud, 2010). After 8 h of growth in phosphatereplete/deplete conditions, biological triplicates of cells were collected by centrifugation at 4000 rpm at 4°C for 10 min. The supernatant was further filtered (0.22 μm pore size) before exoproteins were precipitated. A final concentration of 0.015% (wt./vol.) sodium deoxycholate (DOC) was added to the filtered supernatant. After a 10 min incubation, a final concentration of 3% (wt./vol.) trichloroacetic acid (TCA) was added followed by a 30 min incubation on ice. Proteins were pelleted at 4000 rpm at 4°C for 15 min. Pellets were re-suspended in water and the addition of DOC and TCA was repeated. Proteins were pelleted at 8000 rpm for 10 min at 4°C, then re-suspended in 1:1 ethanol:ether. Samples were further centrifuged at 13 000 rpm for 15 min at 4°C, after which the pellets were left to dry at room temperature overnight. Peptide isolation continued by SDS-PAGE and trypsin digestion, as described for the cellular fraction.

Proteomics analysis

Extracted peptides were analysed by nanoLC-ESI-MS/ MS using the Ultimate 3000 RSLCnano system and Thermo Orbitrap Fusion mass spectrometer (Dionex/ Thermo Fisher Scientific). The proteome used for peptide analysis was concatenated from protein sequences of the B. cenocepacia K56-2 Valvano genome from the JGI IMG genome portal (Genome ID: 2562617106). Further data analysis was carried out using MaxQuant and Perseus software (Tyanova et al., 2016a; Tyanova et al., 2016b). Peptides without triplicate measures were filtered out. From these data sheets, BlastKOALA was used to assign KEGG to significantly differentially expressed proteins. Clusters of Orthologous Groups of proteins were assigned manually from IMG/M. Details of these methods can be found in the Supplementary Materials.

Swarming assays

Overnight cultures of *B. cenocepacia* in LB medium were collected by centrifugation at 4000 rpm for 10 min at 4°C. Pellets were re-suspended in 10 ml of modified minimal

A media to OD_{600} of 0.15 with a final concentration of 0 or 1 mM Na₂HPO₄ for the Pi-deplete or replete conditions respectively. These cultures were incubated at 37°C at 180 rpm until cultures reached mid-exponential phase (0.6–0.8 OD_{600}). Agarose plates for the swarming assay were made using modified minimal A media with 0.4% (wt./vol.) agarose (Sigma Aldrich) for swarming with no added phosphate. Each plate had 20 ml of medium with agarose and the plates were dried for 30 min in the centre of a laminar flow hood. On each plate, a 5-µl droplet of bacterial culture was plated on the surface. Plates were incubated for 18 h at 37°C before the diameter (mm) of the swarm was measured.

Acknowledgements

We are very grateful to the University of Warwick's Proteomics Research Technology Platform for processing the LC– MS/MS samples. We acknowledge the BBSRC MIBTP (BB/M01116X/1) doctoral training centre and MRC doctoral training centre for funding to H.S. and R.A.J. respectively.

References

- Agnoli, K., Lowe, C.A., Husnain, S.I., and Thomas, M.S. (2006) The ornibactin biosynthesis and transport genes of *Burkholderia cenocepacia* are regulated by an extracytoplasmic function sigma factor which is a part of the fur regulon. *J Bacteriol* **188**: 3631–3644.
- Antelmann, H., Scharf, C., and Hecker, M. (2000) Phosphate starvation-inducible proteins of *Bacillus subtilis*: proteomics and transcriptional analysis. *J Bacteriol* **182**: 4478–4490.
- Baek, J.H., and Lee, S.Y. (2006) Novel gene members in the Pho regulon of *Escherichia coli. FEMS Microbiol Lett* **264**: 104–109.
- Bains, M., Fernández, L., and Hancock, R.E.W. (2012) Phosphate starvation promotes swarming motility and cytotoxicity of *Pseudomonas aeruginosa*. *Appl Environ Microbiol* **78**: 6762–6768.
- Ball, G., Antelmann, H., Imbert, P.R.C., Gimenez, M.R., Voulhoux, R., and Ize, B. (2016) Contribution of the twin arginine translocation system to the exoproteome of *Pseudomonas aeruginosa*. Sci Rep 6: 1–14.
- Banerjee, S.K., Bhatt, K., Misra, P., and Chakraborti, P.K. (2000) Involvement of a natural transport system in the process of efflux-mediated drug resistance in *Mycobacterium smegmatis*. *Mol Gen Genet* **262**: 949–956.
- Batenburg, J.J. (1992) Surfactant phospholipids: synthesis and storage. *Am J Physiol Cell Mol Physiol* **262**: L367–L385.
- Bazzini, S., Udine, C., Sass, A., Pasca, M.R., Longo, F., Emiliani, G., *et al.* (2011) Deciphering the role of RND efflux transporters in *Burkholderia cenocepacia*. *PLoS One* **6**: e18902.
- Bernier, S.P., Son, S., and Surette, M.G. (2018) The Mla pathway plays an essential role in the intrinsic resistance of *Burkholderia cepacia* complex species to antimicrobials and host innate components. *J Bacteriol* **200**: e00156-18.

- Bertrand, N., Houle, S., LeBihan, G., Poirier, É., Dozois, C. M., and Harel, J. (2010) Increased Pho regulon activation correlates with decreased virulence of an avian pathogenic *Escherichia coli* O78 strain. *Infect Immun* **78**: 5324–5331.
- Bozue, J.A., Chaudhury, S., Amemiya, K., Chua, J., Cote, C. K., Toothman, R.G., et al. (2016) Phenotypic characterization of a novel virulence-factor deletion strain of *Burkholderia mallei* that provides partial protection against inhalational glanders in mice. *Front Cell Infect Microbiol* 6: 21. https://doi.org/10.3389/fcimb.2016.00021.
- Brown, K.J., Formolo, C.A., Seol, H., Marathi, R.L., Duguez, S., An, E., *et al.* (2012) Advances in the proteomic investigation of the cell secretome. *Expert Rev Proteomics* **9**: 337–345.
- Burkholder, W.H. (1950) Sour skin, a bacterial rot of onion bulbs. *Phytopathology* **40**: 115–117.
- Burtnick, M.N., Brett, P.J., and DeShazer, D. (2014) Proteomic analysis of the *Burkholderia pseudomallei* type II secretome reveals hydrolytic enzymes, novel proteins, and the deubiquitinase TssM. *Infect Immun* 82: 3214–3226.
- Cafiso, V., Stracquadanio, S., Lo Verde, F., Gabriele, G., Mezzatesta, M.L., Caio, C., *et al.* (2019) Colistin resistant *A. baumannii*: genomic and transcriptomic traits acquired under colistin therapy. *Front Microbiol* **9**: 1–20.
- Choi, S., Choi, E., Cho, Y.J., Nam, D., Lee, J., and Lee, E.J. (2019) The *Salmonella* virulence protein MgtC promotes phosphate uptake inside macrophages. *Nat Commun* **10**: 3326.
- Christie-Oleza, J.A., and Armengaud, J. (2010) In-depth analysis of exoproteomes from marine bacteria by shotgun liquid chromatography-tandem mass spectrometry: the *Ruegeria pomeroyi* DSS-3 case-study. *Mar Drugs* 8: 2223–2239.
- Chuanchuen, R., Narasaki, C.T., and Schweizer, H.P. (2002) The MexJK efflux pump of *Pseudomonas aeruginosa* requires OprM for antibiotic efflux but not for efflux of triclosan. *J Bacteriol* **184**: 5036–5044.
- Chuang, Y.P., Peng, Z.R., Tseng, S.F., Lin, Y.C., Sytwu, H. K., and Hsieh, Y.C. (2015) Impact of the *glpQ2* gene on virulence in a *Streptococcus pneumoniae* serotype 19a sequence type 320 strain. *Infect Immun* 83: 682–692.
- Chugani, S.A., and Greenberg, E.P. (2007) The influence of human respiratory epithelia on *Pseudomonas aeruginosa* gene expression. *Microb Pathog* **42**: 29–35.
- Coulon, P.M.L., Groleau, M.C., and Déziel, E. (2019) Potential of the *Burkholderia cepacia* complex to produce 4-hydroxy-3-methyl-2-alkyquinolines. *Front Cell Infect Microbiol* **9**: 33.
- Crépin, S., Chekabab, S.M., Le Bihan, G., Bertrand, N., Dozois, C.M., and Harel, J. (2011) The Pho regulon and the pathogenesis of *Escherichia coli*. *Vet Microbiol* **153**: 82–88.
- Diercks, H., Semeniuk, A., Gisch, N., Moll, H., Duda, K.A., and Hölzl, G. (2015) Accumulation of novel glycolipids and ornithine lipids in *Mesorhizobium loti* under phosphate deprivation. *J Bacteriol* **197**: 497–509.
- Diniz, M.M.P., Goulart, C.L., Barbosa, L.C., Farache, J., Lery, L.M.S., Pacheco, A.B.F., *et al.* (2011) Fine-tuning control of PhoBR expression in *Vibrio cholerae* by binding

of PhoB to multiple Pho boxes. *J Bacteriol* **193**: 6929–6938.

- Drevinek, P., Holden, M.T.G., Ge, Z., Jones, A.M., Ketchell, I., Gill, R.T., and Mahenthiralingam, E. (2008) Gene expression changes linked to antimicrobial resistance, oxidative stress, iron depletion and retained motility are observed when *Burkholderia cenocepacia* grows in cystic fibrosis sputum. *BMC Infect Dis* **8**: 1–16.
- Eriksson, S., Lucchini, S., Thompson, A., Rhen, M., and Hinton, J.C.D. (2003) Unravelling the biology of macrophage infection by gene expression profiling of intracellular *Salmonella enterica*. *Mol Microbiol* **47**: 103–118.
- Fernández, M., Duque, E., Pizarro-Tobías, P., Van Dillewijn, P., Wittich, R.M., and Ramos, J.L. (2009) Microbial responses to xenobiotic compounds. Identification of genes that allow *Pseudomonas putida* KT2440 to cope with 2,4,6-trinitrotoluene. *J Microbial Biotechnol* 2: 287–294.
- Franco, M., D'haeseleer, P.M., Branda, S.S., Liou, M.J., Haider, Y., Segelke, B.W., and El-Etr, S.H. (2018) Proteomic profiling of *Burkholderia thailandensis* during host infection using bio-orthogonal noncanonical amino acid tagging (BONCAT). *Front Cell Infect Microbiol* **8**: 370.
- Frisk, A., Schurr, J.R., Wang, G., Bertucci, D.C., Marrero, L., Hwang, S.H., *et al.* (2004) Transcriptome analysis of *Pseudomonas aeruginosa* after interaction with human airway epithelial cells. *Infect Immun* **72**: 5433–5438.
- Gagic, D., Ciric, M., Wen, W.X., Ng, F., and Rakonjac, J. (2016) Exploring the secretomes of microbes and microbial communities using filamentous phage display. *Front Microbiol* **7**: 1–19.
- Gangaiah, D., Kassem, I.I., Liu, Z., and Rajashekara, G. (2009) Importance of polyphosphate kinase 1 for *Campylobacter jejuni* viable-but-nonculturable cell formation, natural transformation, and antimicrobial resistance. *Appl Environ Microbiol* **75**: 7838–7849.
- Gilbert, J.A., Thomas, S., Cooley, N.A., Kulakova, A., Field, D., Booth, T., *et al.* (2009) Potential for phosphonoacetate utilization by marine bacteria in temperate coastal waters. *Environ Microbiol* **11**: 111–125.
- Godoy, P., Molina-Henares, A.J., De La Torre, J., Duque, E., and Ramos, J.L. (2010) Characterization of the RND family of multidrug efflux pumps: in silico to in vivo confirmation of four functionally distinct subgroups. *J Microbial Biotechnol* **3**: 691–700.
- Goldsmith, D.B., Crosti, G., Dwivedi, B., McDaniel, L.D., Varsani, A., Suttle, C.A., *et al.* (2011) Development of *phoH* as a novel signature gene for assessing marine phage diversity. *Appl Environ Microbiol* **77**: 7730–7739.
- Haddad, A., Jensen, V., Becker, T., and Häussler, S. (2009) The Pho regulon influences biofilm formation and type three secretion in *Pseudomonas aeruginosa*. *Environ Microbiol Rep* **1**: 488–494.
- Hanuszkiewicz, A., Pittock, P., Humphries, F., Moll, H., Rosales, A.R., Molinaro, A., *et al.* (2014) Identification of the flagellin glycosylation system in *Burkholderia cenocepacia* and the contribution of glycosylated flagellin to evasion of human innate immune responses. *J Biol Chem* 289: 19231–19244.
- Hong, T., Kong, A., Lam, J., and Young, L. (2007) Periplasmic alkaline phosphatase activity and abundance in

Escherichia coli B23 and C29 during exponential and stationary phase. *J Exp Microbiol Immunol* **11**: 8–13.

- Hurley, B.P., and McCormick, B.A. (2008) Multiple roles of phospholipase A2 during lung infection and inflammation. *Infect Immun* **76**: 2259–2272.
- Inhülsen, S., Aguilar, C., Schmid, N., Suppiger, A., Riedel, K., and Eberl, L. (2012) Identification of functions linking quorum sensing with biofilm formation in *Burkholderia cenocepacia* H111. *Microbiology* 1: 225–242.
- Ishige, T., Krause, M., Bott, M., Wendisch, V.F., and Sahm, H. (2003) The phosphate starvation stimulon of *Corynebacterium glutamicum* determined by DNA microarray analyses. *J Bacteriol* **185**: 4519–4529.
- Jensen, V., Löns, D., Zaoui, C., Bredenbruch, F., Meissner, A., Dieterich, G., et al. (2006) RhIR expression in *Pseudomonas aeruginosa* is modulated by the *Pseudomonas* quinolone signal via PhoB-dependent and -independent pathways. J Bacteriol **188**: 8601–8606.
- Jiang, J., Yu, K., Qi, L., Liu, Y., Cheng, S., Wu, M., *et al.* (2018) A proteomic view of *Salmonella Typhimurium* in response to phosphate limitation. *Proteomes* **6**: 19.
- Jorge, A.M., Schneider, J., Unsleber, S., Xia, G., Mayer, C., and Peschel, A. (2018) *Staphylococcus aureus* counters phosphate limitation by scavenging wall teichoic acids from other staphylococci via the teichoicase GlpQ. *J Biol Chem* **293**: 14916–14924.
- Joshi, C., Patel, P., Palep, H., and Kothari, V. (2019) Validation of the anti-infective potential of a polyherbal "Panchvalkal" preparation, and elucidation of the molecular basis underlining its efficacy against *Pseudomonas aeruginosa*. *BMC Complement Altern Med* **19**: 1–15.
- Kim, S.K., Park, S.J., Li, X.H., Choi, Y.S., Im, D.S., and Lee, J.H. (2018) Bacterial ornithine lipid, a surrogate membrane lipid under phosphate-limiting conditions, plays important roles in bacterial persistence and interaction with host. *Environ Microbiol* **20**: 3992–4008.
- Korbsrisate, S., Tomaras, A.P., Damnin, S., Ckumdee, J., Srinon, V., Lengwehasatit, I., *et al.* (2007) Characterization of two distinct phospholipase C enzymes from *Burkholderia pseudomallei*. *Microbiology* **153**: 1907–1915.
- Kumar, B., Sorensen, J.L., and Cardona, S.T. (2018) A c-di-GMP-modulating protein regulates swimming motility of *Burkholderia cenocepacia* in response to arginine and glutamate. *Front Cell Infect Microbiol* **8**: a56.
- Lamarche, M.G., Kim, S.H., Crépin, S., Mourez, M., Bertrand, N., Bishop, R.E., *et al.* (2008a) Modulation of hexa-acyl pyrophosphate lipid A population under *Escherichia coli* phosphate (Pho) regulon activation. *J Bacteriol* **190**: 5256–5264.
- Lamarche, M.G., Wanner, B.L., Crépin, S., and Harel, J. (2008b) The phosphate regulon and bacterial virulence: a regulatory network connecting phosphate homeostasis and pathogenesis. *FEMS Microbiol Rev* **32**: 461–473.
- Lee, K.I., Whang, J., Choi, H.G., Son, Y.J., Jeon, H.S., Back, Y.W., *et al.* (2016) *Mycobacterium avium* MAV2054 protein induces macrophage apoptosis by targeting mitochondria and reduces intracellular bacterial growth. *Sci Rep* **6**: 1–16.
- Leitão, J.H., Feliciano, J.R., Sousa, S.A., Pita, T., and Guerreiro, S.I. (2017) *Burkholderia cepacia* complex infections among cystic fibrosis patients: perspectives and

challenges. In *Progress in Understanding Cystic Fibrosis*, pp. 74–99. London, UK: IntechOpen Limited.

- Li, X.Z., and Nikaido, H. (2004) Efflux-mediated drug resistance in bacteria. *Drugs* 64: 159–204.
- Lidbury, I.D.E.A., Murphy, A.R.J., Scanlan, D.J., Bending, G. D., Jones, A.M.E., Moore, J.D., *et al.* (2016) Comparative genomic, proteomic and exoproteomic analyses of three *Pseudomonas* strains reveals novel insights into the phosphorus scavenging capabilities of soil bacteria. *Environ Microbiol* 18: 3535–3549.
- Liu, H., Chen, L., Si, W., Wang, C., Zhu, F., Li, G., and Liu, S. (2017) Physiology and pathogenicity of *cpdB* deleted mutant of avian pathogenic *Escherichia coli*. *Res Vet Sci* **111**: 21–25.
- Liu, H., Chen, L., Wang, X., Si, W., Wang, H., Wang, C., et al. (2015) Decrease of colonization in the chicks' cecum and internal organs of *Salmonella enterica* serovar *Pullorum* by deletion of *cpdB* by Red system. *Microb Pathog* **80**: 21–26.
- Loutet, S.A., Di Lorenzo, F., Clarke, C., Molinaro, A., and Valvano, M.A. (2011) Transcriptional responses of *Burkholderia cenocepacia* to polymyxin B in isogenic strains with diverse polymyxin B resistance phenotypes. *BMC Genomics* **12**: 472.
- Lubin, E.A., Henry, J.T., Fiebig, A., Crosson, S., and Laub, M.T. (2016) Identification of the PhoB regulon and role of PhoU in the phosphate starvation response of *Caulobacter crescentus. J Bacteriol* **198**: 187–200.
- Luu, L.D.W., Octavia, S., Zhong, L., Raftery, M.J., Sintchenko, V., and Lan, R. (2018) Proteomic adaptation of Australian epidemic *Bordetella pertussis*. *Proteomics* **18**: 1–12.
- Maddocks, S.E., and Oyston, P.C.F. (2008) Structure and function of the LysR-type transcriptional regulator (LTTR) family proteins. *Microbiology* **154**: 3609–3623.
- Mahenthiralingam, E., Urban, T.A., and Goldberg, J.B. (2005) The multifarious, multireplicon *Burkholderia cepacia* complex. *Nat Rev Microbiol* **3**: 144–156.
- Matteoli, F.P., Passarelli-Araujo, H., Reis, R.J.A., Da Rocha, L.O., De Souza, E.M., Aravind, L., *et al.* (2018) Genome sequencing and assessment of plant growthpromoting properties of a *Serratia marcescens* strain isolated from vermicompost. *BMC Genomics* **19**: 1–19.
- Memišević, V., Zavaljevski, N., Pieper, R., Rajagopala, S.V., Kwon, K., Townsend, K., et al. (2013) Novel Burkholderia mallei virulence factors linked to specific host-pathogen protein interactions. Mol Cell Proteomics 12: 3036–3051.
- Memišević, V., Zavaljevski, N., Rajagopala, S.V., Kwon, K., Pieper, R., DeShazer, D., et al. (2015) Mining host-pathogen protein interactions to characterize *Burkholderia mallei* infectivity mechanisms. *PLoS Comput Biol* **11**: 1–28.
- Mira, N.P., Madeira, A., Moreira, A.S., Coutinho, C.P., and Sá-Correia, I. (2011) Genomic expression analysis reveals strategies of *Burkholderia cenocepacia* to adapt to cystic fibrosis patients' airways and antimicrobial therapy. *PLoS One* **6**: e28831.
- Miryala, S.K., Anbarasu, A., and Ramaiah, S. (2020) Role of SHV-11, a class A β -lactamase, gene in multidrug resistance among *Klebsiella pneumoniae* strains and understanding its mechanism by gene network analysis. *Microb Drug Resist* **26**: 1–9.

- Mulcahy, H., Charron-Mazenod, L., and Lewenza, S. (2010) *Pseudomonas aeruginosa* produces an extracellular deoxyribonuclease that is required for utilization of DNA as a nutrient source. *Environ Microbiol* **12**: 1621–1629.
- Nocek, B., Kochinyan, S., Proudfoot, M., Brown, G., Evdokimova, E., Osipiuk, J., *et al.* (2008) Polyphosphatedependent synthesis of ATP and ADP by the family-2 polyphosphate kinases in bacteria. *Proc Natl Acad Sci* USA **105**: 17730–17735.
- Novotna, J., Vohradsky, J., Berndt, P., Gramajo, H., Langen, H., Li, X.M., et al. (2003) Proteomic studies of diauxic lag in the differentiating prokaryote Streptomyces coelicolor reveal a regulatory network of stress-induced proteins and central metabolic enzymes. *Mol Microbiol* **48**: 1289–1303.
- Nowak-Lovato, K.L., Hickmott, A.J., Maity, T.S., Bulyk, M.L., Dunbar, J., and Hong-Geller, E. (2012) DNA binding site analysis of *Burkholderia thailandensis* response regulators. *J Microbiol Methods* **90**: 46–52.
- O'Grady, E.P., Viteri, D.F., Malott, R.J., and Sokol, P.A. (2009) Reciprocal regulation by the CepIR and CciIR quorum sensing systems in *Burkholderia cenocepacia*. *BMC Genomics* **10**: 441.
- Podnecky, N.L., Rhodes, K.A., and Schweizer, H.P. (2015) Efflux pump-mediated drug resistance in *Burkholderia*. *Front Microbiol* **6**: 305.
- Pratt, J.T., McDonough, E.K., and Camilli, A. (2009) PhoB regulates motility, biofilms, and cyclic di-GMP in *Vibrio cholerae*. *J Bacteriol* **191**: 6632–6642.
- Prunty, M.P., Noone, D., and Devine, K.M. (2018) The distinct PhoPR mediated responses to phosphate limitation in *Bacillus subtilis* subspecies *subtilis* and spizizenii stem from differences in wall teichoic acid composition and metabolism. *Mol Microbiol* **109**: 23–40.
- Racki, L.R., Tocheva, E.I., Dieterle, M.G., Sullivan, M.C., Jensen, G.J., and Newman, D.K. (2017) Polyphosphate granule biogenesis is temporally and functionally tied to cell cycle exit during starvation in *Pseudomonas* aeruginosa. Proc Natl Acad Sci USA **114**: E2440–E2449.
- Regan, K., and Bhatt, J. (2019) Eradication therapy for *Burkholderia cepacia* complex in people with cystic fibrosis. *Cochrane Database Syst Rev*(4), CD009876. https://doi.org/10.1002/14651858.CD009876.pub4.
- Richards, G.R., and Vanderpool, C.K. (2012) Induction of the Pho regulon suppresses the growth defect of an *Escherichia coli sgrS* mutant, connecting phosphate metabolism to the glucose-phosphate stress response. *J Bacteriol* **194**: 2520–2530.
- Richter, A.M., Fazli, M., Schmid, N., Shilling, R., Suppiger, A., Givskov, M., et al. (2019) Key players and individualists of cyclic-di-GMP signaling in *Burkholderia* cenocepacia. Front Microbiol **10**: 1–15.
- Rodríguez-García, A., Barreiro, C., Santos-Beneit, F., Sola-Landa, A., and Martín, J.F. (2007) Genome-wide transcriptomic and proteomic analysis of the primary response to phosphate limitation in *Streptomyces coelicolor* M145 and in a $\Delta phoP$ mutant. *Proteomics* **7**: 2410–2429.
- Rosenberg, H., Gerdes, R.G., and Chegwidden, K. (1977) Two systems for the uptake of phosphate in *Escherichia coli. J Bacteriol* **131**: 505–511.
- Rüberg, S., Pühler, A., and Becker, A. (1999) Biosynthesis of the exopolysaccharide galactoglucan in *Sinorhizobium*

meliloti is subject to a complex control by the phosphatedependent regulator PhoB and the proteins ExpG and MucR. *Microbiology* **145**: 603–611.

- Ryan, K.A., Karim, N., Worku, M., Penn, C.W., and O'Toole, P.W. (2005) *Helicobacter pylori* flagellar hookfilament transition is controlled by a FliK functional homolog encoded by the gene HP0906. *J Bacteriol* **187**: 5742–5750.
- Santos-Beneit, F. (2015) The Pho regulon: a huge regulatory network in bacteria. *Front Microbiol* **6**: 402.
- Schmidl, S.R., Otto, A., Lluch-Senar, M., Piñol, J., Busse, J., Becher, D., and Stülke, J. (2011) A trigger enzyme in *Mycoplasma pneumoniae*: impact of the glycerophosphodiesterase *glpq* on virulence and gene expression. *PLoS Pathog* **7**: e1002263.
- Srikumar, S., Kröger, C., Hébrard, M., Colgan, A., Owen, S. V., Sivasankaran, S.K., *et al.* (2015) RNA-seq brings new insights to the intra-macrophage transcriptome of Salmonella typhimurium. *PLoS Pathog* **11**: 1–26.
- Stone, J.K., DeShazer, D., Brett, P.J., and Burtnick, M.N. (2014) Melioidosis: molecular aspects of pathogenesis. *Expert Rev Anti Infect Ther* **12**: 1487–1499.
- Subramoni, S., Nguyen, D.T., and Sokol, P.A. (2011) *Burkholderia cenocepacia* ShvR-regulated genes that influence colony morphology, biofilm formation, and virulence. *Infect Immun* **79**: 2984–2997.
- Taylor, C.J., Anderson, A.J., and Wilkinson, S.G. (1998) Phenotypic variation of lipid composition in *Burkholderia cepacia*: a response to increased growth temperature is a greater content of 2-hydroxy acids in phosphatidylethanolamine and ornithine amide lipid. *Microbiology* **144**: 1737–1745.
- Tolman, J.S., and Valvano, M.A. (2012) Global changes in gene expression by the opportunistic pathogen *Burkholderia cenocepacia* in response to internalization by murine macrophages. *BMC Genomics* **13**: 63.
- Torriani, A. (1990) From cell membrane to nucleotides: the phosphate regulon in *Escherichia coli*. *Bioessays* **12**: 371–376.
- Trülzsch, K., Roggenkamp, A., Pelludat, C., Rakin, A., Jacobi, C.A., and Heesemann, J. (2001) Cloning and characterization of the gene encoding periplasmic 2',3'cyclic phosphodiesterase of *Yersinia enterocolitica* O:8. *Microbiology* 147: 203–213.
- Tuanyok, A., Tom, M., Dunbar, J., and Woods, D.E. (2006) Genome-wide expression analysis of *Burkholderia pseudomallei* infection in a hamster model of acute melioidosis. *Infect Immun* **74**: 5465–5476.
- Tunpiboonsak, S., Mongkolrob, R., Kitudomsub, K., Thanwatanaying, P., Kiettipirodom, W., Tungboontina, Y., and Tungpradabkul, S. (2010) Role of a *Burkholderia pseudomallei* polyphosphate kinase in an oxidative stress response, motilities, and biofilm formation. *J Microbiol* **48**: 63–70.
- Tyanova, S., Temu, T., and Cox, J. (2016a) The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat Protoc* **11**: 2301–2319.
- Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., *et al.* (2016b) The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat Methods* **13**: 731–740.

- Valvano, M.A. (2015) Intracellular survival of *Burkholderia cepacia* complex in phagocytic cells. *Can J Microbiol* **61**: 607–615.
- Van Bogelen, R.A., Olson, E.R., Wanner, B.L., and Neidhardt, F.C. (1996) Global analysis of proteins synthesized during phosphorus restriction in *Escherichia coli*. *J Bacteriol* **178**: 4344–4366.
- Vences-Guzmán, M.Á., Geiger, O., and Sohlenkamp, C. (2012) Ornithine lipids and their structural modifications: from A to E and beyond. *FEMS Microbiol Lett* **335**: 1–10.
- Vial, L., Chapalain, A., Groleau, M.C., and Déziel, E. (2011) The various lifestyles of the *Burkholderia cepacia* complex species: a tribute to adaptation. *Environ Microbiol* **13**: 1–12.
- Vial, L., Groleau, M.C., Dekimpe, V., and Deziel, E. (2007) *Burkholderia* diversity and versatility: an inventory of the extracellular products. *J Microbiol Biotechnol* **17**: 1407–1429.
- Von Krüger, W.M.A., Humphreys, S., and Ketley, J.M. (1999) A role for the PhoBR regulatory system homologue in the *Vibrio cholerae* phosphate-limitation response and intestinal colonization. *Microbiology* **145**: 2463–2475.
- Von Krüger, W.M.A., Santos Lery, L.M., Soares, M.R., Saloum De Neves-Manta, F., Batista E Silva, C.M., Da Costa Neves-Ferreira, A.G., *et al.* (2006) The phosphatestarvation response in *Vibrio cholerae* O1 and *phoB* mutant under proteomic analysis: disclosing functions involved in adaptation, survival and virulence. *Proteomics* **6**: 1495–1511.
- Wagley, S., Hemsley, C., Thomas, R., Moule, M.G., Vanaporn, M., Andreae, C., *et al.* (2014) The twin arginine translocation system is essential for aerobic growth and full virulence of *Burkholderia thailandensis*. *J Bacteriol* **196**: 407–416.
- Wanner, B.L. (1990) Phosphorus assimilation and its control of gene expression in *Escherichia coli*. In *The Molecular Basis of Bacterial Metabolism*, pp. 152–163. Berlin Heidelberg: Springer-Verlag.
- Wanner, B.L. (1993) Gene regulation by phosphate in enteric bacteria. *J Cell Biochem* **51**: 47–54.
- Wanner, B.L. (1996) Phosphorus assimilation and control of the phosphate regulon. In *Escherichia coli and Salmonella: Cellular and Molecular Biology*, pp. 1357–1381. Washington, DC: ASM Press.
- Williams, A.W., Yamaguchi, S., Togashi, F., Aizawa, S.I., Kawagishi, I., and Macnab, R.M. (1996) Mutations in *fliK* and *flhB* affecting flagellar hook and filament assembly in *Salmonella typhimurium. J Bacteriol* **178**: 2960–2970.
- Wilton, M., Halverson, T.W.R., Charron-Mazenod, L., Parkins, M.D., and Lewenza, S. (2018) Secreted phosphatase and deoxyribonuclease are required by *Pseudomonas aeruginosa* to defend against neutrophil extracellular traps. *Infect Immun* **86**: 1–12.
- Yabuuchi, E., Kosako, Y., Oyaizu, H., Yano, I., Hotta, H., Hashimoto, Y., et al. (1992) Proposal of Burkholderia gen. nov. and transfer of seven species of the genus Pseudomonas homology group II to the new genus, with the type species Burkholderia cepacia (Palleroni and Holmes 1981) comb. nov. Microbiol Immunol **36**: 1251–1275.
- Yamada, M., Makino, K., Amemura, M., Shinagawa, H., and Nakata, A. (1989) Regulation of the phosphate regulon of

Escherichia coli: analysis of mutant *phoB* and *phoR* genes causing different phenotypes. *J Bacteriol* **171**: 5601–5606.

- Yang, C., Huang, T.W., Wen, S.Y., Chang, C.Y., Tsai, S.F., Wu, W.F., and Chang, C.H. (2012) Genome-wide PhoB binding and gene expression profiles reveal the hierarchical gene regulatory network of phosphate starvation in *Escherichia coli. PLoS One* **7**: e47314.
- Yoshida, Y., Sugiyama, S., Oyamada, T., Yokoyama, K., and Makino, K. (2010) Identification and characterization of novel phosphate regulon genes, ecs0540-ecs0544, in *Escherichia coli* O157:H7. *Mol Genet Genomics* **284**: 197–205.
- Yuan, Z., Zaheer, R., and Finan, T.M. (2006a) Regulation and properties of PstSCAB, a transport system of *Sinorhizobium meliloti. J Bacteriol* **188**: 1089–1102.
- Yuan, Z.C., Zaheer, R., Morton, R., and Finan, T.M. (2006b) Genome prediction of PhoB regulated promoters in *Sinorhizobium meliloti* and twelve proteobacteria. *Nucleic Acids Res* **34**: 2686–2697.
- Zaborin, A., Romanowski, K., Gerdes, S., Holbrook, C., Lepine, F., Long, J., *et al.* (2009) Red death in *Caenorhabditis elegans* caused by *Pseudomonas*

aeruginosa PAO1. Proc Natl Acad Sci U S A 106: 6327–6332.

- Zadjelovic, V., Chhun, A., Quareshy, M., Silvano, E., Hernandez-Fernaud, J.R., Aguilo-Ferretjans, M.M., *et al.* (2020) Beyond oil degradation: enzymatic potential of *Alcanivorax* to degrade natural and synthetic polyesters. *Environ Microbiol* **22**: 1356–1369.
- Zavaleta-Pastor, M., Sohlenkamp, C., Gao, J.-L., Guan, Z., Zaheer, R., Finan, T.M., *et al.* (2010) *Sinorhizobium meliloti* phospholipase C required for lipid remodeling during phosphorus limitation. *Proc Natl Acad Sci U S A* **107**: 302–307.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Supplementary Table 1. The list of proteins that are detected in the cellular proteome and extracellular proteome. **Appendix S1:** Supplementary material.