

# Weight loss with bariatric surgery or behaviour modification and the impact on female obesity-related urine incontinence: A comprehensive systematic review and meta-analysis

William Sheridan<sup>1</sup> | Ana Sofia Da Silva<sup>2</sup> | Bianca M Leca<sup>3</sup> | Eduard Ostrijas<sup>4</sup> |  
 Ameet G Patel<sup>5</sup> | Simon JB Aylwin<sup>6</sup> | Royce P Vincent<sup>1,7</sup> |  
 Spyros Panagiotopoulos<sup>5</sup> | Shamsi El-Hasani<sup>8</sup> | Carel W le Roux<sup>9</sup> |  
 Alexander D Miras<sup>10</sup> | Linda Cardozo<sup>2</sup> | Georgios K Dimitriadis<sup>1,6</sup> 

<sup>1</sup>Faculty of Life Sciences and Medicine, School of Life Course Sciences, King's College London, London, UK

<sup>2</sup>Department of Urogynaecology, King's College Hospital NHS Foundation Trust, Denmark Hill, London, UK

<sup>3</sup>University Hospitals Coventry and Warwickshire NHS Trust, Coventry, UK

<sup>4</sup>Institute for Translational Medicine, University of Pecs Medical School, Pécs, Hungary

<sup>5</sup>Department of Minimal Access Surgery, King's College Hospital NHS Foundation Trust, Denmark Hill, London, UK

<sup>6</sup>Department of Endocrinology ASO/EASO COM, King's College Hospital NHS Foundation Trust, Denmark Hill, London, UK

<sup>7</sup>Department of Clinical Biochemistry, King's College Hospital NHS Foundation Trust, Denmark Hill, London, UK

<sup>8</sup>Minimal Access and Bariatric Unit, Princess Royal University Hospital, King's College Hospital NHS Foundation Trust, Orpington, UK

<sup>9</sup>Diabetes Complication Research Centre, School of Medicine and Medical Science, UCD Conway Institute, University College Dublin, Belfield, Ireland

## Summary

Women with obesity are at risk of pelvic floor dysfunction with a 3-fold increased incidence of urge urinary incontinence (UUI) and double the risk of stress urinary incontinence (SUI). The National Institute for Health and Care Excellence (NICE) and European Association of Urology (EAU) recommend that women with a body mass index  $\geq 30 \text{ kg/m}^2$  should consider weight loss prior to consideration for incontinence surgery. This systematic review and meta-analysis will assess this recommendation to aid in the counselling of women with obesity-related urinary incontinence (UI). Medical Literature Analysis and Retrieval System online (MEDLINE), EMBASE, Cochrane, ClinicalTrials.gov, and SCOPUS were systematically and critically appraised for all peer reviewed manuscripts that suitably fulfilled the inclusion criteria established a priori and presented original, empirical data relevant to weight loss intervention in the management of urinary incontinence. Thirty-three studies and their outcomes were meta-analysed. Weight loss interventions were associated in a decreased prevalence in UI (OR 0.222, 95% CI [0.147, 0.336]), SUI (OR 0.354, 95% CI [0.256, 0.489]), UUI (OR 0.437, 95% CI [0.295, 0.649]) and improved quality of life (PFDI-20, SMD -0.774 (95% CI [-1.236, -0.312])). This systematic review and meta-analysis provide evidence that weight loss interventions are effective in reducing the prevalence of obesity-related UI symptoms in women. Bariatric surgery in particular shows greater sustained weight loss and improvements in UI prevalence. Further large scale,

**Abbreviations:** BMI, body mass index; CENTRAL, Cochrane Central Register of Controlled Trials; EAU, European Association of Urology; EMBASE, Excerpta Medica; ICIQ-SF, International Consultation on Incontinence Questionnaire Short Form; IU, Urinary incontinence; KHQ, Kings Health Questionnaire; LAGB, Laparoscopic Adjustable Gastric Band; LVSG, Laparoscopic Vertical Sleeve Gastrectomy; MEDLINE, Medical Literature Analysis and Retrieval System online; MeSH, Medical Subject Heading; National Institute for PFDI-20, Pelvic Floor Distress Inventory; NICE, National Institute for Health and Care Excellence; PRIDE, Program to Reduce Incontinence by Diet and Exercise; PRISMA-P, Preferred Reporting Items for Systematic review and Meta-Analysis Protocols; PROSPERO, International Prospective Register of Systematic Reviews; QoL, Quality of Life; RYBG, roux-en-Y Gastric Bypass.; SUI, Stress urinary incontinence; UUI, Urge urinary incontinence.

William Sheridan, Ana Sofia Da Silva and Linda Cardozo, Georgios K Dimitriadis denotes co-first and co-last authors, respectively

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Clinical Obesity* published by John Wiley & Sons Ltd on behalf of World Obesity Federation.

<sup>10</sup>Department of Metabolism, Digestion and Reproduction, Imperial College London, London, UK

#### Correspondence

Georgios K Dimitriadis, Department of Endocrinology ASO/EASO COM, King's College Hospital NHS Foundation Trust, Denmark Hill, London, SE5 9RS UK.  
Email: georgios.dimitriadis@kcl.ac.uk

randomized control trials assessing the effect of bariatric surgery on women with obesity-related UI are needed to confirm this study's findings.

#### KEY WORDS

bariatric surgery, behaviour modification, meta-analysis, obesity, systematic review, urinary incontinence, weight loss

## 1 | INTRODUCTION

Obesity is a growing pandemic, with the World Health Organization (WHO) reporting that obesity has nearly tripled worldwide since 1975.<sup>1</sup> In 2016, more than 1.9 billion adults were overweight (body mass index [BMI]  $\geq 25 \text{ kg/m}^2$ ), and of these, over 650 million were obese (BMI  $\geq 30 \text{ kg/m}^2$ ). Obesity is associated with an increased prevalence in multiple non-communicable diseases.<sup>1</sup>

Urinary incontinence (UI) is defined as the involuntary loss of urine<sup>2</sup> and is a common health condition that can significantly affect an individual's quality of life (QoL).<sup>3</sup> Reportedly, approximately 50% of adult women may experience UI,<sup>4</sup> double the prevalence of UI documented in men.<sup>5</sup> This difference in prevalence is due to the structure of the female urinary tract, as well as risk factors such as pregnancy, childbirth, and hysterectomy that can damage the pelvic floor musculature and connective tissue.<sup>5</sup> Obesity is an independent risk factor for urinary storage (leakage) and UI symptoms<sup>6</sup>; such that a 5 points increase of BMI is associated with a 20%-70% increased risk of UI.<sup>6</sup>

Urge urinary incontinence (UUI) is the involuntary leakage of urine associated with urgency whilst stress urinary incontinence (SUI) presents as involuntary leakage with sneezing or coughing.<sup>2</sup> Women with obesity have a 3-fold increased incidence of urge urinary incontinence (UUI) and double the risk of stress urinary incontinence (SUI).<sup>7,8</sup>

In women with obesity, weight loss interventions have been shown to reduce the frequency of UI.<sup>9-11</sup> Currently, clinical guidance bodies such as the National Institute for Health and Care Excellence (NICE) and the European Association of Urology (EAU) recommend life-style intervention as the first-line treatment for women with UI and a BMI  $\geq 30 \text{ kg/m}^2$ .<sup>12</sup> This encompasses the advice of weight loss through dietary, pharmacological, behavioural therapy interventions, or a combination of these.<sup>12</sup>

Bariatric surgery, including Roux-en-Y gastric bypass (RYBG), Laparoscopic Adjustable Gastric Band (LAGB) and Laparoscopic Vertical Sleeve Gastrectomy (LVSG), is a common intervention to treat morbid obesity (BMI  $\geq 40 \text{ kg/m}^2$ ),<sup>13</sup> but can be an appropriate measure for individuals with a BMI  $\geq 35 \text{ kg/m}^2$  and related co-morbidities such as diabetes and hypertension. Although behavioural intervention studies have shown to produce statistically significant weight loss in participants,<sup>14</sup> bariatric surgery has proven to lead to significantly greater sustained weight loss in patients with obesity compared to non-surgical alternatives.<sup>15</sup>

### What is known on the subject?

- Obesity is a recognized risk factor for UI
- Weight loss has proven to reduce UI and is currently recommended in the form of behavioural interventions.
- Bariatric surgery is known to lead to greater, sustained weight loss in individuals compared to conservative weight loss methods.

### What does this study add?

- This study aims to clarify the effects of surgically induced weight loss on urinary incontinence and put forward recommendations for further areas of research.

This systematic review and meta-analysis were carried out in order to compare and understand better the effects of surgical and non-surgical weight loss interventions for female obesity-related UI.

## 2 | METHODOLOGY

This protocol was developed according to the Preferred Reporting Items for Systematic review and Meta-Analysis Protocols (PRISMA-P), and followed methods outlined in The Cochrane Handbook for Systematic Reviews of Interventions.<sup>16</sup> This systematic review has been registered with PROSPERO (International Prospective Register of Systematic Reviews) with ID number 222714.

### 2.1 | Search strategy

Two reviewers conducted systematic searches of the following databases: Medical Literature Analysis and Retrieval System Online (MEDLINE), Excerpta Medica (EMBASE), Cochrane Central Register of Controlled Trials (CENTRAL), ClinicalTrials.gov, SCOPUS databases. Our key MeSH (Medical Subject Heading) search terms were; "weight loss surgery" OR "bariatric surgery" OR "metabolic surgery" OR "weight

loss" OR "obesity surgery" OR "obesity treatment" OR "gastric sleeve" OR "sleeve gastrectomy" OR "gastric bypass" OR "gastric band" OR "duodenal switch" OR "biliopancreatic diversion" OR "LSG" OR "LVSG" OR "RYGB" OR "LAGB" OR "diet induced weight loss" OR "lifestyle intervention" OR "obesity medication" OR "weight loss medicine" OR "weight loss drug" OR "weight loss device" AND "urinary incontinence" OR "lower urinary tract symptoms" OR "LUTS" OR "stress urinary incontinence" OR "overactive bladder" OR "detrusor overactivity" OR "urgency urinary incontinence" OR "Prolapse" OR "POP".

Moreover, reference lists of selected articles and other literature sources were browsed to ensure a comprehensive literature search was completed. None of the database searches filtered results based on year of publication date, and the last search was carried out in September 2020.

## 2.2 | Study selection

Single case reports, expert opinion manuscripts, letters to the editor, commentaries, conference papers, animal studies, meta-analyses, narrative review articles and articles not in English were excluded. Data were only included on adult (18 years or older), non-pregnant women. Articles were included if they examined the effect of any weight loss intervention on UI. No restrictions were made regarding the intervention type, where a study took place, the number of participants or the duration of follow up. Covidence Software was used to manage the study selection process. Publications were initially screened for any duplicates before being assessed independently and in parallel by two reviewers. Any conflicts regarding the inclusion of a study were met with discussion and consensus. If an agreement had not been reached, arbitration by a third reviewer would have been required.

## 2.3 | Data extraction

Data were extracted independently by three reviewers following Cochrane Public Health Group Data Extraction and Assessment Template to construct our own data extraction template that was pilot-tested and systematically used for each article. Data extracted included; study description (eg, title, primary author, publication year, type of study, number of participants, type of weight loss intervention, and follow-up duration), participant demographics (eg, mean age), pre- and post- weight loss intervention anthropometric parameters (mean weight, mean BMI), UI assessment (eg, types of UI assessed, validated questionnaire[s] used), overall diagnosis (urodynamic tests and UI severity rating), quality of life assessments (scores of validated questionnaires). The primary outcomes were the prevalence of total UI, SUI and UUI before and after weight loss intervention.

## 2.4 | Quality assessment

Each study was assessed for bias using the Newcastle-Ottawa scale. Studies were evaluated on eight factors, categorized into three

groups: selection (including whether the cohort is representative of the population), comparability (assessed on grounds of study design and the analysis performed) and outcome (ie, the assessment of outcome, follow-up rate and adequacy follow-up period).

Stars were awarded per category, with a maximum of four, two and three stars possible for the "selection", 'comparability' and 'outcome' categories respectively. By following the guidelines of the Newcastle-Ottawa scale,<sup>17</sup> two reviewers assessed the studies to be of poor (3 stars or less), fair (4-6 stars), or good (7-9 stars) quality.

## 2.5 | Data synthesis and statistical analysis

All meta-analytical calculations were performed using MedCalc, the same software was used for generating forest plots. Pooled odds ratios (OR) were calculated with 95% confidence intervals (CI) from the extracted data with count data, while pooled standardized mean differences were calculated from the extracted continuous variables, also with 95% CI. The DerSimonian-Laird random-effects model was used in all calculations. Heterogeneity was assessed using Cochran's Q and  $I^2$  statistics. Subgroup analyses with respect to the type of weight loss intervention were also performed for BMI and weight change differences as well as the count data variables where at least two subgroup data were available. Publication bias was assessed by using Egger's and Begg's tests along with a visual evaluation of the funnel plots when at least ten studies were available.

## 3 | RESULTS

Of the 702 references imported for screening, 157 duplicates were removed, leaving 545 articles to be screened against title and abstract, which resulted in 69 papers progressing to a full-text assessment. Due to one sample of patients from the PRIDE study being reported in 5 different articles<sup>10,11,18-20</sup> only Subak et al.<sup>18</sup> was included in the meta-analysis as this paper reported the outcomes of interest of the present study from the first follow up date. The same decision was made for other articles that reported on one sample of patients, including Treacy et al.<sup>21</sup> and Mazoyer et al.,<sup>22</sup> along with Leshem et al.<sup>22</sup> and Shimonov et al.<sup>23</sup> The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA-P) flow diagram is shown in Figure 1, outlining the outcomes of the screening process.

The 33 studies included in the present study, comprise studies from 12 different countries, amassing 5616 participants, with an age range of 18-78 across all intervention types, with sample size ranging from 10 to 1565. Table 1 (25 surgical intervention studies) and Table 2 (8 behavioural intervention studies) provide an overview of the studies included in this systematic review; including the assessment of outcomes used in the study, a summary of findings for change in total UI, and weekly UI episodes as well as Newcastle-Ottawa scale quality assessment scores.




FIGURE 1 PRISMA flow diagram of systematic reviews modified from Moher et al.<sup>25</sup>

### 3.1 | Meta-analysis of outcomes

#### 3.1.1 | Changes in BMI and weight

The standardized mean difference (SMD) in BMI pre- and post-intervention, of both intervention types, was calculated from 18 studies as  $-1.948$  (95% CI  $[-2.363, -1.532]$ ,  $p < .001$ ,  $I^2 = 94.10\%$ ), as shown in Figure 2(A). After subgroup analysis, surgical interventions showed a statistically significant SMD with the value of  $-2.120$  (95% CI  $[-2.543, -1.696]$ ,  $p < .001$ ,  $I^2 = 94.07\%$ ), whilst the two non-surgical interventions yielded statistically non-significant difference in pre- and post- BMI (SMD  $-0.402$ , 95% CI  $[-0.880, 0.0761]$ ,  $p = 0.098$ ,  $I^2 = 0.00\%$ ), as shown in Figures 2(B), (C) respectively. For the six studies that reported weight change, SMD for all intervention types was  $-0.803$  (95% CI  $[-1.160, -0.446]$ ,  $p < 0.001$ ,  $I^2 = 83.74\%$ ), see Figure 3. Out of that pool, weight change was reported in two surgical intervention studies (SMD  $-1.615$ , 95% CI  $[-2.684, -0.546]$ ,  $p = .003$ ,  $I^2 = 78.76\%$ ) and in four behavioural intervention studies (SMD  $-0.545$ , 95% CI  $[-0.683, -0.407]$ ,  $p < .001$ ,  $I^2 = 0.00\%$ ).

#### 3.1.2 | Changes in total UI prevalence

A total of 23 studies reported changes in prevalence of UI before and after weight loss interventions. They yielded a significant pooled odds ratio (OR) of  $0.222$  (95% CI  $[0.147, 0.336]$ ,  $p < .001$ ,  $I^2 = 89.76\%$ ), as shown in Figure 4(A). In the subgroup analysis, 18 surgical studies yielded a statistically significant result (OR  $0.209$ , 95% CI  $[0.148, 0.296]$ ,  $p < .001$ ,  $I^2 = 64.96\%$ ), while the result for five non-surgical interventions did not achieve statistical significance (OR  $0.746$ , 95% CI  $[0.545, 1.020]$ ,  $p = .066$ ,  $I^2 = 52.17\%$ ), shown in Figures 4(B), (C) respectively.

#### 3.1.3 | Change in UI prevalence by subtype

Analysis of the total pool of 20 studies that reported prevalence of SUI pre- and post- weight loss intervention, produced a statistically significant pooled OR of  $0.354$  (95% CI  $[0.256, 0.489]$ ,  $p < .001$ ,  $I^2 = 82.38\%$ ), Figure 5(A). The subgroup analysis of 17 surgical

**TABLE 1** Characteristics of Surgical Intervention Studies

Study	Country	Type of Surgery	Study design and primary outcome	Follow-up (Months)	Sample size (n)	Mean (SD) BMI Pre- and Post-Intervention (kg/m <sup>2</sup> )	Mean (SD) Weight Pre- and Post-Intervention (kg)	Assessment of outcome	Change in Urinary Incontinence Prevalence and/or No. of Episodes	NOS Star Award and Assessment of Quality
Castro 2012 <sup>26</sup>	Brazil	Not Specified.	Prospective Cohort study on the probability of UI before and after surgery	12	17	45.96 (5.77) to 29.97 (3.48)		Clinical interview, questionnaires on UI, KHQ, Modified Oxford Scale, perineometry, physical exam	At baseline, 17/17 Incontinent women. Post-operatively, 5/17 were incontinent. 12 women became continent. p<0.001.	8- Good
Olivera 2012 <sup>27</sup>	United States	RYGB, LAGB, LVSG (not specified)	Prospective cohort study on the impact on UI	37.8	36	45.76 (6.48) to 31.55 p<0.0001		PFIQ, PISQ-12, FSFI	No prevalence data	7- Good
Kim 2017 <sup>28</sup>	Korea	RYGB (57)	Prospective Cohort study on benefit of surgery on LUTS (UI).	12	57	37.5 (5.9) to 28 (4.9) p<0.001		IPSS, OABSS, PPBC, Sandvik questionnaire, BDI	At baseline, 40.74% were incontinent at Post-operatively, 18.51% had incontinence.	7- Good
Romero Talamás 2016 <sup>29</sup>	United States	RYGB (65), LVSG (5), LAGB (2)	Prospective cohort study assessing prevalence of UI, SUI, UUI and other PFDs.	12	72	47.5 (9.4) to 32.7 (8.1) p<0.001		PFDI-20, PFIQ-7, PISQ-12, screening questions, urodynamics, POP-Q	SUI was identified in 60 (83.3%) patients at baseline and 32 (44.4%) at follow-up, p=0.001. UI prevalence at baseline and follow up was 54- and 27 respectively, p>0.001. Urodynamic evaluation confirmed decreased in SUI prevalence after surgery (from 76.9% to 30.8%, P = 0.01). No confirmation in UUI.	7- Good
Wasserberg 2009 <sup>30</sup>	United States	RYGB (21), Duodenal switch (22), LVSG (3)	Prospective Cohort study on impact on PFD (mainly UI).	18.6	46	45 to 75	128 to 79	PFIDI-20, PFIQ-7	At baseline, 71% had incontinence. Post-operatively, incontinence decreased to 39% (P = 0.003)	5- Fair
Whitcomb 2012 <sup>31</sup>	United States	LVSG (3), LAGB (95)	Prospective cohort study on SUI and UUI prevalence.	12	69	39.7 (6.2) to 34 (5.6)		EPIQ, PISQ, FSFI, SF-36	At baseline: Prevalence of SUI was 22/69 (32 %). At 6 months follow up, this decreased to 10/69 (15 %). (p = 0.006) and at 12 months, was 14/69 (20 %) at (p = 0.027)	6- Fair
Knepfle 2016 <sup>32</sup>	France	RYGB (67), LVSG (2)	Prospective cohort study on UI incidence.	11.3	70	44.5 (6.31) to 31.83 (5.83)		PFIDI-20	UI incidence reduced from 57/70 (81%) at baseline to 41/70 (59%) post-intervention (P = 0.003). The number of continent patients increased from 13 at baseline, to 29 at follow up. (P = 0.003)	5- Fair
Bump 1992 <sup>33</sup>	United States	RYGB (13)	Prospective cohort study on impact on UI following bariatric surgery.	14.5		49.4 (7.9) to 33.1 (6.7) p=0.0000001	131.5 (119.9) to 88.1 (17)	Urogynaecologic history, genitourinary physical and neurologic examination, Voiding diary, fluid loss quantitation test, urodynamic evaluation	UI prevalence was 12/13 at baseline, and 3/13 post-surgery (p = 0.004). Number of incontinence episodes/ week before: 13.4 (SD 2.9), and after weight loss: 0.9 (SD 0.5), p = 0.001. Number of incontinence pads used/ day: 2.0 (SD 0.30), decreased to 0.08 (SD 0.08), p = 0.00008	7- Good

(Continues)

TABLE 1 (Continued)

Study	Country	Type of Surgery	Study design and primary outcome	Follow-up (Months)	Sample size (n)	Mean (SD) BMI Pre- and Post- Intervention (kg/m <sup>2</sup> )	Mean (SD) Weight Pre- and Post- Intervention (kg)	Assessment of outcome	Change in Urinary Incontinence Prevalence and/or No. of Episodes	NOS Star Award and Assessment of Quality
Daucher 2010 <sup>34</sup>	United States	RYGB (34)	Prospective Cohort study on the impact on urinary function and other PFDs.	6	34	46 (6) to 33 (6)	PFDI, PFIQ, PISQ-12, POP-Q, 3-day bladder diary	At baseline: 12/34 were incontinent. Median number of incontinent episodes of 6.5 per day (range, 3-34; mean, 2). Postoperatively, 6 incontinent women became continent. Weekly episodes decreased by a median of 2.5 (range, 1-11; mean, 0 p = 0.05) in the 6 women who remained incontinent.	At baseline: 12/34 were incontinent. Median number of incontinent episodes of 6.5 per day (range, 3-34; mean, 2). Postoperatively, 6 incontinent women became continent. Weekly episodes decreased by a median of 2.5 (range, 1-11; mean, 0 p = 0.05) in the 6 women who remained incontinent.	8- Good
Cucicchi 2013 <sup>35</sup>	Italy	RYGB (87)	Prospective cohort study assessing change in prevalence of UI.	12	87	34 (6.7) to 30 (5.9)	[CIQ-SF, PFDI, PFIQ-7, POPDI-6]	At baseline: 58.6% were incontinent. Post-operatively, prevalence decreased to 17.2% and 9.2% at 6- and 12-months respectively (P < 0.0001). At 12 months, 43 (84.3%) of 51 patients became continent. The remaining: 4 had improved UI, 2 had no change, and 2 reported worsening.	At baseline: 58.6% were incontinent. Post-operatively, prevalence decreased to 17.2% and 9.2% at 6- and 12-months respectively (P < 0.0001). At 12 months, 43 (84.3%) of 51 patients became continent. The remaining: 4 had improved UI, 2 had no change, and 2 reported worsening.	7- Good
Ranasinghe 2011 <sup>36</sup>	Australia	LAGB	Retrospective study on the change in UI (and subtypes) symptoms.	120	143	43.5 (6.65) to 35.5 (6.8) p<0.0001	[CIQ-SF, IPSS]	SUI was 27/97 (p= 0.0164) UII prevalence worsened; 17/97 (p= 0.1088)	At baseline, SUI prevalence: 38/97 and UII prevalence: 11/97. Post-operatively, 11 women became continent.	4- Fair
Fujisaki 2019 <sup>37</sup>	Japan	RYGB (8), LVSG (30),	Prospective Cohort study on UI incidence	12	55	Not reported	Pad usage, [CIQ-UI-SF,	SUI prevalence: At baseline = 35, post-surgery = 11 (p = 0.001). UI prevalence: At baseline = 11 post-surgery= 4 (p= ns)	At baseline, SUI prevalence: At baseline = 35, post-surgery = 11 (p = 0.001). UI prevalence: At baseline = 11 post-surgery= 4 (p= ns)	4- Fair
Laungani 2009 <sup>38</sup>	United States	Gastric bypass	Prospective cohort stay on impact on UI symptoms following surgery.	12	58	48 (7) to 32 (4)	[CIQ-SF	At baseline, 0/58 patients never leaked. At 12 months, 16/25 participants had UI resolved and never leaked (p<0.001). The number of patients with resolved or improved UI symptoms was 92% (23/25) at follow up. SUI baseline: 46/58 (79%), follow-up: 0/525 (p=0.001)	6- Fair	

TABLE 1 (Continued)

Study	Country	Type of Surgery	Study design and primary outcome	Follow-up (Months)	Sample size (n)	Mean (SD) BMI Pre- and Post-Intervention (kg/m <sup>2</sup> )	Mean (SD) Weight Pre- and Post-Intervention (kg)	Assessment of outcome	Change in Urinary Incontinence Prevalence and/or No. of Episodes	NOS Star Award and Assessment of Quality
Shimonov 2017 <sup>24</sup>	Israel	LVSG	Prospective cohort study on UI prevalence after surgery.	6	77	42.2 to 33		[ICIQ-Ul-SF, BFLUTSSF, PFDI-20, PISQ-12]	Total UI prevalence at baseline: 29 (37.7%) Total UI prevalence at follow-up: 14. Change: 15 (51.7%) The number of UI episodes decreased by 47% in the intervention group, compared with 28% in the control group.	6- Fair
Scozzari 2013 <sup>39</sup>	Italy	RYGB (18), LVSG (2), LAGB (12)	Prospective cohort study on UI incidence following surgery.	15.6	32	46.3 (6.3) to 31.3 p<0.001	122 to 83.2	PFDI-20	At baseline, Total UI prevalence: 59.4% At follow-up: 50% (p = ns) SUI: no significant decrease UUI-Baseline: 14 (43.8%) Follow-up: 5 (15.6%) (p=0.029)	8- Good
Durigon Keller 2019 <sup>40</sup>	Brazil	RYGB (83)	Prospective Cohort study on weight loss impact on UI severity	12	26	44.96 (6.49) to 29.98 (6.23) p<0.0001		% Weight loss	Change in Urinary Incontinence Prevalence and/or No. of Episodes	NOS Star Award and Assessment of Quality
Bulbulier 2017 <sup>41</sup>	Turkey	LVSG (120)	Prospective Cohort study on impact on UI, SUI, UUI.	6	120	46.17 (5.35) to 31.6 (4.37)		ICIQ-SF, KHQ	At baseline: Total UI prevalence: 26 (100%); Continent 0, mild 7, moderate 12, severe 6, very severe 1. Post-operative UI prevalence: 14; continent 15, mild 6, moderate 3, severe 2.	7- Good
Kuruba 2007 <sup>42</sup>	United States	RYGB (42), LAGB (3)	Prospective cohort study on the impact of bariatric surgery on UI severity.	12	45		EBWL: 64% (range 19 to 91%)	UI severity index by Sandvik	At Baseline: 44 women (100%) Preoperatively, the patients reported slight (4%), moderate (47%), or severe (49%) urinary incontinence. At follow up: - 50% of the patients had resolution of urinary incontinence, 18% had slight, 18% had moderate, 13% had severe symptoms	5- Fair
Mazoyer 2019 <sup>22</sup>	France	RYGB (30), LVSG (24)	Prospective Monocentric Pilot Study on the incidence of urinary symptoms.	12	54		TWI: 33% (Range 9-51)	USP-PFDI-20 (POPD, CRADI, UDI)	95% reported subjective improvement or resolution of UI, 5% reported unresolved UI	7- Good

(Continues)

TABLE 1 (Continued)

Study	Country	Type of Surgery	Study design and primary outcome	Follow-up (Months)	Sample size (n)	Mean (SD) BMI Pre- and Post-intervention (kg/m <sup>2</sup> )	% Weight loss	Assessment of outcome	Change in Urinary Incontinence Prevalence and/or No. of Episodes	NOS Star Award and Assessment of Quality
Subak 2015 <sup>43</sup>	United States	RYGB (1111), Duodenal switch (12), LVSG (33), LAGB (387)	Prospective Cohort study on the change in UI prevalence (after weight loss).	12	1565	-29.5%	Validated participant reported UUI	Baseline total UI prevalence: 772; Follow-up total UI prevalence by subtype: SUI at baseline: 646, Post-operatively: 188 UUI at baseline: 505, Post-operatively: 165	7- Good	
Burgio 2007 <sup>44</sup>	United States	RYGB (101)	Prospective cohort study on the prevalence of UI following bariatric surgery.	6 and 12	101	48.9 (7.2) to 30.2 (5.7)	MESA, UDI, HQ	At baseline: Total UI prevalence: 66.7%. At follow-up: Prevalence decreased to 37% at 12 months (56% reduction). p <0.001 (CI 95% 18.6-40.0). Women with UI who lost at least 18 BMI points, 71% had regained urinary continence 12 months after surgery. SUI prevalence at baseline was 59.8% and decreased to 31.5% at 12 months. (p <0.001) (CI 17.2-39.3). UOJ prevalence at baseline was 35.9% and decreased to 15.2% at 12 months. (p <0.001) (CI 9.3% 10.4-30.9). MUJ prevalence at baseline was 31.6% and decreased to 9.9% at 12 months. (p<0.001) (CI 95% 11.0-30.7).	6- Fair	
O'Boyle 2016 <sup>45</sup>	Ireland	RYGB (57), LVSG (24), LAGB (1)	Prospective cohort study on prevalence of UI.	15	82	50 (6.3) to 34 (6.8)	ICIQ-SF	At baseline: 82(100%) women had UI. 75% of patients complained of moderate to very severe incontinence. At follow up: 55- 83% patients reported reduced symptom severity. 33% reported complete symptom resolution. 37% had moderate to very severe incontinence.	6- Fair	
AitSaid 2017 <sup>46</sup>	France	LVSG and RYGB	Prospective Cohort study on prevalence of SUI and UUI.	12	83	43.7 (7.3) to 30.2 (7)	USP, ICIQ, - SF Wexner scale, QOL	Baseline prevalence: SUI - 49.4%; OAB - 43.3% Follow-up prevalence: SUI - 21.7%, OAB: 7.3%	7- Good	

No correlation between the degree of weight loss and improvement of SUI ( $r = 0.007$ ,  $p = 0.55$ ) and UUI ( $r = 0.05$ ,  $p = 0.64$ ).

TABLE 1 (Continued)

Study	Country	Type of Surgery	Study design and primary outcome	Follow-up (Months)	Sample size (n)	Mean (SD) BMI Pre- and Post-intervention ( $\text{kg}/\text{m}^2$ )	Assessment of outcome	NOS Star Award and Assessment of Quality
Anglim 2018 <sup>47</sup>	Ireland	RYGB (40), LVSG (21)	Prospective Cohort study on impact on UI, and cure rate.	12	61	51 (7) to 33 (9)	ICIQ-SF	At baseline: Total UI prevalence at baseline: 61 6-Fair
Leshem 2018 <sup>48</sup>	Israel	RYGB (7), LAGB (94)	Prospective Cohort study on UI prevalence.	24	101	41.6 (4.6) to 27.5 (4.4)	ICIQ-SF, BFLUTS, PFDI-20, PISQ-12	At follow-up: 27 SUI change in prevalence: 22 to 9 UI (OAB) change: 13 to 5 MUI change: 27 to 13 At follow-up, prevalence of UI was 18.13 had SUI, 2 had MUI, 3 had UUJ.

Abbreviations: RYBG, Roux-en-Y Gastric Bypass; LAGB, Laparoscopic Adjustable Gastric Band; LVSG, Laparoscopic Vertical Sleeve Gastrectomy; BMI, Body Mass Index; EBWL, Excess Body Weight Loss; TWL, Total Weight Loss; KHQ, Kings Health Questionnaire; PFIQ-7, Pelvic floor Impact Questionnaire; PISQ-12, Pelvic Organ Prolapse/Urinary Incontinence Sexual Function Index; FSFI, Female Sexual Function Index; IPSS, International Prostate Symptom Score; OABSS, Overactive Bladder Symptom Score; PPBC, Patient perception of bladder condition; BDI, Beck Depression Inventory; PFDI-20, Pelvic Floor Distress Inventory; POP-Q, Pelvic Organ Prolapse Qualification; EPIC, Epidemiology of Prolapse and Incontinence Questionnaire; SF-36, Short-Form 36 Health Survey; POPDI-6, Pelvic Organ Prolapse Distress Inventory; ICIQ-SF, International Consultation on Incontinence Questionnaire Short Form; MESA, The Medical, Epidemiological, and Social Aspects of Ageing Questionnaire; CRADI, Colorectal-anal Distress Inventory; USP, Urinary Symptoms Profile; UI, Urinary Incontinence; SUI, Stress Urinary Incontinence; UUI, Urge Urinary Incontinence; BFLUTS, Bristol Female Lower Urinary Tract Symptoms; PGII, Patient Global Impression of Improvement; PFM-S, Pelvic Floor Muscle Strength.


**TABLE 2** Characteristics of Behavioural Intervention studies

Study	Country	Description of intervention	Study design and primary outcome	Follow-up (Months)	Sample size (n)	Mean (SD) Weight Pre- and Post-Intervention (kg)	Assessment of outcome	Change in Urinary Incontinence Prevalence and/or No. of Episodes	NOS Star Award and Assessment of Quality	
Subak 2005 <sup>5</sup>	United States	Behavioural modification; intensive group-based medical and behavioural weight loss program. Low-calorie liquid diet (800 kcal per day or less), increased exercise (60 mins daily), taught standard cognitive & behavioural skills. Weekly group sessions led by a nutritionist, exercise physiologist or behavioural therapist.	Randomised controlled trial assessing change in weekly UI episodes.	6	19	99 to 84	A 7-day voiding diary, HQ, UDI, SF-36, Urodynamic studies	Median (IQR) Episodes per week: Total UI: At baseline: 21 (11-33); At follow up: 4 (2-28); % improvement: 50% (30-89); p = 0.005. SUI: At baseline: 8 (2-13); At follow up: 0 (0-4); % improvement: 92% (66-100); p = 0.003. UO: At baseline: 8 (4-20); At follow up: 2 (0-10); % improvement: 70% (16-100), p = 0.03	9- Good	
Phelan 2012 <sup>49</sup>	United States	The Look AHEAD trial promoted weight loss of at least 7% weight loss by the first year through a low calorie, low fat, and portion-controlled diet as well as weekly exercise (175 minutes).	Randomised controlled trial on the change of weekly UI episodes.	12	1385	II -7.7 (7.0); Control 0.7(5.0)	Validated self-report questions	Weight loss intervention was associated with a 20% reduction in the odds of having UI (OR 0.80, 95% CI 0.65-0.98, p = 0.03), compared with control group.	9- Good	
Subak 2009 <sup>6</sup>	United States	Behavioural modification; PRIDE: Program modelled on Look AHEAD. Calorie and fat restriction of 1200 to 1800 kcal daily, depending on initial weight, with less than 30% of calories from fat. Encouraged to increase exercise to 200 mins per week. Behavioural skills, including self-monitoring, stimulus control, and problem-solving, were emphasised.	Randomised controlled trial on the percentage change in number of incontinence episodes.	6	226	98.17 (17) to 90 (17)	7-day diary of voiding	At baseline, SUI was prevalent in 44 women and UUI prevalence was 104, (p<0.001). At follow-up, 32 women (27% reduction) had SUI, and 94 (19% reduction) had UUI. 41% of women in intervention group had >70% reduction in any UI episodes, compared to 22% of control group	At baseline, SUI was prevalent in 44 women and UUI prevalence was 104, (p<0.001). At follow-up, 32 women (27% reduction) had SUI, and 94 (19% reduction) had UUI. 41% of women in intervention group had >70% reduction in any UI episodes, compared to 22% of control group	8- Good
Brown 2006 <sup>50</sup>	United States	Behavioural Modification: Included 3 groups; I, Metformin at 850 mg 2x daily, or placebo twice daily. Through a low-fat diet and moderate-intensity physical activity for at least 150 min each week, the I, aimed to lose a minimum 7% of initial body weight.	Randomised controlled trial on the change in UI prevalence as a result of lifestyle intervention	36	660	-3.4 (8.2)	Self-reported questionnaire	Prevalence of weekly incontinence episodes at baseline was 48.1%. At follow-up, reduced to 38.3% (p<0.001). Intensive lifestyle intervention was associated with a 20% reduction in the odds (OR 0.80 [95% CI 0.64-1.01]) of weekly SUI episodes compared with placebo.	8- Good	


TABLE 2 (Continued)

Study	Country	Description of intervention	Study design and primary outcome	Follow-up (Months)	Sample size (n)	Mean (SD) BMI Pre- and Post-intervention ( $\text{kg}/\text{m}^2$ )	Mean (SD) Weight Pre- and Post-intervention (kg)	Assessment of outcome	Change in Urinary Incontinence Prevalence and/or No. of Episodes	NOS Star Award and Assessment of Quality
Auwad 2008 <sup>51</sup>	UK	Behavioural modification: All participants were offered a commercial run programme of diet and exercise. Anti-obesity medication Orlistat was offered to those who did not lose 5% of starting weight within 9 months.	Prospective Cohort study on the incidence of UI episodes.	20	42	36.2 to 31.9		24-h pad test; KHQ, Perineometry, Modified Oxford score.	At Baseline: 6 had mild incontinence, 26 had moderate incontinence and ten had severe incontinence (before weight loss). At Follow-up: 21 had mild leakage, 15 had moderate leakage, and only four had severe leakage (p < 0.001).	8- Good
Fjerbaek 2020 <sup>52</sup>	Denmark	Behavioural modification: Four individual dietary counselling session (one group received 5 sessions), two training sessions per week for 12 weeks.	Prospective Cohort study assessing UI impact on quality of life after weight loss.	9	24	32.7 (4.54) to 31.2 (4.23) p<0.0001	89.29 (12.58) to 85.5 (12.31)	ICIQ-SF, PGI-I, 24-hour pad test, PFMS	Mean 24hr pad test results (by weight): 38.75g at baseline; 18.5g at follow up (p <0.001). Objective urodynamic baseline UI prevalence: SU-20, UIU-8, MU-8. At follow-up, 8 women (40%) with an initial diagnosis of SUI had become continent.	6- Fair
Subak 2002 <sup>53</sup>	United States	Behavioural modification: Weight loss program - low calorie liquid diet or reduced-calorie solid diet and exercise.	Prospective cohort study on change in incontinence frequency	3	10	38 (10) to 33 (7) p=0.03	106 (32) to 91 (22)	7-day urinary diary	At Baseline: UI episodes per week: 13(SD 10). At Follow-up: UI episodes per week: 8 (SD 10), with a median reduction of 6 episodes per week. Pad use at baseline was 8 and following weight loss there was a 60% decrease in pad use (p <0.04).	7- Good
Gozukara 2014 <sup>54</sup>	Turkey	Behavioural modification: Aimed to produce an average loss of >7% within 6 months	Randomised controlled trial on change in number of UI episodes.	6	163	85.1 (9.7) to 78.3 (10.4)		3-day voiding diary, PFDI, POP-Q, UDI, POPDI, CRADI	At baseline: SUI episodes per 3 days: 7.96 UI episodes per 3 days: 2.85	9- Good

Abbreviations: AHFAD, Action for Health in Diabetes; DSF, Diabetes Support and Education; ILI, Intensive Lifestyle Intervention; PRIDE, Program to Reduce Incontinence by Diet and Exercise.



**FIGURE 2** Forest plot of standardized mean difference in pre- and post- BMI from all interventions (A), Surgical intervention studies (B) and behavioural (non-surgical) intervention studies (C)




**FIGURE 3** Forest plot of standardized mean difference in pre- and post-weight change

interventions for changes in prevalence of SUI was statistically significant for both surgical interventions ( $OR\ 0.312, 95\% CI\ [0.235, 0.414], p < .001, I^2 = 61.68\%$ ) and non-surgical interventions ( $OR\ 0.754, 95\% CI\ [0.614, 0.926], p = .007, I^2 = 0.00\%$ ), shown in Figures 5(B), (C) respectively.

The calculated OR for UUI with regard to any weight loss intervention was statistically significant ( $OR\ 0.437, 95\% CI\ [0.295, 0.649], p = .001, I^2 = 77.39\%$ ), shown in Figure 6(A). When considering UUI changes by intervention type; meta-analysis of the 15 surgical intervention studies showed a statistically significant reduction in UUI prevalence ( $OR\ 0.378, 95\% CI\ [0.259, 0.552], p < .001, I^2 = 65.49\%$ ), shown in Figure 6(B), whilst the two non-surgical ones did not ( $OR\ 0.943, 95\% CI\ [0.661, 1.344], p = .745, I^2 = 0.00\%$ ), shown in Figure 6(C).

Meanwhile combined analysis of six studies demonstrated changes in mixed UI prevalence ( $OR\ 0.330, 95\% CI\ [0.189, 0.576], p < .001, I^2 = 59.10\%$ ), all of which were surgical interventions, shown in Figure 7.



**FIGURE 4** Forest plot of OR values of UI prevalence from all interventions (A), Surgical intervention studies (B) and behavioural (non-surgical) intervention studies (C)

### 3.1.4 | Change in questionnaire scores

Several questionnaires were used to assess improvement of urinary function and QoL in the observed studies. Statistically significant results were obtained while calculating SMDs in all questionnaires of interest. For Pelvic Floor Distress Inventory (PFDI-20), the calculated SMD was  $-0.774$  (95% CI  $[-1.236, -0.312]$ ,  $p = .001$ ,  $I^2 = 93.42\%$ ), shown in Figure 8(A). For International Consultation on Incontinence Questionnaire Short Form (ICIQ-Urinary SF), SMD was  $-1.110$  (95% CI  $[-1.497, -0.723]$ ,  $p < .001$ ,  $I^2 = 89.68\%$ ), as shown in Figure 8(B). Finally, for Urinary Distress Inventory (UDI), SMD was  $-0.742$  (95% CI  $[-0.911, -0.572]$ ,  $p < .001$ ,  $I^2 = 40.10\%$ ), shown in Figure 8(C).


## 4 | DISCUSSION

The results of this meta-analysis demonstrate that the prevalence of UI before and after weight loss was significantly reduced (OR 0.222,

95% CI [0.147, 0.336]). Subgroup analysis of interventions showed surgical weight loss interventions were associated with greater reductions in the odds of patients experiencing UI (OR 0.209, 95% CI [0.148, 0.296]) compared with behavioural intervention studies (OR 0.746, 95% CI [0.545, 1.020]).

SUI and UUI prevalence also significantly improved after weight loss (OR 0.354, 95% CI [0.256, 0.489] and OR 0.437, 95% CI [0.295, 0.649], respectively). In addition, comparison of interventions showed patients who underwent weight loss through surgical means, experienced significantly decreased odds ratios in both SUI and UUI prevalence at follow-up (OR 0.312, 95% CI [0.235, 0.414] and OR 0.378, 95% CI [0.259, 0.552] respectively), whilst behavioural interventions showed a weaker improvement in SUI (OR 0.754, 95% CI [0.614, 0.926]) and a non-significant improvement in UUI (OR 0.943, 95% CI [0.661, 1.344]).

The evidence from this meta-analysis showed the studies where bariatric surgery had been performed led to greater reductions in BMI (SMD  $-2.120$ , 95% CI  $[-2.543, -1.696]$ ), which may explain the greater improvement in UI, SUI, and UUI prevalence, compared with



**FIGURE 5** Forest plot of OR values of stress urinary incontinence prevalence from all interventions (A), Surgical intervention studies (B) and behavioural (non-surgical) intervention studies (C)

behavioural weight loss interventions that demonstrated less significant reductions in BMI ( $SMD -0.402$ , 95% CI  $[-0.880, 0.0761]$ ).


The Patient Reported Outcome Questionnaires included in this study were PFDI-20, ICIQ-UI-SF, and UDI, commonly used to assess UI, the severity of incontinence symptoms and their effects on quality of life. Pooled analysis of these scores showed statistically significant reductions in standardized mean differences of the scores from baseline to follow-up, suggesting that weight loss was associated with a significant improvement in UI prevalence and patient's quality of life.

Whilst the underlying pathophysiology of UI is multifactorial and its aetiology not completely known, obesity is a recognized risk factor,<sup>6,33,55,56</sup> and a dose-response relationship between increased weight and UI incidence has been observed.<sup>7</sup> A study by Noblet et al. showed a strong correlation between BMI, intra-abdominal pressure and intravesical (bladder) pressure, with a Pearson correlation coefficient value of 0.76 and 0.71, respectively, both  $p < .0001$ .<sup>57</sup> It was hypothesised that this increased intra-abdominal pressure increases stress on pelvic floor organs and supporting structures, and that such stress is responsible for the symptoms presented in SUI.<sup>57</sup> Similarly,


an increase in abdominal pressure as a result of increased abdominal adiposity, may also be responsible for detrusor instability (detrusor muscle overactivity) which can cause the symptoms of urgency that present in UUI.<sup>9</sup>

A cohort study examining the effects of surgical weight loss interventions highlighted that reduced abdominal adiposity was associated with significant decreases in urinary bladder pressure.<sup>58</sup> The effects of reduced bladder pressure and SUI were observed by Bump et al., in whose study, patients' intravesical pressure readings were significantly improved at follow-up and associated with a resolution in SUI in 7 of the 10 patients who complained of SUI at baseline.<sup>33</sup>

Traditional management of UUI can include bladder retraining and lifestyle changes, which typically advise avoidance of bladder irritants and reducing overall fluid intake.<sup>59</sup> Both carbonated drinks and caffeinated beverages are considered to be bladder irritants.<sup>60,61</sup> Following bariatric surgery, patients are advised to avoid bladder irritants and drinking fluids with meals.<sup>62</sup> In a similar way, the dietary changes promoted in behavioural studies involve reduced calorie intake, which is likely to be associated with reduced fluid intake. The




**FIGURE 6** Forest plot of OR values of urge urinary incontinence prevalence from all interventions (A), Surgical intervention studies (B) and behavioural (non-surgical) intervention studies (C)



**FIGURE 7** Forest plot of OR values of MUI prevalence from all interventions

changes in the amount of fluid intake or type of fluids being consumed may be contributing to the observed improvement in UUI symptoms from the weight loss intervention studies, and not due to weight loss alone. Current understanding of the effects of these dietary factors on urgency symptoms would benefit from further research.

The impact on UI severity following weight loss was only reported in four of the included studies assessing bariatric surgery as an intervention. In each, the number of participants experiencing severe UI decreased significantly.<sup>35,40,42,45</sup> For example, in the study conducted by Kuruba et al, 49% of women complained of severe UI at baseline compared with only 13% at 12 months post-surgery. Furthermore, compared with surgical procedures that treat UI symptoms alone, bariatric surgery can provide further health benefits to obesity-related co-morbidities such as diabetes and hypertension, as well as improved mortality.<sup>63</sup> Despite the paucity in evidence that shows



**FIGURE 8** Forest plot of standardized mean difference in validated questionnaire scores; PFDI-20 (A), ICIQ-UI-SF(B) and UDI (C)

bariatric surgery-induced weight loss leads to significant improvements in UI severity, it is tenable that women with UI that are diagnosed with severe obesity ( $BMI \geq 35 \text{ kg/m}^2$ ), could be counselled on the plethora of benefits of bariatric surgery.

Adverse effects were not reported in any behavioural studies and only discussed in one study assessing bariatric surgery.<sup>30</sup> Adverse effects of bariatric surgery are well established and vary depending on the type of procedure carried out. Identifying all associated adverse effects is beyond the scope of this study but should nevertheless be taken into account if bariatric surgery is to be considered as a treatment option. In addition, the short-term cost of bariatric surgery, although beyond the scope of this review, is something to be considered. However, by preventing long-term complications of UI and other obesity-related comorbidities, it would likely prove cost effective in the long-term.<sup>64</sup>

The limitations of the present study include a disproportionately small number of controlled studies assessing surgical weight loss interventions on UI, compared to a large number of behavioural weight loss intervention studies most of which were randomized control trials. This highlights a paucity of randomized controlled trials evaluating the effects of surgical weight loss interventions on female obesity

related IU. The lack of comparison groups suggests the evidence provided by these manuscripts may be of low certainty.

Furthermore, there was a lack of consistency among the use of Patient Reported Outcome Questionnaires to evaluate patients' UI symptoms and QoL. This meant pooled analysis of scores from infrequently used questionnaires such as the IPSS, BFLUTS, SF-36, USP, and Sandvik severity scale could not be completed. Regarding heterogeneity, meta-analysis of study outcomes showed inconsistency was substantial for many of them ( $I^2 > 75\%$ ), despite using a random effects model. This heterogeneity was particularly evident across the outcomes of the surgical intervention studies and may be a result of different bariatric surgery techniques being used. Alternatively, this could be a result of inconsistent follow-up times across studies, some of which were likely too short for maximum weight loss to occur, hence limiting the significance of comparability.

The majority of studies used subjective measures for improvement in symptoms such as the common use of Patient Reported Outcome questionnaires<sup>9,22,24,26-32,34-52,54</sup> and voiding diaries,<sup>9,18,33,34,53,54</sup> which are arguably the most relevant for clinical care. However, compared with objective urodynamic measurements,<sup>9,26,29,33,51</sup> the results are

subject to patient response bias and their ability to accurately remember and complete these forms of assessment.<sup>65</sup>

## 5 | CONCLUSION

This is the first meta-analysis to study both bariatric surgery and behavioural weight loss interventions and their effects on UI, SUI, and UUI. Across all studies, weight loss was observed to be beneficial in improving UI. Specifically, bariatric surgery was associated with significantly reduced UI prevalence and sustained weight loss compared with behavioural interventions. The implication of these results from this study is that weight loss is an effective treatment for UI management in women with obesity. The authors of this paper recommend that more large-scale trials, with more robust study designs, are carried out to determine the relationship (linear or otherwise) between the degree of weight loss and impact on UI, and assess whether bariatric surgery should be offered to women with obesity-related UI, if other non-surgical UI management options have failed.

## ACKNOWLEDGEMENTS

The authors are grateful to Francesco Rubino and Dudley Robinson for their support. We are grateful to Alexander Miras for providing access to Covidence systematic review software. The data that support the findings of this manuscript are available by the corresponding author upon reasonable request.

## CONFLICT OF INTEREST

The authors declare no potential conflict of interest.

## ORCID

Georgios K Dimitriadis  <https://orcid.org/0000-0002-6662-804X>

## REFERENCES

- WHO. Obesity and overweight. Accessed Aug 31, 2020
- Haylen BT, de Ridder D, Freeman RM, et al. An International Urogynecological Association (IUGA)/International Continence Society (ICS) joint report on the terminology for female pelvic floor dysfunction. *NeurourolUrodyn*. 2010;21:5-26. <https://doi.org/10.1002/nau>.
- Tang DH, Colayco DC, Khalaf KM, et al. Impact of urinary Incontinence on healthcare resource utilization, health-related quality of life and productivity in patients with overactive bladder. *BJU Int*. 2014; 113(3):484-491. <https://doi.org/10.1111/bju.12505>.
- Minassian VA, Stewart WF, Wood GC. Urinary Incontinence in women: variation in prevalence estimates and risk factors. *Obstet Gynecol*. 2008;111(2 Pt 1):324-331. <https://doi.org/10.1097/01.AOG.0000267220.48987.17>.
- Abrams, P; Cardozo, Linda; Wagg, Adrian; Wein, A. J. (Alan J); International continence society. *Incontinence: 6th International Consultation on Incontinence*, Tokyo, International Continence Society; 2016:1-141.
- Subak LL, Richter HE, Hunskaar SO. Urinary incontinence: epidemiology and clinical research update. *J Urol*. 2009;182(6, Supplement):S2-S7. <https://doi.org/10.1016/j.juro.2009.08.071>.
- Hunskaar SA. Systematic review of overweight and obesity as risk factors and targets for clinical Intervention for urinary Incontinence in women. *NeurourolUrodyn*. 2008;27(8):749-757. <https://doi.org/10.1002/nau.20635>.
- Vaughan Camille P, Anssi A, Rufus C, et al. Impact of obesity on urinary storage symptoms: results from the FINNO study. *J Urol*. 2013; 189(4):1377-1382. <https://doi.org/10.1016/j.juro.2012.10.058>.
- Subak LL, Whitcomb E, Shen H, Saxton J, Vittinghoff E, Brown JS. Weight loss: a novel and effective treatment for urinary incontinence. *J Urol*. 2005;174(1):190-195. <https://doi.org/10.1097/01.ju.0000162056.30326.83>.
- Wing RR, West DS, Grady D, et al. Program to reduce Incontinence by diet and exercise group. Effect of weight loss on urinary Incontinence in overweight and obese women: results at 12 and 18 months. *J Urol*. 2010;184(3):1005-1010. <https://doi.org/10.1016/j.juro.2010.05.031>.
- Wing RR, Creasman JM, Smith West D, et al. Improving urinary Incontinence in overweight and obese women through modest weight loss. *Obstet Gynecol*. 2010;116(2 Pt 1):284-292. <https://doi.org/10.1097/AOG.0b013e3181e8fb60>.
- National Institute for Health and Care Excellence. Overview|Urinary Incontinence and Pelvic Organ Prolapse in Women: Management|Guidance|NICE. 2019.
- Buchwald H, Oien DM. Metabolic/bariatric surgery worldwide 2011. *Obes Surg*. 2013;23(4):427-436. <https://doi.org/10.1007/s11695-012-0864-0>.
- Norris SL, Zhang X, Avenell A, Gregg E, Schmid CH, Lau J. Long-term non-pharmacological weight loss interventions for adults with predabetes. *Cochrane Database Syst Rev*. 2005. <https://doi.org/10.1002/14651858.cd005270>.
- Adams TD, Davidson LE, Litwin SE, et al. Weight and metabolic outcomes 12 years after gastric bypass. *N Engl J Med*. 2017;377(12): 1143-1155. <https://doi.org/10.1056/NEJMoa1700459>.
- Cumpston J, Li T, Page MJ, et al. Updated guidance for trusted systematic reviews: a new edition of the Cochrane Handbook for Systematic Reviews of Interventions. *Cochrane Database of Systematic Reviews*. 2019. <https://doi.org/10.1002/14651858.ED000142>.
- Wells G, Shea B, O'Connell D, Peterson J. The Newcastle-Ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. *Ott Ott Hosp Res Inst*. 2000.
- Subak LL, Wing R, DS W, et al. Weight loss to treat urinary incontinence in overweight and obese women. *N Engl J Med*. 2009;360(5 PG-481-90):481-490. <https://doi.org/10.1056/NEJMoa0806375>.
- Myers DL, Sung VW, Richter HE, Creasman J, Subak LL. Program to Reduce Incontinence by Diet and Exercise (PRIDE). *Female Pelvic Medicine & Reconstructive Surgery*. 2012;18(1):55-59. <https://doi.org/10.1097/SPV.0b013e31824171f9>.
- Breyer BN, Creasman JM, Richter HE, et al. Behavioral weight loss program and nonurinary Incontinence lower urinary tract symptoms in overweight and obese women with urinary Incontinence: a secondary data analysis of PRIDE. *J Urol*. 2018;199(1):215-222. <https://doi.org/10.1016/j.juro.2017.07.087>.
- Treacy P-J, Mazoyer C, Falagario U, Iannelli A. Sexual activity after bariatric surgery: a prospective monocentric study using the PISQ-IR questionnaire. *J Sex Med*. 2019;16(12):1930-1937. <https://doi.org/10.1016/j.jsxm.2019.09.004>.
- Mazoyer C, Treacy P, Turchi L, Lehur PA, Benizri E, Iannelli A. Laparoscopic roux-En-Y gastric bypass versus sleeve gastrectomy on pelvic floor disorders in morbidly obese women: a prospective monocentric pilot study. *Obes Surg*. 2019;29(2):609-616. <https://doi.org/10.1007/s11695-018-3577-1>.
- Leshem A, Shimonov M, Amir H, Gordon D, Groutz A. Effects of bariatric surgery on female pelvic floor disorders. *Urology*. 2017;105:42-47. <https://doi.org/10.1016/j.urology.2017.03.003>.
- Shimonov M, Groutz A, Schachter P, Gordon DI. Bariatric surgery the answer to urinary Incontinence in obese women? *Neurourol Urodyn*. 2017;36(1):184-187. <https://doi.org/10.1002/nau.22909>.

25. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6(7):e1000097. <https://doi.org/10.1371/journal.pmed.1000097>.
26. Castro LA, Sobottka W, Baretta G, Freitas AC. Effects of bariatric surgery on pelvic floor function. *Arq Bras Cir Dig.* 2012;25(4):263-268. <https://doi.org/10.1590/S0102-67202012000400010>.
27. Olivera CK, Herron DM, Kini SU, et al. Long-term quality of life and pelvic floor dysfunction after bariatric surgery. *Am. J. Obstet. Gynecol.* 2012;207(5):431.e1-431.e4. <https://doi.org/10.1016/j.ajog.2012.06.041>.
28. Kim JH, Sun HY, Lee HY, et al. Improvement of voiding characteristics in morbidly obese women after bariatric surgery: a single-center study with a 1-year follow-up. *Surg Obes Relat Dis.* 2017;13(5):836-841. <https://doi.org/10.1016/j.soard.2017.01.047>.
29. Romero-Talamás H, Unger CA, Aminian A, Schauer PR, Barber M, Brethauer S. Comprehensive evaluation of the effect of bariatric surgery on pelvic floor disorders. *Surg Obes Relat Dis.* 2016;12(1):138-143. <https://doi.org/10.1016/j.soard.2015.08.499>.
30. Wasserberg N, Petrone P, Haney M, Crookes PF, Kaufman HS. Effect of surgically induced weight loss on pelvic floor disorders in morbidly obese women. *Ann Surg.* 2009;249(1):72-76. <https://doi.org/10.1097/SLA.0b013e31818c7082>.
31. Whitcomb EL, Horgan S, Donohue MC, Lukacz ES. Impact of surgically induced weight loss on pelvic floor disorders. *Int Urogynecology J.* 2012;23(8):1111-1116. <https://doi.org/10.1007/s00192-012-1756-5>.
32. Knepfler T, Valero E, Triki E, Chilintseva N, Koensgen S, Rohr S. Bariatric surgery improves female pelvic floor disorders. *J Visc Surg.* 2016;153(2):95-99. <https://doi.org/10.1016/j.jviscsurg.2015.11.011>.
33. Bump RC, Sugerman HJ, Fantl JA, McClish DK. Obesity and lower urinary tract function in women: effect of surgically induced weight loss. *Am J Obstet Gynecol.* 1992;167(2):392-399. [https://doi.org/10.1016/S0002-9378\(11\)91418-5](https://doi.org/10.1016/S0002-9378(11)91418-5).
34. Daucher JA, Ellison RE, Lowder JL. Pelvic support and urinary function improve in women after surgically induced weight reduction. *Female Pelvic Med Reconstr Surg.* 2010;16(5):263-267. <https://doi.org/10.1097/SPV.0b013e3181eeedb30>.
35. Cuicchi D, Lombardi R, Cariani S, Leuratti L, Lecce F, Cola B. Clinical and instrumental evaluation of pelvic floor disorders before and after bariatric surgery in obese women. *Surg Obes Relat Dis.* 2013;9(1):69-75. <https://doi.org/10.1016/j.soard.2011.08.013>.
36. Ranasinghe WKB, Wright T, Attia J, et al. Effects of bariatric surgery on urinary and sexual function. *BJU Int.* 2011;107(1):88-94. <https://doi.org/10.1111/j.1464-410X.2010.09509.x>.
37. Fujisaki A, Mizumura Y, Haruyama I, Seki Y, Kasama K, Yoshimura Y. Time-dependent changes in lower urinary tract symptoms in Japanese patients who undergo bariatric and metabolic surgery. *Int J Urol.* 2019;26(4):524-526. <https://doi.org/10.1111/iju.13925>.
38. Laungani RG, Seleno N, Carlin AM. Effect of laparoscopic gastric bypass surgery on urinary Incontinence in morbidly obese women. *Surg Obes Relat Dis.* 2009;5(3):334-338. <https://doi.org/10.1016/j.soard.2008.12.003>.
39. Scovazzi G, Rebecchi F, Giaccone C, Chiaro P, Misstrangelo M, Morino M. Bariatric surgery improves urinary Incontinence but not anorectal function in obese women. *Obes Surg.* 2013;23(7):931-938. <https://doi.org/10.1007/s11695-013-0880-8>.
40. Durigon Keller, K.; La Rosa, V. L.; Marques Cerentini, T.; Machado de Souza, C.; Langlois Costa, F.; Viana da Rosa, P.; da Silva Klahr, P.; de Almeida Pereira, E.; Telles da Rosa, L. H. Quality of life and urinary Incontinence symptoms in women undergoing bariatric surgery. *Female Pelvic Med Reconstr Surg.* 2020;26(11):e62-e67. <https://doi.org/10.1097/SPV.0000000000000757>.
41. Bulbulla N, Habibi M, Yuksel M, et al. Effects of bariatric surgery on urinary Incontinence. *Ther Clin Risk Manag.* 2017;13:95-100. <https://doi.org/10.2147/TCRM.S125781>.
42. Kuruba R, Almahmeed T, Martinez F, et al. Bariatric surgery improves urinary Incontinence in morbidly obese individuals. *Surg Obes Relat Dis.* 2007;3(6):586-590. <https://doi.org/10.1016/j.soard.2007.08.007>.
43. Subak LL, King WC, Belle SH, et al. Urinary Incontinence before and after bariatric surgery. *JAMA Intern Med.* 2015;175(8):1378-1387. <https://doi.org/10.1001/jamainternmed.2015.2609>.
44. Burgio KL, Richter HE, Clements RH, Redden DT, Goode PS. Changes in urinary and fecal Incontinence symptoms with weight loss surgery in morbidly obese women. *Obstet Gynecol.* 2007;110(5):1034-1040. <https://doi.org/10.1097/01.AOG.0000285483.22898.9c>.
45. O'Boyle CJ, O'Sullivan OE, Shabana H, Boyce M, O'Reilly BA. The effect of bariatric surgery on urinary Incontinence in women. *Obes Surg.* 2016;26(7):1471-1478. <https://doi.org/10.1007/s11695-015-1969-z>.
46. Ait Said K, Leroux Y, Menahem B, Doerfler A, Alves A, Tillou X. Effect of bariatric surgery on urinary and fecal Incontinence: prospective analysis with 1-year follow-up. *Surg Obes Relat Dis.* 2017;13(2):305-312. <https://doi.org/10.1016/j.soard.2016.08.019>.
47. Anglim B, O'Boyle CJ, O'Sullivan OE, O'Reilly BA. The long-term effects of bariatric surgery on female urinary Incontinence. *Eur J Obstet Gynecol Reprod Biol.* 2018;231:15-18. <https://doi.org/10.1016/j.ejogrb.2018.10.011>.
48. Leshem A, Groutz A, Amir H, Gordon D, Shimonov M. Surgically induced weight loss results in a rapid and consistent improvement of female pelvic floor symptoms. *Scand J Urol.* 2018;52(3):219-224. <https://doi.org/10.1080/21681805.2018.1447600>.
49. Phelan S, Kanaya AM, Subak LL, et al. Weight loss prevents urinary Incontinence in women with type 2 diabetes: results from the look AHEAD trial. *J Urol.* 2012;187(3):939-944. <https://doi.org/10.1016/j.juro.2011.10.139>.
50. Brown JS, Wing R, Barrett-Connor E, et al. Lifestyle Intervention is associated with lower prevalence of urinary Incontinence: the diabetes prevention program. *Diabetes Care.* 2006;29(2):385-390. <https://doi.org/10.2337/diacare.29.02.06 dc05-1781>.
51. Auwad W, Steggles P, Bombieri L, Waterfield M, Wilkin T, Freeman R. Moderate weight loss in obese women with urinary Incontinence: a prospective longitudinal study. *Int Urogynecol J Pelvic Floor Dysfunct.* 2008;19(9 PG-1251-9):1251-1259. <https://doi.org/10.1007/s00192-008-0616-9>.
52. Fjerbæk A, Søndergaard L, Andreasen J, Glavind K. Treatment of urinary Incontinence in overweight women by a multidisciplinary Lifestyle Intervention. *Arch Gynecol Obstet.* 2020;301(2):525-532. <https://doi.org/10.1007/s00404-019-05371-x>.
53. Subak LL, Johnson C, Johnson C, et al. Does weight loss improve Incontinence in moderately obese women? *Int Urogynecology J.* 2002;13(1):40-43. <https://doi.org/10.1007/s001920200008>.
54. Gozukara YM, Akalan G, Tok EC, Aytan H, Ertunc D. The improvement in pelvic floor symptoms with weight loss in obese women Does not correlate with the changes in pelvic anatomy. *Int Urogynecology J.* 2014;25(9):1219-1225. <https://doi.org/10.1007/s00192-014-2368-z>.
55. Zache MM, Giarenis I, Thiagamoorthy G, Robinson D, Cardozo L. Is there an association between aspects of the metabolic syndrome and overactive bladder? A prospective cohort study in women with lower urinary tract symptoms. *Eur J Obstet Gynecol Reprod Biol.* 2017;217:1-5. <https://doi.org/10.1016/j.ejogrb.2017.08.002>.
56. Aoki Y, Brown HW, Brubaker L, Cornu JN, Daly JO, Cartwright R. Urinary Incontinence in women. *Nat Rev Primer.* 2017;3:17042. <https://doi.org/10.1038/nrdp.2017.42.Urinary>.

57. Noblett KL, Jensen JK, Ostergard DR. The relationship of body mass index to intra-abdominal pressure as measured by multichannel cystometry. *Int Urogynecol J Pelvic Floor Dysfunct.* 1997;8(6):323-326. <https://doi.org/10.1007/BF02765589>.
58. Sugerman H, Windsor A, Bessos M, Kellum J, Reines H, DeMaria E. Effects of surgically induced weight loss on urinary bladder pressure, sagittal abdominal diameter and obesity co-morbidity. *Int J Obes.* 1998;22(3):230-235. <https://doi.org/10.1038/sj.ijo.0800574>.
59. Faiena I, Patel N, Parihar JS, Calabrese M, Tunuguntla H, Robert R. Conservative Management of Urinary Incontinence in Women. *Rev Urol.* 2015;17(3):129-139. <https://doi.org/10.3909/riu0651>.
60. Dallosso HM, McGrother CW, Matthews RJ, Donaldson MMK. The Association of Diet and Other Lifestyle Factors with overactive bladder and stress Incontinence: a longitudinal study in women. *BJU Int.* 2003;92(1):69-77. <https://doi.org/10.1046/j.1464-410X.2003.04271.x>.
61. Creighton SM, Stanton SLC. Does it affect your bladder? *Br J Urol.* 1990;66(6):613-614. <https://doi.org/10.1111/j.1464-410X.1990.tb07192.x>.
62. Dagan SS, Goldenshluger A, Globus I, et al. Nutritional recommendations for adult bariatric surgery patients: clinical practice. *Adv Nutr.* 2017;8(2):382-394. <https://doi.org/10.3945/an.116.014258>.
63. Adams TD, Gress RE, Smith SC, et al. After gastric bypass surgery. *Surv Anesthesiol.* 2007;57(8):753-761. <https://doi.org/10.1097/01.san.0000307862.68683.00>.
64. Keating CL, Bulfone L, Dixon JB, et al. Cost-effectiveness of surgically induced weight loss for the management of Type 2 diabetes: modeled lifetime analysis. *Diabetes Care.* 2009;32(4):567-574. <https://doi.org/10.2337/dc08-1749>.
65. Yap TL, Cromwell DA, Brown C, van der Meulen J, Emberton M. The relationship between objective frequency-volume chart data and the I-PSS in men with lower urinary tract symptoms. *Eur Urol.* 2007;52(3):811-818. <https://doi.org/10.1016/j.euro.2007.01.013>.

**How to cite this article:** Sheridan W, Da Silva AS, Leca BM, et al. Weight loss with bariatric surgery or behaviour modification and the impact on female obesity-related urine incontinence: A comprehensive systematic review and meta-analysis. *Clin Obes.* 2021;e12450. <https://doi.org/10.1111/cob.12450>