

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/151071>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Nonexistence of spectral gaps in Hölder spaces for continuous time dynamical systems

Ian Melbourne* Nicolò Paviato[†] Dalia Terhesiu[‡]

27 February 2021

Abstract

We show that there is a natural restriction on the smoothness of spaces where the transfer operator for a continuous dynamical system has a spectral gap. Such a space cannot be embedded in a Hölder space with Hölder exponent greater than $\frac{1}{2}$ unless it consists entirely of coboundaries.

1 Introduction

Decay of correlations (rates of mixing) and strong statistical properties are well-understood for Axiom A diffeomorphisms since the work of [2, 9, 10]. Mixing rates are computed with respect to any equilibrium measure with Hölder potential. Up to a finite cycle, such diffeomorphisms have exponential decay of correlations for Hölder observables. In the one-sided (uniformly expanding) setting, this is typically proved by establishing quasicompactness and a spectral gap for the associated transfer operator L . Such a spectral gap yields a decay rate $\|L^n v - \int v\| \leq C_v e^{-an}$ for v Hölder, where $\|\cdot\|$ is a suitable Hölder norm and a, C_v are positive constants. Decay of correlations for Hölder observables is an immediate consequence of the decay for L^n . This philosophy has been extended to large classes of nonuniformly expanding dynamical systems with exponential [13] and subexponential decay of correlations [14].

For continuous time dynamical systems, the usual techniques [5, 7, 8] bypass spectral gaps; the only exceptions that we know of being Tsujii [11, 12]. However, the result in [11] is for suspension semiflows over the doubling map with a C^3 roof function, where the smoothness of the roof function is crucial and very restrictive. A spectral gap for contact Anosov flows is obtained in [12]; unfortunately it seems nontrivial to extend this to nonuniformly hyperbolic contact flows (or uniformly hyperbolic

*Mathematics Institute, University of Warwick, Coventry, CV4 7AL, UK

[†]Mathematics Institute, University of Warwick, Coventry, CV4 7AL, UK

[‡]Mathematisch Instituut, University of Leiden, Niels Bohrweg 1, 2333 CA Leiden, Netherlands

contact flows with unbounded distortion), see [1] which proves exponential decay of correlations for billiard flows with a contact structure but does not establish a spectral gap. Indeed, apart from [11, 12], there are no results on spectral gaps of transfer operators for semiflows and flows.

The results of Tsujii [11, 12] provide a spectral gap in an anisotropic Banach space. In this paper we obtain a restriction on the Banach spaces that can yield a spectral gap. We work in the following general setting:

Let (Λ, d) be a bounded metric space with Borel probability measure μ , and let $F_t : \Lambda \rightarrow \Lambda$ be a measure-preserving semiflow. We suppose that $t \rightarrow F_t$ is Lipschitz a.e. on Λ . Let $L_t : L^1(\Lambda) \rightarrow L^1(\Lambda)$ denote the transfer operator corresponding to F_t (so $\int_{\Lambda} L_t v w d\mu = \int_{\Lambda} v w \circ F_t d\mu$ for all $v \in L^1(\Lambda)$, $w \in L^\infty(\Lambda)$, $t > 0$). Let $v \in L^\infty(\Lambda)$ and define $v_t = \int_0^t v \circ F_r dr$ for $t \geq 0$.

Theorem 1.1 *Let $\eta \in (\frac{1}{2}, 1)$. Suppose that $L_t v \in C^\eta(\Lambda)$ for all $t > 0$ and that $\int_0^\infty \|L_t v\|_\eta dt < \infty$. Then v_t is a coboundary:*

$$v_t = \chi \circ F_t - \chi \quad \text{for all } t \geq 0, \text{ a.e. on } \Lambda$$

where $\chi = \int_0^\infty L_t v dt \in C^\eta(\Lambda)$. In particular, $\sup_{t \geq 0} |v_t|_\infty < \infty$.

Here, $|g|_\infty = \text{ess sup}_\Lambda |g|$ and $\|g\|_\eta = |g|_\infty + \sup_{x \neq y} |g(x) - g(y)|/d(x, y)^\eta$.

Theorem 1.1 implies that any Banach space admitting a spectral gap and embedded in $C^\eta(\Lambda)$ for some $\eta > \frac{1}{2}$ is cohomologically trivial. However, for (non)uniformly expanding semiflows and (non)uniformly hyperbolic flows of the type in the aforementioned references, coboundaries are known to be exceedingly rare, see for example [3, Section 2.3.3]. Hence, Theorem 1.1 can be viewed as an “anti-spectral gap” result for such continuous time dynamical systems.

2 Proof of Theorem 1.1

Let $v \in L^\infty(\Lambda)$, with $L_t v \in C^\eta(\Lambda)$ for all $t > 0$ and $\int_0^\infty \|L_t v\|_\eta dt < \infty$ where $\eta \in (\frac{1}{2}, 1)$. Following Gordin [6] we consider a martingale-coboundary decomposition. Define $\chi = \int_0^\infty L_t v dt \in C^\eta(\Lambda)$, and

$$v_t = \int_0^t v \circ F_r dr, \quad m_t = v_t - \chi \circ F_t + \chi,$$

for $t \geq 0$. Let \mathcal{B} denote the Borel σ -algebra on Λ .

Proposition 2.1 *(i) $t \rightarrow m_t$ is C^η a.e. on Λ .*

(ii) $\mathbb{E}(m_t | F_t^{-1} \mathcal{B}) = 0$ for all $t \geq 0$.

Proof (i) For $0 \leq s \leq t \leq 1$ and $x \in \Lambda$,

$$\begin{aligned} |m_s(x) - m_t(x)| &\leq |v_s(x) - v_t(x)| + |\chi(F_s x) - \chi(F_t x)| \\ &\leq |s - t| |v|_\infty + |\chi|_\eta d(F_s x, F_t x)^\eta. \end{aligned}$$

Since $t \mapsto F_t$ is a.e. Lipschitz, it follows that $t \mapsto m_t$ is a.e. C^η .

(ii) Let $U_t v = v \circ F_t$, and recall that $L_t U_t = I$ and $\mathbb{E}(\cdot | F_t^{-1} \mathcal{B}) = U_t L_t$. Then

$$\begin{aligned} L_t m_t &= L_t(v_t - U_t \chi + \chi) = \int_0^t L_t U_r v dr - \chi + \int_0^\infty L_t L_r v dr \\ &= \int_0^t L_{t-r} v dr - \chi + \int_0^\infty L_{t+r} v dr = \int_0^t L_r v dr - \chi + \int_t^\infty L_r v dr = 0. \end{aligned}$$

Hence $\mathbb{E}(m_t | F_t^{-1} \mathcal{B}) = U_t L_t m_t = 0$. ■

Proof of Theorem 1.1 Fix $T > 0$, and define

$$M_T(t) = m_T - m_{T-t} = m_t \circ F_{T-t}, \quad t \in [0, T].$$

Also, define the filtration $\mathcal{G}_{T,t} = F_{T-t}^{-1} \mathcal{B}$. It is immediate that $M_T(t) = m_t \circ F_{T-t}$ is $\mathcal{G}_{T,t}$ -measurable. Also, for $s < t$ we have $M_T(t) - M_T(s) = m_{T-s} - m_{T-t} = m_{t-s} \circ F_{T-t}$, so

$$\mathbb{E}(M_T(t) - M_T(s) | \mathcal{G}_{T,s}) = \mathbb{E}(m_{t-s} \circ F_{T-t} | F_{T-s}^{-1} \mathcal{B}) = \mathbb{E}(m_{t-s} | F_{t-s}^{-1} \mathcal{B}) \circ F_{T-t} = 0$$

by Proposition 2.1(ii). Hence M_T is a martingale for each $T > 0$. Next,

$$|M_T(t)|_\infty = |m_t \circ F_{T-t}|_\infty \leq |m_t|_\infty \leq |v_t|_\infty + 2|\chi|_\infty \leq T|v|_\infty + 2|\chi|_\infty.$$

Hence $M_T(t)$, $t \in [0, T]$, is a bounded martingale.

By Proposition 2.1(i), M_T has C^η sample paths. Since $\eta > \frac{1}{2}$, it follows from general martingale theory that $M_T \equiv 0$ a.e. Taking $t = T$, we obtain $m_T = 0$ a.e. Hence $v_T = \chi \circ F_T - \chi$ a.e. for all $T > 0$ as required.

For completeness, we include the argument that $M_T \equiv 0$ a.e. We require two standard properties of the quadratic variation process $t \mapsto [M_T](t)$; a reference for these is [4, Theorem 4.1]. First, $[M_T](t)$ is the limit in probability as $n \rightarrow \infty$ of

$$S_n(t) = \sum_{j=1}^n (M_T(jt/n) - M_T((j-1)t/n))^2.$$

Second (noting that $M_T(0) = 0$),

$$[M_T](t) = M_T(t)^2 - 2 \int_0^t M_T dM_T,$$

where the stochastic integral has expectation zero. In particular, $\mathbb{E}([M_T]) \equiv \mathbb{E}(M_T^2)$.

Since M_T has Hölder sample paths with exponent $\eta > \frac{1}{2}$, we have a.e. that

$$|S_n(t)| = O(t^\eta n^{-(2\eta-1)}) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Hence $[M_T] \equiv 0$ a.e. It follows that $\mathbb{E}(M_T^2) \equiv 0$ and so $M_T \equiv 0$ a.e. ■

References

- [1] V. Baladi, M. F. Demers and C. Liverani. Exponential decay of correlations for finite horizon Sinai billiard flows. *Invent. Math.* **211** (2018) 39–177.
- [2] R. Bowen. *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*. Lecture Notes in Math. **470**, Springer, Berlin, 1975.
- [3] I. Chevyrev, P. K. Friz, A. Korepanov, I. Melbourne and H. Zhang. Multiscale systems, homogenization, and rough paths. *Probability and Analysis in Interacting Physical Systems: In Honor of S.R.S. Varadhan, Berlin, August, 2016*” (P. Friz et al., ed.), Springer Proceedings in Mathematics & Statistics **283**, Springer, 2019, pp. 17–48.
- [4] K. L. Chung and R. J. Williams. *Introduction to stochastic integration*, second ed., Probability and its Applications, Birkhäuser Boston, Inc., Boston, MA, 1990.
- [5] D. Dolgopyat. On the decay of correlations in Anosov flows. *Ann. of Math.* **147** (1998) 357–390.
- [6] M. I. Gordin. The central limit theorem for stationary processes. *Soviet Math. Dokl.* **10** (1969) 1174–1176.
- [7] C. Liverani. On contact Anosov flows. *Ann. of Math.* **159** (2004) 1275–1312.
- [8] M. Pollicott. On the rate of mixing of Axiom A flows. *Invent. Math.* **81** (1985) 413–426.
- [9] D. Ruelle. *Thermodynamic Formalism*. Encyclopedia of Math. and its Applications **5**, Addison Wesley, Massachusetts, 1978.
- [10] Y. G. Sinai. Gibbs measures in ergodic theory. *Russ. Math. Surv.* **27** (1972) 21–70.
- [11] M. Tsujii. Decay of correlations in suspension semi-flows of angle-multiplying maps. *Ergodic Theory Dynam. Systems* **28** (2008) 291–317.
- [12] M. Tsujii. Quasi-compactness of transfer operators for contact Anosov flows. *Nonlinearity* **23** (2010) 1495–1545.
- [13] L.-S. Young. Statistical properties of dynamical systems with some hyperbolicity. *Ann. of Math.* **147** (1998) 585–650.
- [14] L.-S. Young. Recurrence times and rates of mixing. *Israel J. Math.* **110** (1999) 153–188.