

Examining vulnerability and resilience in maternal, newborn and child health through a gender lens in low-income and middle-income countries: a scoping review

Fatima Abdulaziz Sule ¹, Olalekan A Uthman ²,
Emmanuel Olawale Olamijuwon ¹, Nchelem Kokomma Ichegbo ¹,
Ifeyanyi C Mgbachi,¹ Babasola Okusanya ³, Olusesan Ayodeji Makinde ^{1,4}

To cite: Sule FA, Uthman OA, Olamijuwon EO, *et al*. Examining vulnerability and resilience in maternal, newborn and child health through a gender lens in low-income and middle-income countries: a scoping review. *BMJ Global Health* 2022;**7**:e007426. doi:10.1136/bmjgh-2021-007426

Handling editor Stephanie M Topp

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjgh-2021-007426>).

Received 14 September 2021
Accepted 20 March 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Olusesan Ayodeji Makinde; sesmak@gmail.com

ABSTRACT

Introduction Gender lens application is pertinent in addressing inequities that underlie morbidity and mortality in vulnerable populations, including mothers and children. While gender inequities may result in greater vulnerabilities for mothers and children, synthesising evidence on the constraints and opportunities is a step in accelerating reduction in poor outcomes and building resilience in individuals and across communities and health systems.

Methods We conducted a scoping review that examined vulnerability and resilience in maternal, newborn and child health (MNCH) through a gender lens to characterise gender roles, relationships and differences in maternal and child health. We conducted a comprehensive search of peer-reviewed and grey literature in popular scholarly databases, including PubMed, ScienceDirect, EBSCOhost and Google Scholar. We identified and analysed 17 published studies that met the inclusion criteria for key gendered themes in maternal and child health vulnerability and resilience in low-income and middle-income countries.

Results Six key gendered dimensions of vulnerability and resilience emerged from our analysis: (1) restricted maternal access to financial and economic resources; (2) limited economic contribution of women as a result of motherhood; (3) social norms, ideologies, beliefs and perceptions inhibiting women's access to maternal healthcare services; (4) restricted maternal agency and contribution to reproductive decisions; (5) power dynamics and experience of intimate partner violence contributing to adverse health for women, children and their families; (6) partner emotional or affective support being crucial for maternal health and well-being prenatal and postnatal.

Conclusion This review highlights six domains that merit attention in addressing maternal and child health vulnerabilities. Recognising and understanding the gendered dynamics of vulnerability and resilience can help develop meaningful strategies that will guide the design and implementation of MNCH programmes in low-income and middle-income countries.

WHAT IS ALREADY KNOWN ON THIS TOPIC?

- ⇒ Socioeconomic inequalities place women and girls in precarious positions that adversely affect their vulnerability and resilience to health shocks.
- ⇒ Research on the gendered dimension of maternal and child health vulnerability and resilience is needed to fully evaluate how gender expectations may result in greater vulnerability for mothers, newborns and children or impact their resilience.

WHAT THIS STUDY ADDS

- ⇒ This study provides new evidence on the gender dynamics of vulnerability and resilience in maternal, newborn and child health (MNCH) and how this impacts health outcomes.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ This research informs programme managers and researchers of the importance of embedding gender perspectives in future design of interventions aimed at addressing MNCH outcomes.
- ⇒ It also calls for further research aimed at generating robust evidence on gender-transformative interventions that best address MNCH vulnerability and resilience across different contexts in low-income and middle-income countries.

INTRODUCTION

Maternal and childhood mortality remains key health challenges in several low-income and middle-income countries (LMICs). In 2019, approximately 5.2 million children died before their fifth birthday, more than 80% of these deaths occurred in sub-Saharan Africa and Central and South Asia.¹ Maternal mortality in sub-Saharan Africa and South Asia bore 86% of the estimated global burden in 2017.² Sub-Saharan Africa's maternal

mortality ratio of 546 per 100 000 live births is estimated to be the highest globally for any region.³

In Maternal, Newborn and Child Health (MNCH), a vulnerable pregnant woman was defined as a woman who is threatened by physical, psychological, cognitive and/or social risk factors in combination with lack of adequate support and/or adequate coping skills.⁴ On the other hand, resilience has been described as the capability of the public health and healthcare systems, communities, and individuals to prevent, protect against, quickly respond to and recover from health emergencies, particularly those whose scale, timing or unpredictability threatens to overwhelm routine capabilities.⁵ Thus in MNCH, vulnerability and resilience are two divergent terms that tend to complement each other by acting as risk or protective factors, respectively, both at the individual level and at the health system. Pregnancy-related morbidity and mortality in LMICs are often preventable or treatable, but poverty, low maternal educational attainment and place of residence, among several other underlying factors, increase women's vulnerability to adverse maternal and child health outcomes.⁶⁻¹¹

Although multiple studies have examined these vulnerabilities, more attention needs to be paid to how they are patterned by gender to influence MNCH outcomes. Similarly, maternal resilience evidenced in women's ability to sustain life satisfaction, self-esteem and purpose amidst emotional, physical and financial difficulties associated with mothering and caregiving has been studied extensively.^{6 8 10-12} However, there has been limited focus on how gender roles and norms may shape these factors.¹²

Institutionalised power, social, political and economic advantages and disadvantages afforded to different genders influence power relations. Gender also intersects with other social determinants of health, including social class, race and ethnicity,¹³ determines the hierarchy of social structure and power dynamics, and influences health outcomes. Health inequalities conditioned by gender are likely to put vulnerable populations at a further disadvantage.¹⁴ Today, there is an increasing need for a critical and systematic assessment of the effect of gender norms, and gender inequality on the constraints faced by and opportunities available to vulnerable populations regarding MNCH. Theoretical and conceptual advances in global health have highlighted the importance of gender expectations, roles and relations in health promotion interventions.¹⁵⁻¹⁷ For example, different gender expectations may result in greater vulnerability to mothers and children. Promising gender-sensitive practices in health have also emerged to address the HIV/AIDS epidemic and influence maternal and child health outcomes.¹⁸⁻²¹

The Sustainable Development Goals (SDG) aims to reduce maternal deaths to less than 70 per 100 000 live births by 2030 (SDG 3.1), neonatal mortality rate to at least as low as 12 per 1000 live births and under-5 mortality rate to at least as low as 25 per 1000 live births (SDG 3.2). These maternal and child health targets may

be impossible to achieve if the critical factors shaping maternal and child health vulnerability and resilience are not well articulated. The SDG agenda must operate with gender as a cross-cutting aspect and therefore integrated within design, resource allocation, implementation, measurement and evaluation. Specifically, understanding how health systems respond to critical factors that shape the health and well-being of mothers, children and newborn is necessary.²²

This scoping review illuminates how gender differences and relations are relevant in providing important insight into how power structures and roles aggravate vulnerability or strengthen resilience in maternal and child health in LMICs. It provides new evidence on gendered dynamics in MNCH research that must be considered as we strive to programme interventions aimed at achieving the SDG targets on maternal and child health.

METHODS

We conducted a scoping review in accordance with Arksey and O'Malley's framework to examine the gendered dimension of vulnerability and resilience in MNCH in LMICs.^{23 24} A scoping review was necessary for a broad and comprehensive analysis without consideration of publication quality. The review followed five stages: (1) identifying the research question; (2) identifying the relevant studies; (3) selecting the studies; (4) charting data and (5) collating, summarising and reporting results.

Identification of relevant peer-reviewed literature

This gender analysis was based on a larger scoping review aimed at developing a framework for vulnerability and resilience in MNCH in LMICs. The initial pool of literature was retrieved from major databases (ie, Medline, Embase, Scopus and Web of Science) based on a comprehensive and exhaustive search strategy that included appropriate keywords (see online supplemental appendix S1). This was supplemented by a grey literature search. The initial search was conducted on 15 January 2021 and updated on 1 March 2021.

The search strategy was structured around three blocks: (1) population (ie, MNCH, health outcomes, healthcare utilisation and social capital), (2) exposure (ie, vulnerability, resilience and high-risk) and (3) setting (ie, low-income and middle-income settings). Critical keywords and thesaurus heading terms were initially tailored to Medline and Embase searches and then adapted in other sources as necessary. Online supplemental appendix S1 shows the full search strategies for Medline and Embase.

We also reviewed reports and technical papers from multilateral and bilateral organisations, foundations, international and local non-governmental organisations, such as the Bill & Melinda Gates Foundation, Jhpiego, Clinton Health Access Initiative, International Centre for Research on Women, Women's Health and Action Research Centre, Gender Watch and pharmacies. To gather as much evidence as possible, including

high-quality literature regarding vulnerable populations in MNCH beyond the traditional sources, we incorporated the research from grey literature into this scoping review. We supplemented the database search with a bibliography search of key articles but found no relevant articles beyond what had already been extracted. We did not apply language restrictions in our search parameters and, thus, engaged translators to translate non-English publications.

Study selection

We developed and validated a high-performance machine learning classifier/algorithm (bidirectional encoder representations from transformers) to identify relevant studies focusing on vulnerability and resilience in MNCH from an initial pool of search results. Previous studies have reported the high predictive ability of machine learning models in title and abstract screening.^{25–27} To train the machine learning algorithm, we randomly selected, screened and annotated the titles and abstracts of 500 records from the database. The performance of the model was evaluated against our classification based on precision, recall, specificity and accuracy scores. Subsequently, we applied the algorithm to review the abstracts and titles of the remaining publications to generate predictions to include or exclude them.

Covidence, an online systematic review software, was used to manage the search outputs and screening of eligible studies (<https://www.covidence.org/>). Two researchers screened the identified manuscripts retained from machine-learning predictions using Covidence. A third researcher reviewed and resolved all conflicts. Titles and abstracts were first screened before a full-text review for possible inclusion in the study. We included studies based on four key criteria. First, if they focus on women (pregnant/lactating and teenage mothers) and/or children (male and female) under 5 years. Second, if they focused on LMICs. We also included studies that focused on vulnerability, frailty or high risk and resilience in LMICs. Lastly, we included all study types including peer-reviewed publications, programmatic reports, and conference abstracts. There were no language restrictions nor exclusions based on the year of publication.

Charting data

To provide a holistic gender analysis, we adapted a conceptual framework for gender analysis in health systems research by Morgan *et al.*²⁸ The framework unifies several other frameworks focusing on health, health systems and development.²⁸ More importantly, the framework's unique focus on how power is constituted and negotiated makes it a valuable resource for understanding gender in terms of power relation and a source of disparity in health systems. The framework had five focal areas, namely, access to resources, division of labour, social norms, rules and decision-making, power negotiation, and structure/environment. All the articles

that met the inclusion criteria for this study were further screened based on these five key gender dimensions.

Relevant data were extracted into a data collection template developed on AirTable. Articles were screened and extracted if they fit any of the five dimensions of gender and power identified in the framework. We extracted the publication metadata (ie, name of the first author, year of publication, publication title and publication country) and additional data (eg, publication type, research design and methods, study context, indices of vulnerability and resilience, and key findings from the research). Categories for the focal areas were not mutually exclusive, which means that a study could belong and be counted in more than one category where evidence of such contributions exists. During the data analysis, we grouped the articles by their specific focus on the different dimensions of gender and power relations. **Table 1** presents the details of the classifications.

Collating, synthesising and reporting the results

This review describes, first, the characteristics of the studies that meet the study inclusion criteria and, second, the findings. We report the summary statistics describing data collection methods, vulnerability/resilience context (eg, maternal or child/newborn health), gender dimension (eg, access to resources, division of labour, social norms, rules and decision making, power negotiation and structure/environment). We did not assess the quality or risk of bias for the included articles as the objective of this review was to scope and describe the breadth of gender dimensions in vulnerability or resilience in MNCH in LMICs. This review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews (PRISMA-ScR) statement guidelines to enhance transparency in reporting scoping reviews.²⁹

Patient and public involvement statement

Patients were not involved in the conduct of this study.

RESULTS

We identified 76 656 records through the database search (**figure 1**). We excluded 57 duplicate records and 73 638 abstracts that were flagged as potentially irrelevant to this study. Thereafter, we screened the remaining titles and abstracts (n=2871), we considered only 96 studies as relevant and selected them for full-text review. Of these, 79 studies did not meet the inclusion criteria and were excluded because of incorrect population, outcomes, setting, and study design or the lack of a gender focus in the analysis. Subsequently, 17 studies met our inclusion criteria for a promising gender-sensitive analysis of vulnerability and resilience in MNCH in LMICs. Online supplemental material S2 provides the details of these studies, including the year of publication, country of publication, context of the study, study design and key findings related to gender as regards vulnerability and resilience in MNCH.

Table 1 Illustrative examples of gender analysis

What constitutes gender power relations		Illustrative gender analysis research question
Access to resources (Who has what?)	Access to resources (education, information, skills, income, employment, services, benefits, time, space, social capital, etc.)	To what extent do women and men have the same access to education, information, income, employment and other resources that contribute to improvement in maternal, newborn, and child health? Do women have sufficient means to make decisions and access healthcare services without financial restrictions?
Division of labour (Who does what?)	Division of labour within and beyond the household and everyday practices	How do women's social roles, such as childbearing, childcare, and infant feeding, affect their economic opportunities and access to health facilities?
Social norms (How are values defined?)	Social norms, ideologies, beliefs and perceptions	How does stigma inhibit women's access to maternal healthcare services and are these available to unmarried women and teenage mothers? How do cultural norms about motherhood put women at risk of adverse health?
Agency and decision making (Who decides?)	Agency and decision making (both formal and informal)	To what extent are women able to advocate for their health needs and contribute to household decisions that shape their and their children's health?
Power negotiation (How is power enacted, negotiated or challenged?)	Critical consciousness, acknowledgement/lack of acknowledgement, agency/apathy, interests, historical and lived experiences, resistance or violence	How is power enacted and negotiated in relation to maternal, newborn, and child health and how does power dynamics or women's experience of intimate partners contribute to adverse health for women, children and their families?

Study characteristics

A total of 17 studies met the inclusion criteria for a gender analysis. Out of these, 13 focused on maternal health and four on child health (figure 2). Eleven studies focused on sub-Saharan African countries (figure 2), of which three were from Kenya. Resilience was a more dominant focus: eight on maternal health and two on child health (online supplemental material S2).

Figure 3 presents the distribution of gender themes across maternal and child health contexts. Access to resources and decision making was the most common focus of the identified studies on both maternal (five) and child (three) health. Three studies examined power negotiation in relation to maternal (two) and child (one) health. Two studies also highlighted partner emotional or mental support in maternal health (two) and another two on the decision-making ability of mothers (two). Only a few studies examined how social norms and division of labour intersect with maternal health.

Access to resources

Access to resources emerged as the dominant gender-focused theme (8 of 17 studies).^{30–37} In most studies, pregnant women or mothers lived in households characterised by low socioeconomic status and had lower levels of education, all of which are potentially related to poor access to maternal and child healthcare services. Among pregnant women living in a community of metropolitan Santiago, Chile,³¹ low socioeconomic status was found to be related to deteriorating reproductive, maternal and neonatal health. Warren *et al* supported this finding and found that most women affected by fistula had secondary education as the highest level of education and a very low

monthly income.³⁶ Most primary caretakers, including mothers, were not income earners and often relied heavily on their spouses or other household members for money.

Access to resources also emerged as an important barrier to child healthcare. For example, Johnson *et al* demonstrated that the classification as orphan and vulnerable children (OVC) directly and indirectly influenced the risk of childhood morbidity (eg, diarrhoea, fever and acute respiratory infection).³⁰ This is because OVCs were more likely to be found in households headed by adults (40 years old), where the mother/caregiver had inadequate access to socioeconomic resources, such as inadequate education, and in urban areas.

Many women were often in precarious positions, relying on their spouses for financial support to access healthcare services even during emergencies. A study in Kenya reported that irrespective of marital status, having male support (eg, husband, brother or uncle), particularly financial support and help in securing transport to hospitals for care, was critical.³⁶ Some women failed to attend clinics because of a lack of support from their husbands. Most husbands did not provide their wives with adequate funds for their needs during delivery.³² The lack of rapid access to money was another important contributing factor to a child's deteriorating condition; it influenced the initiation of a treatment-seeking action, including where and by whom (all households) the action was performed. For instance, women in a study made many references to 'waiting to talk to my husband', 'waiting to be sent money from my husband,' and waiting for 'his permission to pursue an action.'³⁷ Such




Figure 1 Summary of the search, selection and inclusion process.* Women (including pregnant/lactating and teenage mothers) AND/OR Children (male and female), 0-4 years OR under 5 years. (Include breastfeeding and childhood immunization studies). **Include peer-reviewed publications, programmatic reports, conference abstracts.

gender-reinforced inequality in access to resources could subsequently affect the health-care-seeking behaviour of mothers and, ultimately, affect childcare, especially in the context of costly maternal healthcare services.

Division of labour

Only one study examined the dimension of the division of labour and how it intersects with maternal and child health.³⁸ This study included 36 Ugandan women who were admitted with obstetric near-miss and revealed that women's need to balance economic activities and reproduction often increased their vulnerability and ability to recover from obstetric complications. In such circumstances, social networks or social capital was generally perceived as an essential component of women's resilience because it provides women with financial, material, and emotional assistance, including those related to household responsibilities, such as childcare.³⁸

Social norms

One study examined the dimensions of social norms in maternal health.³⁹ It explored how values related to motherhood are defined and how this definition shapes or inhibits women's access to maternal healthcare services

or places women at risk of adverse health. An in-depth case study of a woman from Burkina Faso suggested that structural impediments, including motherhood and childbearing, limit individual resilience.³⁹ This case study noted that the high level of social pressure on women to bear children as soon as possible, even when they are not physically or mentally capable, and the stigma associated with childlessness exacerbate maternal mortality and morbidity risks.³⁹ These conditions contributed to the death of women in the case study, who could not be rescued from dying from childbirth-related complications despite having access to skilled birth attendance and emergency obstetric care.³⁹

Agency and decision making

Two studies underscored the ability of women and mothers to make informed choices and contribute to decisions related to maternal and child healthcare.^{40 41} For example, Prates *et al* showed women's inability to adequately plan the timing of childbirth because of poor socioeconomic status and inequalities in gender power, all of which contribute to multiparity.⁴¹ More importantly, the existing power imbalance motivates male partner



Figure 2 Geographical distribution of the 17 studies included in the scoping review.

resistance to condom use as a means of family planning.⁴¹ Additionally, Den Hollander *et al* in Ghana underscored women's low negotiating ability and autonomy in healthcare decision making.⁴⁰ The study reported wide power differences between health providers and women, especially in a context shaped by authority. Women were generally uninformed about their basic health information. A high level of therapeutic misconceptions was also observed in this study. Women were also reported to rely more often on a medical professional's opinion rather than being guided by their motivation.⁴⁰

Power negotiation

Power negotiation also emerged as a dominant gender dimension of vulnerability and resilience in maternal and child health. This dimension refers to how power is enacted and negotiated in relation to MNCH and how power dynamics or women's experience of intimate partner violence contributes to adverse health for women, children and their families. Our analysis found

two studies that examined power negotiation in maternal health^{42 43} and one in child health.⁴⁴

Although seropositive status disclosure is a crucial aspect of HIV programming, women living with HIV were generally reluctant in disclosing their HIV status to their partner to avoid negative reactions from the latter, including intimate partner physical violence.⁴⁴ Men were often not in favour of having their wives tested, fearing the indirect disclosure of their own infection.⁴⁴ Nonetheless, partner involvement is crucial for prevention of mother-to-child transmission (PMTCT), especially because this might require mothers to use antiretroviral therapy and formula feeding for infants. The authors recommended couple counselling and partner involvement in PMTCT programmes, as only testing women can increase their susceptibility to violence despite careful counselling.

Furthermore, women's exposure to intimate partner violence could also affect other aspects of their health. For example, Vivilaki *et al* observed that the lack of or disappointment with partner support, poor marital




Figure 3 Summary of the 17 studies by gender focus and maternal or child health context.

relationship and emotional/physical abuse had been associated with high levels of postpartum anxiety and depression.⁴³ Likewise, McNaughton Reyes *et al* found that women exposed to intimate partner violence may be likely to experience persistent poor mental health across the antenatal and postnatal periods.⁴²

Partner emotional or affective support

The two studies on partner emotional or affective support were primarily related to maternal health.^{45 46} Families and partners often reacted negatively by rejecting unwed pregnant teenagers or teenage mothers.⁴⁶ These rejections were expressed differently, including avoiding pregnant teenagers or verbal abuse.⁴⁶ The analysis suggested that low-resilient women with threatened premature labour reported higher pressures from child support concerns after delivery, less active coping, less positive affect and more negative affect.⁴⁵

DISCUSSION

This scoping review illuminates the gendered dynamics of vulnerability and resilience in MNCH research. Based on the 17 studies reviewed, we found that gender norms, roles and relationships significantly influence and reinforce vulnerability and resilience in maternal and child health. The role of gender-transformative interventions cannot be overemphasised in addressing these societal structures and widely held social values that perpetuate the gender inequities identified in this review. Our work highlights some promising gender-transformative interventions that should be prioritised in addressing vulnerabilities in MNCH (see [table 2](#) for the summary). These

are potential interventions based on the problems identified. Most importantly, women should have unhindered access to maternal and child healthcare services regardless of education, level of wealth, age or marriage.

As highlighted in this review, access to resources was a dominant theme in 8 of the 17 reviewed studies.^{30–37} Mothers in most of the studies reported having to wait for their husbands or other relatives for funds before they could access healthcare services. This process could pose a significant threat to them and their children's health and well-being, especially during emergencies. Women's access to healthcare services is compounded by socio-cultural stereotypes that impede maternal access to healthcare services, including marriage and adolescent motherhood. Multiple studies have highlighted how cultural stereotypes and stigma may hinder healthcare access for the same people who need the service the most.⁴⁷ In some cultural settings, unmarried women and adolescent mothers are unable to access care, partly because of the emphasis on marriage and motherhood in many African societies. Many women in search of assistance have fallen victims of human trafficking rings in baby factories where their babies are sold, and then they have been held against their will, thereby compounding their woes.^{48 49}

However, these barriers to healthcare access can be alleviated through a multisectoral intervention that addresses sociocultural stereotypes and the high costs of access to health services, including the cost of registration, treatment and care. For example, in Nigeria, the removal of user fees and increased community engagement for the most vulnerable is associated with a higher

Table 2 Implications and recommendations for programme and policy

Gender aspect	Recommendations
Access to resources (Who has what?)	<ul style="list-style-type: none"> ▶ Eliminate financial barriers in accessing maternal and child health services. ▶ Empower mothers through formal and informal education to enhance their health awareness and consciousness, efficacy and ability to make informed decisions about their health and their child/children's health.
Division of labour (Who does what?)	<ul style="list-style-type: none"> ▶ Provide adequate support and affordable childcare for mothers to enhance their productivity and participation in the labour force. ▶ Incentivise programmes that motivate the involvement of men in childcare and house chores.
Social norms (How are values defined?)	<ul style="list-style-type: none"> ▶ Address issues regarding cultural stereotypes that impede maternal access to healthcare services, including those related to marriage and adolescent motherhood. This could be in the form of providing a friendly and safe environment for adolescent and unmarried mothers to access healthcare. ▶ Engage community leaders in alleviating social norms that put women and girls at risk of poor health. This includes social norms that limit the contributions of women beyond motherhood.
Agency and decision making (Who decides?)	<ul style="list-style-type: none"> ▶ Provide universal access to safe and effective means of contraception, irrespective of the level of education and wealth. ▶ Strengthen the capacity of women and girls through education and job creation to contribute significantly to household decision making. ▶ Empower women to make decisive decisions about whether they want to have a/another baby and when they want to do so.
Power negotiation (How is power enacted, negotiated or challenged?)	<ul style="list-style-type: none"> ▶ Develop effective systems and strategies for the reporting and management of intimate partner violence and abuse.

level of maternal health-seeking behaviour.⁵⁰ Similar findings have been reported in other LMICs, including China, Zambia, Jamaica and India.^{51 52}

Although the abolition of user fee policies is necessary to achieve universal access to quality healthcare, multiple studies have underscored that such policies are not sufficient to improve maternal healthcare utilisation.^{53 54} The removal of user fees may increase uptake but may not reduce mortality proportionally if the quality of facility-based care is poor.⁵⁵ This may especially be salient in settings where healthcare access is limited by structural barriers related to the distance of health facilities or cost of transportation, waiting times and other additional costs.^{56 57} Masiye *et al* emphasised that the cost of transportation is mainly responsible for limiting the protective effect of user fee removal on catastrophic healthcare among the poorest households.⁵⁷ This finding is supported by Dahab and Sakellariou, who identified transportation barriers as among the most important barriers to maternal health in low-income African countries.⁵⁶ In fact, one study in our review reported that receiving financial support and helping in securing transport to hospitals for healthcare is critical.³⁶

Previous studies have also highlighted that poorly implemented user fee removal policies benefit more well-off women than poor ones, and in cases where there are significant immediate effects on the uptake of facility delivery, this trend is not sustained over time.^{58 59} Given these findings, there is an overarching need for comprehensive and multisectoral approaches to achieve sustainable improvements in maternal health. In some studies, women who received financial incentives as a part of

neonatal care or conditional cash transfers reported better healthcare-seeking behaviours than those who did not.⁶⁰ Morgan *et al* emphasised that financial incentives can increase the quantity and quality of maternal health services and address health systems and financial barriers that prevent women from accessing and providers from delivering quality and lifesaving maternal healthcare.⁶⁰ There is also an increasing consensus on the need to engage the community and religious leaders in challenging many of the cultural impediments to healthcare access. Countries in which these have been attempted have reported huge successes in improving healthcare access and service utilisation.

In several LMICs, women are tasked with the responsibility of childbearing and child-rearing; both could significantly affect women's economic productivity. Empowering women through skill acquisition could also offer a viable financial alternative and alleviate the high cost of accessing healthcare services, especially for women in low socioeconomic strata. Adequate incentives and support for mothers of children could also significantly ease the pressure on women to balance motherhood and economic activities. Some studies have reported the positive effects of programmes that help women with childcare.^{61 62} Such empowerment programmes could also be extended to single women and women in sole-based or female-headed households, because these family types are characterised by low levels of education and household wealth.

Another important gender dimension is the need for women and mothers to make decisions about their health and well-being. As highlighted in our review, women have

limited contribution to decision-making processes that are related to healthcare and family planning.^{40 41} This limitation is complicated by power imbalances between women and their spouses and between women and healthcare workers.^{40 41} One study found that women are only aware of condoms as a means of contraception and that their male partners resist to use condom. However, they are unwilling to use other means of contraception, perhaps because of the known or perceived side effects. Family planning services must be integrated into existing maternal and child health programmes, so that women are adequately equipped with sexual and reproductive health information and have the autonomy to choose their preferred means of contraception with minimal effects on pleasure.

Male partner involvement is also crucial for PMTCT of HIV, especially because this requires mothers to use antiretroviral therapy and feed the child using formula feeding.⁴⁴ Although the involvement of the spouse during childbirth and child-rearing could alleviate some of the economic implications of motherhood, unfortunately, many male partners are not usually involved in childcare.⁶³ A few studies in our review reported on women's experiences of intimate partner violence and how this intersects with maternal and child vulnerabilities.⁴²⁻⁴⁴ Women's exposure to intimate partner violence is associated with high levels of postpartum anxiety and depression and their experience of persistent poor mental health across the antenatal and postnatal periods.^{42 43} The fear of intimate partner violence has also been reported to influence women's disclosure of HIV status to their spouses.⁴⁴ This occurs especially because men are often not favouring having their wives tested, fearing the indirect disclosure of their own infection. As recommended by Gaillard *et al*⁴⁴ and other scholars,⁶⁴ the continued counselling of women alone may not eliminate some of the maternal risks of intimate partner violence. However, MNCH programmes could alleviate these risks through couple-counselling and partner involvement in PMTCT programmes.

Aside from increasing male partner involvement in reducing maternal risks of intimate partner violence, the development of effective systems and strategies for the reporting and management of intimate partner violence and abuse is important. Many LMICs have legal structures for seeking redress for intimate partner violence; however, reporting the same has not been effective. Multiple studies have examined women's motivation to remain in violent unions.⁶⁵⁻⁶⁸ The findings of these studies, among several others, have highlighted the subsistence and stereotypes associated with being divorced, among others. As a result, strong systems may especially be important for women in low socioeconomic status who must remain in violent marriages for survival. Altogether, these findings have pointed to the need for context and a women-centric perspective in developing strategies to eliminate violence against women, as such strategies may be inefficient if they do not address some

of the bottlenecks for combating violence against women. Some studies have reported the effectiveness of women's social empowerment combined with economic empowerment in reducing women's vulnerabilities to intimate partner violence.⁶⁹ Such interventions may also provide women with resources to access healthcare services and alleviate maternal experiences of intimate partner violence. However, these interventions could aggravate experiences of intimate partner violence, especially in settings where maternal empowerment is perceived to threaten established gender norms.⁷⁰⁻⁷² Nonetheless, multiple studies in Tanzania have reported that maternal empowerment has led to considerable reductions in physical intimate partner violence and posed no additional adverse health risks.⁶⁹

Watts and Mayhew⁷³ and García-Moreno *et al*⁷⁴ recommended a more active approach, that is, to integrate health systems response into maternal and child healthcare. Today, there is a global consensus to strengthen healthcare professionals' ability to identify victims of intimate partner violence and provide first-line supportive care and referral to other care services.⁷⁴ A functional and well-financed health system is also important to prevent violence against women and respond to victims and survivors in a consistent, safe, and effective manner to enhance their health and well-being.⁷⁴ Health providers could probe women about their experiences of violence or evaluate them for any potential indicator of partner violence, such as any history of unexplained injury or maternal bleeding, preterm labour or birth, and foetal injury or death.⁷³ The healthcare system can also provide women with a safe environment in which they can confidentially disclose experiences of violence and receive a supportive response.

Although our review addresses an important gap in the literature, it is not without limitation. The first is that the inclusion of articles in this review is based solely on their focus on vulnerability or resilience in LMICs. Therefore, studies on vulnerability or resilience outside of LMICs, in locations where pockets of vulnerable populations occur in high-income nations have not been captured. Additionally, while we made every attempt to find all accessible material, it is possible that we omitted some publications with distinct perspectives that were not represented in the review's evidence from grey literature particularly, given how broad it is.

CONCLUSION

Only a few studies have examined vulnerability and resilience in maternal and child health, especially in LMICs. We have identified some gendered dynamics of vulnerability and resilience in MNCH through this scoping review. Findings from this scoping review suggest that there is a great need to continue to empower women and mothers to access resources, contribute to decisions about their own health, and eliminate structural or social stereotypes that limit their agency.

Author affiliations

¹Department of Research and Development, Viable Helpers Development Organization, Abuja, Federal Capital Territory, Nigeria

²Division of Health Sciences, University of Warwick, Coventry, UK

³Department of Obstetrics and Gynaecology, University of Lagos College of Medicine, Lagos, Nigeria

⁴Department of Research and Development, Viable Knowledge Masters, Gwarinpa, Federal Capital Territory, Nigeria

Twitter Emmanuel Olawale Olamijuwon @eolamijuwon, Babasola Okusanya @Babsokusanya and Olusesan Ayodeji Makinde @sesmak

Acknowledgements We would like to thank the Bill and Melinda Gates Foundation for funding this work. We would like to acknowledge Ms Claire-Helene Mershon, the Program Officer at the Foundation, who guided us through this piece of work.

Contributors OAM conceptualised the review, developed the initial search strategy for the study, screened studies for eligibility, reviewed draft manuscript and supervised the overall research. OAU developed the search strategy and machine learning programme for study screening and screened studies for eligibility. EOO and FAS drafted the manuscript. NKI, ICM and BO screened studies for eligibility, performed data extraction and reviewed the draft manuscript. All authors read the manuscript, contributed to the revisions as required and approved the final manuscript. OAM takes responsibility for the overall content as the guarantor.

Funding This work was supported by the Bill & Melinda Gates Foundation under investment INV-015806 awarded to the Viable Helpers Development Organisation.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study did not receive nor require ethics approval, as it does not involve human and animal participants.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Fatima Abdulaziz Sule <http://orcid.org/0000-0001-8684-0630>

Olalekan A Uthman <http://orcid.org/0000-0002-8567-3081>

Emmanuel Olawale Olamijuwon <http://orcid.org/0000-0002-6109-8131>

Nchelem Kokomma Ichehgo <http://orcid.org/0000-0003-1325-8759>

Babasola Okusanya <http://orcid.org/0000-0003-2381-873X>

Olusesan Ayodeji Makinde <http://orcid.org/0000-0002-1575-3588>

REFERENCES

- Sharrow D, Hug L, Liu Y. Levels and trends in child mortality: report, 2020. New York: UNICEF. Available: <https://www.unicef.org/media/79371/file/UN-IGME-child-mortality-report-2020.pdf.pdf> [Accessed 13 Feb 2022].
- WHO. Maternal mortality, 2019. Available: <https://www.who.int/news-room/fact-sheets/detail/maternal-mortality> [Accessed 11 Dec 2020].
- Sharrow D, Liu S, Liu Y, et al. Levels and trends in child mortality: report 2020, 2020. New York: UNICEF. Available: <https://www.unicef.org/media/79371/file/UN-IGME-child-mortality-report-2020.pdf.pdf> [Accessed 13 Feb 2022].
- Scheele J, Harmsen van der Vliet-Torij HW, Wingelaar-Loomans EM, et al. Defining vulnerability in European pregnant women, a Delphi study. *Midwifery* 2020;86:102708.
- Bayntun C, Rockenschaub G, Murray V. Developing a health system approach to disaster management: a qualitative analysis of the core literature to complement the who toolkit for assessing health-system capacity for crisis management. *PLoS Curr* 2012;4:e5028b6037259a.
- Andriano L, Monden CWS. The causal effect of maternal education on child mortality: evidence from a Quasi-Experiment in Malawi and Uganda. *Demography* 2019;56:1765–90.
- Bolarinwa OA, Ameen HA, Durowade KA. Mapping knowledge management resources of maternal, newborn and child health (MNCH) among people living in rural and urban settings of Ilorin, Nigeria. *Pan Afr Med J* 2014;17.
- Dolan CB, BenYishay A, Grépin KA, et al. The impact of an insecticide treated bednet campaign on all-cause child mortality: a geospatial impact evaluation from the Democratic Republic of Congo. *PLoS One* 2019;14:e0212890.
- Gakidou E, Cowling K, Lozano R, et al. Increased educational attainment and its effect on child mortality in 175 countries between 1970 and 2009: a systematic analysis. *Lancet* 2010;376:959–74.
- Kinney MV, Kerber KJ, Black RE, et al. Sub-Saharan Africa's mothers, newborns, and children: where and why do they die? *PLoS Med* 2010;7:e1000294.
- Pickbourn L, Ndikumana L. Does health aid reduce infant and child mortality from diarrhoea in sub-Saharan Africa? *J Dev Stud* 2019;55:2212–31.
- Seely HD, Mickelson KD. Maternal resilience as a protective factor between financial stress and child outcomes. *J Fam Issues* 2019;40:1604–26.
- Creighton G, Oliffe JL. Theorising masculinities and men's health: A brief history with a view to practice. *Health Sociology Review* 2010;19:409–18.
- Braveman P, Gruskin S. Defining equity in health. *J Epidemiol Community Health* 2003;57:254–8.
- Greaves L, Jategaonkar N. Tobacco policies and vulnerable girls and women: toward a framework for gender sensitive policy development. *J Epidemiol Community Health* 2006;60:ii57–65.
- Hemings N, Greaves L. Gender norms, roles and relations and Cannabis-Use patterns: a scoping review. *Int J Environ Res Public Health* 2020;17:17030947. doi:10.3390/ijerph17030947
- Ostlin P, Eckermann E, Mishra US, et al. Gender and health promotion: a multisectoral policy approach. *Health Promot Int* 2006;21:25–35.
- Wechsberg WM, Browne FA, Bonner CP, et al. Current interventions for people living with HIV who use alcohol: why gender matters. *Curr HIV/AIDS Rep* 2021;18:351–64.
- Dunkle KL, Jewkes R. Effective HIV prevention requires gender-transformative work with men. *Sex Transm Infect* 2007;83:173–4.
- Ruane-McAteer E, Gillespie K, Amin A, et al. Gender-transformative programming with men and boys to improve sexual and reproductive health and rights: a systematic review of intervention studies. *BMJ Glob Health* 2020;5:e002997.
- Rasanathan K, Damji N, Atsbeha T, et al. Ensuring multisectoral action on the determinants of reproductive, maternal, newborn, child, and adolescent health in the post-2015 era. *BMJ* 2015;351:h4213.
- Morgan R, Dhatt R, Kharel C, et al. A patchwork approach to gender equality weakens the SDGs: time for cross-cutting action. *Glob Health Promot* 2020;27:3–5.
- Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005;8:19–32.
- Peters MDJ, Marnie C, Tricco AC, et al. Updated methodological guidance for the conduct of scoping reviews. *JBI Evid Synth* 2020;18:2119–26.
- Bannach-Brown A, Przybyła P, Thomas J, et al. Machine learning algorithms for systematic review: reducing workload in a preclinical review of animal studies and reducing human screening error. *Syst Rev* 2019;8:23.
- Hempel S, Shetty KD, Shekelle PG. *Machine learning methods in systematic reviews: identifying quality improvement intervention evaluations*. Rockville, MD: Agency for Healthcare Research and Quality (US), 2012. <http://www.ncbi.nlm.nih.gov/books/NBK109711/>
- Marshall IJ, Wallace BC. Toward systematic review automation: a practical guide to using machine learning tools in research synthesis. *Syst Rev* 2019;8:163.

- 28 Morgan R, George A, Ssali S, *et al.* How to do (or not to do)... gender analysis in health systems research. *Health Policy Plan* 2016;31:1069–78.
- 29 Tricco AC, Lillie E, Zarin W, *et al.* PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018;169:467–73.
- 30 Amoako Johnson F, Padmadas SS, Smith PWF. Orphanhood and vulnerability: a conduit to poor child health outcomes in Rwanda. *AIDS Care* 2010;22:314–23.
- 31 Arcos E, Munoz LA, Sanchez X. [Vulnerability of pregnant women living in a community of Metropolitan Santiago] 2011;139:739–47.
- 32 Lavender T, Bedwell C, Blaikie K, *et al.* Journey of vulnerability: a mixed-methods study to understand intrapartum transfers in Tanzania and Zambia. *BMC Pregnancy Childbirth* 2020;20:292.
- 33 Pourette D. [Haitian women living with HIV/AIDS in Guadeloupe: pregnancy between medical stakes, social benefits and administrative vulnerability. A qualitative study]. *Bull Soc Pathol Exot* 2012;105:143–9.
- 34 Rotheram-Borus MJ, Christodoulou J, Hayati Rezvan P, *et al.* Maternal HIV does not affect resiliency among uninfected/HIV exposed South African children from birth to 5 years of age. *AIDS* 2019;33:S5–16.
- 35 Scheidell JD, Beau De Rochars VM, Séraphin MN, *et al.* Socioeconomic vulnerability and sexually transmitted infection among pregnant Haitian women. *Sex Transm Dis* 2018;45:626–31.
- 36 Warren CE, Sripad P, Mwangi A. “Sickness of shame”: investigating challenges and resilience among women living with obstetric fistula in Kenya. In: Choudhury S, Erausquin JT, Withers M, eds. *Global perspectives on women’s sexual and reproductive health across the lifecourse*. Cham: Springer International Publishing, 2018: 91–109.
- 37 Zakayo SM, Njeru RW, Sanga G. Vulnerability and agency across treatment-seeking journeys for acutely ill children: how family members navigate complex healthcare before, during and after hospitalisation in a rural Kenyan setting. *a* 2020;19:136.
- 38 Kaye DK, Kakaire O, Nakimuli A, *et al.* Survivors’ understanding of vulnerability and resilience to maternal near-miss obstetric events in Uganda. *Int J Gynaecol Obstet* 2014;127:265–8.
- 39 Storeng KT, Drabo S, Filippi V. Too poor to live? A case study of vulnerability and maternal mortality in Burkina Faso. *Glob Health Promot* 2013;20:33–8.
- 40 Den hollander GC, Browne JI, Arhinful D, *et al.* Power difference and risk perception: mapping vulnerability within the decision process of pregnant women towards clinical trial participation in an urban middle-income setting. *Dev World Bioeth* 2018;18:68–75.
- 41 Prates CdeS, Abib GMdeC, de Oliveira DLLC. [Gender power, poverty and contraception: multiparous women’s experiences]. *Rev Gaucha Enferm* 2008;29:604–11.
- 42 McNaughton Reyes HL, Maman S, Groves AK, *et al.* Intimate partner violence and postpartum emotional distress among South African women: moderating effects of resilience and vulnerability factors. *Glob Public Health* 2020;15:1157–67.
- 43 Vivilaki VG, Dafermos V, Patelarou E, *et al.* Maternal screening for early postnatal vulnerability. *Psychiatriki* 2016;27:169–81.
- 44 Gaillard P, Melis R, Mwanyumba F, *et al.* Vulnerability of women in an African setting: lessons for mother-to-child HIV transmission prevention programmes. *AIDS* 2002;16:937–9.
- 45 Nie C, Dai Q, Zhao R, *et al.* The impact of resilience on psychological outcomes in women with threatened premature labor and spouses: a cross-sectional study in Southwest China. *Health Qual Life Outcome* 2017;15:26.
- 46 Nor Jana S, Ghazinour M, Richter J. Mental health predicted by coping, social support, and resilience among young unwed pregnant Malaysian women and mothers living in shelter homes. *Int J Hum Rights Healthc* 2016;9:185–97.
- 47 Brewis A, Wutich A. Why we should never do it: stigma as a behaviour change tool in global health. *BMJ Glob Health* 2019;4:e001911.
- 48 Makinde OA, Olaleye O, Makinde OO, *et al.* Baby factories in Nigeria: starting the discussion toward a national prevention policy. *Trauma Violence Abuse* 2017;18:98–105 <https://journals.sagepub.com/doi/10.1177/1524838015591588> doi:10.1177/1524838015591588
- 49 Makinde OA. Infant trafficking and baby factories: a new tale of child abuse in Nigeria. *Child Abuse Rev* 2016;25:433–43.
- 50 Ajayi AI, Akpan W. Maternal health care services utilisation in the context of ‘Abiye’ (safe motherhood) programme in Ondo State, Nigeria. *BMC Public Health* 2020;20:362.
- 51 Liu X, Yan H, Wang D. The evaluation of ‘Safe Motherhood’ program on maternal care utilization in rural western China: a difference in difference approach. *BMC Public Health* 2010;10:566.
- 52 Li Z, Li M, Fink G, *et al.* User-fee-removal improves equity of children’s health care utilization and reduces families’ financial burden: evidence from Jamaica. *J Glob Health* 2017;7.
- 53 Hadley M. Does increase in utilisation rates alone indicate the success of a user fee removal policy? A qualitative case study from Zambia. *Health Policy* 2011;103:244–54.
- 54 Plouffe V, Bicaba F, Bicaba A, *et al.* User fee policies and women’s empowerment: a systematic scoping review. *BMC Health Serv Res* 2020;20:982.
- 55 Hatt LE, Makinen M, Madhavan S, *et al.* Effects of user fee exemptions on the provision and use of maternal health services: a review of literature. *J Health Popul Nutr* 2013;31:S67–80.
- 56 Dahab R, Sakellariou D. Barriers to accessing maternal care in low income countries in Africa: a systematic review. *Int J Environ Res Public Health* 2020;17:17124292. doi:10.3390/ijerph17124292
- 57 Masiye F, Kaonga O, Kirigia JM. Does user fee removal policy provide financial protection from catastrophic health care payments? Evidence from Zambia. *PLoS One* 2016;11:e0146508.
- 58 Lépine A, Lagarde M, Le Nestour A. How effective and fair is user fee removal? Evidence from Zambia using a pooled synthetic control. *Health Econ* 2018;27:493–508.
- 59 Obare F, Abuya T, Matanda D, *et al.* Assessing the community-level impact of a decade of user fee policy shifts on health facility deliveries in Kenya, 2003–2014. *Int J Equity Health* 2018;17:65.
- 60 Morgan L, Stanton ME, Higgs ES. Financial incentives and maternal health: where do we go from here? *J Health Popul Nutr* 2013;31:S8–22.
- 61 Todd PE. *Effectiveness of interventions aimed at improving women’s employability and quality of work: a critical review*. Washington, DC: World Bank, 2012.
- 62 Clark S, Kabiru CW, Laszlo S, *et al.* The impact of childcare on poor urban women’s economic empowerment in Africa. *Demography* 2019;56:1247–72.
- 63 Erzse A, Goldstein S, Tugendhaft A, *et al.* The roles of men and women in maternal and child nutrition in urban South Africa: a qualitative secondary analysis. *Matern Child Nutr* 2021;17:e13161.
- 64 Abramsky T, Kapinga I, Mshana G, *et al.* Couples data from north-western Tanzania: insights from a survey of male partners of women enrolled in the MAISHA cluster randomized trial of an intimate partner violence prevention intervention. *PLoS One* 2020;15:e0240112.
- 65 Oduaran A, Chukwudeh O. Trap in the closet: intra-ethnic marriage and intimate partner violence in sub-Saharan Africa. *Soc Sci* 2021;10:31.
- 66 Zust BL, Housley J, Klatke A. Evangelical christian pastors’ lived experience of counseling victims/survivors of domestic violence. *Pastoral Psychol* 2017;66:675–87.
- 67 Sichimba F, Nakazwe KC, Phiri T. Untold stories of women living in violence: lived realities of why women stay: a case study of Ngombe and Kanyama compounds in Lusaka. *J Aggress Maltreat Trauma* 2020;29:767–84.
- 68 Saraswati R. Shame and Indonesian women victims of domestic violence in making the decision to divorce. *Identities* 2020;27:557–73.
- 69 Kapiga S, Harvey S, Mshana G, *et al.* A social empowerment intervention to prevent intimate partner violence against women in a microfinance scheme in Tanzania: findings from the MAISHA cluster randomised controlled trial. *Lancet Glob Health* 2019;7:e1423–34.
- 70 Abramsky T, Watts CH, Garcia-Moreno C, *et al.* What factors are associated with recent intimate partner violence? findings from the WHO multi-country study on women’s health and domestic violence. *BMC Public Health* 2011;11:109.
- 71 Bourey C, Williams W, Bernstein EE, *et al.* Systematic review of structural interventions for intimate partner violence in low- and middle-income countries: organizing evidence for prevention. *BMC Public Health* 2015;15:1165.
- 72 Robinson AB, Stephenson R, Merrill KG. Missed opportunities for addressing intimate partner violence through Microcredit: qualitative findings from Bangladesh. *Violence Against Women* 2020;1077801220963906.
- 73 Watts C, Mayhew S. Reproductive health services and intimate partner violence: shaping a pragmatic response in sub-Saharan Africa. *Int Fam Plan Perspect* 2004;30:207–13.
- 74 Garcia-Moreno C, Hegarty K, d’Oliveira AFL, *et al.* The health-systems response to violence against women. *Lancet* 2015;385:1567–79.

Appendix 1: Database search strategies

Maternal Health

'maternal health'.ti,ab,kw.
('pregnancy'.ti,ab,kw. OR 'adolescent pregnancy'.ti,ab,kw. OR 'first trimester pregnancy'.ti,ab,kw. OR 'multiple pregnancy'.ti,ab,kw. OR 'twin pregnancy'.ti,ab,kw. OR 'second trimester pregnancy'.ti,ab,kw. OR 'third trimester pregnancy'.ti,ab,kw. OR 'unplanned pregnancy'.ti,ab,kw. OR 'unwanted pregnancy'.ti,ab,kw. ((pregnan*) adj3 (adolescen* OR teen* OR multip* OR twin* OR quad* OR quin* OR trip* OR trimester OR early OR late OR unplanned OR unintended OR unwanted OR undesired OR intrauterine OR maintenance)) (twining* OR 'multiple birth' OR quadruplet* OR quintuplet* OR triplet* OR superfetation OR 'child bearing' OR childbearing OR gestation OR gravidity OR 'labor presentation' OR ((trimester) adj3 (first OR second OR third OR 1st OR 2nd OR 3rd OR mid*))).ti,ab,kw.
Pregnancy in Adolescence/
(pregnan* adj2 (adolescen* or teen* or schoolchild*)).ti,ab,kw.
Pregnancy, unplanned/ or Pregnancy, unwanted/
pregnan* adj3 (prevent* or interrupt* or unplanned or unwanted or mistimed)).ti,ab,kw.
Prenatal Care/
(((antenatal or ante-natal or prenatal or pre-natal or antepartum or ante-partum) adj3 (care or service* or counsel* or test*)) or (birth adj3 prepar*)).ti,ab,kw.
Maternal Health Services/
((maternal or mother*) adj3 (health or service* or care)).ti,ab,kw.
Midwifery/
(midwi* or skilled birth or skilled attendan*).ti,ab,kw.
Postnatal Care/
Perinatal Care/
Postpartum Period/
((postnatal or post-natal or perinatal or peri-natal or postpartum or post-partum) adj2 (care or service*)).ti,ab,kw.
Maternal Nutritional Physiological Phenomena/
Prenatal Nutritional Physiological Phenomena/
Breast Feeding/
(breast feed* or breast fed or breastfeed* or breastfed).ti,ab,kw.

Child health

(Infant health)) OR (Infant Services, Health)) OR (Service, Child Health)) OR (Infant Health Service)) OR (Neonate)) OR (Neonate)) OR (Neonates)) OR (Newborn Infants)) OR (Infants, Newborn) (Infant, Premature/ or Infant, Newborn/ or Infant, Low Birth Weight/ or Infant, Extremely Low Birth Weight/ or Infant, Small for Gestational Age/ or Infant/ or Infant, Very Low Birth Weight/ or Infant, Postmature/ or Infant, Extremely Premature/) and Early Diagnosis/ (early adj1 diagnos* adj2 (infant* or neonat* or newborn*).ti,ab,kw. diagnosis.fs. and (infant* or neonat* or newborn*).ti,ab,kw. (Immunization/ or Immunization, passive/ or Immunization schedule/ or Immunization, secondary/ or Immunization Programs/ or Vaccination/ or Mass vaccination/) and (Infant, Premature/ or Infant, Newborn/ or Infant, Low Birth Weight/ or Infant, Extremely Low Birth Weight/ or Infant, Small for Gestational Age/ or Infant/ or Infant, Very Low Birth Weight/ or Infant, Postmature/ or Infant, Extremely Premature/ or Child/ or Child, Preschool/ or Adolescent/ or Pregnancy/) ((immuniz* or immunis* or vaccinat*) and (infant* or newborn* or neonat* or child* or adolescen* or teen*).ti,ab,kw. Child health services/ or Maternal-child health services/ exp child nutrition disorders/ or exp infant nutrition disorders/ or (exp nutrition disorders/ and (exp Infant/ or Child, Preschool/)) 'Delivery of Health Care, Integrated'/ ((integrat* adj3 (health care or healthcare or management or treat* or service*) adj3 (child* or schoolchild* or infant* or neonat* or newborn or adolescen* or teen*)) or IMCI or IMNCI).ti,ab,kw. (Guideline Adherence/ or Quality Assurance, Health Care/) and (exp Infant/ or Child, Preschool/ (((guideline* or protocol*) adj3 (adher* or observ*)) or "prescribed care") and (infant* or newborn* or neonat* or child*).ti,ab,kw. (Diarrhea/di, dt, ep, pc, th, tm or Diarrhea, Infantile/di, dt, ep, pc, th, tm) and (Infant, Premature/ or Infant, Newborn/ or Infant, Low Birth Weight/ or Infant, Extremely Low Birth Weight/ or Infant, Small for Gestational Age/ or Infant/ or Infant, Very Low Birth Weight/ or Infant, Postmature/ or Infant, Extremely Premature/ or Child/ or Child, Preschool/ or Adolescent/) (diarrh* and (infant* or newborn* or neonat* or child* or schoolchild* or adolescen* or teen*).ti,ab,kw. Fluid Therapy/ and (Infant, Premature/ or Infant, Newborn/ or Infant, Low Birth Weight/ or Infant, Extremely Low Birth Weight/ or Infant, Small for Gestational Age/ or Infant/ or Infant, Very Low Birth Weight/ or Infant, Postmature/ or Infant, Extremely Premature/ or Child/ or Child, Preschool/) Child Development/ or Adolescent Development/

Health utilisation

'health care system'.ti,ab,kw.
'health care access'.ti,ab,kw.
'health care availability'.ti,ab,kw.
'health care disparity'.ti,ab,kw.
'health care distribution'.ti,ab,kw.
'health care need'.ti,ab,kw.
'health care planning'.ti,ab,kw.
'healthcare system'.ti,ab,kw.
'health disparity'.ti,ab,kw.
'health care delivery'.ti,ab,kw.
'ambulatory care'.ti,ab,kw.
'hospital care'.ti,ab,kw.
'primary health care'.ti,ab,kw.
'secondary health care'.ti,ab,kw.
'tertiary care center'.ti,ab,kw.
'health care quality'.ti,ab,kw.
((health* OR service*) adj3 (access* OR availab* OR disparat* OR equit* OR inequal* OR plan* OR priorit* OR resource* OR suppl* OR deliver*)).ti,ab,kw.

Health Outcomes

Obstetric Labor Complications/
 Pregnancy Complications/
 stillbirth.ti,ab,kw.
 stillborn.ti,ab,kw.
 ((obstetric* or pregnan* or labour or labor or parturition) adj3 (emergenc* or complication*)).ti,ab,kw.
 ('maternal morbidity'.ti,ab,kw. OR 'maternal mortality'.ti,ab,kw. OR 'perinatal death'.ti,ab,kw. OR 'perinatal morbidity'/exp OR 'newborn morbidity'.ti,ab,kw. OR 'newborn death'.ti,ab,kw. OR 'child death'.ti,ab,kw. OR 'infant mortality'.ti,ab,kw. OR 'small for date infant'.ti,ab,kw. OR 'low birth weight'.ti,ab,kw. OR 'extremely low birth weight'.ti,ab,kw. OR 'very low birth weight'.ti,ab,kw. OR 'apgar score'.ti,ab,kw.
 ((maternal OR mother OR perinatal OR neonat* OR infant*) NEAR/3 (death OR mortality OR morbidity OR disease)) OR 'small for gestational age' OR ((SGA OR 'small for gestational age' OR 'small for date' OR 'small for age') adj3 (infant* OR neonat* OR newborn OR baby))
 (('ELBW' OR 'VLBW' OR 'LBW' OR (('birth weight' OR birthweight OR 'LBW' OR underweight) adj3 (infant* OR neonat* OR newborn OR low))
 ((apgar OR APGAR) adj3 (classif* OR coeff* OR index* OR rating OR scale OR test OR timer OR score)).ti,ab,kw.
 ((child* or schoolchild* or adolescen* or teen*) adj2 (develop* or progress*)).ti,ab,kw.
 Pregnancy in Adolescence/
 Kangaroo-Mother Care Method/
 (kangaroo adj2 (mother or infant or care)).ti,ab,kw.
 (Anemia/dt, pc or Anemia, Hypochromic/dt, pc or Anemia, Iron-Deficiency/dt, pc) and Pregnancy/
 ((maternal or mother* or pregnan*) adj2 (nutrition* or folate or folic or iron or anaemi* or anemi*)).ti,ab,kw.
 (Malaria/di, dt, pc or Malaria, Falciparum/di, dt, pc or Malaria, Vivax/di, dt, pc) and (Pregnancy/ or Pregnancy Complications, Parasitic/
 ((malaria* or falciparum or vivax) adj3 (pregnan* or mother* or maternal or postpartum or postpartum)).ti,ab,kw.
 Mental health/ or Mental disorders/ or Mental health services/ or Community mental health services/
 Maternal behavior/ or Mother-child relations/ or Parenting/ or Paternal behavior/
 Depression, Postpartum/
 (((mental or behavio*) adj3 (health or disorder*)) or postpartum depression or post-partum depression).ti,ab,kw.

Social capital

'social determinants of health'.ti,ab,kw.
'neighborhood'.ti,ab,kw. OR 'community'.ti,ab,kw. OR 'religion'.ti,ab,kw. OR 'occupation'.ti,ab,kw.
'employment'/exp OR 'income'.ti,ab,kw. OR 'ethnicity'.ti,ab,kw. OR 'race'.ti,ab,kw. OR 'education'.ti,ab,kw.
'social status'/exp OR 'lowest income group'.ti,ab,kw. OR 'poverty'.ti,ab,kw. OR 'social background'.ti,ab,kw.
'social class'.ti,ab,kw. OR 'social network'.ti,ab,kw. OR 'social capital'.ti,ab,kw. OR 'health literacy'.ti,ab,kw. OR
'locus of control'.ti,ab,kw.
'attitude to health'.ti,ab,kw. OR 'health insurance'.ti,ab,kw. OR 'private insurance'.ti,ab,kw. OR 'public health
insurance'.ti,ab,kw. OR
'social stigma'.ti,ab,kw. OR 'stigma'.ti,ab,kw.
'risk perception' OR 'medically uninsured'.ti,ab,kw. OR 'risk factor'.ti,ab,kw. OR 'relative risk'
((health*) adj3 (knowledge OR attitude* OR practice OR accept* OR barrier))
'HLC' OR 'health locus of control' OR uninsured OR 'insurance status'
neighborhood OR neighbourhood OR 'poverty areas' OR 'religious belief' OR spiritual* OR profession OR
'employment status'
income OR 'social network' OR 'sociocultural class' OR 'socioeconomic class' OR residenc*
(education*) adj3 (achievement OR attain* OR level))
((social*) adj3 (determin* OR factor))
((underserve*) adj3 (neighborhood OR neighbourhood OR area))
((social* OR socio* OR religious OR gender OR politic* OR econom*) NEAR/3 (barrier OR threshold)).ti,ab,kw.)

Resource limited setting

'developing countries'.ti. OR 'low-income countries'.ti. OR 'middle-income country'.ti. OR Imic.ti. OR Imics.ti. OR 'less-developed countries'.ti. OR 'under-developed nations'.ti. OR 'third-world country'.ti. OR 'third-world nations'.ti.

(afghanistan OR albania OR algeria OR american samoa OR angola OR "antigua and barbuda"
OR antigua OR barbuda OR argentina OR armenia OR armenian OR aruba OR azerbaijan OR bahrain OR bangladesh OR barbados OR republic
of belarus OR belarus OR byelarus OR belorussia OR byelorussian OR belize OR british honduras OR benin OR dahomey OR bhutan OR bolivia OR "bosnia and herzegovina"
OR bosnia OR herzegovina OR botswana OR bechuanaland OR brazil OR brasil OR bulgaria OR burkina faso OR burkina fasso OR upper volta OR burundi OR urundi OR cabo verde OR
cape verde OR cambodia OR kampuchea OR khmer republic OR cameroon OR cameron OR cameroun OR central african republic OR ubangi shari OR chad OR chile OR china OR colombia OR comoros OR comoro islands
OR iles comores OR mayotte OR democratic republic of the congo OR democratic republic congo OR congo OR zaire OR costa rica OR "cote d'ivoire" OR "cote d'ivoire" OR cote d'ivoire OR cote d'ivoire OR ivory coast
OR croatia OR cuba OR cyprus OR czech republic OR czechoslovakia OR djibouti OR french somaliland OR dominica OR dominican republic OR ecuador OR egypt OR united arab republic OR el salvador OR equatorial guinea
OR spanish guinea OR eritrea OR estonia OR eswatini OR swaziland OR ethiopia OR fiji OR gabon OR gabonese republic OR gambia OR "georgia (republic)" OR georgian OR ghana OR gold coast
OR gibraltar OR greece OR grenada OR guam OR guatemala OR guinea OR guinea bissau OR guyana OR british guiana OR haiti OR hispaniola OR honduras OR hungary OR india OR indonesia OR timor OR iran OR iraq OR isle of
man OR jamaica OR jordan OR kazakhstan OR kazakh OR kenya OR "democratic people's republic of korea" OR republic of korea OR north korea OR
south korea OR korea OR kosovo OR kyrgyzstan OR kirghizia OR kirgizstan OR kyrgyz republic OR kirghiz OR laos OR lao pdr OR "lao people's democratic republic" OR latvia OR lebanon OR lebanese republic
OR lesotho OR basutoland OR liberia OR libya OR libyan arab jamahiriya OR lithuania OR macau OR macao OR republic of north macedonia OR macedonia OR madagascar OR malagasy republic
OR malawi OR nysalaland OR malaysia OR malaya federation OR malaya federation OR maldives OR indian ocean islands OR indian ocean OR mali OR malta OR micronesia OR federated states
of micronesia OR kiribati OR marshall islands OR nauru OR northern mariana islands
OR palau OR tuvalu OR mauritania OR mauritius OR mexico OR moldova OR moldovian OR mongolia OR montenegro OR morocco OR ifni OR mozambique OR portuguese east africa OR myanmar OR burma OR namibia OR nep
al OR netherlands antilles OR nicaragua OR niger OR nigeria OR oman OR muscat OR pakistan OR panama OR papua new guinea OR new guinea
OR paraguay OR peru OR philippines OR philippines OR philippines OR philippines OR poland OR "polish people's republic" OR portugal OR portuguese republic OR puerto rico OR romania OR russia OR russian federation
OR ussr OR soviet union OR union of soviet socialist republics OR rwanada OR ruanda OR samoa OR pacific islands OR polynesia OR samoa islands OR navigator island OR navigator islands OR "sao tome and principe"
OR saudi arabia OR senegal OR serbia OR seychelles OR sierra leone OR slovakia OR slovak republic OR slovenia OR melanesia OR solomon island OR solomon islands OR norfolk island OR norfolk islands OR somalia OR
south africa OR south sudan OR sri lanka OR ceylon OR "saint kitts and nevis" OR "st. kitts and nevis" OR saint lucia OR "st. lucia" OR "saint vincent and the grenadines" OR saint vincent OR "st. vincent" OR grenadines
OR sudan OR suriname OR surinam OR dutch guiana OR netherlands guiana OR syria OR syrian arab republic
OR tajikistan OR tadjikistan OR tadjikistan OR tadjik OR tanzania OR tanganyika OR thailand OR siam OR timor leste OR east timor OR togo OR togolese republic OR tonga OR "trinidad and tobago"
OR trinidad OR tobago OR tunisia OR turkey OR turkmenistan OR turkmen OR uganda OR ukraine OR uruguay OR uzbekistan OR uzbek OR vanuatu OR new hebrides OR venezuela OR vietnam OR viet nam OR middle east OR
west bank OR gaza OR palestine OR yemen OR yugoslavia OR zambia OR zimbabwe OR northern rhodesia OR global south OR africa south of the sahara OR sub-saharan africa OR subsaharan africa OR africa, central OR
central africa OR africa, northern OR north africa OR northern africa OR magreb OR maghrib OR sahara OR africa, southern OR southern africa OR africa, eastern OR east africa OR eastern africa OR africa, western OR
west africa OR western africa OR west indies OR indian ocean islands OR caribbean OR central america OR latin america OR "south and central america" OR south america OR asia, central OR central asia OR asia, northern OR
north asia OR northern asia OR asia, southeastern OR southeastern asia OR south eastern asia OR southeast asia OR south east asia OR asia, western OR western asia OR europe, eastern OR east europe OR eastern europe OR
developing country OR developing countries OR developing nation? OR developing population? OR developing world OR less developed countr* OR less developed nation? OR less developed population? OR less developed
world OR lesser developed countr* OR lesser developed nation? OR lesser developed population? OR lesser developed world OR under developed countr* OR under developed nation? OR under developed population?
OR under developed world OR underdeveloped countr* OR underdeveloped nation? OR underdeveloped population? OR underdeveloped world OR middle income countr* OR middle income nation? OR middle
income population? OR low income countr* OR low income nation? OR low income population? OR lower income countr* OR lower income nation? OR lower income population? OR underserved countr* OR underserved
nation? OR underserved population? OR underserved world OR under served countr* OR under served nation? OR under served population? OR under served world OR deprived countr* OR deprived nation? OR deprived
population? OR deprived world OR poor countr* OR poor nation? OR poor population? OR poor world OR poorer countr* OR poorer nation? OR poorer population? OR poorer world OR developing econom* OR less
developed econom* OR lesser developed econom* OR under developed econom* OR underdeveloped econom* OR middle income econom* OR low income econom* OR lower income econom* OR low gdp OR low gnp OR
low gross domestic OR low gross national OR lower gdp OR lower gnp OR lower gross domestic OR lower gross national OR Imic OR Imics OR third world OR lami countr* OR transitional countr* OR emerging economies OR
emerging nation?),ti,ab,sh,kf.

(afghan OR afghans OR afghani OR albanian? OR algerian? OR american samoan? OR angolan? OR antiguan? OR barbudan? OR argentine? OR argentinian? OR argentinean? OR armenian? OR aruban? OR azerbaijani?
OR bahraini? OR bangladeshi? OR bangalees OR bajan? OR belarusian? OR byelorussian? OR belizean? OR beninese? OR bhutanese OR bolivian? OR bosnian? OR botswana OR batswana OR brazilian? OR brasilian?
OR bulgarian? OR burkinabe OR burkinese OR burundian? OR cape verdean? OR cabo verdean? OR cambodian? OR khmer OR cameroonian? OR central african? OR chadian? OR chilean? OR chinese OR colombian?
OR comorian? OR congolese OR costa rican? OR ivorian? OR croatian? OR cuban? OR cyriot? OR czech? OR djiboutian? OR dominican? OR ecuadorian? OR egyptian? OR salvadoran? OR equatorial guinean?
OR equatoguinean? OR eritrean? OR estonian? OR swazi? OR swati? OR ethiopian? OR fiji OR gabonese OR gabonaise OR gambian? OR georgian? OR ghanaian? OR gibraltarian? OR greek? OR grenadian? OR guamanian?
OR guatemalan? OR guinean? OR bissau guinean? OR guyanese OR haitian? OR honduran? OR indian? OR indonesian? OR iranian? OR iraqian? OR iraqi? OR manx OR jamaican? OR jordanian? OR kazakhstanian?
OR kenyan? OR kirabati OR kirabatian? OR north korean? OR korean? OR kosovar? OR kosovan? OR kyrgyz* OR lao OR laotian? OR latvian? OR lebanese OR lesothan? OR lesothonian? OR mosotho OR basotho OR liberian?
OR libyan? OR lithuanian? OR macanese OR macedonian? OR malagasy OR madagascan? OR malawian? OR malaysian? OR maldivian? OR malian? OR maltese OR marshallese? OR mauritanian? OR mauritian? OR mexican?
OR micronesian? OR moldovan? OR mongolian? OR mongol OR montenegrin? OR moroccan? OR mozambican? OR burmese OR myanma OR namibian? OR nauruan? OR nepali OR nepalese OR netherlands antillean?
OR nicaraguan? OR nigerien? OR nigerian? OR northern mariana islander? OR mariana? OR omani? OR pakistani? OR palauan? OR panamanian? OR papua new guinean? OR paraguayian? OR peruvian? OR philippine?
OR philippine? OR philippine? OR philippine? OR filipino? OR filipina? OR polish OR pole OR poles OR portuguese OR puerto rican? OR romanian? OR russian? OR soviet people OR soviet population OR rwanadan?
OR rwandese OR ruanadan? OR ruandese OR samoan? OR sao tomean? OR santomean? OR saudi arabian? OR saudi? OR seychellois OR seychelloise? OR sierra leonean?

Supplementary file 2: Full details of the 17 studies included in the scoping review

S/N	Author Year	Publication Title	Country [Context]	Main-Theme	Main Findings
1	Rotheram-Borus et al. (2009)	Maternal HIV does not affect resiliency among uninfected/HIV exposed South African children from birth to 5 years of age	South Africa [Child health]	Access to resources [Who has what?]	Resiliency was significantly associated with lower income, food security, not having a live-in partner, and the absence of maternal risk (i.e. not being depressed, using alcohol, or being a victim of intimate partner violence).
2	Jana et al. (2013)	Mental health predicted by coping, social support, and resilience among young unwed pregnant Malaysian women and mothers living in shelter homes	Malaysia [Maternal health]	Partner emotional or mental support/acceptance	Families and partners often reacted to the pregnancy by rejection of the unwed pregnant teenagers or teenage mothers. These rejections took different expressions in terms of avoiding them or verbal harassment.
3	Vivilaki et al. (2016)	Maternal screening for early postnatal vulnerability	Crete [Maternal health]	Power negotiation [How is power enacted, negotiated, or challenged?]	The lack of or disappointment with partner support, poor marital relationship, and emotional/physical abuse have been associated with high levels of postpartum anxiety and depression.
4	Nie et al. (2017)	The impact of resilience on psychological outcomes in women with threatened premature labor and spouses: a cross-sectional study in Southwest China	China [Maternal health]	Partner emotional or mental support/acceptance	The analysis suggested that low resilient women with threatened premature labour reported higher pressures from concerns of child support after delivery, less active coping, less positive affect, and more negative affect. High resilient women reported more social support.
5	Scheidell et al. (2018)	Socioeconomic vulnerability and sexually transmitted infection among pregnant Haitian women	Haiti [Maternal health]	Access to resources [Who has what?]	The study findings highlighted how socioeconomic vulnerabilities, such as low levels of women's education, increase their risk of early sexual debut.

S/N	Author Year	Publication Title	Country [Context]	Main-Theme	Main Findings
6	Kaye et al. (2014)	Survivors' understanding of vulnerability and resilience to maternal near-miss obstetric events in Uganda	Uganda [Maternal health]	Division of labour [Who does what?]	The study highlighted how women's need to balance economic activities and reproduction increased their vulnerability and ability to recover from obstetric complications. In such circumstances, social networks or social capital was generally perceived as an essential component of women's resilience because it provided women with the financial, material, and emotional assistance, including those related to household responsibilities, such as childcare.
7	Johnson et al. (2010)	Orphanhood and vulnerability: a conduit to poor child health outcomes in Rwanda	Rwanda [Child health]	Access to resources [Who has what?]	The study showed that the orphan and vulnerable children (OVC) status directly and indirectly influenced the risk of childhood morbidity (e.g., diarrhoea, fever, and acute respiratory infection [ARI] symptoms). This is because OVCs were more likely to be found in households headed by older adults (40 years old) and females, where the mother/carer has inadequate access to socioeconomic infrastructures (e.g. inadequate education), and that are more likely to be in urban areas.
8	Gaillard et al. (2002)	Vulnerability of women in an African setting: lessons for mother-to-child HIV transmission prevention programmes	Kenya [Child health]	Power negotiation [How is power enacted, negotiated, or challenged?]	Women were generally reluctant about disclosing their HIV status to their partner. Violence against women living with HIV was reported in this study. This was primarily linked to women's disclosure of their seropositivity status. Men were often not in favour of having their wives tested, fearing the indirect disclosure of their own infection. The authors recommended couple-counselling and partner involvement in mother-to-child-transmission (MTCT) prevention programmes, as only testing women can increase their susceptibility to violence despite careful counselling. Partner support is crucial for preventing MTCT, especially because this requires mothers to use antiretroviral therapy and feed the child using formula feeding.

S/N	Author Year	Publication Title	Country [Context]	Main-Theme	Main Findings
9	Zakayo et al. (2020)	Vulnerability and agency across treatment-seeking journeys for acutely ill children: how family members navigate complex healthcare before, during and after hospitalisation in a rural Kenyan setting	Kenya [Child health]	Access to resources [Who has what?]	The lack of rapid access to money was an important contributing factor to a child's deteriorating condition; it influenced the initiation of a treatment-seeking. For instance, women made many references to "waiting to talk to my husband," "waiting to be sent money from my husband," and waiting for "his permission to pursue an action." Most primary caretakers, including mothers, were not income earners, and often relied heavily on their spouses or other household providers for money. At times, this social support was reported to be a burden and to cause delays in care-seeking or create an additional financial burden for households.
10	Storeng et al. (2012)	Too poor to live? A case study of vulnerability and maternal mortality in Burkina Faso	Burkina Faso [Maternal health]	Social norms [How are values defined?]	The study highlighted how structural impediments, including those related to motherhood and childbearing, limit individual resilience. For example, the social importance of bearing a child as soon as possible [amidst being physically or mentally capable] and the stigma associated with childlessness contributed to the death of the study actor who, despite having access to skilled birth attendance and emergency obstetric care, could not be rescued from maternal mortality.
11	Den Hollander et al. (2018)	Power difference and risk perception: mapping vulnerability within the decision process of pregnant women towards clinical trial participation in an urban middle-income setting	Ghana [Maternal health]	Agency and decision-making [Who decides?]	This study found a wide power difference between health providers and patients, and a different perception of risk through externalisation of responsibility of risk management within a religious context and a context shaped by authority. Therapeutic misconception was also observed, leading women to rely on the opinion of the medical professional, rather than being guided by their own motivation to participate. On their decision to participate in clinical trials, four women said that they would want to consult their husbands (or other family members living with them) before agreeing to participate. They

S/N	Author Year	Publication Title	Country [Context]	Main-Theme	Main Findings
					claimed that they would not be able to say they would participate without discussing it with the partner.
12	Arcos et al. (2011)	Vulnerability of pregnant women living in a community of metropolitan Santiago	Chile [Maternal health]	Access to resources [Who has what?]	It was observed that reproductive, neonatal, and mental and family health deteriorates when women have precarious jobs, with pregnancies without social security, informal and insecure habitability of housing, low schooling, psychosocial risk, absence of a partner in the home, and high dependency ratio.
13	Reyes et al. (2020)	Intimate partner violence and postpartum emotional distress among South African women: moderating effects of resilience and vulnerability factors	South Africa [Maternal health]	Power negotiation [How is power enacted, negotiated, or challenged?]	Consistent with the research hypothesis, this study found that the strength of the association between exposure to intimate partner violence and postpartum distress was stronger for women who reported greater levels of distress during pregnancy and weaker for women with greater socioeconomic assets. Intimate partner exposure impacts were greater for women reporting higher levels of emotional distress during pregnancy suggests that IPV-exposed women may be particularly likely to experience persistence (vs. discontinuity) of poor mental health across the antenatal and postnatal periods.
14	Warren et al. (2018)	“Sickness of shame”: investigating challenges and resilience among women living with obstetric fistula in Kenya	Kenya [Maternal health]	Access to resources [Who has what?]	Most women affected by fistula had secondary education and a very low monthly income. It was reported that irrespective of marital status, having male support (e.g. husband, brother, or uncle), particularly financial support and help in securing transport to hospitals for repair care, was critical. This was seen especially in cases where facilities denied care to women who were unaccompanied by their husbands.
15	Lavender et al. (2020)	Journey of vulnerability: a mixed-methods study to understand intrapartum transfers in Tanzania and Zambia	Tanzania and Zambia [Maternal health]	Access to resources [Who has what?]	Women were usually not in control of their own finances, making them reliant on their partners for support; this made it difficult for them to be ready for emergency situations. Some women failed to attend clinics as a result of the lack of support from their husbands.

S/N	Author Year	Publication Title	Country [Context]	Main-Theme	Main Findings
					*Most husbands did not give their wives adequate funds for the needs during delivery.
16	Pourette (2012)	Haitian women living with HIV in Guadeloupe: pregnancy between medical issues, social benefits and administrative vulnerability: a qualitative study	Guadeloupe [Maternal health]	Access to resources [Who has what?]	In one interview with a respondent, the woman received financial support (e.g. rent fee, feeding) from her sexual partner.
17	Prates Cde et al. (2008)	Gender power, poverty and contraception: multiparous experiences	Brazil [Maternal health]	Agency and decision-making [Who decides?]	The authors illuminated the challenges of multiparity in poorest sections of the population, where precarious living conditions, including inequalities in gender power, resulted in reduced potential empowerment of women to organise their lives, including planning childbirth. *Male resistance to condom use appeared to be prominent in the testimonials. The decrease in this autonomy affected the reproductive life, reducing its power to control one's fertility, which resulted in multiparity.