

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/168901>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

SCHOTTKY SPACES AND UNIVERSAL MUMFORD CURVES OVER \mathbb{Z}

JÉRÔME POINEAU AND DANIELE TURCHETTI

ABSTRACT. For every integer $g \geq 1$ we define a universal Mumford curve of genus g in the framework of Berkovich spaces over \mathbb{Z} . This is achieved in two steps: first, we build an analytic space \mathcal{S}_g that parametrizes marked Schottky groups over all valued fields. We show that \mathcal{S}_g is an open, connected analytic space over \mathbb{Z} . Then, we prove that the Schottky uniformization of a given curve behaves well with respect to the topology of \mathcal{S}_g , both locally and globally. As a result, we can define the universal Mumford curve \mathcal{C}_g as a relative curve over \mathcal{S}_g such that every Schottky uniformized curve can be described as a fiber of a point in \mathcal{S}_g . We prove that the curve \mathcal{C}_g is itself uniformized by a universal Schottky group acting on the relative projective line $\mathbb{P}_{\mathcal{S}_g}^1$. Finally, we study the action of the group $\text{Out}(F_g)$ of outer automorphisms of the free group with g generators on \mathcal{S}_g , describing the quotient $\text{Out}(F_g) \backslash \mathcal{S}_g$ in the archimedean and non-archimedean cases. We apply this result to compare the non-archimedean Schottky space with constructions arising from geometric group theory and the theory of moduli spaces of tropical curves.

CONTENTS

1. Introduction	1
2. Berkovich spaces over \mathbb{Z}	7
3. Schottky groups	13
4. The Schottky space over \mathbb{Z}	20
5. Outer automorphisms and connectedness of \mathcal{S}_g	28
6. Schottky uniformization for families of curves	36
References	40

1. INTRODUCTION

The uniformization of Riemann surfaces is one of the most central results in the theory of analytic curves. Independently proven in 1907 by P. Koebe and H. Poincaré, this theorem is the culmination of almost a century of work in complex geometry¹, and states that the universal cover of a connected compact complex analytic curve of genus g is analytically isomorphic to:

- the projective line if $g = 0$;
- the affine line if $g = 1$;
- the open unit disc if $g \geq 2$.

This fact bears important consequences in numerous fields of mathematics, such as differential equations, number theory (special functions, modular forms, elliptic curves), Kleinian and Fuchsian group representations, and the theory of algebraic curves and their fundamental groups. After proving the aforementioned theorem, Koebe went on to show several related results that clarified

2020 *Mathematics Subject Classification*. 14H15, 14G22, 30F10, 32G15, 32P05.

Key words and phrases. Schottky group, Schottky space, uniformization, Mumford curve, Berkovich space over \mathbb{Z} , outer space.

¹A complete account of the results and the mathematicians that made this breakthrough possible is given with detailed proofs in the impressive collective work [dSG10].

many aspects of uniformization theory. The most notorious one is the *retrosection theorem*, stating that every connected compact complex analytic curve is the quotient of an open dense subset $O \subset \mathbb{C}$ by the action of a free, finitely generated subgroup Γ of $\mathrm{PGL}_2(\mathbb{C})$ for which O is the region of discontinuity. Any group Γ arising in this way is called a *Schottky group*, and the resulting theory is usually referred to as *Schottky uniformization* of Riemann surfaces.

The quest for a non-archimedean analogue of uniformization theory is at the heart of the establishment of non-archimedean analytic geometry. In fact, J. Tate developed his theory of rigid analytic geometry to show that elliptic curves with split multiplicative reduction over a non-archimedean field k are always of the form $\mathbb{G}_m/q^{\mathbb{Z}}$ for some element $q \in k$ such that $0 < |q| < 1$. In the early '70s, D. Mumford had the intuition that Tate's uniformization could be extended to higher genus curves by building a non-archimedean theory of Schottky uniformization. In his celebrated paper [Mum72], he defined a notion of non-archimedean Schottky group and, using tools from formal geometry and Bruhat-Tits theory, showed that any such group acts on a suitable open subset of the projective line, and that the quotient identifies to (the analytification of) a projective curve. Finally, he characterized the curves that admit Schottky uniformization according to their reduction type. A further development of this theory in the context of rigid geometry was subsequently carried out by several authors, most notably by L. Gerritzen and M. van der Put in their book [GvdP80].

The non-archimedean theory is remarkable in many aspects, and particularly for the ways it addresses the issue of the lack of a nice topological structure: non-archimedean fields are totally disconnected, hence it is highly nontrivial even to define what an open dense subset of the projective line in this context should be. The solution of Tate's rigid geometry, in the late '50s, is to consider non-archimedean spaces with a Grothendieck topology instead of a classical topology. A more modern approach, developed by V. Berkovich in the late '80s, consists in defining non-archimedean analytic spaces as spaces of absolute values. Since they contain many points beyond the classical ones, they are not easily described explicitly, but they may be endowed with a structure of topological space in the classical sense. Given a non-archimedean field k , the group $\mathrm{PGL}_2(k)$ acts naturally on the Berkovich projective line $\mathbb{P}_k^{1,\mathrm{an}}$ and one can describe Schottky uniformization of Mumford curves in complete analogy with the complex case. A study of classical results on Schottky groups and Mumford curves in this new framework was initiated in Berkovich's first monograph (see [Ber90, §4.4]) and has been expanded in recent work by the authors [PT21], whose appendix contains a description of striking applications of these results.

In this paper, we are interested in the interplay between the archimedean and non-archimedean theories of Schottky uniformization. In order to construct a rigorous and coherent common framework for these two theories, we adopted the viewpoint of "Berkovich spaces over \mathbb{Z} ". Let us introduce it in a few words.

Although the theory of Berkovich analytic geometry was originally geared towards non-archimedean spaces, that is to say spaces over non-archimedean valued fields, it is worth noting that Berkovich's original definitions apply under less restrictive assumptions, and allow to make sense of analytic spaces over arbitrary Banach rings. For example, one may consider the base ring $(\mathbb{C}, |\cdot|_{\infty})$, where $|\cdot|_{\infty}$ denotes the usual absolute value. The Berkovich analytic spaces obtained in this way are nothing but the familiar complex analytic spaces.

Note that the field \mathbb{C} may be endowed with other absolute values. By considering a power $|\cdot|_{\infty}^{\varepsilon}$ of the usual absolute value, with $\varepsilon \in (0, 1]$, one still obtains a theory completely parallel to the complex analytic theory, but with a different normalization. Passing to the limit when ε tends to 0, one is led to consider the field \mathbb{C} endowed with the trivial absolute $|\cdot|_0$ (defined by $|a|_0 = 0$ if $a = 0$ and $|a|_0 = 1$ otherwise), which is a non-archimedean field, hence belongs to the realm of the usual theory of Berkovich spaces.

By using the theory of Berkovich spaces over Banach rings, one may fit all the preceding spaces into a common one. To be more precise, let us endow the field \mathbb{C} with the (non-multiplicative) hybrid norm $\|\cdot\|_{\text{hyb}} := \max(|\cdot|_0, |\cdot|_\infty)$. Its spectrum in the sense of Berkovich is $\mathcal{M}_{\text{hyb}} := \{|\cdot|_0\} \cup \{|\cdot|_\infty^\varepsilon, 0 < \varepsilon \leq 1\}$, and every Berkovich space over $(\mathbb{C}, \|\cdot\|_{\text{hyb}})$ admits a natural morphism pr_{hyb} to \mathcal{M}_{hyb} . As one may expect, the fibers $\text{pr}_{\text{hyb}}^{-1}(|\cdot|_\infty^\varepsilon)$ are complex analytic spaces, whereas the fiber $\text{pr}_{\text{hyb}}^{-1}(|\cdot|_0)$ is a non-archimedean Berkovich space. As a result, such a hybrid space witnesses complex analytic spaces converging towards a non-archimedean space. This rough idea actually leads to concrete results and has allowed to investigate precisely various properties of degeneration of families of complex spaces such as mixed Hodge structures [Ber09], volume forms [BJ17], equilibrium measures of endomorphisms [Fav20], etc. It also found a striking arithmetic application to uniform Manin-Mumford bounds for a family of genus 2 curves in [DKY20]. From this same point of view, in this paper, we will observe complex curves with a Schottky uniformization converging to non-archimedean curves with a Mumford uniformization.

It is possible to push further this line of thought and consider not only families of complex spaces and their non-archimedean limits (lying over a trivially valued field), but even p -adic spaces and, more generally, spaces over arbitrary valued fields. To this end, one starts with the base ring $(\mathbb{Z}, |\cdot|_\infty)$ and considers the space $\mathcal{M}(\mathbb{Z})$ of all absolute values (and more generally multiplicative seminorms) on \mathbb{Z} . Thanks to Ostrowski's theorem, up to a power, the latter are known to be exactly the usual one $|\cdot|_\infty$, the p -adic one $|\cdot|_p$, where p is a prime number, the trivial one $|\cdot|_0$ and the multiplicative seminorm $|\cdot|_{p,0}$ induced by the trivial absolute value on \mathbb{F}_p , where p is a prime number and \mathbb{F}_p is the finite field with p elements. By the same reasoning as in the hybrid case, Berkovich analytic spaces over \mathbb{Z} are global objects that naturally admit a morphism $\text{pr}_{\mathbb{Z}}$ to $\mathcal{M}(\mathbb{Z})$. The fibers of $\text{pr}_{\mathbb{Z}}$ may either be complex analytic spaces, p -adic analytic spaces, or spaces over trivially valued fields. This is the framework we will use to carry out our study of analytic uniformization of curves in a uniform way. It will allow us to build a space parametrizing all uniformizable curves (or, equivalently, Schottky groups) over all possible valued fields: archimedean or not, of arbitrary characteristic and arbitrary residue characteristic.

The foundations of the theory of analytic spaces over Banach rings were laid by Berkovich at the beginning of his manuscript [Ber90], but he soon switched more specifically to non-archimedean spaces. Over base rings of a special type (including, among others, \mathbb{Z} , rings of integers of numbers fields, and the hybrid \mathbb{C}), the theory was then further developed by the first-named author in [Poi10a] (case of the affine line), [Poi13] (local algebraic properties such as Noetherianity of the stalks or coherence of the structure sheaf) and [LP20] with Th. Lemanissier (definition of the category of analytic spaces, local path-connectedness of the spaces, cohomological vanishing on disks). As an example of application, let us mention that Berkovich spaces over \mathbb{Z} were used in [Poi10b] to give a geometric proof of a result of D. Harbater [Har88] solving the inverse Galois problem over a ring of convergent arithmetic power series (the subring of $\mathbb{Z}[[t]]$ consisting of power series that converge on the complex open unit disc).

Let us go back to the construction of a common framework for Schottky uniformization. Our first main new contribution in this direction is the definition of a moduli space \mathcal{S}_g of Schottky groups of rank g as a Berkovich space over \mathbb{Z} . This comes as a generalization of the *Schottky spaces* that have been extensively investigated both over the complex numbers and over non-archimedean fields. In order to simplify the notation, let us assume that $g \geq 2$ (the case $g = 1$ is analogous in many aspects, though much simpler). The complex Schottky space is defined by L. Bers in [Ber75] as a complex submanifold of \mathbb{C}^{3g-3} and parametrizes complex Schottky groups together with the choice of a basis up to conjugation by elements of $\text{PGL}_2(\mathbb{C})$. Its properties, generalizations, and various applications have been studied ever since, for example in [JMM79], [GH88], and [Her91]. In analogy with the complex case, Gerritzen [Ger81], [Ger82] gave a definition of Schottky space over a

non-archimedean field k as a rigid analytic subspace of the affine space of dimension $3g - 3$ over k , *i.e.* $\bar{k}^{3g-3}/\text{Gal}(\bar{k}/k)$, where \bar{k} is an algebraic closure of k . Both parametrizations are based on the notion of *Koebe coordinates*: a hyperbolic element γ of $\text{PGL}_2(\mathbb{C})$ or $\text{PGL}_2(k)$ is uniquely determined by the datum of its attracting fixed point α , its repelling fixed point α' , and its multiplier β , and, conversely, any ordered triplet (α, α', β) of elements of \mathbb{C} or k satisfying

$$\begin{cases} \alpha \neq \alpha' \\ 0 < |\beta| < 1 \end{cases}$$

gives rise to a hyperbolic element of $\text{PGL}_2(\mathbb{C})$ or $\text{PGL}_2(k)$. A Schottky group of rank g admits a basis consisting of g elements, which gives rise to $3g$ Koebe coordinates. Using a Möbius transformation to send the first three fixed points to $0, 1$ and ∞ , which amounts to some normalization, reduces the number to $3g - 3$ coordinates.

The space \mathcal{S}_g defined in this paper is a subset of an affine Berkovich space over \mathbb{Z} , namely $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$. Recall that each point x in $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$ naturally determines a complete valued field $\mathcal{H}(x)$, its *complete residue field*, as well as $3g - 3$ elements in it (obtained by evaluating the coordinate functions at x). The space \mathcal{S}_g then consists of the points x in $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$ whose associated $(3g - 3)$ -tuple of elements in $\mathcal{H}(x)$ corresponds, up to normalization, to the Koebe coordinates of a basis of a Schottky group of rank g in $\text{PGL}_2(\mathcal{H}(x))$ (see Definition 4.2.1 for more precision). This definition allows us to retrieve both the complex Schottky space and the Berkovich analogue of Gerritzen's space, as the complete residue fields $\mathcal{H}(x)$ may be complex, p -adic, etc. We first establish some topological properties of the Schottky space over \mathbb{Z} .

Theorem 1 (Theorems 4.3.4 and 5.2.1). *The space \mathcal{S}_g is open in $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$ and path-connected.*

As a result of its openness, the space \mathcal{S}_g inherits a structure of analytic space. In other words, the Mumford and Schottky uniformizations naturally fit together into a well-behaved family. Under this new perspective, Mumford's construction appears to be more than a mere analogue of the complex construction: both are particular instances of a more general theory.

The properties of the space \mathcal{S}_g allow to move continuously between the archimedean and the non-archimedean constructions, as in the case of hybrid spaces mentioned before. Similarly, there is a phenomenon of continuous degeneration from p -adic Schottky spaces to Schottky spaces in characteristic p . We believe that these features could be exploited to build interesting partial compactifications of Schottky spaces and to establish analogies between the theory of Mumford curves in mixed characteristic and in positive characteristic.

We then turn to the question of the connection between the space \mathcal{S}_g and the moduli space of curves. Let $\text{Out}(F_g)$ be the group of outer automorphisms of the free group with g generators. There is a natural action of $\text{Out}(F_g)$ on \mathcal{S}_g , whose orbits consist of unmarked Schottky groups (*i.e.* with no chosen basis). Since a Mumford curve over a non-archimedean field k determines a unique conjugacy class of Schottky groups, it is natural to consider the quotient by this action of the non-archimedean part of the Schottky space as a global space of Mumford curves. In order to do so, we let $\mathcal{S}_g^{\text{na}}$ be the set of non-archimedean points of \mathcal{S}_g . We prove the following result.

Theorem 2 (Corollaries 5.3.7 and 5.3.8). *The action of $\text{Out}(F_g)$ on $\mathcal{S}_g^{\text{na}}$ is proper and has finite stabilizers. For each $a \in \mathcal{M}(\mathbb{Z})$, the quotient space $\text{Mumf}_{g,a} := \text{Out}(F_g) \backslash (\mathcal{S}_g \cap \text{pr}_{\mathbb{Z}}^{-1}(a))$ inherits a structure of $\mathcal{H}(a)$ -analytic space.*

The action of $\text{Out}(F_g)$ is proper also over the space \mathcal{S}_g^{a} of archimedean points of \mathcal{S}_g . The proof of this fact boils down to a globalized version of the analogous result over the complex numbers, which is already known and can be proven using Teichmüller theory, once one provides the connection between the Schottky space and the Teichmüller space (see [HS07] for a detailed discussion of this

connection). In the non-archimedean framework, our proof is inspired by the work of Gerritzen [Ger82], and consists of applying Serre’s theory of free groups acting on trees [Ser77, §3] to the case of Schottky groups acting on the Berkovich projective line. This strategy is not easily adapted to the archimedean case, and we are not able to prove the properness of the action of $\text{Out}(F_g)$ on the entire space \mathcal{S}_g . This would be a first step in providing the global quotient $\text{Out}(F_g)\backslash\mathcal{S}_g$ with a structure of an analytic space over \mathbb{Z} , a result that we believe to hold true.

Even without a \mathbb{Z} -analytic structure on the global space of uniformizable curves, the theory of Schottky spaces over \mathbb{Z} allows us to construct a *universal* Mumford curve, in the weak sense of a relative curve over \mathcal{S}_g that encodes all possible archimedean and non-archimedean uniformizations at once, even though the same curve can appear more than once.

Theorem 3. *There is a smooth proper morphism of analytic spaces over \mathbb{Z}*

$$\mathcal{C}_g \longrightarrow \mathcal{S}_g$$

that is of relative dimension 1 and satisfies the following: given a point $x \in \mathcal{S}_g$, its preimage in \mathcal{C}_g is a curve over $\mathcal{H}(x)$ uniformized by the Schottky group Γ_x .

It is important to note that the uniformization property which lies at the basis of the theories of Schottky and Mumford carries over to our global setting. Indeed, the universal Mumford curve \mathcal{C}_g can be uniformized by an open subset of the relative projective line over \mathcal{S}_g . The precise statement is as follows.

Theorem 4 (Corollary 6.1.3). *There exists an open subset Ω_g of $\mathbb{P}_{\mathcal{S}_g}^1$ and a morphism $\Omega_g \rightarrow \mathcal{C}_g$ over \mathcal{S}_g that is a local isomorphism. Over each point x in \mathcal{S}_g , the restriction of the morphism gives back the uniformization by the Schottky group Γ_x .*

To contextualize our result, let us mention a different construction of families of Mumford curves over \mathbb{Z} , due to T. Ichikawa. In [Ich00], he introduces objects called *generalized Tate curves*: stable curves defined over a \mathbb{Z} -algebra mixing polynomials and power series. A typical example is the curve C_0 , which, by specialization on the base, gives back Mumford curves whose reduction graph is a rose with g petals. It is defined over the ring $R_0 := \mathbb{Z}[\underline{x}, \prod_{i \neq j} \frac{1}{x_i - x_j}][[\underline{y}]]$, where the multivariables \underline{x} and \underline{y} are deformation parameters related to Schottky uniformization. A similar curve C_Δ exists for each stable graph Δ of genus g and is defined over a ring R_Δ analogous to R_0 . Ichikawa’s construction makes no use of Schottky spaces and is rather a generalization of Mumford’s strategy to the case of a nonlocal base ring. More recently, in the preprint [Ich20b], which appeared online after the first version of the present work, Ichikawa has shown that the generalized Tate curves may be glued into a universal Mumford curve, defined over a formal scheme over \mathbb{Z} . Ichikawa’s approach has allowed him to study objects of arithmetic interest, such as Teichmüller modular forms [Ich18], periods of Mumford curves [Ich20a], and p -adic solutions of certain systems of partial differential equations, such as the Korteweg-deVries hierarchy [Ich01].

There are several advantages in our construction of universal Schottky uniformization over \mathbb{Z} . First of all, the curve \mathcal{C}_g admits a nice description as a global object: the Schottky space \mathcal{S}_g is a subspace of the affine space and therefore has canonical global coordinates. Moreover, the presence of the analytic topology provides a finer description of degenerations of families of Schottky groups, as well as a concrete way of studying natural group actions on the universal Mumford curve. Finally, Berkovich geometry brings out the connections between our construction and objects in neighboring theories such as tropical geometry and geometric group theory.

A convenient way to study these connections is the notion of *skeleton* of a non-archimedean analytic space, a combinatorial object that plays an important role in Berkovich’s theory and that can be interpreted as a “tropical shadow” of such a space. In the one-dimensional case, skeletons are

finite graphs that capture fundamental properties of the curve they represent.² They appear in this work as invariants that define strata in the Schottky space, bringing out connections with tropical moduli spaces as follows. Let us fix a non-archimedean point $a \in \mathcal{M}(\mathbb{Z})$, and recall that $\text{Mumf}_{g,a}$ is the space of Mumford curves over extensions of $\mathcal{H}(a)$, as in Theorem 2. Then, by assigning to each Mumford curve its skeleton, we can build a map of topological spaces

$$\psi : \text{Mumf}_{g,a} \rightarrow M_g^{\text{trop}},$$

where M_g^{trop} denotes the moduli space of tropical curves, parametrizing stable graphs of genus g with weights assigned at vertices. The image of the map ψ consists of unweighted metric graphs, and can be characterized as the quotient of the Culler-Vogtmann outer space CV'_g by the action of $\text{Out}(F_g)$. The outer space is a fundamental object of geometric group theory, and has been used to show many group theoretic properties of $\text{Out}(F_g)$ that were previously unknown. The reader can find more information in the original paper [CV86], as well as in [Vog14], which discusses more recent advances. The space CV'_g parametrizes graphs with an extra structure called a marking, and the action of the group $\text{Out}(F_g)$ is nontrivial only at the level of the marking, in a way that is reminiscent of the action of $\text{Out}(F_g)$ on \mathcal{S}_g . It is therefore a very natural problem to ask for a comparison between the two actions, which we are able to provide with the following result.

Theorem 5 (Theorem 6.2.2). *Let $a \in \mathcal{M}(\mathbb{Z})$ be a non-archimedean point and let $\mathcal{S}_{g,a} = \mathcal{S}_g \cap \text{pr}_{\mathbb{Z}}^{-1}(a)$ be the fiber of the Schottky space over a . Then, there is a continuous surjective function*

$$\phi : \mathcal{S}_{g,a} \longrightarrow CV'_g \times_{M_g^{\text{trop}}} \text{Mumf}_{g,a},$$

which is not injective for $g \geq 2$.

The interplay between non-archimedean Schottky spaces, tropical geometry, and the outer space is also the object of very recent work of M. Ulirsch. In [Uli21], which appeared online at the same time as the present work, he defines a non-archimedean analytic Deligne-Mumford stack $\overline{\mathcal{T}}_g$, over a fixed algebraically closed non-archimedean complete valued field K , which provides an analogue of Teichmüller space in non-archimedean analytic geometry. The construction proceeds in two steps. He first considers the logarithmic algebraic stack $\mathcal{T}_g^{\text{log}} = \mathcal{M}_g^{\text{log}} \times_{M_g^{\text{trop}}} \mathcal{T}_g^{\text{trop}}$, where $\mathcal{M}_g^{\text{log}}$ is the logarithmic algebraic stack of stable curves and $\mathcal{T}_g^{\text{trop}}$ is the tropical Teichmüller space defined in [CMV13]. The desired space $\overline{\mathcal{T}}_g$ is then obtained by taking the algebraic stack underlying $\mathcal{T}_g^{\text{log}}$, base changing it to the value group of K , and applying Raynaud's generic fiber functor.

The space $\overline{\mathcal{T}}_g$ parametrizes Berkovich stable curves over valued extensions of K , together with a choice of a basis of their topological fundamental group. It contains a natural locus of marked Mumford curves, which is related to the Schottky space $\mathcal{S}_g \times_{\mathbb{Z}} K$. Even though the techniques of the present paper significantly differ from those of [Uli21], we can describe the extent of this relation by considering the connections of the two constructions with the spaces M_g^{trop} and CV'_g . This is done in Remark 6.2.4, as well as in Remark 5 in the introduction of [Uli21].

Finally, let us note that other moduli spaces of arithmetic significance have been extensively studied, over both archimedean and non-archimedean valued fields. It seems natural to try to define a version over \mathbb{Z} of these objects, the closest to the present work being the moduli space $\text{Mumf}_{g,n}$ of pointed Mumford curves and the moduli space \mathcal{A}_g of principally polarized abelian varieties. While the case of $\text{Mumf}_{g,n}$ descends easily from the results already established by the authors, the space \mathcal{A}_g and a version of the Torelli map over \mathbb{Z} would require a bigger effort. We plan to address these issues and related applications in future work.

²For more details and a discussion of higher dimensional skeletons, we refer the reader to the excellent survey by A. Werner [Wer16].

The paper is structured as follows. Section 2 is a self-contained introduction to the affine and projective Berkovich spaces over the Banach ring $(\mathbb{Z}, |\cdot|_\infty)$, providing all the definitions and basic material on the subject needed for the rest of the paper. In Section 3, we formulate classical definitions and results of the theory of Schottky groups in a way that applies to any valued field, archimedean and non-archimedean alike. We adapt classical notions in the non-archimedean theory to the framework of Berkovich spaces, relying on previous work [PT21] by the authors. In Section 4, we define the space \mathcal{S}_g as a Berkovich space over \mathbb{Z} and we prove that it is open in $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$. In Section 5, we study the natural action of $\text{Out}(F_g)$ on \mathcal{S}_g : we establish some results that allow us to complete the proof of Theorem 1 and determine the properties of the quotient as stated in Theorem 2. Finally, in Section 6, we prove the universal uniformization theorem (Theorems 3 and 4) and a result clarifying the connections with the outer space and the moduli space of tropical curves (Theorem 5).

All the results proven in this paper remain valid if we replace $(\mathbb{Z}, |\cdot|_\infty)$ with the ring of integers of a number field.

Acknowledgements. We warmly thank Melody Chan, Frank Herrlich, Fabian Ruoff, Thibaud Lemanissier, Marco Maculan, and the anonymous referee for stimulating mathematical exchanges that improved a preliminary version of this paper. We are especially grateful to Sam Payne who raised to us the question answered by Theorem 6.2.2 and to Martin Ulirsch for sharing with us a preliminary version of his work [Uli21].

2. BERKOVICH SPACES OVER \mathbb{Z}

2.1. Analytic spaces over Banach rings. Let $(A, \|\cdot\|)$ be a Banach ring. In this section, we recall Berkovich's definition of analytic spaces over A (see [Ber90, Section 1.5]).

We start with the affine analytic space of dimension n over A , denoted by $\mathbb{A}_A^{n, \text{an}}$. It is a locally ringed space and we define it in three steps: underlying set, topology and structure sheaf.

The set underlying $\mathbb{A}_A^{n, \text{an}}$ is the set of bounded multiplicative seminorms on $A[T_1, \dots, T_n]$ that are bounded on A , *i.e.* the set of maps

$$|\cdot|: A[T_1, \dots, T_n] \longrightarrow \mathbb{R}_{\geq 0}$$

that satisfy the following properties:

- (i) $|0| = 0$ and $|1| = 1$;
- (ii) $\forall P, Q \in A[T_1, \dots, T_n], |P + Q| \leq |P| + |Q|$;
- (iii) $\forall P, Q \in A[T_1, \dots, T_n], |P + Q| = |P| |Q|$;
- (iv) $\forall a \in A, |a| \leq \|a\|$.

We set $\mathcal{M}(A) := \mathbb{A}_A^{0, \text{an}}$ and call it the *spectrum* of A . Note that we have a projection map $\text{pr}_A: \mathbb{A}_A^{n, \text{an}} \rightarrow \mathcal{M}(A)$ induced by the morphism $A \rightarrow A[T_1, \dots, T_n]$.

Let x be a point of $\mathbb{A}_A^{n, \text{an}}$. Denote by $|\cdot|_x$ the multiplicative seminorm associated to it. The ring $A[T_1, \dots, T_n]/\ker(|\cdot|_x)$ is a domain and we can consider its field of fractions. The seminorm $|\cdot|_x$ induces an absolute value on the later it and we can consider its completion, which we denote by $\mathcal{H}(x)$. We simply denote by $|\cdot|$ the absolute value on $\mathcal{H}(x)$ induced by $|\cdot|_x$ since no confusion may result.

We have a natural morphism $\chi_x: A[T_1, \dots, T_n] \rightarrow \mathcal{H}(x)$. For each $P \in A[T_1, \dots, T_n]$, we set $P(x) := \chi_x(P)$. Note that, by definition, we have $|P(x)| = |P|_x$.

The set $\mathbb{A}_A^{n, \text{an}}$ is endowed with the coarsest topology such that, for each $P \in A[T_1, \dots, T_n]$, the map

$$x \in \mathbb{A}_A^{n, \text{an}} \longmapsto |P(x)| \in \mathbb{R}_{\geq 0}$$

is continuous. The resulting topological space is Hausdorff and locally compact. The spectrum $\mathcal{M}(A)$ is even compact. The projection map pr_A is continuous.

For each open subset V of $\mathbb{A}_A^{n,\text{an}}$, we denote by S_V the set of elements of $A[T_1, \dots, T_n]$ that do not vanish on V and set $K(V) := S_V^{-1}A[T_1, \dots, T_n]$.

Let U be an open subset of $\mathbb{A}_A^{n,\text{an}}$. We define $\mathcal{O}(U)$ to be the set of maps

$$f: U \longrightarrow \bigsqcup_{x \in U} \mathcal{H}(x)$$

such that

- (i) for each $x \in U$, $f(x) \in \mathcal{H}(x)$;
- (ii) each $x \in U$ has an open neighborhood V on which f is a uniform limit of elements of $K(V)$.

One may now define arbitrary analytic spaces over A as locally ringed spaces that are locally isomorphic to some $(V(\mathcal{I}), \mathcal{O}_U/\mathcal{I})$, where U is an open subset of $\mathbb{A}_A^{n,\text{an}}$ and \mathcal{I} is a sheaf of ideals of \mathcal{O}_U .

A point x of an analytic space X over A is said to be archimedean or non-archimedean if the associated absolute value on $\mathcal{H}(x)$ is. We denote by X^{a} (resp. X^{na}) the set of archimedean (resp. non-archimedean) points of x and call it the archimedean (resp. non-archimedean) part of X . It is well-known that an absolute value on a field is archimedean if, and only if, its restriction to the prime field is. It follows that we have

$$X^{\text{a}} = \{x \in X : |2(x)| > 1\} \text{ and } X^{\text{na}} = \{x \in X : |2(x)| \leq 1\}$$

(and 2 could be replaced by any integer bigger than 1). In particular, the archimedean and non-archimedean parts of X are respectively open and closed subsets of X .

To go further, one should define the category of analytic spaces over A . When A is a complete non-archimedean valued field, this has been achieved by V. Berkovich in [Ber90, Ber93] (with a more general notion of analytic space). In [LP20, §2.1], Th. Lemanissier and the first-named author gave a definition over an arbitrary Banach ring. However, the category is shown to enjoy nice properties only under additional assumptions, for instance when A is a discrete valuation ring (with some mild extra hypotheses) or the ring of integers of a number field (see Section 2.3 for some definitions related to this setting). For future use, we note that, in those cases, fiber products exist (see [LP20, Théorème 4.3.8]).

2.2. Relative projective line. In the rest of the text, we will not only need affine spaces, but also projective spaces and, more precisely, relative projective lines over affine spaces. We explain here how to construct them in a down-to-earth way. Let $(A, \|\cdot\|)$ be a Banach ring. Let $n \in \mathbb{N}$ and denote by S the analytic space $\mathbb{A}_A^{n,\text{an}}$ with coordinates T_1, \dots, T_n .

Let U (resp. V) be the affine space $\mathbb{A}_A^{n+1,\text{an}}$ with coordinates T_1, \dots, T_n, Z (resp. T_1, \dots, T_n, Z') and denote by U_0 (resp. V_0) the open subset defined by the inequality $Z \neq 0$ (resp. $Z' \neq 0$). The morphism

$$\begin{array}{ccc} A[T_1, \dots, T_n, Z, Z^{-1}] & \longrightarrow & A[T_1, \dots, T_n, Z', Z'^{-1}] \\ T_i & \longmapsto & T_i \\ Z & \longmapsto & Z'^{-1} \end{array}$$

induces an isomorphism $U_0 \xrightarrow{\sim} V_0$.

We denote by \mathbb{P}_S^1 the analytic space obtained by glueing U and V along U_0 and V_0 via the previous isomorphism. It comes with a natural projection morphism $\pi: \mathbb{P}_S^1 \rightarrow S$. For any open subset S' of S , we denote by $\mathbb{P}_{S'}^1$ the analytic space $\pi^{-1}(S')$.

When $n = 0$, we will denote $\mathbb{P}_{\mathcal{M}(A)}^1$ by $\mathbb{P}_A^{1,\text{an}}$. Note that, for each $s \in S$, the fiber $\pi^{-1}(s)$ is identified with $\mathbb{P}_{\mathcal{H}(s)}^{1,\text{an}}$.

Let $M := \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathcal{O}(S))$. We may associate to it an endomorphism of \mathbb{P}_S^1 by the usual expression in coordinates

$$Z \mapsto \frac{aZ + b}{cZ + d}.$$

In this way, we get an action of $\mathrm{GL}_2(\mathcal{O}(S))$ on \mathbb{P}_S^1 . It factors through $\mathrm{PGL}_2(\mathcal{O}(S))$. The image of M in $\mathrm{PGL}_2(\mathcal{O}(S))$ will be denoted by $[M] = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Note that the action restricts to an action on each fiber of π , hence also on $\mathbb{P}_{S'}^1$, for any open subset S' of S .

2.3. Berkovich spaces over \mathbb{Z} . In this section, we consider the special case where $(A, \|\cdot\|) = (\mathbb{Z}, |\cdot|_\infty)$, where $|\cdot|_\infty$ denotes the usual absolute value. We refer to [Poi10a], and especially Section 3.1 there, for more details.

The spectrum $\mathcal{M}(\mathbb{Z})$ is easily described using Ostrowski's theorem. It contains the following points:

- a point a_0 , associated to the trivial absolute value $|\cdot|_0$, with residue field \mathbb{Q} ;
- for each $\varepsilon \in (0, 1]$, a point a_∞^ε associated to the absolute value $|\cdot|_\infty^\varepsilon$, with residue field \mathbb{R} ;
- for each prime number p and each $\varepsilon \in (0, +\infty)$, a point a_p^ε associated to the absolute value $|\cdot|_p^\varepsilon$, with residue field \mathbb{Q}_p ;
- for each prime number p , a point $a_{p,0}$ associated to the seminorm on \mathbb{Z} induced by the trivial absolute value on $\mathbb{Z}/p\mathbb{Z}$, with residue field $\mathbb{Z}/p\mathbb{Z}$.

We will sometimes drop the exponent 1 and write a_p and a_∞ instead of a_p^1 and a_∞^1 . The archimedean part of $\mathcal{M}(\mathbb{Z})$ is the open subset $\mathcal{M}(\mathbb{Z})^a = \{a_\infty^\varepsilon \mid \varepsilon \in (0, 1]\}$.

The topology of $\mathcal{M}(\mathbb{Z})$ is quite simple. First, the branches are all homeomorphic to segments: for each prime number p , the map

$$b_p: \eta \in [0, 1] \mapsto \begin{cases} a_{p,0} & \text{if } \eta = 0; \\ a_p^{-\log(\eta)} & \text{if } \eta \in (0, 1); \\ a_0 & \text{if } \eta = 1 \end{cases}$$

is a homeomorphism and the map

$$\beta_\infty: \varepsilon \in [0, 1] \mapsto \begin{cases} a_0 & \text{if } \varepsilon = 0; \\ a_\infty^\varepsilon & \text{if } \varepsilon \in (0, 1] \end{cases}$$

is a homeomorphism too. Moreover, a subset U of $\mathcal{M}(\mathbb{Z})$ containing a_0 is open if, and only if, the intersection of U with each $b_p([0, 1])$ and $\beta_\infty([0, 1])$ is open and only finitely many of those sets are not contained entirely in U . In other words, $\mathcal{M}(\mathbb{Z})$ is homeomorphic to the Alexandroff one-point compactification of the disjoint union of the $b_p([0, 1])$'s and $\beta_\infty([0, 1])$, the point at infinity being a_0 .

We will often think about an analytic space over \mathbb{Z} as a family of analytic spaces over the different valued fields associated to the points of $\mathcal{M}(\mathbb{Z})$. The spaces over \mathbb{Q}_p , \mathbb{Q} , $\mathbb{Z}/p\mathbb{Z}$ (the last two being endowed with the trivial absolute value) are then usual Berkovich spaces. Recall that the analytic spaces over \mathbb{R} in the sense of Berkovich are the quotients of the corresponding usual analytic spaces over \mathbb{C} by the complex conjugation.

Let us be more precise in the case of an affine space $\mathbb{A}_{\mathbb{R}, |\cdot|_\infty^\varepsilon}^{n, \mathrm{an}}$, for some $\varepsilon \in (0, 1]$. The complex conjugation induces an automorphism of \mathbb{C}^n given by

$$c: z = (z_1, \dots, z_n) \in \mathbb{C}^n \mapsto (\bar{z}_1, \dots, \bar{z}_n) \in \mathbb{C}^n$$




FIGURE 1. The analytic spectrum $\mathcal{M}(\mathbb{Z})$.

and we have a homeomorphism

$$\rho_{\varepsilon}: z \in \mathbb{C}^n / \langle c \rangle \mapsto v_{z,\varepsilon} \in \mathbb{A}_{\mathbb{R},|\cdot|_{\infty}}^{n,\text{an}},$$

where $v_{z,\varepsilon}: P(T) \in \mathbb{R}[T] \mapsto |P(z)|_{\infty}^{\varepsilon}$. It follows that all the archimedean fibers are the same. More precisely, the map

$$\Phi: (v, \varepsilon) \in \mathbb{A}_{\mathbb{R},|\cdot|_{\infty}}^{n,\text{an}} \times (0, 1] \mapsto \rho_{\varepsilon} \circ \rho_1^{-1}(v) \in (\mathbb{A}_{\mathbb{Z}}^{n,\text{an}})^{\text{a}}$$

is a homeomorphism. Note that $\Phi(v, \varepsilon)$ may also be defined explicitly as the seminorm defined by

$$\Phi(v, \varepsilon): P(T) \in \mathbb{R}[T] \mapsto |P(v)|^{\varepsilon}.$$

In particular, the seminorms v and $\Phi(v, \varepsilon)$ are equivalent.

As regards topology, analytic spaces over \mathbb{Z} are known to be locally path-connected thanks to [LP20, Théorème 7.2.17]. As one can expect, surprising phenomena occur when passing from archimedean to the non-archimedean part. We illustrate this by giving two examples of continuous sections of the projection $\text{pr}_{\mathbb{Z}}: \mathbb{A}_{\mathbb{Z}}^{1,\text{an}} \rightarrow \mathcal{M}(\mathbb{Z})$.

Example 2.3.1. Let α be an element of \mathbb{C} that is transcendental over \mathbb{Q} . For each $a \in \mathcal{M}(\mathbb{Z})^{\text{na}}$, denote by $\eta_{a,1}$ the Shilov boundary of the disc of center 0 and radius 1, *i.e.* the Gauß point, in the fiber $\text{pr}_{\mathbb{Z}}^{-1}(a)$. The map

$$\sigma: a \in \mathcal{M}(\mathbb{Z}) \mapsto \begin{cases} \eta_{a,1} & \text{if } a \text{ is non-archimedean;} \\ \rho_{\varepsilon}(\alpha) & \text{if } a = a_{\infty}^{\varepsilon} \text{ with } \varepsilon \in (0, 1]. \end{cases}$$

is a continuous section of $\text{pr}_{\mathbb{Z}}: \mathbb{A}_{\mathbb{Z}}^{1,\text{an}} \rightarrow \mathcal{M}(\mathbb{Z})$. For this it is enough to show that $\rho_{\varepsilon}(\alpha)$ tends to $\eta_{a_0,1}$ when ε goes to 0. Remark that the point $\eta_{a_0,1}$ corresponds to the trivial absolute value on $\mathbb{Z}[T]$ and that, for each $P \in \mathbb{Z}[T] - \{0\}$, we have

$$|P(\rho_{\varepsilon}(\alpha))| = |P(\alpha)|_{\infty}^{\varepsilon} \xrightarrow{\varepsilon \rightarrow 0} 1$$

since α is transcendental over \mathbb{Q} . The result follows.

Example 2.3.2. Let $r \in (0, 1)$. For each $a \in \mathcal{M}(\mathbb{Z})^{\text{na}}$, denote by $\eta_{a,r}$ the Shilov boundary of the disc of center 0 and radius r in the fiber $\text{pr}_{\mathbb{Z}}^{-1}(a)$. The map

$$\tau_r : a \in \mathcal{M}(\mathbb{Z}) \mapsto \begin{cases} \eta_{a,r} & \text{if } a \text{ is non-archimedean;} \\ \rho_\varepsilon(r^{1/\varepsilon}) & \text{if } a = a_\infty^\varepsilon \text{ with } \varepsilon \in (0, 1]. \end{cases}$$

is a continuous section of $\text{pr}_{\mathbb{Z}} : \mathbb{A}_{\mathbb{Z}}^{1,\text{an}} \rightarrow \mathcal{M}(\mathbb{Z})$. It is enough to show that $\rho_\varepsilon(r^{1/\varepsilon})$ tends to $\eta_{a_0,r}$ when ε goes to 0. This is clear since, for each $\varepsilon \in (0, 1]$, we have

$$|T(\rho_\varepsilon(r^{1/\varepsilon}))| = |r^{1/\varepsilon}|_\infty^\varepsilon = r$$

and $\eta_{a_0,r}$ is the only point of the fiber $\text{pr}_{\mathbb{Z}}^{-1}(a_0)$ where T has absolute value r .

One can build a similar theory replacing \mathbb{Z} by the ring of integers \mathcal{O}_K of a number field K . To be more precise, let us denote by Σ_K the set of complex embeddings of K up to conjugation and endow \mathcal{O}_K with the norm

$$\|\cdot\|_K := \max(|\sigma(\cdot)|_\infty, \sigma \in \Sigma_K).$$

Then, the spectrum $\mathcal{M}(\mathcal{O}_K)$ looks very similar to $\mathcal{M}(\mathbb{Z})$: it is a tree with one point associated to the trivial absolute value and, for each place of K , one branch emanating from it.

Remark that the restriction of seminorms induces a map $\mathcal{M}(\mathcal{O}_K) \rightarrow \mathcal{M}(\mathbb{Z})$, and more generally a map $\mathbb{A}_{\mathcal{O}_K}^{n,\text{an}} \rightarrow \mathbb{A}_{\mathbb{Z}}^{n,\text{an}}$. Those maps are continuous and open.

Note also that $\mathcal{M}(\mathcal{O}_K)$ is an analytic space over \mathbb{Z} in the sense of Section 2.1. In particular, by [LP20, Théorème 4.3.8], it makes sense to consider the fiber product of an analytic space over \mathbb{Z} by $\mathcal{M}(\mathcal{O}_K)$ over $\mathcal{M}(\mathbb{Z})$. We obtain canonical identifications $\mathbb{A}_{\mathcal{O}_K}^{n,\text{an}} = \mathbb{A}_{\mathbb{Z}}^{n,\text{an}} \times_{\mathcal{M}(\mathbb{Z})} \mathcal{M}(\mathcal{O}_K)$, $\mathbb{P}_{\mathcal{O}_K}^{1,\text{an}} = \mathbb{P}_{\mathbb{Z}}^{1,\text{an}} \times_{\mathcal{M}(\mathbb{Z})} \mathcal{M}(\mathcal{O}_K)$, etc.

2.4. Some useful inequalities. In this section, we fix a complete valued field $(k, |\cdot|)$, archimedean or not. We state a few results that will be useful later.

Lemma 2.4.1. *Let $a, b \in k$. We have*

$$|a + b| \leq \max(|2|, 1) \max(|a|, |b|).$$

If $|a| > \max(|2|, 1) |b|$, then we have

$$|a + b| \geq \frac{|a|}{\max(|2|, 1)}.$$

Proof. If $(k, |\cdot|)$ is non-archimedean, then $\max(|2|, 1) = 1$, and those inequalities are well-known.

Assume that $(k, |\cdot|)$ is archimedean. Then $(k, |\cdot|)$ embeds isometrically into $(\mathbb{C}, |\cdot|_\infty)$ for some $\varepsilon \in (0, 1]$ and it is enough to prove the result for the latter. In this case, we have $\max(|2|, 1) = 2^\varepsilon$. For any $a, b \in \mathbb{C}$, we have $|a + b|_\infty \leq 2 \max(|a|_\infty, |b|_\infty)$ and the first result follows by raising the inequality to the power ε .

The inequality applied to $a + b$ and $-b$ gives $|a| \leq |2| \max(|a + b|, |b|)$. As a consequence, if we have $|a| > |2| |b|$, we must have $|a| \leq |2| |a + b|$. \square

It will be useful to introduce a notation for discs. We consider here the Berkovich affine line $\mathbb{A}_k^{1,\text{an}}$ over k with coordinate T .

Notation 2.4.2. For $a \in k$ and $r \in \mathbb{R}_{>0}$, we set

$$D^+(a, r) := \{x \in \mathbb{A}_k^{1,\text{an}} \mid |T(x) - a| \leq r\},$$

$$D^-(a, r) := \{x \in \mathbb{A}_k^{1,\text{an}} \mid |T(x) - a| < r\}.$$

Lemma 2.4.3. *Let $a, b \in k$ and $\rho_a, \rho_b \in \mathbb{R}_{>0}$. If $|a - b| > \max(|2|, 1) \max(\rho_a, \rho_b)$, then the closed discs $D^+(a, \rho_a)$ and $D^+(b, \rho_b)$ are disjoint.*

If $|\cdot|$ is non-archimedean, then the closed discs $D^+(a, \rho_a)$ and $D^+(b, \rho_b)$ are disjoint if, and only if, $|a - b| > \max(\rho_a, \rho_b)$.

Proof. If there exists a point x in $D^+(a, \rho_a) \cap D^+(b, \rho_b)$, then we have $|T(x) - a| \leq \rho_a$ and $|T(x) - b| \leq \rho_b$ in $\mathcal{H}(x)$, hence

$$|a - b| = |(a - T(x)) + (T(x) - b)| \leq \max(|2|, 1) \max(\rho_a, \rho_b)$$

by Lemma 2.4.1. The first part of the result follows.

The converse implication in the non-archimedean setting is well-known. \square

Lemma 2.4.4. *Let $a, b \in k$. If $|a + b|^2 > \max(|4|, 1) |ab|$, then $|a| \neq |b|$.*

If $|\cdot|$ is non-archimedean, then we have $|a| \neq |b|$ if, and only if, $|ab| < |a + b|^2$.

Proof. If $|a| = |b|$, then, by Lemma 2.4.1, we have $|a + b| \leq \max(|2|, 1) |a|$, hence

$$|a + b|^2 \leq \max(|2|^2, 1) |a|^2 = \max(|4|, 1) |ab|.$$

The first part of the result follows.

Let us now assume that $|\cdot|$ is non-archimedean. Assume that $|a| \neq |b|$. Then, we have $|a + b| = \max(|a|, |b|) > \min(|a|, |b|)$, hence

$$|a + b|^2 = \max(|a|, |b|)^2 > \max(|a|, |b|) \min(|a|, |b|) = |a| |b|.$$

The converse implication follows directly from the first part of the statement. \square

2.5. Metric structure. In this section, we fix a complete non-archimedean valued field $(k, |\cdot|)$. In the following, we will often encounter the projective line $\mathbb{P}_k^{1, \text{an}}$ and we gather here a few metric properties.

First recall that $\mathbb{P}_k^{1, \text{an}}$ has the structure of a real tree (see [Duc, (3.4.20)]). In particular, for any two distinct points $x, y \in \mathbb{P}_k^{1, \text{an}}$, there exists a unique segment $[xy]$ joining x to y . Recall also that each segment consisting of points of type 2 or 3 carries a multiplicative length (or modulus) that is invariant under isomorphisms of $\mathbb{P}_k^{1, \text{an}}$, *i.e.* under Möbius transformations (see [Duc, (3.6.23)]). To define the length of such a segment I , one may proceed as follows.

Notation 2.5.1. For $a \in k$ and $r \in \mathbb{R}_{>0}$, we denote by $\eta_{a,r}$ the unique point in the Shilov boundary of the closed disc $D^+(a, r)$.

There exist a finite extension k' of k , a coordinate T on $\mathbb{P}_k^{1, \text{an}}$, $a \in k'$ and $r \leq s \in \mathbb{R}_{>0}$ such that I is the image of the segment $[\eta_{a,r}, \eta_{a,s}]$ by the projection map $\mathbb{P}_{k'}^{1, \text{an}} \rightarrow \mathbb{P}_k^{1, \text{an}}$. We then set

$$\ell(I) := \frac{s}{r} \in [1, +\infty).$$

It is independent of the choices made. It will be convenient to set $\ell(\emptyset) := 1$.

Lemma 2.5.2. *Let a, b, c, d be distinct points of $\mathbb{P}^1(k)$ and denote their cross-ratio by $[a, b; c, d]$.*

Set $I := [ab] \cap [cd]$. It is either a segment consisting of points of type 2 or 3 or the empty set. If I is a non-trivial segment and if going from a to b and from c to d induces the same orientation on I , then we set $\varepsilon := -1$. In all other cases, we set $\varepsilon := 1$.

Then, we have

$$|[a, b; c, d]| = \ell(I)^\varepsilon.$$

Proof. Since the cross-ratio is invariant under Möbius transformations, we may assume that $b = 1$, $c = 0$ and $d = \infty$. Assume that $|a| < 1$. Then $[ab] \cap [cd] = [\eta_{0,|a|}, \eta_{0,1}]$ and going from a to b and from c to d induces the same orientation on it, hence $\varepsilon = -1$. We have

$$\ell([ab] \cap [cd])^{-1} = |a| = |[a, b; c, d]|$$

as desired. The other cases are dealt with similarly. □

3. SCHOTTKY GROUPS

The notion of Schottky group is classical over \mathbb{C} (see [MSW15]) and even over a complete non-archimedean valued field (see [GvdP80]). The definitions, results and proofs that appear in this section are adaptations of the standard ones to a relative setting. We will start the section by describing a typical geometric situation giving rise to Schottky groups before stating the general definition (Definition 3.5.1).

We would also like to refer the reader to the recent text [PT21] by the authors, which provides a detailed exposition of the theory non-archimedean Schottky groups and Schottky uniformization in the setting of Berkovich spaces and fixes some gaps in the existing literature. Our treatment here follows this reference quite closely.

3.1. Geometric situation. Let S be an analytic space over a Banach ring (archimedean or not). As in Section 2.2, consider the analytic space \mathbb{P}_S^1 and the projection morphism $\pi: \mathbb{P}_S^1 \rightarrow S$. In this section, we describe geometric properties of the action of some groups of automorphisms of \mathbb{P}_S^1 . It follows the strategy of [GvdP80, I, 4.1], see also [PT21, §6.4.1].

Definition 3.1.1. Let $(\gamma_1, \dots, \gamma_g) \in \mathrm{PGL}_2(\mathcal{O}(S))^g$. Let $\mathcal{B} = (B^+(\gamma_i^\varepsilon), 1 \leq i \leq g, \varepsilon = \pm 1)$ be a family of closed subsets of \mathbb{P}_S^1 that are disjoint. For each $i \in \{1, \dots, g\}$ and $\varepsilon \in \{-1, 1\}$, set

$$B^-(\gamma_i^\varepsilon) := \gamma_i^\varepsilon(\mathbb{P}_S^1 - B^+(\gamma_i^{-\varepsilon})).$$

For each $s \in S$, $i \in \{1, \dots, g\}$, $\varepsilon \in \{-1, 1\}$ and $\sigma \in \{-, +\}$, set $B_s^\sigma(\gamma_i^\varepsilon) := B^\sigma(\gamma_i^\varepsilon) \cap \pi^{-1}(s)$.

We say that \mathcal{B} is a *Schottky figure* adapted to $(\gamma_1, \dots, \gamma_g)$ if, for each $s \in S$, $i \in \{1, \dots, g\}$ and $\varepsilon \in \{-1, 1\}$, $B_s^+(\gamma_i^\varepsilon)$ is a closed disc in $\pi^{-1}(s) \simeq \mathbb{P}_{\mathcal{H}(s)}^1$ and $B_s^-(\gamma_i^\varepsilon)$ is a maximal open disc inside it.

Remark 3.1.2. Let $\varphi \in \mathrm{PGL}_2(\mathcal{O}(S))$. If $\mathcal{B} = (B^+(\gamma_i^\varepsilon), 1 \leq i \leq g, \varepsilon = \pm 1)$ is a Schottky figure adapted to $(\gamma_1, \dots, \gamma_g)$, then $(\varphi(B^+(\gamma_i^\varepsilon)), 1 \leq i \leq g, \varepsilon = \pm 1)$ is a Schottky figure adapted to $(\varphi\gamma_1\varphi^{-1}, \dots, \varphi\gamma_g\varphi^{-1})$.

In this section, we assume that we are in the situation of Definition 3.1.1. For $\sigma \in \{-, +\}$, we set

$$F^\sigma := \mathbb{P}_S^1 - \bigcup_{\substack{1 \leq i \leq g \\ \varepsilon = \pm 1}} B^{-\sigma}(\gamma_i^\varepsilon).$$

Note that, for $\gamma_0 \in \{\gamma_1^{\pm 1}, \dots, \gamma_g^{\pm 1}\}$ and $\sigma \in \{-, +\}$, $B^\sigma(\gamma_0)$ is the unique disc among the $B^\sigma(\gamma)$'s containing $\gamma_0 F^\sigma$.

Set $\Delta := \{\gamma_1, \dots, \gamma_g\}$. Denote by F_g the free group over the alphabet Δ and by Γ the subgroup of $\mathrm{PGL}_2(\mathcal{O}(B))$ generated by Δ . We have a natural morphism $\varphi: F_g \rightarrow \Gamma$ sending each γ in Δ to γ . It induces an action of F_g on \mathbb{P}_S^1 .

We now define subsets of \mathbb{P}_S^1 associated to the elements of F_g . As usual, we will identify those elements with the words over the alphabet $\Delta^\pm := \{\gamma_1^{\pm 1}, \dots, \gamma_g^{\pm 1}\}$.

Notation 3.1.3. For a non-empty reduced word $w = w'\gamma$ over Δ^\pm and $\sigma \in \{-, +\}$, we set

$$B^\sigma(w) := w' B^\sigma(\gamma).$$

The following result is stated and proved in [PT21, Lemma 6.4.6].

Lemma 3.1.4. *Let u be a non-empty reduced word over Δ^\pm . Then we have $uF^+ \subseteq B^+(u)$.*

Let v be a non-empty reduced word over Δ^\pm . If there exists a word w over Δ^\pm such that $u = vw$, then we have $uF^+ \subseteq B^+(u) \subseteq B^+(v)$. If, moreover, $u \neq v$, then we have $B^+(u) \subseteq B^-(v)$.

Conversely, if we have $B^+(u) \subseteq B^+(v)$, then there exists a word w over Δ^\pm such that $u = vw$. \square

Corollary 3.1.5. *The morphism φ is an isomorphism and the group Γ is free on the generators $\gamma_1, \dots, \gamma_g$. \square*

As a consequence, we will now identify Γ with F_g and express the elements of Γ as words over the alphabet Δ^\pm . In particular, we will allow us to speak of the length of an element γ of Γ . We will denote it by $|\gamma|$.

Set

$$O_n := \bigcup_{|\gamma| \leq n} \gamma F^+.$$

Since the complement of F^+ is the disjoint union of the open disks $B^-(\gamma)$ with $\gamma \in \Delta^\pm$, it follows from the description of the action that, for each $n \geq 0$, we have

$$\mathbb{P}_S^1 - O_n = \bigsqcup_{|\gamma|=n+1} B^-(\gamma).$$

It follows from Lemma 3.1.4 that, for each $n \geq 0$, O_n is contained in the interior of O_{n+1} . We set

$$O := \bigcup_{n \geq 0} O_n = \bigcup_{\gamma \in \Gamma} \gamma F^+.$$

3.2. Over a valued field. Let $(k, |\cdot|)$ be a complete valued field. In this section, we will focus on the particular case $S = \mathcal{M}(k)$. In this setting, the material of this section is classical: see [MSW15, Project 4.5] and [GvdP80, I, 4.1.3] (or [PT21, §6.4.1]) in the archimedean and non-archimedean case respectively.

We still assume that we are in the situation of Definition 3.1.1. Set $\iota := \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \in \mathrm{PGL}_2(k)$. It corresponds to the map $z \mapsto 1/z$ on $\mathbb{P}_k^{1, \mathrm{an}}$. The first result follows from an explicit computation.

Lemma 3.2.1. *Let $\alpha \in k^*$ and $\rho \in [0, |\alpha|)$. Then, we have*

$$\iota D^+(\alpha, \rho) = \begin{cases} D^+\left(\frac{\bar{\alpha}}{|\alpha|^2 - \rho^2}, \frac{\rho}{|\alpha|^2 - \rho^2}\right) & \text{if } k \text{ is archimedean;} \\ D^+\left(\frac{1}{\alpha}, \frac{\rho}{|\alpha|^2}\right) & \text{otherwise.} \end{cases}$$

\square

Lemma 3.2.2. *Let $r > 0$ and let $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ in $\mathrm{PGL}_2(k)$ such that $\gamma D^+(0, r) \subseteq \mathbb{A}_k^{1, \mathrm{an}}$. Then, we have $|d| > r|c|$ and*

$$\gamma D^+(0, r) = \begin{cases} D^+\left(\frac{b\bar{d} - a\bar{c}r^2}{|d|^2 - |c|^2 r^2}, \frac{|ad - bc|r}{|d|^2 - |c|^2 r^2}\right) & \text{if } k \text{ is archimedean;} \\ D^+\left(\frac{b}{\bar{d}}, \frac{|ad - bc|r}{|d|^2}\right) & \text{otherwise.} \end{cases}$$

Proof. Let us first assume that $c = 0$. Then, we have $d \neq 0$, so the inequality $|d| > r|c|$ holds, and γ is affine with ratio a/d . The result follows.

Let us now assume that $c \neq 0$. In this case, we have $\gamma^{-1}(\infty) = -\frac{d}{c}$, which does not belong to $D(0, r)$ if, and only if, $|d| > r|c|$. Note that we have the following equality in $k(T)$:

$$\frac{aT + b}{cT + d} = \frac{a}{c} - \frac{ad - bc}{c^2} \frac{1}{T + \frac{d}{c}}.$$

By Lemma 3.2.1, there exist $\beta \in k$ and $\sigma > 0$ such that $\iota D^+(\frac{d}{c}, r) = D^+(\beta, \sigma)$. Then, we have $\gamma D^+(0, r) = D^+(\frac{a}{c} - \frac{ad-bc}{c^2} \beta, |\frac{ad-bc}{c^2}| \sigma)$ and the result follows from an explicit computation. \square

Lemma 3.2.3. *Let $D' \subseteq D$ be closed concentric discs in $\mathbb{A}_k^{1, \text{an}}$. Let $\gamma \in \text{PGL}_2(k)$ such that $\gamma D' \subseteq \gamma D \subseteq \mathbb{A}_k^{1, \text{an}}$. Then, we have*

$$\frac{\text{radius of } \gamma(D')}{\text{radius of } \gamma(D)} \leq \frac{\text{radius of } D'}{\text{radius of } D},$$

with an equality if k is non-archimedean.

Proof. Let p be the center of D and D' and let τ be the translation sending p to 0. Up to changing D into τD , D' into $\tau D'$, γ into $\gamma\tau^{-1}$ and γ' into $\gamma'\tau^{-1}$, we may assume that D and D' are centered at 0. The result then follows from Lemma 3.2.2. \square

Proposition 3.2.4. *Assume that $\infty \in F^-$. Then, there exist $R > 0$ and $c \in (0, 1)$ such that, for each $\gamma \in \Gamma - \{\text{id}\}$, $B^+(\gamma)$ is a closed disc of radius at most $Rc^{|\gamma|}$.*

Proof. Let $\delta, \delta' \in \Delta^\pm$. By Lemma 3.1.4, we have an inclusion of discs $B^+(\delta'\delta) \subseteq B^+(\delta')$. There exists $f_{\delta, \delta'} \in \text{PGL}_2(k)$ that sends those discs to concentric disks inside $\mathbb{A}_k^{1, \text{an}}$. In the non-archimedean case, the discs are already concentric, so one may take $f_{\delta, \delta'} = \text{id}$, while, in the archimedean case, there is some work to be done, for which we refer to [MSW15, Project 3.4]. Set

$$c_{\delta, \delta'} := \frac{\text{radius of } f_{\delta, \delta'}(B^+(\delta'\delta))}{\text{radius of } f_{\delta, \delta'}(B^+(\delta'))} \in (0, 1).$$

For each $\gamma \in \Gamma$ such that $\gamma\delta'$ is a reduced word, by Lemma 3.2.3, we have

$$\frac{\text{radius of } B^+(\gamma\delta'\delta)}{\text{radius of } B^+(\gamma\delta')} = \frac{\text{radius of } \gamma f_{\delta, \delta'}^{-1} f_{\delta, \delta'}(B^+(\delta'\delta))}{\text{radius of } \gamma f_{\delta, \delta'}^{-1} f_{\delta, \delta'}(B^+(\delta'))} \leq c_{\delta, \delta'}.$$

Set

$$R := \max(\{\text{radius of } B^+(\gamma) \mid \gamma \in \Delta^\pm\})$$

and

$$c := \max(\{c_{\gamma, \gamma'} \mid \gamma, \gamma' \in \Delta^\pm, \gamma' \neq \gamma^{-1}\}).$$

By induction, for each $\gamma \in \Gamma - \{\text{id}\}$, we have

$$\text{radius of } B^+(\gamma) \leq R c^{|\gamma|}.$$

\square

Let us mention an easy consequence of that result (see [PT21, Corollary 6.4.12] for details).

Corollary 3.2.5. *Every element of $\Gamma - \{\text{id}\}$ is loxodromic.* \square

We now investigate the set O .

Corollary 3.2.6. *Let $w = (w_n)_{n \geq 0}$ be a sequence of reduced words over Δ^\pm such that the associated sequence of discs $(B^+(w_n))_{n \geq 0}$ is strictly decreasing. Then, the intersection $\bigcap_{n \geq 0} B^+(w_n)$ is a single k -rational point p_w . Moreover, the discs $B^+(w_n)$ form a basis of neighborhoods of p_w in $\mathbb{P}_k^{1, \text{an}}$.*

Proof. We include here the idea of the proof and refer the interested reader to [PT21, Corollary 6.4.13] for the complete details (in the non-archimedean case, but the same arguments apply when k is archimedean).

One checks that it is enough to prove the result after extending the scalars to a finite extension k_0 of k . As a result, we may assume that $F^- \cap \mathbb{P}_k^{1,\text{an}}(k) \neq \emptyset$, and then, up to changing coordinates, that $\infty \in F^-$. By Proposition 3.2.4, the radius of $B^+(w_n)$ tends to 0 when n goes to ∞ , and the result follows. \square

Corollary 3.2.7. *The set O is dense in $\mathbb{P}_k^{1,\text{an}}$ and its complement is contained in $\mathbb{P}_k^{1,\text{an}}(k)$. \square*

3.3. Schottky uniformization. We return to the general case of Definition 3.1.1 with an arbitrary analytic space S . Here again, we follow [PT21, §6.4.1].

Definition 3.3.1. We say that a point $x \in \mathbb{P}_S^1$ is a *limit point* if there exist $x_0 \in \mathbb{P}_S^1$ and a sequence $(\gamma_n)_{n \geq 0}$ of distinct elements of Γ such that $\lim_{n \rightarrow \infty} \gamma_n(x_0) = x$.

The *limit set* L of Γ is the set of limit points of Γ .

Following [Bou71, III, §4, Définition 1], we say that the action of Γ on a subset E of \mathbb{P}_S^1 is *proper* if the map

$$\begin{aligned} \Gamma \times E &\longrightarrow E \times E \\ (\gamma, x) &\longmapsto (x, \gamma \cdot x) \end{aligned}$$

is proper, where Γ is endowed with the discrete topology. By [Bou71, III, §4, Proposition 7], it is equivalent to requiring that, for every $x, y \in E$, there exist neighborhoods U_x and U_y of x and y respectively such that the set $\{\gamma \in \Gamma : \gamma U_x \cap U_y \neq \emptyset\}$ is finite. By [Bou71, III, §4, Proposition 3], in this case, the quotient $\Gamma \backslash E$ is Hausdorff.

We denote by C the set of points $x \in \mathbb{P}_S^1$ that admit a neighborhood U_x satisfying $\{\gamma \in \Gamma : \gamma U_x \cap U_x \neq \emptyset\} = \{\text{id}\}$. Then C is an open subset of \mathbb{P}_S^1 and the quotient map $(\mathbb{P}_S^1 - C) \rightarrow \Gamma \backslash (\mathbb{P}_S^1 - C)$ is a local homeomorphism. In particular, the topological space $\Gamma \backslash (\mathbb{P}_S^1 - C)$ is naturally endowed with a structure of analytic space *via* this map.

Proposition 3.3.2. *We have $O = C = \mathbb{P}_S^1 - L$. Moreover, the action of Γ on O is free and proper and the quotient map $\Gamma \backslash O \rightarrow S$ is proper.*

Proof. The proof is the same as that of [PT21, Theorem 6.4.18]. We include it for the convenience of the reader.

Let $x \in L$. By definition, there exist $x_0 \in \mathbb{P}_S^1$ and a sequence $(\gamma_n)_{n \geq 0}$ of distinct elements of Γ such that $\lim_{n \rightarrow \infty} \gamma_n(x_0) = x$. Assume that $x \in F^+$. Since F^+ is contained in the interior of O_1 , there exists $N \geq 0$ such that $\gamma_N(x_0) \in O_1$, hence we may assume that $x_0 \in O_1$. Lemma 3.1.4 then leads to a contradiction. It follows that L does not meet F^+ , hence, by Γ -invariance, L is contained in $\mathbb{P}_S^1 - O$.

Let $y \in \mathbb{P}_S^1 - O$. By definition, there exists a sequence $(w_n)_{n \geq 0}$ of reduced words over Δ^\pm such that, for each $n \geq 0$, $|w_n| \geq n$ and $y \in B^-(w_n)$. Let $y_0 \in F^-$. By Lemma 3.1.4, for each $n \geq 0$, we have $w_n(y_0) \in B^-(w_n)$ and the sequence of discs $(B^+(w_n))_{n \geq 0}$ is strictly decreasing. By Corollary 3.2.6, $(w_n(y_0))_{n \geq 0}$ tends to y , hence $y \in L$. It follows that $\mathbb{P}_S^1 - O = L$.

Set

$$U := F^+ \cup \bigcup_{\gamma \in \Delta^\pm} \gamma F^- = \mathbb{P}_S^1 - \bigsqcup_{|\gamma|=2} B^+(\gamma).$$

It is an open subset of \mathbb{P}_S^1 and it follows from the properties of the action (see Lemma 3.1.4) that we have $\{\gamma \in \Gamma \mid \gamma U \cap U \neq \emptyset\} = \{\text{id}\} \cup \Delta^\pm$. Using the fact that the stabilizers of the points of U are trivial, we deduce that $U \subseteq C$. Letting Γ act, it follows that $O \subseteq C$. Since no limit point may belong to C , we deduce that this is actually an equality.

We have already seen that the action is free on O . Let us prove that it is proper. Let $x, y \in O$. There exists $n \geq 0$ such that x and y belong to the interior of O_n . By Lemma 3.1.4, the set $\{\gamma \in \Gamma \mid \gamma O_n \cap O_n \neq \emptyset\}$ is made of element of length at most $2n + 1$. In particular, it is finite. We deduce that the action of Γ on O is proper.

Let K be a compact subset of S . Since $\pi: \mathbb{P}_S^1 \rightarrow S$ is proper, $\pi^{-1}(K)$ is compact, hence its closed subset $F^+ \cap \pi^{-1}(K)$ is compact too. Since $F^+ \cap \pi^{-1}(K)$ contains a point of every orbit of every element of $\pi^{-1}(K)$, we deduce that $\Gamma \backslash (O \cap \pi^{-1}(K))$ is compact. \square

3.4. Koebe coordinates. Let $(k, |\cdot|)$ be a complete valued field and let γ be a loxodromic element of $\mathrm{PGL}_2(k)$. The eigenvalues of γ belong to a quadratic extension of k and have distinct absolute values. If k is archimedean, it follows immediately that they both belong to k , hence γ admits exactly two fixed points $\alpha, \alpha' \in \mathbb{P}^{1, \mathrm{an}}(k)$. If k is non-archimedean, then the result still holds by the same argument in characteristic different from 2 and, in general, as a consequence of Hensel's lemma (see [GvdP80, I, 1.4]).

We can choose α so that the associated eigenvalue has minimal absolute value. In this case, α and α' will be respectively the attracting and repelling fixed points of the Möbius transformation associated to γ . Denote by β the multiplier of γ , *i.e.* the ratio of the eigenvalues such that $|\beta| < 1$. For $\varepsilon \in \mathrm{PGL}_2(k)$ such that $\varepsilon(0) = \alpha$ and $\varepsilon(\infty) = \alpha'$, we have $\varepsilon^{-1}\gamma\varepsilon(z) = \beta z$. It follows that the parameters α, α' and β determine uniquely the transformation γ . They are called the *Koebe coordinates* of γ .

Conversely, given $(\alpha = [u: v], \alpha' = [u': v'], \beta) \in (\mathbb{P}_k^{1, \mathrm{an}})^3$ with $\alpha \neq \alpha'$ (*i.e.* $wv' \neq u'v$) and $0 < |\beta| < 1$, it is not difficult to determine explicitly the element of $\mathrm{PGL}_2(k)$ that has those Koebe coordinates. It is given by

$$M(\alpha, \alpha', \beta) = \begin{bmatrix} uv' - \beta u'v & (\beta - 1)uu' \\ (1 - \beta)vv' & \beta uv' - u'v \end{bmatrix} \in \mathrm{PGL}_2(k).$$

In the rest of the paper, we will sometimes abuse notation and allow ourselves to use $M(\alpha, \alpha', \beta)$ in different contexts, for example when α, α', β belong to a ring (provided the conditions $\alpha \neq \alpha'$ and $0 < |\beta| < 1$ are satisfied at each point of its spectrum). This should not cause any trouble.

The following interpretation of $|\beta|$ will be useful later.

Lemma 3.4.1. *Assume that k is non-archimedean. Let D be an open disc of $\mathbb{P}_k^{1, \mathrm{an}}$ containing α' and not α . Then $\gamma(D)$ is an open disc containing α' and the segment joining the boundary point of D to that of $\gamma(D)$ consists of points of type 2 or 3 and has length equal to $|\beta|^{-1}$.*

Proof. Möbius transformations preserve open discs, their boundary points, and the length of segments. Since $\varepsilon^{-1}\gamma\varepsilon(\varepsilon^{-1}(D)) = \varepsilon^{-1}(\gamma(D))$, it is enough to prove the result for $\varepsilon^{-1}\gamma\varepsilon$ and $\varepsilon^{-1}(D)$. In this case, it is clear. \square

We now check that the Koebe coordinates depend analytically on the entries of the corresponding matrix. In fact, this is true not only over a valued field, but even over \mathbb{Z} . To prove this, let us introduce some notation. Set

$$K_{\mathbb{Z}} := \{(\alpha, \alpha', \beta) \in (\mathbb{P}_{\mathbb{Z}}^{1, \mathrm{an}})^3 : \alpha \neq \alpha', 0 < |\beta| < 1\}.$$

It is an open subset of $(\mathbb{P}_{\mathbb{Z}}^{1, \mathrm{an}})^3$. We also consider $\mathbb{P}_{\mathbb{Z}}^{3, \mathrm{an}}$ and write its elements in coordinates in the form $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ instead of the usual $[a: b: c: d]$. Denote by $L_{\mathbb{Z}}$ the set of elements $x \in \mathbb{P}_{\mathbb{Z}}^{3, \mathrm{an}}$ such that the matrix

$$\begin{bmatrix} a(x) & b(x) \\ c(x) & d(x) \end{bmatrix} \in \mathrm{PGL}_2(\mathcal{H}(x))$$

is loxodromic.

Lemma 3.4.2. *The subset $L_{\mathbb{Z}}$ is open in $\mathbb{P}_{\mathbb{Z}}^{3,\text{an}}$.*

Proof. Let us first consider the archimedean part $L_{\mathbb{Z}}^{\text{a}}$ of $L_{\mathbb{Z}}$. By Section 2.3, it is enough to prove that its intersection with the fiber over the point a_{∞}^1 , corresponding to the usual absolute value, is open. This allows to translate the statement into a statement about $\mathbb{P}^3(\mathbb{C})$ (since the set is clearly stable by complex conjugation), where it is a consequence of the continuity of the roots of a (degree 2) polynomial.

Let us now handle the non-archimedean part $L_{\mathbb{Z}}^{\text{na}}$. By Lemma 2.4.4, we have

$$L_{\mathbb{Z}}^{\text{na}} = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in (\mathbb{P}_{\mathbb{Z}}^{3,\text{an}})^{\text{na}} : |ad - bc| < |a + d|^2 \right\}.$$

Let $x \in L_{\mathbb{Z}}^{\text{na}}$. There exists $r > 1$ such that $r|ad - bc| < |a + d|^2$. The open subset of $\mathbb{P}_{\mathbb{Z}}^{3,\text{an}}$ defined by the inequality

$$r \max(|4|, 1) |ad - bc| < |a + d|^2$$

contains x and sits inside $L_{\mathbb{Z}}$, by Lemma 2.4.4 again. The result follows. \square

Proposition 3.4.3. *The morphism*

$$M: (\alpha, \alpha', \beta) \in K_{\mathbb{Z}} \mapsto M(\alpha, \alpha', \beta) \in L_{\mathbb{Z}}$$

is an isomorphism of analytic spaces over \mathbb{Z} . Its inverse is the map that associates to a loxodromic matrix its Koebe coordinates.

Proof. The map M is clearly analytic and it follows from the discussion above that it is a bijection. Let us prove that its inverse is also analytic.

Let $m \in L_{\mathbb{Z}}$. We may work in an affine chart of $\mathbb{P}_{\mathbb{Z}}^{3,\text{an}}$ containing m and identify it to $\mathbb{A}_{\mathbb{Z}}^{3,\text{an}}$. As a result, we may assume that the coefficients $a(m), b(m), c(m), d(m)$ of m are well-defined. Denote by $\lambda(m)$ and $\lambda'(m)$ the eigenvalues of the matrix associated to m , chosen so that $|\lambda(m)| < |\lambda'(m)|$. Remark that the inequality on the absolute values implies that $(a + d)(m) \neq 0$. The elements $\lambda(m)$ and $\lambda'(m)$ are then the two roots of the characteristic polynomial of the matrix:

$$X^2 - (a + d)(m)X + (ad - bc)(m) = (a + d)^2(m) \left(Y^2 - Y + \frac{ad - bc}{(a + d)^2}(m) \right),$$

where $Y = \frac{1}{(a + d)(m)} X$.

Note that the polynomial $P(Y) := Y^2 - Y + (ad - bc)/(a + d)^2$ (which is actually well-defined on the whole $L_{\mathbb{Z}}$) has analytic coefficients. We claim that λ and λ' are analytic functions of m . If m is archimedean and the discriminant $\Delta(m)$ of $P(m)(Y)$ is not real, this follows from the fact that there exists a branch of the square-root that is analytic in a neighborhood of $\Delta(m)$. If m is archimedean and $\Delta(m)$ is real, then $\Delta(m) > 0$, since otherwise $\lambda(m)$ and $\lambda'(m)$ would be complex conjugates hence would have the same absolute value. The result then follows from the fact that there exists a branch of the square-root that is analytic in a neighborhood of $\Delta(m)$ and commutes with the complex conjugation.

Assume that m is non-archimedean. The stalk \mathcal{O}_m of the structure sheaf is a local ring (whose maximal ideal is the set of elements that vanish at m). Denote its residue field by $\kappa(m)$ and set

$$\kappa(m)^{\circ} := \{f \in \kappa(m) : |f(m)| \leq 1\}$$

and

$$\kappa(m)^{\circ\circ} := \{f \in \kappa(m) : |f(m)| < 1\}.$$

The set $\kappa(m)^{\circ}$ is a local ring with maximal ideal $\kappa(m)^{\circ\circ}$. We denote its residue field by $\tilde{\kappa}(m)$. By Lemma 2.4.4, the image of $P(Y)$ in $\kappa(m)[Y]$ has coefficients in $\kappa(m)^{\circ}$ and its reduction is $Y^2 - Y$.

The roots $\lambda(m)$ and $\lambda'(m)$ of $P(m)(Y)$ reduce respectively to the roots 0 and 1 of $Y^2 - Y$. By [Poi13, Corollaire 5.3] and [Poi10a, Corollaire 2.5.2], $\kappa(m)^\circ$ and \mathcal{O}_m are henselian, and it follows that λ and λ' are analytic in the neighborhood of m .

It is now clear that $\beta = \lambda/\lambda'$ is analytic in the neighborhood of m . Note that we can also recover α and α' from λ and λ' since they correspond to the associated eigenline. More precisely, we have $\alpha(m) = [b(m) : (\lambda - a)(m)]$ if $\lambda(m) \neq a(m)$ and $\alpha(m) = [(\lambda - d)(m) : c(m)]$ otherwise, and similarly for α' . It follows that α and α' are analytic in the neighborhood of m . \square

3.5. Group theory. Let $(k, |\cdot|)$ be a complete valued field. In this section, we give the general definition of Schottky group over k and explain how it relates to the geometric situation considered in Section 3.1. Here, we borrow from [PT21, §6.4.2 and §6.4.3].

Definition 3.5.1. A discrete subgroup Γ of $\mathrm{PGL}_2(k)$ is said to be a *Schottky group over k* if

- (i) it is free and finitely generated;
- (ii) all its non-trivial elements are loxodromic;
- (iii) there exists a non-empty Γ -invariant connected open subset of $\mathbb{P}_k^{1,\mathrm{an}}$ on which the action of Γ is free and proper (cf. Section 3.3).

A *marked Schottky group over k* is the datum of a Schottky group Γ over k together with a basis of Γ (as a free group).

Remark 3.5.2. The notion of Schottky group over k is left unchanged if one replaces the absolute value on k by an equivalent one.

Remark 3.5.3. Let $(k', |\cdot|')$ be an extension of $(k, |\cdot|)$. A subgroup Γ of $\mathrm{PGL}_2(k)$ is a Schottky group over k if, and only if, it is a Schottky group over k' .

Schottky groups arise naturally from Schottky figures, as the following proposition shows.

Proposition 3.5.4. *Let Γ be a discrete subgroup of $\mathrm{PGL}_2(k)$ generated by finitely many elements $\gamma_1, \dots, \gamma_g$. If there exists a Schottky figure adapted to $(\gamma_1, \dots, \gamma_g)$, then Γ is a Schottky group, and $(\gamma_1, \dots, \gamma_g)$ is a basis of it.*

Proof. The group Γ satisfies (i) by Corollary 3.1.5, (ii) by Corollary 3.2.5 and (iii) by Corollary 3.2.7. The fact that $(\gamma_1, \dots, \gamma_g)$ is a basis follows from Corollary 3.1.5 again. \square

Definition 3.5.5. Let Γ be a Schottky group of rank g . We say that a basis $B = (\gamma_1, \dots, \gamma_g)$ of Γ is a *Schottky basis* if there exists a Schottky figure that is adapted to it.

If k is archimedean, it is a classical result of Marden [Mar74a] that there exist Schottky groups with no Schottky bases. On the contrary, in the non-archimedean case, a theorem of Gerritzen ensures that Schottky bases always exist (see [Ger74, §2, Satz 1]). We rephrase it here using the notation of the present paper (see [PT21, Corollary 6.4.32] for a proof using Berkovich geometry). We first introduce some terminology.

Definition 3.5.6. Let $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{PGL}_2(k)$, with $c \neq 0$, be a loxodromic matrix and let $\lambda \in \mathbb{R}_{>0}$ be a positive real number. We call open and closed *twisted Ford discs* associated to (γ, λ) the sets

$$D_{(\gamma, \lambda)}^- := \left\{ z \in k \mid \lambda |\gamma'(z)| = \lambda \frac{|ad - bc|}{|cz + d|^2} > 1 \right\}$$

and

$$D_{(\gamma, \lambda)}^+ := \left\{ z \in k \mid \lambda |\gamma'(z)| = \lambda \frac{|ad - bc|}{|cz + d|^2} \geq 1 \right\}.$$

Lemma 3.5.7. Let $\alpha, \alpha', \beta \in k$ with $|\beta| < 1$ and let $\lambda \in \mathbb{R}_{>0}$. Set $\gamma = M(\alpha, \alpha', \beta) = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. The twisted Ford discs $D_{(\gamma, \lambda)}^-$ and $D_{(\gamma, \lambda)}^+$ have center

$$\frac{\alpha' - \beta\alpha}{1 - \beta} = -\frac{d}{c}$$

and radius

$$\rho = \frac{(\lambda|\beta|)^{1/2}|\alpha - \alpha'|}{|1 - \beta|} = \frac{(\lambda|ad - bc|)^{1/2}}{|c|}.$$

The twisted Ford discs $D_{(\gamma^{-1}, \lambda^{-1})}^-$ and $D_{(\gamma^{-1}, \lambda^{-1})}^+$ have center

$$\frac{\alpha - \beta\alpha'}{1 - \beta} = \frac{a}{c}$$

and radius $\rho' = \rho/\lambda$. □

Lemma 3.5.8. For every loxodromic $\gamma \in \mathrm{PGL}_2(k)$ that does not fix ∞ and every $\lambda \in \mathbb{R}_{>0}$, we have $\gamma(D_{(\gamma, \lambda)}^+) = \mathbb{P}_k^{1, \mathrm{an}} - D_{(\gamma^{-1}, \lambda^{-1})}^-$. □

Theorem 3.5.9 (Gerritzen). Assume that k is non-archimedean. Let Γ be a Schottky group over k whose limit set does not contain ∞ . Then, there exist a basis $(\delta_1, \dots, \delta_g)$ of Γ and positive real numbers $\lambda_1, \dots, \lambda_g \in \mathbb{R}_{>0}$ such that the family of twisted Ford discs $(D_{\delta_1, \lambda_1}^+, \dots, D_{\delta_g, \lambda_g}^+, D_{\delta_1^{-1}, \lambda_1^{-1}}^-, \dots, D_{\delta_g^{-1}, \lambda_g^{-1}}^-)$ is a Schottky figure adapted to $(\delta_1, \dots, \delta_g)$. □

Despite the different behaviour of Schottky groups in the archimedean and non-archimedean cases, we will see in the next section that a uniform construction of a parameter space over \mathbb{Z} comprising the two worlds still enjoys nice topological properties.

4. THE SCHOTTKY SPACE OVER \mathbb{Z}

In this section, we define a parameter space for marked Schottky groups of a given rank (where the marking is given by the choice of a basis, see Definition 3.5.1) over all complete valued fields, archimedean and non-archimedean. Already for Schottky groups of rank one, one gets an interesting construction, but most uniformization phenomena that are at the center of our interest become apparent only when the rank is at least two.

4.1. The space \mathcal{S}_1 .

Let $\Gamma = \langle \gamma \rangle$ be a Schottky group of rank one over a complete valued field $(k, |\cdot|)$. Then, γ is conjugated in $\mathrm{PGL}_2(k)$ to a unique matrix of the form $M(0, \infty, \beta)$ with $0 < |\beta| < 1$ (which corresponds to the multiplication by β as an endomorphism of $\mathbb{P}_k^{1, \mathrm{an}}$; see Section 3.4 for the notation).

Consider the affine line $\mathbb{A}_{\mathbb{Z}}^{1, \mathrm{an}}$ with coordinate Y and set

$$\mathcal{S}_1 := \{x \in \mathbb{A}_{\mathbb{Z}}^{1, \mathrm{an}} : 0 < |Y(x)| < 1\}.$$

With each point $x \in \mathcal{S}_1$, one can canonically associate a Schottky group of rank one

$$\Gamma_x := \langle M(0, \infty, Y(x)) \rangle \subset \mathrm{PGL}_2(\mathcal{H}(x)).$$

The condition imposed on \mathcal{S}_1 ensures that $M(0, \infty, Y(x))$ is a loxodromic transformation of $\mathbb{P}_{\mathcal{H}(x)}^{1, \mathrm{an}}$ having 0 as attracting point and ∞ as repelling point.

Given a Schottky group of rank one over \mathbb{Q}_p , we can retrieve it as $\langle M(0, \infty, Y(x)) \rangle$ for a unique $x \in \mathcal{S}_1$ with $\mathcal{H}(x) = \mathbb{Q}_p$. For a general valued field $(k, |\cdot|)$ and an element $\beta \in k$ such that $0 < |\beta| < 1$, the group generated by $M(0, \infty, \beta)$ can be retrieved as above from a point of $\mathcal{S}_1 \times_{\mathbb{Z}} k$. This can be seen as a consequence of Lemma 4.2.5 below in the special case where $g = 1$.

4.2. Construction of \mathcal{S}_g and equivalent definitions.

Fix $g \geq 2$. Consider the space $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$ and denote its coordinates by $X_3, \dots, X_g, X'_2, \dots, X'_g, Y_1, \dots, Y_g$. For notational convenience, set $X_1 := 0$, $X_2 := 1$ and $X'_1 := \infty$ (seen as morphisms from $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$ to $\mathbb{P}_{\mathbb{Z}}^{1, \text{an}}$). Denote by $\text{pr}_{\mathbb{Z}}: \mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}} \rightarrow \mathcal{M}(\mathbb{Z})$ the projection morphism. Let U_g be the open subset of $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$ defined by the inequalities

$$\begin{cases} 0 < |Y_i| < 1 \text{ for } 1 \leq i \leq g; \\ X_i^{\sigma_i} \neq X_j^{\sigma_j} \text{ for } i, j \in \{1, \dots, g\} \text{ and } \sigma_i, \sigma_j \in \{\emptyset, '\}. \end{cases}$$

For $i \in \{1, \dots, g\}$, consider the transformations

$$M_i := M(X_i, X'_i, Y_i) \in \text{PGL}_2(\mathcal{O}(U_g)).$$

Definition 4.2.1. The *Schottky space of rank g* over \mathbb{Z} , denoted by \mathcal{S}_g , is the set of points $x \in U_g$ such that the subgroup Γ_x of $\text{PGL}_2(\mathcal{H}(x))$ defined by

$$\langle M_1(x), M_2(x), \dots, M_g(x) \rangle$$

is a Schottky group of rank g .

Notation 4.2.2. Recall that a Schottky group over a complete valued field $(k, |\cdot|)$ gives rise to a unique k -analytic curve by means of the uniformization described in Section 3.3. Given $x \in \mathcal{S}_g$, we denote by Γ_x the marked Schottky group of ordered basis $(M_1(x), M_2(x), \dots, M_g(x))$, and by \mathcal{C}_x the $\mathcal{H}(x)$ -analytic curve obtained via Schottky uniformization by Γ_x .

In the non-archimedean case, the curve \mathcal{C}_x has semi-stable reduction, and results of Berkovich ([Ber90, Section 4.3]) then assert that the dual graph of the stable model of \mathcal{C}_x can be canonically realized as a subset of \mathcal{C}_x . Such a subset, denoted by Σ_x , is a graph of Betti number g , called the *skeleton* of \mathcal{C}_x . If $\mathcal{B} = (D_{i,\varepsilon}^+, 1 \leq i \leq g, \varepsilon = \pm 1)$ is a Schottky figure adapted to a Schottky basis of Γ_x and F^+ is the associated ‘‘closure of a fundamental domain’’ (see Definition 3.1.1 and following paragraph), then there is an isomorphism $\mathcal{C}_x \cong F^+/\Gamma_x$ and the skeleton Σ_x of the Mumford curve \mathcal{C}_x is obtained through pairwise identification, for every $i = 1, \dots, g$, of the points of the Shilov boundaries of $D_{i,1}^+, D_{i,-1}^+$ in the tree corresponding to the skeleton of F^+ (see Figure 2 for an example with $g = 2$). We refer the reader to [PT21, Theorem 6.4.18] for a proof of this fact.

If $k = \mathbb{C}$, it is a deep theorem that in general there does not exist a Schottky basis for Γ_x (See [Mar74b] for an abstract proof and [Yam91] for an explicit example), but an analogue of the set F^+ can be built by replacing the Schottky figure \mathcal{B} with a $2g$ -uple of Jordan curves $(C_{i,\varepsilon}, 1 \leq i \leq g, \varepsilon = \pm 1)$ in $\mathbb{P}_{\mathbb{C}}^1$ with disjoint interiors and such that $\gamma_i(C_{i,1}) = C_{i,-1}$ for every $i = 1, \dots, g$. More specifically, the existence of such a $2g$ -uple is the condition originally used to define complex Schottky groups. B. Maskit [Mas67] successively showed the equivalence between this definition and the group-theoretical Definition 3.5.1. In the context of complex Schottky groups, the closed set F^+ is the complement of the interiors of these Jordan curves. The quotient map $F^+ \rightarrow \mathcal{C}_x$ sends the Jordan curves $C_{1,1}, \dots, C_{g,1}$ into smooth simple non-intersecting curves $\alpha_1, \dots, \alpha_g$ inside \mathcal{C}_x . The complex description allows to easily treat the case $k = \mathbb{R}$: every real Schottky group can be seen as a complex Schottky group, and the corresponding uniformization gives rise to a curve defined over \mathbb{R} .

Note that the identification of $\mathcal{H}(x)$ with a valued extension of $\mathcal{H}(\text{pr}_{\mathbb{Z}}(x))$ is not canonical, and to different immersions $\mathcal{H}(\text{pr}_{\mathbb{Z}}(x)) \hookrightarrow \mathcal{H}(x)$ one associates different Schottky groups, yielding curves \mathcal{C}_x that might not be isomorphic. To lift the ambiguity from this situation one has then to consider the points of the base-change $\mathcal{S}_g \times_{\mathbb{Z}} \mathcal{H}(x)$, as made more precise in the following remark.




FIGURE 2. The domain F^+ of a Schottky figure for the Schottky group Γ_x in the non-Archimedean case is depicted on the left. The group Γ_x identifies the ends of the skeleton Σ_{F^+} , so that the corresponding Mumford curve (on the right) contains the finite graph Σ_x .

Remark 4.2.3. Let $(A, \|\cdot\|)$ be a Banach ring. Starting with $\mathcal{M}(A)$ instead of $\mathcal{M}(\mathbb{Z})$, one can define a Schottky space $\mathcal{S}_{g,A}$ over A . It is related to the Schottky space over \mathbb{Z} in the following way. If we denote by $\pi_A: \mathbb{A}_A^{3g-3, \text{an}} \rightarrow \mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$ the projection map, it follows from Remark 3.5.3 that we have $\mathcal{S}_{g,A} = \pi_A^{-1}(\mathcal{S}_g)$. In other words, assuming that the suitable categories and fiber products are defined, we have $\mathcal{S}_{g,A} = \mathcal{S}_g \times_{\mathcal{M}(\mathbb{Z})} \mathcal{M}(A)$. Moreover, the projection map π_A respects all the data: for each $x \in \mathcal{S}_{g,A}$, the group Γ_x is the image of $\Gamma_{\pi(x)}$ by the inclusion $\text{PGL}_2(\mathcal{H}(\pi(x))) \subseteq \text{PGL}_2(\mathcal{H}(x))$, the limit set of $\Gamma_{\pi(x)}$ in $\mathbb{P}_{\mathcal{H}(\pi(x))}^{1, \text{an}}$ is the preimage of the limit set of Γ_x in $\mathbb{P}_{\mathcal{H}(x)}^{1, \text{an}}$, and so on.

In the special case where $A = \mathbb{C}$, we obtain a subset $\mathcal{S}_{g,\mathbb{C}}$ of \mathbb{C}^{3g-3} . As one may expect, this is a classical object that has already been closely investigated. By [Hej75, Lemma 5.11], one has a covering map $\mathcal{T}_{g,\mathbb{C}} \rightarrow \mathcal{S}_{g,\mathbb{C}}$ from the Teichmüller space to the complex Schottky space. In particular, one deduces that $\mathcal{S}_{g,\mathbb{C}}$ is a connected subset of \mathbb{C}^{3g-3} . See also [Ber75, Proposition 2] for a direct proof.

Recall that, for $\varepsilon \in (0, 1]$, we denote by a_∞^ε the point of $\mathcal{M}(\mathbb{Z})$ associated with $|\cdot|_\infty^\varepsilon$ and that we have an isomorphism $\mathcal{H}(a_\infty^\varepsilon) \simeq \mathbb{R}$. We will use the canonical map $\rho_\varepsilon: \mathbb{C}^{3g-3} = \mathbb{A}_{\mathbb{C}}^{3g-3, \text{an}} \rightarrow \mathbb{A}_{\mathbb{R}}^{3g-3, \text{an}}$, where \mathbb{R} and \mathbb{C} are endowed with $|\cdot|_\infty^\varepsilon$. See Section 2.3 for details.

Lemma 4.2.4. *For $\varepsilon \in (0, 1]$, we have $\mathcal{S}_g \cap \text{pr}_{\mathbb{Z}}^{-1}(a_\infty^\varepsilon) = \rho_\varepsilon(\mathcal{S}_{g,\mathbb{C}})$. The set $\mathcal{S}_g \cap (\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}})^{\text{a}}$ is a connected open subset of $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$.*

Proof. By Remarks 3.5.2 and 3.5.3, $\rho_\varepsilon^{-1}(\mathcal{S}_g \cap \text{pr}_{\mathbb{Z}}^{-1}(a_\infty^\varepsilon))$ coincides with the usual complex Schottky space $\mathcal{S}_{g,\mathbb{C}}$. In other words, $\mathcal{S}_g \cap \text{pr}_{\mathbb{Z}}^{-1}(a_\infty^\varepsilon)$ is the quotient of $\mathcal{S}_{g,\mathbb{C}}$ by the complex conjugation. In particular, it is a connected open subset of $\mathbb{A}_{\mathbb{R}}^{3g-3, \text{an}}$.

Recall the homeomorphism

$$\Phi: \mathbb{A}_{\mathbb{R}}^{3g-3, \text{an}} \times (0, 1] \rightarrow (\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}})^{\text{a}}$$

from Section 2.3. It follows from Remark 3.5.2 that it induces a bijection between $(\mathcal{S}_g \cap \text{pr}_{\mathbb{Z}}^{-1}(a_\infty^\varepsilon)) \times (0, 1]$ and \mathcal{S}_g^{a} . As a consequence, \mathcal{S}_g^{a} is connected and open, since $(\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}})^{\text{a}}$ is open in $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$. \square

Let F_g be the free group of rank g with basis e_1, \dots, e_g . For each complete valued field k , we denote by $\text{Hom}_S(F_g, \text{PGL}_2(k))$ the set of group homomorphisms $\varphi: F_g \rightarrow \text{PGL}_2(k)$ that satisfy the following conditions:

- (i) $\varphi(e_1)$ is loxodromic with attracting fixed point 0 and repelling fixed point ∞ ;
- (ii) $\varphi(e_2)$ is loxodromic with attracting fixed point 1;
- (iii) the image of φ is a Schottky group of rank g .

Each point x of \mathcal{S}_g gives rise to an element φ_x of $\text{Hom}_S(F_g, \text{PGL}_2(\mathcal{H}(x)))$ that sends e_i to $M_i(x)$.

To state a converse result, we need to introduce an equivalence relation similar to that of [Ber90, Remark 1.2.2 (ii)]. We say that two elements $\varphi_1 \in \text{Hom}_S(F_g, \text{PGL}_2(k_1))$ and $\varphi_2 \in \text{Hom}_S(F_g, \text{PGL}_2(k_2))$ are equivalent if there exists an element $\varphi \in \text{Hom}_S(F_g, \text{PGL}_2(k))$ and isometric embeddings $k \hookrightarrow k_1$ and $k \hookrightarrow k_2$ that make the following diagram commute:

$$\begin{array}{ccc}
 & & \text{PGL}_2(k_1) \\
 & \nearrow^{\varphi_1} & \nearrow \\
 F_g & \xrightarrow{\varphi} & \text{PGL}_2(k) \\
 & \searrow_{\varphi_2} & \searrow \\
 & & \text{PGL}_2(k_2)
 \end{array}$$

We denote by $\text{Hom}_S(F_g, \text{PGL}_2)$ the set of classes of this equivalence relation.

Lemma 4.2.5. *The map $x \mapsto \varphi_x$ is a bijection between the underlying set of \mathcal{S}_g and $\text{Hom}_S(F_g, \text{PGL}_2)$.*

Proof. If $\varphi_x = \varphi_y$ then $\mathcal{H}(x) = \mathcal{H}(y)$ and $\varphi_x(F_g) = \varphi_y(F_g)$ coincide as marked Schottky groups in $\text{PGL}_2(\mathcal{H}(x))$. Hence $x = y$ in \mathcal{S}_g and $x \mapsto \varphi_x$ is injective. Conversely, for a valued field k and a map $\varphi \in \text{Hom}_S(F_g, \text{PGL}_2(k))$, we can consider the point $y \in \mathbb{A}_k^{3g-3, \text{an}}$ given by the Koebe coordinates of the marked Schottky group $\varphi(F_g)$. The image of y under the canonical projection $\mathbb{A}_k^{3g-3, \text{an}} \rightarrow \mathbb{A}_Z^{3g-3, \text{an}}$ is a point $x \in \mathcal{S}_g$ and there is an isometric embedding $\mathcal{H}(x) \hookrightarrow k$ realizing φ_x as canonical representative of the class of φ in $\text{Hom}_S(F_g, \text{PGL}_2)$. Hence $x \mapsto \varphi_x$ is surjective. \square

Remark 4.2.6. Every marked Schottky group of rank g over a complete valued field k is conjugated in $\text{PGL}_2(k)$ to a unique marked Schottky group with the property that its Koebe coordinates are of the form $\{(0, \infty, \beta_1), (1, \alpha'_2, \beta_2), \dots, (\alpha_g, \alpha'_g, \beta_g)\}$. The combination of this observation with Lemma 4.2.5 implies that every Schottky group over k can be retrieved, up to conjugation, as Γ_x for some suitable point $x \in \mathcal{S}_{g,k}$.

4.3. Openness of \mathcal{S}_g .

Definition 4.3.1. Let S be an analytic space. Consider the relative affine line \mathbb{A}_S^1 with coordinate Z .

For $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ in $\text{PGL}_2(\mathcal{O}(S))$ and $\lambda \in \mathbb{R}_{>0}$, set

$$D_{(\gamma, \lambda)}^+ := \{x \in \mathbb{A}_S^1 : |(cZ + d)(x)|^2 \leq \lambda |(ad - bc)(x)|\}$$

and

$$D_{(\gamma, \lambda)}^- := \{x \in \mathbb{A}_S^1 : |(cZ + d)(x)|^2 < \lambda |(ad - bc)(x)|\}.$$

We call such sets closed and open *relative twisted Ford discs* respectively.

We now generalize Gerritzen's theorem 3.5.9 to the relative setting.

Theorem 4.3.2. *Let x be a non-archimedean point of \mathcal{S}_g such that ∞ is not a limit point of Γ_x . There exist an open neighborhood W of x in U_g , an automorphism $\tau \in \text{Aut}(F_g)$ and positive real numbers $\lambda_1, \dots, \lambda_g \in \mathbb{R}_{>0}$ such that, denoting*

$$(N_1, \dots, N_g) := \tau \cdot (M_1, \dots, M_g) \in \text{GL}_2(\mathcal{O}(U_g))^g,$$

the family of relative twisted Ford discs over W

$$(D_{N_1, \lambda_1}^+, \dots, D_{N_g, \lambda_g}^+, D_{N_1^{-1}, \lambda_1^{-1}}^+, \dots, D_{N_g^{-1}, \lambda_g^{-1}}^+)$$

is a Schottky figure adapted to the family (N_1, \dots, N_g) of $\text{PGL}_2(\mathcal{O}(W))$.

Proof. By Theorem 3.5.9, we can find a basis $(\delta_1, \dots, \delta_g)$ of Γ_x and positive real numbers $\lambda_1, \dots, \lambda_g$ such that the family of twisted Ford discs $(D_{\delta_1, \lambda_1}^+, \dots, D_{\delta_g, \lambda_g}^+, D_{\delta_1^{-1}, \lambda_1^{-1}}^+, \dots, D_{\delta_g^{-1}, \lambda_g^{-1}}^+)$ is a Schottky figure adapted to $(\delta_1, \dots, \delta_g)$.

Denote by τ the automorphism of Γ_x sending $M_i(x)$ to δ_i . Identifying F_g with Γ_x by sending e_i to $M_i(x)$, we get an automorphism of F_g that we still denote by τ . Set

$$(N_1, \dots, N_g) := \tau \cdot (M_1, \dots, M_g) \in \text{GL}_2(\mathcal{O}(U_g))^g$$

and write

$$N_i = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix}$$

for $i \in \{1, \dots, g\}$. Note that the coefficients of the N_i 's are rational functions in the X_j 's, X_j' 's and Y_j 's. Denote by U' the open subset of U_g where they are all defined.

Let V be the open subset of U' defined by

$$|\text{tr}(N_i)|^2 > \max(|4|, 1) |\det(N_i)| \text{ for } 1 \leq i \leq g.$$

By Lemma 2.4.4, V contains x and, for each $y \in V$ and each $i \in \{1, \dots, g\}$, the matrix N_i is loxodromic.

Let $i \neq j \in \{1, \dots, g\}$. Since ∞ is not a limit point of Γ_x , it cannot be a fixed point of $N_i(x)$ or $N_j(x)$, hence $c_i(x)c_j(x) \neq 0$. There exists a neighborhood $W_{i,j}$ of x in V such that $c_i c_j$ does not vanish on $W_{i,j}$. In this case, for each $y \in W_{i,j}$, ∞ is not a fixed point of $N_i(y)$ or $N_j(y)$ and, by Lemma 3.5.7, we have

$$D_{N_i(y), \lambda_i}^+ = D^+ \left(-\frac{d_i}{c_i}(y), \left| \frac{a_i d_i - b_i c_i}{c_i^2}(y) \right|^{1/2} \lambda_i^{1/2} \right)$$

and

$$D_{N_j(y), \lambda_j}^+ = D^+ \left(-\frac{d_j}{c_j}(y), \left| \frac{a_j d_j - b_j c_j}{c_j^2}(y) \right|^{1/2} \lambda_j^{1/2} \right).$$

By assumption, the discs at x are disjoint and Lemma 2.4.3 ensures that we have

$$\left| \frac{d_i}{c_i}(x) - \frac{d_j}{c_j}(x) \right| > \max \left(\left| \frac{a_i d_i - b_i c_i}{c_i^2}(x) \right|^{1/2} \lambda_i^{1/2}, \left| \frac{a_j d_j - b_j c_j}{c_j^2}(x) \right|^{1/2} \lambda_j^{1/2} \right).$$

Since x is non-archimedean, we have $\max(|2(x)|, 1) = 1$, hence, up to shrinking $W_{i,j}$, we may assume that, for each $y \in W$, we have

$$\left| \frac{d_i}{c_i}(y) - \frac{d_j}{c_j}(y) \right| > \max(|2(y)|, 1) \max \left(\left| \frac{a_i d_i - b_i c_i}{c_i^2}(y) \right|^{1/2} \lambda_i^{1/2}, \left| \frac{a_j d_j - b_j c_j}{c_j^2}(y) \right|^{1/2} \lambda_j^{1/2} \right),$$

which implies that $D_{N_i(y), \lambda_i}^+$ and $D_{N_j(y), \lambda_j}^+$ are disjoint, by Lemma 2.4.3. Similar arguments show that, up to shrinking $W_{i,j}$, we may ensure that the discs $D_{N_i(y), \lambda_i}^+$, $D_{N_i^{-1}(y), \lambda_i^{-1}}^+$, $D_{N_j(y), \lambda_j}^+$ and $D_{N_j^{-1}(y), \lambda_j^{-1}}^+$ are all disjoint.

The result now holds with $W := \bigcap_{i \neq j} W_{i,j}$. \square

Corollary 4.3.3. *Let x be a non-archimedean point of \mathcal{S}_g . There exist an open neighborhood W of x in U_g , an automorphism τ of F_g and a family of closed subsets of \mathbb{P}_W^1 that is a Schottky figure adapted to $\tau \cdot (M_1, \dots, M_g)$.*

Proof. We will distinguish two cases.

- Assume that the extension $\mathcal{H}(x)/\mathcal{H}(\text{pr}_{\mathbb{Z}}(x))$ is finite.

Then, there exists an algebraic integer that does not belong to $\mathcal{H}(x)$, in the sense that there exists a polynomial $P \in \mathbb{Z}[T]$ with no roots in $\mathcal{H}(x)$. If $\text{pr}_{\mathbb{Z}}(x) = a_0$, so that $\mathcal{H}(\text{pr}_{\mathbb{Z}}(x)) = \mathbb{Q}$, we may moreover assume that P is totally real: all of its complex roots are real. Let K be a number field containing a root ω of P . Note that $\omega \in \mathcal{O}_K$.

We will work over the Schottky space $\mathcal{S}_{g, \mathcal{O}_K}$ over $\mathcal{M}(\mathcal{O}_K)$ defined in Remark 4.2.3. Denote by $p_K: \mathcal{S}_{g, \mathcal{O}_K} \rightarrow \mathcal{S}_g$ the projection morphism. Let $x_K \in p_K^{-1}(x)$ and set $U'_g := p_K^{-1}(U_g)$.

Let $A = \begin{pmatrix} 0 & 1 \\ 1 & -\omega \end{pmatrix} \in \text{GL}_2(\mathcal{O}(U'_g))$. We have $A(\omega) = \infty$. If $\text{pr}_{\mathbb{Z}}(x) = a_0$, then, for each $z \in (U'_g)^a$, we have $A(z) \in \text{GL}_2(\mathbb{R})$. For $i \in \{1, \dots, g\}$, set $M_{\infty, i} := A M_i A^{-1}$ in $\text{GL}_2(\mathcal{O}(U'_g))$. Denote by Γ_{∞, x_K} the subgroup of $\text{PGL}_2(\mathcal{H}(x_K))$ generated by $M_{\infty, 1}(x_K), \dots, M_{\infty, g}(x_K)$. By Corollary 3.2.7 and Remark 4.2.3, ω is not a limit point of Γ_{x_K} , hence ∞ is not a limit point of Γ_{∞, x_K} . By Theorem 4.3.2, there exists an open neighborhood W' of x_K in U'_g , an automorphism τ of F_g and $\lambda_1, \dots, \lambda_g \in \mathbb{R}_{>0}$ such that, denoting

$$(N_{\infty, 1}, \dots, N_{\infty, g}) := \tau \cdot (M_{\infty, 1}, \dots, M_{\infty, g}) \in \text{GL}_2(\mathcal{O}(U'_g))^g,$$

the family of twisted isometric discs

$$(D_{N_{\infty, 1}, \lambda_1}^+, \dots, D_{N_{\infty, g}, \lambda_g}^+, D_{N_{\infty, 1}^{-1}, \lambda_1^{-1}}^+, \dots, D_{N_{\infty, g}^{-1}, \lambda_g^{-1}}^+)$$

is a Schottky figure adapted to the family $(N_{\infty, 1}, \dots, N_{\infty, g})$ of $\text{PGL}_2(\mathcal{O}(W'))$. If $p(x) \neq a_0$, then x belongs to the interior of the non-archimedean part of \mathcal{S}_g and we may assume that $W' \subseteq \mathcal{S}_g^{\text{na}}$.

For each $i \in \{1, \dots, g\}$, set $N_i := A^{-1} N_{\infty, i} A$. Note that we have

$$(N_1, \dots, N_g) := \tau \cdot (M_1, \dots, M_g) \in \text{GL}_2(\mathcal{O}(U'_g))^g$$

and that the family $(A^{-1}(D_{N_{\infty, 1}, \lambda_1}^+), \dots, A^{-1}(D_{N_{\infty, g}, \lambda_g}^+), A^{-1}(D_{N_{\infty, 1}^{-1}, \lambda_1^{-1}}^+), \dots, A^{-1}(D_{N_{\infty, g}^{-1}, \lambda_g^{-1}}^+))$ is a Schottky figure adapted to the family (N_1, \dots, N_g) of $\text{PGL}_2(\mathcal{O}(W'))$ (see Remark 3.1.2). Set $W := p_K(W')$. By [LP20, Corollaire 5.6.4], it is an open subset of U_g .

Let us still denote by p_K the projection morphism $\mathbb{P}_{\mathcal{S}_{g, \mathcal{O}_K}}^1 \rightarrow \mathbb{P}_{\mathcal{S}_g}^1$. For $i \in \{1, \dots, g\}$ and $\varepsilon \in \{-1, 1\}$, set $B^+(N_i^\varepsilon) := p_K(A^{-1}(D_{N_{\infty, i}, \lambda_i^\varepsilon}^+))$. It is a closed subset of \mathbb{P}_W^1 and, by construction, for each $y' \in W'$, we have $B^+(N_i^\varepsilon) \cap \pi^{-1}(y) = p_K(A^{-1}(D_{N_{\infty, i}, \lambda_i^\varepsilon}^+) \cap \pi^{-1}(y'))$. To prove that the family $(B^+(N_i^\varepsilon), 1 \leq i \leq g, \varepsilon = \pm 1)$ is a Schottky figure adapted to the family (N_1, \dots, N_g) of $\text{PGL}_2(\mathcal{O}(W))$, it is enough to prove that for each $y \in W$, $i \in \{1, \dots, g\}$ and $\varepsilon \in \{-1, 1\}$, there exists a closed disc E_y in $\pi^{-1}(y)$ such that, for each $y' \in p_K^{-1}(y) \cap W'$, $A^{-1}(D_{N_{\infty, i}, \lambda_i^\varepsilon}^+) \cap \pi^{-1}(y') = p_K^{-1}(E_y)$.

Let $y \in W$, $i \in \{1, \dots, g\}$ and $\varepsilon \in \{-1, 1\}$. Let $y' \in p_K^{-1}(y) \cap W'$. Assume that y is non-archimedean. The set $A^{-1}(D_{N_{\infty, i}, \lambda_i^\varepsilon}^+) \cap \pi^{-1}(y')$ is an open disc over $\mathcal{H}(y')$ that contains a fixed point of $N_i(y')$. Since $N_i(y')$ is defined over $\mathcal{H}(y)$, its fixed points come from $\mathcal{H}(y)$ -rational points by base change to $\mathcal{H}(y')$ and we deduce that $A^{-1}(D_{N_{\infty, i}, \lambda_i^\varepsilon}^+) \cap \pi^{-1}(y')$ is the preimage by p_K of a closed

disc in $\pi^{-1}(y)$. Assume that y is archimedean. Note that, in this case, we have $\text{pr}_{\mathbb{Z}}(x) = a_0$. If $\mathcal{H}(y') = \mathcal{H}(y)$, then the result holds. Otherwise, we have $\mathcal{H}(y') = \mathbb{C}$ and $\mathcal{H}(y) = \mathbb{R}$ and the result follows from the fact that $A(y') \in \text{GL}_2(\mathbb{R})$.

- Assume that the extension $\mathcal{H}(x)/\mathcal{H}(\text{pr}_{\mathbb{Z}}(x))$ is infinite.

We deduce that the field $\mathcal{H}(x)$ is not locally compact, hence $\mathbb{P}^1(\mathcal{H}(x))$ is not compact. It follows from Corollary 3.2.7 that there exists a point $\omega \in \mathbb{P}^1(\mathcal{H}(x))$ that is not a limit point of Γ_x . Moreover, by definition, the image $\kappa(x)$ of $\text{Frac}(\mathcal{O}_x)$ in $\mathcal{H}(x)$ is dense. By Corollary 3.2.7 again, the limit set of Γ_x is closed, hence we may assume that ω belongs to $\mathbb{P}^1(\kappa(x))$.

If $\omega = \infty$, then the result follows directly from Theorem 4.3.2, so let us assume that this is not the case. Then, there exists an open neighborhood V of x in U_g and an element $\Omega \in \mathcal{O}(V)$ whose image in $\kappa(x)$ is ω . Let $A = \begin{pmatrix} 0 & 1 \\ 1 & -\Omega \end{pmatrix} \in \text{GL}_2(\mathcal{O}(V))$. By construction, we have $A(x)(\omega) = \infty$ in $\mathbb{P}^1(\mathcal{H}(x))$. The same arguments as in the proof of the first case apply (without having to worry about a base-change), and the result follows. \square

We have now collected all the results necessary to prove the openness of \mathcal{S}_g .

Theorem 4.3.4. *The Schottky space \mathcal{S}_g is an open subset of $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$.*

Proof. Let $x \in \mathcal{S}_g$. We want to prove that there exists an open subset of $\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$ containing x that is contained in \mathcal{S}_g . This follows from Lemma 4.2.4 when x is archimedean and from Corollary 4.3.3 and Proposition 3.5.4 when x is non-archimedean. \square

4.4. The space of Schottky bases. Let us assume $g \geq 2$ and fix a complete non-archimedean valued field $(k, |\cdot|)$. We denote by $\mathcal{SB}_{g,k}$ the subspace of $\mathcal{S}_{g,k}$ consisting of Schottky bases (see Remark 4.2.3 and Definition 3.5.5 for these notions).

Notation 4.4.1. Let A be a finite subset of $\mathbb{P}_k^{1, \text{an}}(k)$ with at least 2 elements. For each $\alpha \in A$, we denote by $D^-(\alpha, A)$ the biggest open disc with center α containing no other element of A and we denote by $p_{\alpha, A}$ its boundary point in $\mathbb{P}_k^{1, \text{an}}$.

Note that, if A contains at least 3 elements, then all the discs $D^-(\alpha, A)$ are disjoint.

Proposition 4.4.2. *Let $\alpha_1, \alpha'_1, \dots, \alpha_g, \alpha'_g$ be distinct elements of $\mathbb{P}_k^{1, \text{an}}(k)$. Set $A := \{\alpha_1, \alpha'_1, \dots, \alpha_g, \alpha'_g\}$. Let β_1, \dots, β_g be elements of k with absolute values in $(0, 1)$. For each $i \in \{1, \dots, g\}$, set $\gamma_i := M(\alpha_i, \alpha'_i, \beta_i) \in \text{PGL}_2(k)$. The following conditions are equivalent:*

- (i) *there exists a Schottky figure adapted to $(\gamma_1, \dots, \gamma_g)$;*
- (ii) *for each $i \in \{1, \dots, g\}$, we have $\ell([p_{\alpha_i, A}, p_{\alpha'_i, A}]) < |\beta_i|^{-1}$;*
- (iii) *for each $i, j, k \in \{1, \dots, g\}$ with $j \neq i, k \neq i$ and $\sigma_j, \sigma_k \in \{\emptyset, '\}$, we have*

$$|\beta_i| \cdot |[\alpha_j^{\sigma_j}, \alpha_k^{\sigma_k}; \alpha_i, \alpha'_i]| < 1.$$

Proof. (i) \implies (ii) Let $\mathcal{B} = (B^+(\gamma_i^\varepsilon), 1 \leq i \leq g, \varepsilon = \pm 1)$ be a Schottky figure adapted to $(\gamma_1, \dots, \gamma_g)$. Let $i \in \{1, \dots, g\}$. Note that we have $\alpha_i \in B^+(\gamma_i)$ and $\alpha'_i \in B^+(\gamma_i^{-1})$, hence $B^+(\gamma_i) \subset D^-(\alpha_i, A)$ and $B^+(\gamma_i^{-1}) \subset D^-(\alpha'_i, A)$. It follows that the segment between the boundary points of $D^-(\alpha_i, A)$ and $D^-(\alpha'_i, A)$ is strictly contained in the segment between the boundary points of $B^+(\gamma_i)$ and $B^+(\gamma_i^{-1})$. Lemma 3.4.1 then provides the desired inequality.

(ii) \implies (iii) Let $i, j, k \in \{1, \dots, g\}$ with $j \neq i, k \neq i$ and $\sigma_j, \sigma_k \in \{\emptyset, '\}$. If $[\alpha_j^{\sigma_j}, \alpha_k^{\sigma_k}] \cap [\alpha_i, \alpha'_i] = \emptyset$, then we have $|[\alpha_j^{\sigma_j}, \alpha_k^{\sigma_k}; \alpha_i, \alpha'_i]| = 1$ and the inequality of the statement holds.

Assume that $I := [\alpha_j^{\sigma_j}, \alpha_k^{\sigma_k}] \cap [\alpha_i, \alpha'_i] \neq \emptyset$. Since $\alpha_j^{\sigma_j}$ and $\alpha_k^{\sigma_k}$ do not belong to the discs $D^-(\alpha_i, A)$ and $D^-(\alpha'_i, A)$, the segment I must be contained in the segment joining the boundary points of

those two discs. It now follows from Lemma 2.5.2 that we have

$$\max(|[\alpha_j^{\sigma_j}, \alpha_k^{\sigma_k}; \alpha_i, \alpha'_i]|, |[\alpha_j^{\sigma_j}, \alpha_k^{\sigma_k}; \alpha_i, \alpha'_i]|^{-1}) = \ell(I) \leq \ell([p_{\alpha_i, A} p_{\alpha'_i, A}]) < |\beta_i|^{-1}.$$

(iii) \implies (i) Let $i \in \{1, \dots, g\}$. We will construct discs $B^+(\gamma_i)$ and $B^+(\gamma_i^{-1})$ that lie in $D^-(\alpha_i, A)$ and $D^-(\alpha'_i, A)$ respectively and such that $\gamma_i(\mathbb{P}_k^{1, \text{an}} - B^+(\gamma_i^{-1}))$ is a maximal open disc inside $B^-(\gamma_i)$ and $\gamma_i^{-1}(\mathbb{P}_k^{1, \text{an}} - B^+(\gamma_i))$ is a maximal open disc inside $B^-(\gamma_i^{-1})$. To do so, we may choose coordinates on $\mathbb{P}_k^{1, \text{an}}$ such that $\alpha_i = 0$ and $\alpha'_i = \infty$. The equalities of the statement then become

$$|\beta_i| \cdot \frac{\alpha_j^{\sigma_j}}{\alpha_k^{\sigma_k}} < 1$$

for $j, k \in \{1, \dots, g\}$ with $j \neq i$, $k \neq i$ and $\sigma_j, \sigma_k \in \{\emptyset, '\}$. It follows that there exists $r_i \in \mathbb{R}_{>0}$ such that

$$|\beta_i| \max(|\alpha_j^{\sigma_j}|, j \neq i, \sigma_j \in \{\emptyset, '\}) < r_i < \min(|\alpha_j^{\sigma_j}|, j \neq i, \sigma_j \in \{\emptyset, '\}).$$

The discs $B^+(\gamma_i) := D^+(0, r_i)$ and $B^+(\gamma_i^{-1}) := \mathbb{P}_k^{1, \text{an}} - D^-(0, |\beta_i|^{-1} r_i)$ then satisfy the required conditions.

The family of discs $(B^+(\gamma_i^\varepsilon), 1 \leq i \leq g, \varepsilon = \pm 1)$ is a Schottky figure adapted to $(\gamma_1, \dots, \gamma_g)$. \square

Corollary 4.4.3. *The topological space $\mathcal{SB}_{g,k}$ is path-connected.*

Proof. Let k' be a complete non-trivially valued extension of k . Denote by $\pi_{k'/k}: \mathbb{A}_{k'}^{3g-3, \text{an}} \rightarrow \mathbb{A}_k^{3g-3, \text{an}}$ the projection map. By Remark 4.2.3, we have $\mathcal{S}_{g,k'} = \pi_{k'/k}^{-1}(\mathcal{S}_{g,k})$ and, by Proposition 4.4.2, $\mathcal{SB}_{g,k'} = \pi_{k'/k}^{-1}(\mathcal{SB}_{g,k})$. Up to replacing k by k' , we may assume that k is not trivially valued.

We will consider the affine spaces $\mathbb{A}_k^{2g-3, \text{an}}$ with coordinates $X_3, \dots, X_g, X'_2, \dots, X'_g$ and $\mathbb{A}_k^{g, \text{an}}$ with the coordinates Y_1, \dots, Y_g . We denote by $\pi_1: \mathbb{A}_k^{3g-3, \text{an}} \rightarrow \mathbb{A}_k^{2g-3, \text{an}}$ and $\pi_2: \mathbb{A}_k^{3g-3, \text{an}} \rightarrow \mathbb{A}_k^{g, \text{an}}$ the corresponding projections.

Let $a, b \in \mathcal{SB}_{g,k}$. Let V be the open subset of $\mathbb{A}_k^{2g-3, \text{an}}$ consisting of the points all of whose coordinates are distinct. It is path-connected and contains $a_1 := \pi_1(a)$ and $b_1 := \pi_1(b)$. Let $\varphi: [0, 1] \rightarrow V$ be a continuous map such that $\varphi(0) = a_1$ and $\varphi(1) = b_1$. The continuous maps $|[X_j^{\sigma_j}, X_k^{\sigma_k}; X_i, X'_i]|$ for $i, j, k \in \{1, \dots, g\}$ with $j \neq i$, $k \neq i$ and $\sigma_j, \sigma_k \in \{\emptyset, '\}$ are all bounded on $\varphi([0, 1])$. Let $M \in \mathbb{R}_{>0}$ be a common upper bound. Since k is not trivially valued, there exists $\beta \in k^*$ such that

$$|\beta| < \min(M^{-1}, |Y_i(a)|, |Y_i(b)|, 1 \leq i \leq g).$$

Let us identify $\pi_1^{-1}(a_1)$ and $\mathbb{A}_{\mathcal{H}(a_1)}^{g, \text{an}}$, so that a may be seen as a point in the latter space. The point $\beta := (\beta, \dots, \beta)$ of $\mathbb{A}_k^{g, \text{an}}$ canonically lifts to a point a_β of $\mathbb{A}_{\mathcal{H}(a_1)}^{g, \text{an}}$ and there exists a continuous path from a to a_β in $\mathbb{A}_{\mathcal{H}(a_1)}^{g, \text{an}}$ along which all the $|Y_i|$'s are non-increasing and remain in $(0, 1)$. By Proposition 4.4.2, the corresponding path in $\mathbb{A}_k^{3g-3, \text{an}}$ stays in $\mathcal{SB}_{g,k}$. We similarly define a point b_β in $\pi_1^{-1}(b_1)$ and a continuous path from b to b_β in $\mathcal{SB}_{g,k}$.

To prove the result, it is now enough to construct a continuous path from a_β to b_β in $\mathcal{SB}_{g,k}$. Note that $\pi_2(a_\beta) = \pi_2(b_\beta) = \beta$, so that a_β and b_β identify to two points of the same fiber $\pi_2^{-1}(\beta) \simeq \mathbb{A}_k^{2g-3, \text{an}} = \mathbb{A}_k^{2g-3, \text{an}}$. We may now use the path defined by φ to go from a_β to b_β . By construction, it stays inside $\mathcal{SB}_{g,k}$. \square

Corollary 4.4.4. *The set $\mathcal{SB}_g^{\text{na}}$ is the subset of U_g^{na} described by the inequalities*

$$|Y_i| \cdot |[X_j^{\sigma_j}, X_k^{\sigma_k}; X_i, X'_i]| < 1$$

for all $i, j, k \in \{1, \dots, g\}$ with $j \neq i$, $k \neq i$ and $\sigma_j, \sigma_k \in \{\emptyset, '\}$. It is a path-connected open subset of $\mathcal{S}_g^{\text{na}}$.

Proof. The first part of the statement follows from Proposition 4.4.2. The fact that $\mathcal{SB}_g^{\text{na}}$ is open in $\mathcal{S}_g^{\text{na}}$ is an immediate consequence.

By Corollary 4.4.3, for each $z \in \mathcal{M}(\mathbb{Z})$, the fiber $\mathcal{SB}_g \cap \text{pr}_{\mathbb{Z}}^{-1}(z)$ is connected and contains the point P_z defined as the unique point in the Shilov boundary of the disc defined by the inequalities

$$\begin{cases} |X_i| \leq 1 \text{ for } 3 \leq i \leq g; \\ |X'_i| \leq 1 \text{ for } 2 \leq i \leq g; \\ |Y_i| \leq \frac{1}{2} \text{ for } 1 \leq i \leq g. \end{cases}$$

The result now follows from the continuity of the map $z \in \mathcal{M}(\mathbb{Z}) \mapsto P_z \in \mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$. \square

5. OUTER AUTOMORPHISMS AND CONNECTEDNESS OF \mathcal{S}_g

In this section, we study a natural action of the group $\text{Out}(F_g)$ of outer automorphisms on the Schottky space \mathcal{S}_g . We show that this action respects the analytic structure of the Schottky space and use this to prove that \mathcal{S}_g is path-connected (see 5.2). Finally, we study the properness of this action. When $g = 1$, $\text{Out}(F_1)$ acts trivially on \mathcal{S}_1 , hence we assume for the rest of the section that $g \geq 2$.

5.1. The action of $\text{Out}(F_g)$ on the Schottky space. Recall from Lemma 4.2.5 that a point of \mathcal{S}_g corresponds to a homomorphism in $\text{Hom}_S(F_g, \text{PGL}_2)$. This identification gives rise to a natural action of $\text{Aut}(F_g)$ on \mathcal{S}_g by letting an element of $\text{Aut}(F_g)$ act on the source of homomorphisms in $\text{Hom}_S(F_g, \text{PGL}_2)$.

More precisely, let $\tau \in \text{Aut}(F_g)$, $x \in \mathcal{S}_g$, and φ_x be the associated homomorphism of $\text{Hom}_S(F_g, \text{PGL}_2)$. Then, the map $\varphi_x \circ \tau$ is a group homomorphism from F_g to $\text{PGL}_2(\mathcal{H}(x))$. Its image, being the same as that of φ_x , is the Schottky group Γ_x . Moreover, there exists a unique Möbius transformation ε , with coefficients in $\mathcal{H}(x)$, that sends the attracting and repelling fixed points of $\varphi_x \circ \tau(e_1)$ to 0 and ∞ respectively and the attracting fixed point of $\varphi_x \circ \tau(e_2)$ to 1. Then, $\varepsilon^{-1}(\varphi_x \circ \tau) \varepsilon$ belongs to $\text{Hom}_S(F_g, \text{PGL}_2(\mathcal{H}(x)))$, hence gives rise to a point of \mathcal{S}_g . We denote it by τx .

Definition 5.1.1. The map $(\tau, x) \in \text{Aut}(F_g) \times \mathcal{S}_g \mapsto \tau x \in \mathcal{S}_g$ defines an action of $\text{Aut}(F_g)$ on \mathcal{S}_g that factors through $\text{Out}(F_g)$.

We now describe the stabilizers of the points of \mathcal{S}_g under the action of $\text{Out}(F_g)$. The corresponding result for rigid Schottky spaces is known (see [Ger81, Satz 3]).

Lemma 5.1.2. *Let $(k, |\cdot|)$ be a complete valued field and let γ be a loxodromic element of $\text{PGL}_2(k)$ with fixed points α and β . Let $\varepsilon_1, \varepsilon_2 \in \text{PGL}_2(k)$ such that $\varepsilon_1^{-1} \gamma \varepsilon_1 = \varepsilon_2^{-1} \gamma \varepsilon_2$. Then, we have $\varepsilon_1^{-1}(\alpha) = \varepsilon_2^{-1}(\alpha)$ and $\varepsilon_1^{-1}(\beta) = \varepsilon_2^{-1}(\beta)$.*

Proof. We may assume that α is the attracting point of γ . Let $P \in \mathbb{P}_k^{1, \text{an}} - \{\varepsilon_1^{-1}(\alpha), \varepsilon_1^{-1}(\beta), \varepsilon_2^{-1}(\alpha), \varepsilon_2^{-1}(\beta)\}$. Then, for each $n \in \mathbb{Z}$, we have

$$\varepsilon_1^{-1}(\gamma^n(\varepsilon_1(P))) = \varepsilon_2^{-1}(\gamma^n(\varepsilon_2(P))).$$

The left-hand side converges to $\varepsilon_1^{-1}(\alpha)$ (resp. $\varepsilon_1^{-1}(\beta)$) when n goes to $+\infty$ (resp. $-\infty$). The right-hand side converges to $\varepsilon_2^{-1}(\alpha)$ (resp. $\varepsilon_2^{-1}(\beta)$) when n goes to $+\infty$ (resp. $-\infty$). The result follows. \square

Proposition 5.1.3. *Let $x \in \mathcal{S}_g$. The stabilizer of x under the action of $\text{Out}(F_g)$ is isomorphic to the quotient $\Gamma_x \backslash N(\Gamma_x)$, where $N(\Gamma_x)$ denotes the normalizer of Γ_x in $\text{PGL}_2(\mathcal{H}(x))$.*

Proof. Let $\varepsilon \in N(\Gamma_x)$. The morphism φ_x induces an isomorphism $\psi_x: F_g \xrightarrow{\sim} \Gamma_x$. Since ε belongs to the normalizer of Γ_x in $\text{PGL}_2(\mathcal{H}(x))$, the conjugation by ε in $\text{PGL}_2(\mathcal{H}(x))$ induces an automorphism c_ε

of Γ_x . It follows from the definitions that $\psi_x^{-1} \circ c_\varepsilon \circ \psi_x$ is an automorphism of F_g stabilizing x . We have just constructed a map $\nu: N(\Gamma_x) \rightarrow \text{Stab}_{\text{Aut}(F_g)}(x)$. It is a morphism of groups.

Let $\varepsilon \in N(\Gamma_x)$. The automorphism $\nu(\varepsilon)$ is inner if, and only if, there exists $w \in F_g$ such that $\nu(\varepsilon) = c_w$, where c_w denotes the automorphism defined by the conjugation by w in F_g . Note that we have $c_w = \psi_x^{-1} \circ c_{\psi_x(w)} \circ \psi_x$. It follows that $\nu(\varepsilon)$ is inner if, and only if, there exists $\delta \in \Gamma_x$ such that $c_\varepsilon = c_\delta$. If the latter condition holds, then, by Lemma 5.1.2, ε^{-1} and δ^{-1} coincide on all the fixed points of the $M_i(x)$'s. Since $g \geq 2$, there are more than two fixed points, hence $\varepsilon = \delta$. The argument shows that ν induces an injective morphism $\nu': \Gamma_x \backslash N(\Gamma_x) \rightarrow \text{Stab}_{\text{Out}(F_g)}(x)$.

To conclude, it remains to prove that ν' is surjective. It is enough to prove that ν is surjective. Let $\tau \in \text{Stab}_{\text{Aut}(F_g)}(x)$. For each $i \in \{1, \dots, g\}$, $\tau(e_i)$ is an element of F_g , that is to say a word $w_i = e_{j_{i,0}}^{n_{i,0}} \cdots e_{j_{i,r_i}}^{n_{i,r_i}}$, for some $r_i \in \mathbb{N}$, $j_{i,0}, \dots, j_{i,r_i} \in \{1, \dots, g\}$, $n_{i,0}, \dots, n_{i,r_i} \in \mathbb{Z}$. Since φ_x is a morphism of groups, we have

$$N_i := \varphi_x \circ \tau(e_i) = M_{j_{i,0}}(x)^{n_{i,0}} \cdots M_{j_{i,r_i}}(x)^{n_{i,r_i}}.$$

In particular, $N_i \in \Gamma_x$. Let $\varepsilon \in \text{PGL}_2(\mathcal{H}(x))$ be the unique element that sends the attracting and repelling fixed points of N_1 to 0 and ∞ respectively and the attracting fixed point of N_2 to 1. By definition of the action, we have

$$\forall i \in \{1, \dots, g\}, \varepsilon^{-1} N_i \varepsilon = M_i(\tau x) = M_i(x).$$

By using the words associated to the morphism τ^{-1} , one may express the $M_j(x)$'s in terms of the N_i 's. It follows that the N_i 's generate the group Γ_x , hence $\varepsilon \in N(\Gamma_x)$. Moreover, we have $\nu(\varepsilon^{-1}) = \tau$. The result follows. \square

Remark 5.1.4. Let $x \in \mathcal{S}_g$. It is easy to check that each element of $N(\Gamma_x)$ preserves the limit set \mathcal{L}_x , hence inducing an automorphism of $\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}} - \mathcal{L}_x$, and that the latter descends to an automorphism of \mathcal{C}_x . This construction gives rise to a group homomorphism $\phi: N(\Gamma_x) \rightarrow \text{Aut}(\mathcal{C}_x)$ with kernel Γ_x .

Let $x \in \mathcal{S}_g^{\text{na}}$ be a non-archimedean point of the Schottky space. Then, the homomorphism ϕ is surjective (see [Mum72, Corollary 4.12] for the first occurrence of this result and [PT21, §6.5.2] for a discussion using Berkovich geometry), and as a result the quotient group $\Gamma_x \backslash N(\Gamma_x)$ considered above is isomorphic to the automorphism group $\text{Aut}(\mathcal{C}_x)$ of the curve \mathcal{C}_x . Every element of $\text{Aut}(\mathcal{C}_x)$ restricts to an isometry of the skeleton Σ_x defined in 4.2.2. This restriction induces an injection of the automorphism group $\text{Aut}(\mathcal{C}_x)$ in the group $\text{Aut}(\Sigma_x)$ of isometric automorphisms of Σ_x (cf. [PT21, Proposition 6.5.9]).

Let $x \in \mathcal{S}_g^{\text{a}}$ be an archimedean point of the Schottky space. In this case the homomorphism ϕ is not surjective in general, and hence we only have an injection of $\Gamma_x \backslash N(\Gamma_x)$ into the group $\text{Aut}(\mathcal{C}_x)$. The question of lifting automorphisms of a Riemann surface to $\Gamma \backslash N(\Gamma)$ for some Schottky group Γ uniformizing such surface has been thoroughly investigated by R. Hidalgo (see for example [Hid05]).

Proposition 5.1.5. *The action of $\text{Out}(F_g)$ on \mathcal{S}_g is analytic and has finite stabilizers. Moreover:*

- (i) *If $g \geq 3$, then this action is faithful;*
- (ii) *If $g = 2$, then the element $\iota \in \text{Out}(F_2)$ defined by $\iota(e_i) = e_i^{-1}$ for $i = 1, 2$ stabilizes every point of \mathcal{S}_2 , and the action of the quotient $\text{Out}(F_2)/\langle \iota \rangle$ on \mathcal{S}_2 is faithful.*

Proof. It is a classical result of Nielsen that $\text{Out}(F_g)$ is generated by the set of four elements $\{\sigma_1, \sigma_2, \sigma_3, \sigma_4\}$ defined by:

$$\begin{cases} \sigma_1(e_1) = e_g, \sigma_1(e_i) = e_{i-1} \forall i > 1 \\ \sigma_2(e_1) = e_2, \sigma_2(e_2) = e_1, \sigma_2(e_i) = e_i \forall i > 2 \\ \sigma_3(e_1) = e_1^{-1}, \sigma_3(e_i) = e_i \forall i > 1 \\ \sigma_4(e_2) = e_1^{-1}e_2, \sigma_4(e_i) = e_i \forall i \neq 2 \end{cases}$$

For $i = 1, 2, 3$ a simple computation shows that every σ_i acts on \mathcal{S}_g by Möbius transformations on the Koebe coordinates. For example, in the case of σ_1 , the point x of \mathcal{S}_g with Koebe coordinates $(\alpha_i, \alpha'_i, \beta_i)$ is sent to a point with basis $(M(\alpha_g, \alpha'_g, \beta_g), M(0, \infty, \beta_1), \dots, M(\alpha_{g-1}, \alpha'_{g-1}, \beta_{g-1}))$. To describe the action in terms of Koebe coordinates, we need to conjugate this basis by the unique Möbius transformation γ such that $\gamma(\alpha_g) = 0$, $\gamma(\alpha'_g) = \infty$, and $\gamma(0) = 1$. This conjugation sends the fixed points of a transformation to their images under γ , while leaving multipliers untouched. Hence, the Koebe coordinates of $\sigma_1(x)$ are

$$\sigma_1(\alpha_i, \alpha'_i, \beta_i) = (\gamma(1), \dots, \gamma(\alpha_{g-1}), \gamma(\infty), \dots, \gamma(\alpha'_{g-1}), \beta_g, \beta_1, \dots, \beta_{g-1}).$$

The cases of σ_2 and σ_3 are completely analogous, and so in these three cases the action is analytic.

Let us show that this is the case for σ_4 as well. Denote by $M := M(\alpha^*, \alpha'^*, \beta^*)$ the matrix representing the product $M(0, \infty, \beta_1)^{-1}M(1, \alpha'_2, \beta_2)$. The Koebe coordinates $(\alpha^*, \alpha'^*, \beta^*)$ are analytic functions in the coefficients of M by virtue of Proposition 3.4.3. Moreover, the coefficients of M are rational functions without poles in the variables $\beta_1, \alpha'_2, \beta_2$ on \mathcal{S}_g , and then analytic as well. Finally, if we want to get a basis with 1 as attracting fixed point of the second generator we have to conjugate every element by multiplication by $\frac{1}{\alpha^*}$. Summarizing, we get the following expression of the action of σ_4 on \mathcal{S}_g :

$$\sigma_4(\alpha_i, \alpha'_i, \beta_i) = \left(\frac{\alpha_3}{\alpha^*}, \dots, \frac{\alpha_g}{\alpha^*}, \frac{\alpha'^*}{\alpha^*}, \frac{\alpha'_3}{\alpha^*}, \dots, \frac{\alpha'_g}{\alpha^*}, \beta_1, \beta^*, \beta_3, \dots, \beta_g \right).$$

Since α^* , α'^* , and β^* are analytic functions of the Koebe coordinates, the action of σ_4 , and hence of $\text{Out}(F_g)$, is analytic on \mathcal{S}_g .

The finiteness of the stabilizers follows from Proposition 5.1.3 and Remark 5.1.4. To prove faithfulness for $g \geq 3$, it is enough to remark that for every valued field there exist Schottky uniformized curves with trivial automorphism groups. For $g = 2$, the outer automorphism ι of order 2 defined by $\iota(e_i) = e_i^{-1}$ for $i = 1, 2$ fixes every point $x \in \mathcal{S}_2$. In fact, writing in Koebe coordinates $x = (\alpha'_2, \beta_1, \beta_2)$, the point $\iota(x)$ corresponds to the ordered basis $(M(\infty, 0, \beta_1), M(\alpha'_2, 1, \beta_2))$. Then, the conjugation of this basis by the Möbius transformation $i : z \mapsto \frac{\alpha'_2}{z}$ produces the ordered basis $(M(0, \infty, \beta_1), M(1, \alpha'_2, \beta_2))$, so that $\iota(x) = x$. Since a generic genus 2 curve that admits Schottky uniformization has an automorphism group of order 2, we can conclude that the action of the quotient $\text{Out}(F_2)/\langle \iota \rangle$ on \mathcal{S}_2 is faithful. \square

Remark 5.1.6. The automorphism ι appearing in part (ii) of Proposition 5.1.5 induces, via the isomorphism given in Proposition 5.1.3, the element $J = \begin{bmatrix} 0 & \alpha'_2 \\ 1 & 0 \end{bmatrix} \in N(\Gamma_x)$. In turn, the class $[J]$ in $\Gamma_x \backslash N(\Gamma_x) \cong \text{Aut}(\mathcal{C}_x)$ induces on \mathcal{C}_x the hyperelliptic involution. If x is a non-archimedean point, this follows from the fact that $[J]$ acts on Σ_x in such a way that the quotient is a tree. If x is an archimedean point, one can argue as follows. Writing $\gamma_1 = M(0, \infty, \beta_1)$ and $\gamma_2 = M(1, \alpha'_2, \beta_2)$, the elements J , $\gamma_1 J$, and $\gamma_2 J$ restrict to the same automorphism of \mathcal{C}_x and give rise to 6 distinct fixed points in $\mathbb{P}_k^{1, \text{an}} - \mathcal{L}_x$. Up to conjugation by elements of Γ_x , one can assume that these points all lie in the same fundamental domain. Hence, the cover of curves $\mathcal{C}_x \rightarrow \mathcal{C}_x / \langle [J] \rangle$ is ramified at least at 6 points and applying Riemann-Hurwitz formula one finds that the genus of the target curve is 0.

5.2. Entr'acte: path-connectedness of \mathcal{S}_g . We now apply the results of the previous section to show that \mathcal{S}_g is a connected topological space.

We already know that the archimedean part \mathcal{S}_g^a of the Schottky space is path-connected, thanks to its relation to the complex Schottky space (see Lemma 4.2.4). This allows to use global arguments to show the connectedness of \mathcal{S}_g .

Theorem 5.2.1. *The Schottky space \mathcal{S}_g is path-connected.*

Proof. Let $x \in \mathcal{S}_g^{\text{na}}$ be a non-archimedean point of the Schottky space over \mathbb{Z} . By Corollary 4.3.3, there is an automorphism $\tau \in \text{Out}(F_g)$ such that $\tau(x) \in \mathcal{SB}_g^{\text{na}}$.

Let $\alpha_3, \dots, \alpha_g, \alpha'_2, \dots, \alpha'_g \in \mathbb{C}$ such that the degree of transcendence of the extension of \mathbb{Q} they generate is maximal (equal to $2g - 3$). Let $r_1, \dots, r_g \in (0, 1)$ whose images in the \mathbb{Q} -vector space $\mathbb{R}_{>0}$ are linearly independent. For $\varepsilon \in (0, 1]$, set

$$\sigma(a_\infty^\varepsilon) := \rho_\varepsilon(\alpha_3, \dots, \alpha_g, \alpha'_2, \dots, \alpha'_g, r_1^{1/\varepsilon}, \dots, r_g^{1/\varepsilon}) \in \text{pr}_{\mathbb{Z}}^{-1}(a_\varepsilon)$$

(see Section 2.3 for this notation). For each $a \in \mathcal{M}(\mathbb{Z})^{\text{na}}$, denote by $\sigma(a)$ the unique point in the Shilov boundary of the disc inside $\text{pr}_{\mathbb{Z}}^{-1}(a) \simeq \mathbb{A}_{\mathcal{H}(a)}^{3g-3}$ defined by the inequalities

$$\begin{cases} |X_i| \leq 1 \text{ for } 3 \leq i \leq g; \\ |X'_i| \leq 1 \text{ for } 2 \leq i \leq g; \\ |Y_i| \leq r_i \text{ for } 1 \leq i \leq g. \end{cases}$$

By comparing the limit of $\sigma(a_\infty^\varepsilon)$ for $\varepsilon \rightarrow 0$ with the non-archimedean valuation $\sigma(a_0)$ over the central point (cf. Examples 2.3.1 and 2.3.2), one shows that the map

$$\sigma: a \in \mathcal{M}(\mathbb{Z}) \mapsto \sigma(a) \in \mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}}$$

is a continuous section of the projection morphism $\text{pr}_{\mathbb{Z}}: \mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}} \rightarrow \mathcal{M}(\mathbb{Z})$. By Corollary 4.4.4, $\sigma(a_0)$ belongs to \mathcal{SB}_g and to the same path-connected component in \mathcal{S}_g as $\tau(x)$.

Since \mathcal{S}_g is open, by Theorem 4.3.4, we deduce that $\sigma(a_\infty^\varepsilon)$ belongs to \mathcal{S}_g for $\varepsilon \in (0, 1]$ small enough. In particular, $\tau(x)$ belongs to the same path-connected component of \mathcal{S}_g as \mathcal{S}_g^a . By Proposition 5.1.5, τ^{-1} acts continuously on \mathcal{S}_g , hence x belongs to the same path-connected component of \mathcal{S}_g as \mathcal{S}_g^a . The result now follows from the path-connectedness of the latter (see Lemma 4.2.4). \square

5.3. Properness of the action of $\text{Out}(F_g)$. Recall that we discussed proper actions at the beginning of Section 3.3. Let us first consider the archimedean case. The following proposition is a consequence of well known facts about the complex Schottky space.

Theorem 5.3.1. *The action of $\text{Out}(F_g)$ on \mathcal{S}_g^a is proper. The quotient space $\text{Out}(F_g) \backslash \mathcal{S}_g^a$ is Hausdorff and reduces locally to a quotient by a finite group.*

Proof. Using the homeomorphism

$$\Phi: \mathbb{A}_{\mathbb{R}}^{3g-3, \text{an}} \times (0, 1] \rightarrow (\mathbb{A}_{\mathbb{Z}}^{3g-3, \text{an}})^a$$

from Section 2.3 as well as Remarks 3.5.2 and 3.5.3 one reduces to the case of the complex Schottky space $\mathcal{S}_{g, \mathbb{C}}$. Recall from the discussion in Remark 4.2.3 the universal covering map $\Psi: \mathcal{T}_{g, \mathbb{C}} \rightarrow \mathcal{S}_{g, \mathbb{C}}$ from the complex Teichmüller space. The mapping class group MCG_g acts properly on $\mathcal{T}_{g, \mathbb{C}}$ ([Gar87, §8, Theorem 6]) in such a way that Ψ is a G -covering for a normal subgroup $G \subset \text{MCG}_g$. The action of $\text{Out}(F_g)$ on $\mathcal{S}_{g, \mathbb{C}}$ considered here comes from the realization of $\text{Out}(F_g)$ as a subgroup of the quotient MCG_g/G (see [HS07, Proposition 5.10]). In particular, it is a proper action.

The fact that the quotient space $\text{Out}(F_g) \backslash \mathcal{S}_g^a$ is Hausdorff follows directly from the properness of the action. The fact that it reduces locally to a quotient by a finite group is a consequence of the finiteness of the stabilizers (see Proposition 5.1.5). \square

Let us now focus on the the non-archimedean case. We follow the strategy outlined in the proof of [Ger82, Proposition 7] and take this opportunity to add details to said proof.

Theorem 5.3.2. *The set*

$$SB = \{\tau \in \text{Out}(F_g) : \tau(\mathcal{SB}_g^{\text{na}}) \cap \mathcal{SB}_g^{\text{na}} \neq \emptyset\}$$

is finite.

Proof. For every point $x \in \mathcal{SB}_g^{\text{na}}$, let us denote by $L_x \subset \mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}}(\mathcal{H}(x))$ the limit set of Γ_x , and by $T_{\Gamma_x} \subset \mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}}$ the infinite tree defined as the skeleton of $\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}} - L_x$.³ The action of Γ_x on the infinite tree T_{Γ_x} is free and without inversions, and gives rise to a universal covering $p_x : T_{\Gamma_x} \rightarrow \Sigma_x$ of the skeleton of the Mumford curve uniformized by Γ_x .

Following Serre [Ser77, §3.1], we call *representative tree of T_{Γ_x}* any subtree of T_{Γ_x} that is a lifting of a spanning tree of Σ_x via p_x . Equivalently, a representative tree is a connected subtree of T_{Γ_x} that has a unique vertex in any given Γ_x -orbit on the set of vertices of T_{Γ_x} . With a representative tree $T \subset T_{\Gamma_x}$, we can associate a generating set of Γ_x as follows. Let us call E_T the set of edges in T_{Γ_x} that have one endpoint in T and the other in its complement $T_{\Gamma_x} - T$. Note that the set E_T consists of $2g$ elements.

Lemma 5.3.3. *The set*

$$G_T = \{\gamma \in \Gamma_x - \{1\} : \exists e \in E_T \text{ with } \gamma(e) \in E_T\}$$

is of the form $B \cup B^{-1}$ with $B = \{\gamma_1, \dots, \gamma_g\}$ a basis of Γ_x and $B^{-1} = \{\gamma_1^{-1}, \dots, \gamma_g^{-1}\}$.

Proof. Let us choose an orientation on the tree T_{Γ_x} compatible with the action of Γ_x , and we consider the set $B \subset \Gamma_x$ consisting of the elements $\gamma \in \Gamma_x$ such that there exists an edge e of T_{Γ_x} starting in T and ending in $\gamma(T)$. By applying a theorem of Serre on free actions on oriented trees [Ser77, §3.3 Théorème 4', a)], we deduce that B is a basis of Γ_x . By construction, the set $B \cup B^{-1}$ is contained in G_T , so it suffices to show that $G_T \subset B \cup B^{-1}$ to conclude. To show this, let us pick $\gamma \in G_T$ and $e_1, e_2 \in E_T$, such that $\gamma(e_1) = e_2$. If we call v_i, w_i the endpoints of e_i in such a way that $v_i \in T$, then v_1 and v_2 can not be in the same orbit, hence $\gamma(v_1) = w_2$ and $\gamma(w_1) = v_2$. As a result, $w_1 \in \gamma(T)$ and $w_2 \in \gamma^{-1}(T)$. Then, depending on the orientation chosen at the beginning, either $\gamma \in B$ or $\gamma \in B^{-1}$, as desired. \square

We denote by \mathfrak{F}_x the set of representative trees of T_{Γ_x} , and by \mathfrak{B}_x the set of generating sets of Γ_x of the form $B \cup B^{-1}$ as in the statement of Lemma 5.3.3.

Lemma 5.3.4. *The function*

$$\begin{aligned} G_x : \mathfrak{F}_x &\rightarrow \mathfrak{B}_x \\ T &\mapsto G_T \end{aligned}$$

is injective. Its image consists of those generating sets $B \cup B^{-1}$ such that B is a Schottky basis.

³Recall from [Ber90, 4.1.3] that the skeleton of an analytic curve C is defined as the subset of C consisting of those points that do not have a neighborhood potentially isomorphic to a disc. If C is the analytification of a smooth proper algebraic curve, its skeleton is a finite graph and this definition coincides with the one at the end of Notation 4.2.2.

Proof. Let B be a Schottky basis, choose a Schottky figure adapted to B , and consider the associated analytic space $F^+ \subset \mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}}$ as in Definition 3.1.1. We claim that the maximal subtree T_B of T_{Γ_x} contained in F^+ has a unique vertex in any Γ_x -orbit on the set of vertices of T_{Γ_x} , and therefore is a representative tree. To prove this, recall from 4.2.2 that the skeleton Σ_x is obtained by pairwise identifying the endpoints of the intersection $F^+ \cap T_{\Gamma_x}$ according to the action of Γ_x . In particular, F^+ contains a fundamental domain for the action of Γ_x on $\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}} - L_x$, so that there is a vertex of T_B in the orbit of v , for every vertex v of T_{Γ_x} . Moreover, if ξ is an endpoint of $F^+ \cap T_{\Gamma_x}$, then the map p_x identifies ξ with only another endpoint of $F^+ \cap T_{\Gamma_x}$, so that $p_x(\xi)$ is a point of degree 2 of Σ_x and hence by definition it is not a vertex of Σ_x . In particular, ξ is not a vertex of T_B , so that all vertices of T_B are contained in a fundamental domain for the action of Γ_x on $\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}} - L_x$. This shows that every element in the orbit of a vertex v of T_B different from itself does not lie in T_B , concluding the proof that T_B is a representative tree.

Note that the endpoints of $F^+ \cap T_{\Gamma_x}$ lie precisely on those edges of T_{Γ_x} that are in E_{T_B} , and by construction of F^+ the elements of Γ_x acting on these endpoints are precisely those lying in $B \cup B^{-1}$. Hence we have $G_{T_B} = B \cup B^{-1}$, showing that $B \cup B^{-1}$ is in the image of G_x . Conversely, if $B = \{\gamma_1, \dots, \gamma_g\}$ is a basis of Γ_x and $B \cup B^{-1} \in \mathfrak{B}_x$ can be written as $G_x(T)$ for some representative tree T , one can build a Schottky figure adapted to B as follows. First one writes the set E_T as $\{e_{-g}, \dots, e_{-1}, e_1, \dots, e_g\}$ in such a way that $\gamma_i(e_{-i}) = e_i$. Then one chooses a set of $2g$ points $\{x_{-g}, \dots, x_{-1}, x_1, \dots, x_g\}$ in $\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}}$ in such a way that $x_{-i} \in e_{-i}$, $x_i \in e_i$ and $\gamma_i(x_{-i}) = x_i$ for every $i = 1, \dots, g$ (here we tacitly identify an element of E_T with the corresponding subset of $\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}}$). Each $x_i \in \mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}}$ is the Shilov boundary of a unique closed disc of $\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}}$ not containing T , that we denote by B_i^+ . The family $\mathcal{B} = \{B_i^+, i \in \{-g, \dots, -1, 1, \dots, g\}\}$ is a Schottky figure adapted to B : In fact, if we denote by B_i^- the unique connected component of $\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}} - \{x_i\}$ such that $T \cap B_i^- = \emptyset$ and $L_x \cap B_i^- \neq \emptyset$, we have that B_i^- is a maximal open disc inside B_i^+ and that

$$B_i^- = \gamma_i(\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}} - B_{-i}^+),$$

where we adopted the convention $\gamma_{-i} = \gamma_i^{-1}$. Then \mathcal{B} is a Schottky basis, constructed in such a way that the representative tree T_B as above coincides with T .

The injectivity of G_x is proved as follows: let $G_x(T_1) = G_x(T_2) = B \cup B^{-1}$. By what we just proved, \mathcal{B} is a Schottky basis and the injectivity of G_x is equivalent to the fact that every Schottky figure associated with \mathcal{B} gives rise to the same representative tree T_B . We can show this by contradiction: suppose that two Schottky figures \mathcal{B}_1 and \mathcal{B}_2 adapted to \mathcal{B} give rise to different representative trees T_1 and T_2 . Then there is a vertex v of T_1 that is not a vertex of T_2 , and there are at least two edges e_i, e_j of E_{T_1} having v as an endpoint that are not edges of T_2 nor they belong to E_{T_2} , for instance those edges departing from v in a direction different from the one of T_2 . As a result, there is a unique connected component of $T_{\Gamma_x} - T_2$ that contains both e_i and e_j . Hence, the two closed discs $B_i^+, B_j^+ \in \mathcal{B}_1$ corresponding to e_i, e_j are contained in a single closed disc $B^+ \in \mathcal{B}_2$. This leads to a contradiction, since every disc in a Schottky figure adapted to \mathcal{B} contains a unique fixed point of a unique element of B , but B^+ contains at least two of them. This shows that $T_1 = T_2$. \square

Thanks to the injectivity of G_x , one can associate with x a unique representative tree $T_x \in \mathfrak{F}_x$, defined as the preimage by G_x of the generating set $\{M_1(x), \dots, M_g(x), M_1^{-1}(x), \dots, M_g^{-1}(x)\}$. Furthermore, with the point x one can also associate the set

$$SB_x = \{\tau \in \text{Out}(F_g) : \tau(x) \in \mathcal{SB}_g\}.$$

Our proof of the theorem then relies on the following lemmas.

Lemma 5.3.5. *Let $x \in \mathcal{SB}_g^{\text{na}}$. Then SB_x is a finite subset of $\text{Out}(F_g)$.*

Proof. Let us fix a vertex $v \in T_x$, and call $\mathfrak{F}_{x,v}$ the subset of \mathfrak{F}_x consisting of those representative trees that contain v . We first prove that every Schottky basis is Γ_x -conjugated to a unique Schottky basis in the image $G_x(\mathfrak{F}_{x,v})$. In fact, if B is a Schottky basis, then $B \cup B^{-1} = G_x(T)$ for some $T \in \mathfrak{F}_x$. For every $\gamma \in \Gamma_x$, $\gamma G_x(T) \gamma^{-1}$ is the image by G_x of the representative tree $\gamma(T)$. Since there exists a unique $\gamma \in \Gamma_x$ such that $v \in \gamma(T)$, there is a unique generating set in $G_x(\mathfrak{F}_{x,v})$ conjugated to $B \cup B^{-1}$ by an element of Γ_x . As a result, the function G_x realizes a bijection between the set $\mathfrak{F}_{x,v}$ and the set of Γ_x -conjugacy classes of generating sets of the form $B \cup B^{-1}$ with B a Schottky basis of Γ_x . As the former is a finite set, the latter is also finite and, in particular, the PGL_2 -conjugacy classes of Schottky bases of Γ_x are finite.

For every $y \in \mathcal{SB}_g^{\mathrm{na}}$, let us define the subset $SB_{x,y} = \{\tau \in \mathrm{Out}(F_g) : \tau(x) = y\}$ of SB_x . If $\tau_1, \tau_2 \in SB_{x,y}$, then $\tau_1 \tau_2^{-1}(x) = x$, so there is an element $\sigma \in \mathrm{Out}(F_g)$ in the stabilizer of x such that $\tau_1 = \sigma \tau_2$. By Proposition 5.1.3, this stabilizer is finite, so the set $SB_{x,y}$ is finite too. We can write the set SB_x as a union

$$SB_x = \bigcup_{y \in \mathcal{SB}_g^{\mathrm{na}}} SB_{x,y}.$$

Note that the set $SB_{x,y}$ is non-empty only if the Schottky basis $(M_1(y), \dots, M_g(y))$ is PGL_2 -conjugated to $\tau(M_1(x), \dots, M_g(x))$. Hence, by what precedes, SB_x is a finite union of finite sets and then it is finite. \square

Given a tree \mathcal{T} , recall that a vertex of degree one of \mathcal{T} is called a *leaf*. If the set of leaves $L(\mathcal{T})$ of \mathcal{T} has cardinality $2g$, we call *leaf labeling* of \mathcal{T} a bijection between $L(\mathcal{T})$ and the set $\{-g, \dots, -1, 1, \dots, g\}$. Let us denote by \mathfrak{L}_g the set of pairs (\mathcal{T}, ℓ) where \mathcal{T} is a finite tree with $2g$ leaves and no vertices of degree 2, and ℓ is a leaf labeling of \mathcal{T} . Since g is fixed, \mathfrak{L}_g is a finite set. For a point $x \in \mathcal{SB}_g^{\mathrm{na}}$, the subtree $T_x \cup E_{T_x}$ of T_{Γ_x} has $2g$ leaves and no vertices of degree 2, and can naturally be endowed with a labeling induced by the writing $E_{T_x} = \{e_{-g}, \dots, e_{-1}, e_1, \dots, e_g\}$ as in the first part of the proof. This assignment defines a map $\lambda : \mathcal{SB}_g^{\mathrm{na}} \rightarrow \mathfrak{L}_g$.

Lemma 5.3.6. *Let x, y be two points of $\mathcal{SB}_g^{\mathrm{na}}$ such that $\lambda(x) = \lambda(y)$. Then $SB_x = SB_y$.*

Proof. Since $\lambda(x) = \lambda(y)$, there exists an isomorphism of finite trees

$$\psi : T_x \cup E_{T_x} \rightarrow T_y \cup E_{T_y}$$

that sends T_x to T_y and respects the leaf labelings. Consider the group isomorphism $\phi : \Gamma_x \rightarrow \Gamma_y$ such that $\phi(M_i(x)) = M_i(y)$. Then there is a unique way to extend ϕ -equivariantly the isomorphism ψ to an isomorphism of infinite trees $\Psi : T_{\Gamma_x} \rightarrow T_{\Gamma_y}$. Namely, for every vertex $v' \in T_{\Gamma_x}$, there is a unique pair (γ, v) with $\gamma \in \Gamma_x$ and v a vertex of T_x such that $v' = \gamma(v)$. The assignment $\Psi(v') = \phi(\gamma)(\Psi(v))$ uniquely determines the isomorphism Ψ .

Let us fix vertices $v_x \in T_x$ and $v_y \in T_y$ such that $\Psi(v_x) = v_y$. Note that we constructed Ψ in such a way to be equivariant, so there is a commutative diagram of the form

$$\begin{array}{ccc} T_{\Gamma_x} & \xrightarrow{\Psi} & T_{\Gamma_y} \\ \downarrow p_x & & \downarrow p_y \\ \Sigma_x & \xrightarrow{\sim} & \Sigma_y \end{array},$$

where the arrow on the bottom is an isomorphism of graphs, and in particular sends spanning trees in Σ_x to spanning trees in Σ_y . As a result, Ψ sends representative trees in T_{Γ_x} to representative trees in T_{Γ_y} . In particular, Ψ restricts to a function

$$\Psi|_{\mathfrak{F}_{x,v_x}} : \mathfrak{F}_{x,v_x} \rightarrow \mathfrak{F}_{y,v_y}$$

satisfying $\Psi|_{\mathfrak{F}_{x,v_x}}(T_x) = T_y$.

Now we consider an element $\tau \in SB_x$. This has a unique representative $\sigma \in \text{Aut}(F_g)$ such that B_σ , the Schottky basis resulting from applying σ to $(M_1(x), \dots, M_g(x))$, satisfies $G_x^{-1}(B_\sigma) \in \mathfrak{F}_{x,v_x}$. If we consider the representative tree $T = \Psi|_{\mathfrak{F}_{x,v_x}}(G_x^{-1}(B_\sigma))$, we have that the generating set $G_y(T)$ of Γ_y is of the form $B \cup B^{-1}$ for some Schottky basis B of Γ_y . By definition, $G_y(T)$ consists of those elements of Γ_y that act on the set E_T . Note that for every vertex $v \in T_{\Gamma_x}$, we have $\Psi((M_i(x))(v)) = (M_i(y))(\Psi(v))$ thanks to the fact that Ψ is ϕ -equivariant. This condition ensures that, if we set $B_\sigma = (M_{j_1,0}^{n_{1,0}}(x) \cdots M_{j_1,r_1}^{n_{1,r_1}}(x), \dots, M_{j_g,0}^{n_{g,0}}(x) \cdots M_{j_g,r_g}^{n_{g,r_g}}(x))$, then B can be taken to be $(M_{j_1,0}^{n_{1,0}}(y) \cdots M_{j_1,r_1}^{n_{1,r_1}}(y), \dots, M_{j_g,0}^{n_{g,0}}(y) \cdots M_{j_g,r_g}^{n_{g,r_g}}(y))$, that is, $\tau(y) \in \mathcal{SB}_g^{\text{na}}$. Hence τ is in SB_y and so $SB_x \subset SB_y$. The same construction applied to the isomorphism Ψ^{-1} shows that $\tau \in SB_y$ implies $\tau \in SB_x$, so that $SB_x = SB_y$. \square

The theorem then follows from the two lemmas above. We write

$$SB = \bigcup_{x \in \mathcal{SB}_g^{\text{na}}} SB_x = \bigcup_{\lambda(x) \in \mathcal{L}_g} SB_x,$$

where the second equality is given by Lemma 5.3.6. The result of Lemma 5.3.5 ensures the finiteness of SB_x for every x , and the finiteness of the set \mathcal{L}_g allows to conclude. \square

Corollary 5.3.7. *The action of $\text{Out}(F_g)$ on $\mathcal{S}_g^{\text{na}}$ is proper. The quotient space $\text{Out}(F_g) \backslash \mathcal{S}_g^{\text{na}}$ is Hausdorff and reduces locally to a quotient by a finite group.*

Proof. Let $x, y \in \mathcal{S}_g^{\text{na}}$. By Corollary 4.3.3, there are $\sigma_x, \sigma_y \in \text{Out}(F_g)$ such that $\sigma_x(x), \sigma_y(y) \in \mathcal{SB}_g^{\text{na}}$. By Corollary 4.4.4, $\mathcal{SB}_g^{\text{na}}$ is open in $\mathcal{S}_g^{\text{na}}$. It follows that $U_x := \sigma_x^{-1}(\mathcal{SB}_g^{\text{na}})$ and $U_y := \sigma_y^{-1}(\mathcal{SB}_g^{\text{na}})$ are open neighborhoods of x and y respectively. By Theorem 5.3.2, the set

$$\{\tau \in \text{Out}(F_g) : \tau(U_x) \cap U_y \neq \emptyset\} = \{\tau \in \text{Out}(F_g) : \sigma_y \tau \sigma_x^{-1}(\mathcal{SB}_g^{\text{na}}) \cap \mathcal{SB}_g^{\text{na}} \neq \emptyset\}$$

is finite. It follows that the action is proper, and that $\text{Out}(F_g) \backslash \mathcal{S}_g^{\text{na}}$ is Hausdorff.

Let us now prove the last part of the statement. Let $x \in \mathcal{S}_g^{\text{na}}$. The previous result applied with $y = x$ ensures that there exists an open neighborhood U_x of x such that

$$T := \{\tau \in \text{Out}(F_g) : \tau(U_x) \cap U_x \neq \emptyset\}$$

is finite. Up to shrinking U_x , we may assume that $T = \text{Stab}(x)$ and that U_x is stable under $\text{Stab}(x)$. We then have a canonical isomorphism $\text{Out}(F_g) \backslash \mathcal{S}_g^{\text{na}} \simeq \text{Stab}(x) \backslash \mathcal{S}_g^{\text{na}}$. The result follows. \square

Corollary 5.3.8. *For each $a \in \mathcal{M}(\mathbb{Z})$, the quotient space $\text{Out}(F_g) \backslash (\mathcal{S}_g \cap \text{pr}_{\mathbb{Z}}^{-1}(a))$ inherits a structure of $\mathcal{H}(a)$ -analytic space.*

Proof. Let $(k, |\cdot|)$ be a complete valued field. Recall that, if X is a k -analytic space and G a finite group acting on X , then the quotient $G \backslash X$ inherits a structure of k -analytic space. The archimedean case reduces to the case of complex analytic spaces, which is handled in [Car57, Théorème 4]. The non-archimedean case is a consequence of [BGR84, Proposition 6.3.3/3].

With these results at hand, the statement follows from Theorem 5.3.1 and Corollary 5.3.7. \square

Remark 5.3.9. Even though the results of this section mostly apply to the local situation, we believe that they should hold globally, which is to say the action of $\text{Out}(F_g)$ on \mathcal{S}_g is proper and the quotient space $\text{Out}(F_g) \backslash \mathcal{S}_g$ inherits a structure of an analytic space over \mathbb{Z} .

6. SCHOTTKY UNIFORMIZATION FOR FAMILIES OF CURVES

6.1. The universal Mumford curve over \mathbb{Z} .

Definition 6.1.1. We call *universal Schottky group* the following subgroup of $\mathrm{PGL}_2(\mathcal{O}(\mathcal{S}_g))$:

$$\mathcal{G}_g := \langle M(0, \infty, Y_1), M(1, X'_2, Y_2), \dots, M(X_g, X'_g, Y_g) \rangle.$$

For $x \in \mathcal{S}_g$, we denote by $L_x \subseteq \pi^{-1}(x) \simeq \mathbb{P}_{\mathcal{H}(x)}^{1, \mathrm{an}}$ the limit set of Γ_x . We call *limit set* of \mathcal{G}_g the set

$$\mathcal{L}_g := \bigcup_{x \in \mathcal{S}_g} L_x \subseteq \mathbb{P}_{\mathcal{S}_g}^1.$$

We set

$$\Omega_g := \mathbb{P}_{\mathcal{S}_g}^1 - \mathcal{L}_g.$$

Theorem 6.1.2. *The limit set \mathcal{L}_g of \mathcal{G}_g is a closed subset of $\mathbb{P}_{\mathcal{S}_g}^1$, the action of \mathcal{G}_g on its complement Ω_g is free and proper and the quotient map $\mathcal{G}_g \backslash \Omega_g \rightarrow \mathcal{S}_g$ is proper.*

Proof. It is enough to prove that every $x \in \mathcal{S}_g$ admits a neighborhood U_x such that $\mathcal{L}_g \cap \pi^{-1}(U_x)$ is a closed subset of $\mathbb{P}_{U_x}^1$, the action of \mathcal{G}_g on $\Omega_g \cap \pi^{-1}(U_x)$ is free and proper and the quotient map $\mathcal{G}_g \backslash (\Omega_g \cap \pi^{-1}(U_x)) \rightarrow U_x$ is proper. Let $x \in \mathcal{S}_g$.

Assume that x is archimedean. Arguing as in Remark 4.2.3 and the proof of Lemma 4.2.4, we may replace \mathcal{S}_g by the complex Schottky space $\mathcal{S}_{g, \mathbb{C}}$. We will add subscripts \mathbb{C} to denote the various objects under consideration in this setting.

By [Ber75, Proposition 3], $\mathcal{L}_{g, \mathbb{C}}$ is closed. The main ingredient in the proof is the existence of an open neighborhood U_0 of 0 in \mathbb{C}^{3g-3} , a neighborhood U of x in $\mathcal{S}_{g, \mathbb{C}}$, an analytic isomorphism $\xi: U_0 \rightarrow U$, and Beltrami coefficients $\mu_1, \dots, \mu_{3g-3}: \mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{C}$ such that, for each $s \in U_0$, we have

$$M_i(\xi(s)) = w^{s\mu} M_i(x) (w^{s\mu})^{-1} \text{ for } 1 \leq i \leq 3g-3,$$

where $w^{s\mu}$ denotes the unique quasiconformal automorphism of $\mathbb{P}^1(\mathbb{C})$ with Beltrami coefficient $s\mu = \sum_{i=1}^{3g-3} s_i \mu_i$ that fixes 0, 1 and ∞ . One may then relate the domain of discontinuity $\Omega_{g, x}$ over x to that over $\xi(s)$, for some $s \in U_0$, by $\Omega_{g, \xi(s)} = w^{s\mu}(\Omega_{g, x})$. It now follows from the continuity properties of $w^{s\mu}$ (see [AB60, Theorem 8]) that Ω_g is open, or, equivalently, that \mathcal{L}_g is closed.

Similarly, let F_x^+ be a compact subset of $\Omega_{g, x}$ intersecting every orbit of the action of $\mathcal{G}_{g, x}$. Let V_0 be a compact neighborhood of 0 in \mathbb{C}^{3g-3} . Then, the set

$$F^+ := \bigcup_{s \in V_0} w^{s\mu}(V_0)$$

is a compact subset of Ω_g that intersects every orbit of the action of $\mathcal{G}_{g, \xi(s)}$ for $s \in V_0$. It follows that $\mathcal{G}_g \backslash (\Omega_g \cap \pi^{-1}(V_0))$ is compact.

Finally, the group $\mathcal{G}_{g, \mathbb{C}}$ acts on $\Omega_{g, \mathbb{C}}$ by preserving the fibers of the morphism $\Omega_{g, \mathbb{C}} \rightarrow \mathcal{S}_{g, \mathbb{C}}$. Since its action is free and proper on each fiber, it is free and proper on the whole $\Omega_{g, \mathbb{C}}$.

Assume that x is non-archimedean. The result then follows from Corollary 4.3.3 and Proposition 3.3.2. \square

It follows from Theorem 6.1.2 that the quotient

$$\mathcal{C}_g := \Omega_g / \mathcal{G}_g$$

defines an analytic space over \mathbb{Z} . We call it the *universal Mumford curve* over \mathbb{Z} . This definition is motivated by the following corollary, summarizing the results of this section.

Corollary 6.1.3. *There is a commutative diagram in the category of analytic spaces over \mathbb{Z} :*

$$\begin{array}{ccc}
 \Omega_g & & \\
 \downarrow \pi & \searrow & \\
 & & \mathcal{C}_g \\
 & \swarrow & \\
 \mathcal{S}_g & &
 \end{array}$$

where the quotient map $\Omega_g \rightarrow \mathcal{C}_g$ is a local isomorphism and the natural map $\mathcal{C}_g \rightarrow \mathcal{S}_g$ is proper and smooth of relative dimension 1.⁴ Given a point $x \in \mathcal{S}_g$, its preimage in \mathcal{C}_g is a curve over $\mathcal{H}(x)$ which is isomorphic to the curve uniformized by the Schottky group Γ_x .

Remark 6.1.4. Note that the adjective *universal* applied to the curve \mathcal{C}_g does not bear any meaning beyond that of Corollary 6.1.3. In fact, while the universal Schottky group of Definition 6.1.1 corresponds to a universal family for the Schottky space, the universal Mumford curve does not correspond to such a family on a suitable moduli space of Mumford curves, at least not without additional structure. Indeed, the same Mumford curve can be obtained through different specializations of the universal Mumford curve.

6.2. Moduli spaces of Mumford curves. The existence of the curve \mathcal{C}_g raises the question of the existence of a moduli space of Mumford curves and its connections with the moduli space of stable curves. Over a non-archimedean fiber this space is obtained as the quotient of \mathcal{S}_g by the action of $\text{Out}(F_g)$ described in section 5.1, which was previously known in the rigid analytic context from work of L. Gerritzen [Ger81] and F. Herrlich [Her84], among others.

Remark 6.2.1. Over an archimedean fiber, the quotient of \mathcal{S}_g by the action of $\text{Out}(F_g)$ captures more than isomorphism classes of Riemann surfaces, as there are different Schottky groups that uniformize the same complex curve. In the case $g = 1$, one gets a trivial action, and the space $\mathcal{S}_{1,\mathbb{C}}$ is the punctured open unit disc over \mathbb{C} , image of the Poincaré open-half plane under the map $z \mapsto e^{2i\pi z}$. As such, $\mathcal{S}_{1,\mathbb{C}}$ is a cover of the modular curves $X_0(N)$ and $X_1(N)$ for every N . Hence a point of $\mathcal{S}_{1,\mathbb{C}}$ not only determines an isomorphism class of an elliptic curve, but also takes into account its N -level structures for every N . It would be interesting to investigate whether the space $\text{Out}(F_g) \backslash \mathcal{S}_g$ over the complex numbers can bear a similar interpretation for curves of higher genus.

In what follows, we consider only the non-archimedean case: let $a \in \mathcal{M}(\mathbb{Z})$ be a non-archimedean point, and consider the fiber $\mathcal{S}_{g,a} = \mathcal{S}_g \cap \text{pr}_{\mathbb{Z}}^{-1}(a)$ over a of the Schottky space. We denote by $\text{Mumf}_{g,a}$ the quotient of $\mathcal{S}_{g,a}$ by the continuous action of $\text{Out}(F_g)$ defined in 5.1.1. This quotient is a $\mathcal{H}(a)$ -analytic space by Corollary 5.3.8. Since any element of $\text{Out}(F_g)$ acts on the marking but does not affect the conjugacy class of a Schottky group, each point $x \in \text{Mumf}_{g,a}$ corresponds to a unique conjugacy class of a Schottky group $\Gamma_x \subset \text{PGL}_2(\mathcal{H}(x))$. Moreover, by [Mum72, Corollary (4.11)], the isomorphism class of a Mumford curve over a non-archimedean field determines the conjugacy class in PGL_2 of its Schottky group. This shows that, for every valued extension k of $\mathcal{H}(a)$ the k -points of $\text{Mumf}_{g,a} \times_{\mathcal{H}(a)} k$ are in 1-to-1 correspondence with isomorphism classes of Mumford curves of genus g defined over k .

6.2.1. Relationship with geometric group theory and tropical moduli. The existence of a faithful action of $\text{Out}(F_g)$ on \mathcal{S}_g with finite stabilizers is reminiscent of Culler-Vogtmann definition of the *outer space* in the context of geometric group theory, as introduced in their seminal paper [CV86].

⁴The property “smooth of relative dimension 1” has been recently defined in [Ber22, Définition 9.2]. More concretely, in our case the fibration ψ is locally isomorphic to the relative line.

This is not a coincidence, and in this section we show that we can indeed relate the topology of \mathcal{S}_g with that of the outer space.

Let us recall the definition of the Culler-Vogtmann outer space. We fix $g \geq 2$ and an abstract graph R_g with one vertex and g edges, identifying its fundamental group $\pi_1(R_g)$ with F_g . A finite connected graph G is said to be *stable* if all its vertices have degree ≥ 3 . A *marking* on a stable graph G of Betti number g is a homotopy equivalence $m : R_g \rightarrow G$ or, equivalently, the choice of a group isomorphism between F_g and the fundamental group $\pi_1(G)$. Two pairs (G, m) and (G', m') each consisting of a stable metric graph and a marking are *equivalent* if there is an isometry $s : G \rightarrow G'$ such that $s \circ m$ is homotopic to m' . For a given marked graph (G, m) , the isomorphism $F_g \cong \pi_1(G)$ determines an action of F_g on the universal cover T of G , a tree naturally endowed with a metric, denoted by d_T . The *translation length function* of (G, m) is the function $\ell_G : F_g \rightarrow \mathbb{R}$ associating to any $\sigma \in F_g$ the quantity $\ell_G(\sigma) := \min_{x \in T} \{d_T(\sigma(x), x)\}$. Let CV_g denote the set of equivalence classes of stable marked graphs endowed with a metric such that the sum of edge lengths is equal to 1, and let \mathcal{C} denote the set of conjugacy classes in F_g . The rule associating with a marked tree its translation length function defines an embedding $CV_g \hookrightarrow \mathbb{R}^{\mathcal{C}}$ into the infinite dimensional real vector space $\mathbb{R}^{\mathcal{C}}$. Thanks to this fact, CV_g inherits a topology from the product topology on $\mathbb{R}^{\mathcal{C}}$. The topological space so obtained is called the *Culler-Vogtmann outer space*, and it is also denoted by CV_g .

The original definition of the outer space can be found in [CV86, §0], where more details about the length functions and the topology of the outer space are given. In what follows, it will be useful to drop the condition that the marked graphs have sum of edge lengths equal to 1. We will then denote by CV'_g the *unprojectivized outer space* $CV_g \times \mathbb{R}_{>0}$, which parametrizes marked graphs with arbitrary edge lengths. There is a natural continuous action of $\text{Out}(F_g)$ on CV_g , which extends to CV'_g using the trivial action on the factor $\mathbb{R}_{>0}$. The quotient space $CV'_g / \text{Out}(F_g)$ has a canonical injection in the moduli space of abstract weighted tropical curves M_g^{trop} , whose image is given by those tropical curves that have weight zero at every vertex. The induced map $CV'_g \rightarrow M_g^{\text{trop}}$ is continuous and corresponds to forgetting the marking on a given metric graph. For more details about equivalent definitions of M_g^{trop} and its properties, we refer to [BMV11, §3], while a comparison between CV_g and M_g^{trop} is discussed in [Cap13, §5.2].

An isomorphism of Mumford curves induces an isometry between their skeletons. This allows to define a continuous function $\text{Mumf}_{g,a} \rightarrow M_g^{\text{trop}}$ sending (the class of) a Mumford curve in (the class of) its skeleton. For a point $x \in \text{Mumf}_{g,a}$, recall that the Schottky uniformization $(\mathbb{P}_{\mathcal{H}(x)}^{1,\text{an}} - \mathcal{L}_x) \rightarrow \mathcal{C}_x$ restricts to a universal cover of the skeleton Σ_x (see [PT21, Theorem 6.4.18]). Via this restriction, the Schottky group $\Gamma_x \cong \pi_1(\mathcal{C}_x)$ can be identified with the topological fundamental group $\pi_1(\Sigma_x)$.

Theorem 6.2.2. *There is a continuous surjective function*

$$\phi : \mathcal{S}_{g,a} \longrightarrow CV'_g \times_{M_g^{\text{trop}}} \text{Mumf}_{g,a}.$$

Proof. Let us consider the following:

- The continuous function $\phi_1 : \mathcal{S}_{g,a} \rightarrow \text{Mumf}_{g,a}$ given by the quotient by the action of $\text{Out}(F_g)$. Note that continuity descends from Proposition 5.1.5;
- The continuous function $\phi_2 : \mathcal{S}_{g,a} \rightarrow CV'_g$ given by assigning to each $y \in \mathcal{S}_{g,a}$ the metric graph corresponding to the skeleton Σ_y of the Mumford curve \mathcal{C}_y and the marking as follows: recall from Lemma 4.2.5 that the point y can be identified with the conjugacy class of a morphism $\varphi_y : F_g \hookrightarrow \text{PGL}_2(\mathcal{H}(y))$, whose image is the fundamental group $\pi_1(\mathcal{C}_y)$, and associate with y the marking corresponding to the isomorphism $F_g \cong \Gamma_y$ induced by φ_y . To prove continuity for ϕ_2 , we prove that the composite function $\mathcal{S}_{g,a} \rightarrow \mathbb{R}^{\mathcal{C}}$ is continuous. This amounts to prove that the following: if $\sigma \in \text{Aut}(F_g)$ is defined by $\sigma(e_i) = e_{j_i,0}^{n_{i,0}} \cdots e_{j_i,r_i}^{n_{i,r_i}}$, for

some $r_i \in \mathbb{N}$, $j_{i,0}, \dots, j_{i,r_i} \in \{1, \dots, g\}$, $n_{i,0}, \dots, n_{i,r_i} \in \mathbb{Z}$, the assignment

$$y \mapsto \ell_{\Sigma_y}(M_{j_{i,0}}(y)^{n_{i,0}} \dots M_{j_{i,r_i}}(y)^{n_{i,r_i}})$$

defines a continuous function $L : \mathcal{S}_{g,a} \rightarrow \mathbb{R}$. By Lemma 3.4.1, the length $\ell_{\Sigma_y}(M)$ for $M \in \Gamma_y$ coincides with $|\beta|^{-1}$, where β is the multiplier of M . The result then follows from Proposition 3.4.3, that ensures that the multiplier of the element $M_{j_{i,0}}(y)^{n_{i,0}} \dots M_{j_{i,r_i}}(y)^{n_{i,r_i}}$ is a continuous function in the Koebe coordinates of y .

The function ϕ_2 is $\text{Out}(F_g)$ -equivariant, and then agrees with ϕ_1 on M_g^{trop} . By the universal property of the fiber product, the pair (ϕ_1, ϕ_2) defines a continuous function

$$\phi : \mathcal{S}_{g,a} \longrightarrow CV'_g \times_{M_g^{\text{trop}}} \text{Mumf}_{g,a}.$$

We now prove that the function ϕ is surjective. Let $([G, m], [C]) \in CV'_g \times_{M_g^{\text{trop}}} \text{Mumf}_{g,a}$ be a pair consisting of an equivalence class of a marked graph and an isomorphism class of a Mumford curve of genus g , such that the graph G is isometric to the skeleton of C . We fix an isometry between G and the skeleton of C , inducing an isomorphism $j : \pi_1(G) \xrightarrow{\sim} \pi_1(C)$. Let us denote by y the point of $\mathcal{S}_{g,a}$ whose underlying Schottky group is $\pi_1(C)$, with marking given by the image of the basis of F_g under the composition of isomorphisms $F_g \xrightarrow{m} \pi_1(G) \xrightarrow{j} \pi_1(C)$. Then $\phi_1(y) = [C]$ and $\phi_2(y) = [G, m]$. Hence ϕ is surjective. \square

Remark 6.2.3. In the proof of the surjectivity of the continuous function ϕ above, a different choice of isometry j between G and the fundamental group $\pi_1(C)$ might determine a different preimage in $\mathcal{S}_{g,a}$ of the pair $([G, m], [C])$. For example, when G is a rose with g loops all of the same length, a permutation of the loops corresponds to a permutation of the basis $\{\gamma_1, \dots, \gamma_g\}$ of the Schottky group $\pi_1(C)$. In most cases the element of $\text{Out}(F_g)$ corresponding to such a permutation does not stabilize a point in $\mathcal{S}_{g,a}$, for instance when two distinct elements $\gamma_i \neq \gamma_j$ have distinct multipliers $\beta_i \neq \beta_j$. This shows in particular that the function ϕ is not injective.

Remark 6.2.4. In [Uli21], Ulirsch constructs a non-archimedean analogue of Teichmüller space $\overline{\mathcal{T}}_g$, using the tropical Teichmüller space that Chan, Melo, and Viviani introduced in [CMV13] and tools from logarithmic geometry. The space $\overline{\mathcal{T}}_g$ is a Deligne-Mumford analytic stack over a non-archimedean algebraically closed field k whose points morally correspond to pairs (C, ϕ) consisting of a stable projective curve C over a valued extension of k and an isomorphism $\phi : \pi_1^{\text{top}}(C^{\text{an}}) \cong F_{b(C)}$, where $b(C)$ is the first Betti number of C^{an} . When restricting this construction on the locus of Mumford curve, one retrieves a space $\mathcal{T}_g^{\text{Mum}}$, and a corollary of Ulirsch's construction is the realization of CV_g as a strong deformation retract of $\mathcal{T}_g^{\text{Mum}}$. Moreover, the fibered product $CV'_g \times_{M_g^{\text{trop}}} \text{Mumf}_{g,a}$ is identified (after a suitable base-change to an algebraically closed field) with the locus of Mumford curves inside the coarse moduli space of $\overline{\mathcal{T}}_g$. As a result, Theorem 6.2.2 and Remark 6.2.3 clarify the relationship between non-archimedean fibers of the Schottky space \mathcal{S}_g over \mathbb{Z} and Ulirsch's $\mathcal{T}_g^{\text{Mum}}$.

REFERENCES

- [AB60] Lars V. Ahlfors and Lipman Bers. Riemann’s mapping theorem for variable metrics. *Ann. Math. (2)*, 72:385–404, 1960.
- [Ber75] Lipman Bers. Automorphic forms for Schottky groups. *Adv. Math.*, 16:332–361, 1975.
- [Ber90] Vladimir G. Berkovich. *Spectral theory and analytic geometry over non-Archimedean fields*, volume 33 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1990.
- [Ber93] Vladimir G. Berkovich. Étale cohomology for non-Archimedean analytic spaces. *Inst. Hautes Études Sci. Publ. Math.*, (78):5–161 (1994), 1993.
- [Ber09] Vladimir G. Berkovich. A non-Archimedean interpretation of the weight zero subspaces of limit mixed Hodge structures. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I*, volume 269 of *Progr. Math.*, pages 49–67. Birkhäuser Boston Inc., Boston, MA, 2009.
- [Ber22] Dorian Berger. Espaces de Berkovich sur \mathbb{Z} : morphismes étales. arXiv, 2022. <https://arxiv.org/abs/2112.02907>.
- [BGR84] Siegfried Bosch, Ulrich Güntzer, and Reinhold Remmert. *Non-Archimedean analysis*, volume 261 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1984. A systematic approach to rigid analytic geometry.
- [BJ17] Sébastien Boucksom and Mattias Jonsson. Tropical and non-Archimedean limits of degenerating families of volume forms. *J. Éc. polytech. Math.*, 4:87–139, 2017.
- [BMV11] Silvia Brannetti, Margarida Melo, and Filippo Viviani. On the tropical Torelli map. *Advances in Mathematics*, 226(3):2546–2586, 2011.
- [Bou71] Nicolas Bourbaki. *Éléments de mathématique. Topologie générale. Chapitres 1 à 4*. Hermann, Paris, 1971.
- [Cap13] Lucia Caporaso. Algebraic and tropical curves: comparing their moduli spaces. In G. Farkas and I. Morrison, editors, *Handbook of Moduli, Volume I*, pages 119–160. Advanced Lectures in Mathematics, Volume XXIV, 2013.
- [Car57] Henri Cartan. Quotient d’un espace analytique par un groupe d’automorphismes. *Princeton Math. Ser.*, 12:90–102, 1957.
- [CMV13] Melody Chan, Margarida Melo, and Filippo Viviani. Tropical Teichmüller and Siegel spaces. In *Algebraic and combinatorial aspects of tropical geometry*, volume 589 of *Contemp. Math.*, pages 45–85. Amer. Math. Soc., Providence, RI, 2013.
- [CV86] Marc Culler and Karen Vogtmann. Moduli of graphs and automorphisms of free groups. *Invent. Math.*, 84(1):91–119, 1986.
- [DKY20] Laura DeMarco, Holly Krieger, and Hexi Ye. Uniform Manin-Mumford for a family of genus 2 curves. *Ann. Math. (2)*, 191(3):949–1001, 2020.
- [dSG10] Henri Paul de Saint-Gervais. *Uniformisation des surfaces de Riemann. Retour sur un théorème centenaire*. ENS Éditions, 2010. 544 pp.
- [Duc] Antoine Ducros. La structure des courbes analytiques. Manuscript available at <http://www.math.jussieu.fr/~ducros/trirss.pdf>.
- [Fav20] Charles Favre. Degeneration of endomorphisms of the complex projective space in the hybrid space. *J. Inst. Math. Jussieu*, 19(4):1141–1183, 2020.
- [Gar87] Frederik P. Gardiner. *Teichmüller theory and quadratic differentials*. John Wiley & Sons, New York, 1987.
- [Ger74] Lothar Gerritzen. Zur nichtarchimedischen Uniformisierung von Kurven. *Math. Ann.*, 210:321–337, 1974.
- [Ger81] Lothar Gerritzen. Zur analytischen Beschreibung des Raumes der Schottky-Mumford-Kurven. *Math. Ann.*, 255(2):259–271, 1981.
- [Ger82] Lothar Gerritzen. p -adic Teichmüller space and Siegel halfspace. *Groupe de travail d’analyse ultramétrique*, 9(2), 1981-1982. talk:26.
- [GH88] L. Gerritzen and F. Herrlich. The extended Schottky space. *J. reine angew. Math.*, 389:190–208, 1988.
- [GvdP80] Lothar Gerritzen and Marius van der Put. *Schottky groups and Mumford curves*, volume 817 of *Lecture Notes in Mathematics*. Springer, Berlin, 1980.
- [Har88] David Harbater. Galois covers of an arithmetic surface. *Amer. J. Math.*, 110(5):849–885, 1988.
- [Hej75] Dennis A Hejhal. On Schottky and Teichmüller spaces. *Advances in Mathematics*, 15(2):133 – 156, 1975.
- [Her84] Frank Herrlich. On the stratification of the moduli space of Mumford curves. *Groupe de travail d’analyse ultramétrique*, 11:1–10, 1983-1984.
- [Her91] Frank Herrlich. Moduli for stable marked trees of projective lines. *Mathematische Annalen*, 291(1):643–661, 1991.
- [Hid05] Rubén A. Hidalgo. Automorphism groups of Schottky type. *Ann. Acad. Sci. Fenn., Math.*, 30(1):183–204, 2005.

- [HS07] Frank Herrlich and Gabriela Schmithüsen. On the boundary of Teichmüller disks in Teichmüller and in Schottky space. In *Handbook of Teichmüller theory I*, pages 293–349. European Mathematical Society, 2007.
- [Ich00] Takashi Ichikawa. Generalized Tate curve and integral Teichmüller modular forms. *Amer. J. Math.*, 122(6):1139–1174, 2000.
- [Ich01] Takashi Ichikawa. Universal periods of hyperelliptic curves and their applications. *Journal of Pure and Applied Algebra*, 163(3):277–288, 2001.
- [Ich18] Takashi Ichikawa. Klein’s formulas and arithmetic of Teichmüller modular forms. *Proceedings of the American Mathematical Society*, 146(12):5105–5112, sep 2018.
- [Ich20a] Takashi Ichikawa. Periods of generalized Tate curves. arXiv, 2020. <https://arxiv.org/abs/1909.10149>.
- [Ich20b] Takashi Ichikawa. The universal Mumford curve and its periods in arithmetic formal geometry. arXiv, 2020. <https://arxiv.org/abs/2010.11517>.
- [JMM79] T. Jørgensen, A. Marden, and B. Maskit. The boundary of classical Schottky space. *Duke Mathematical Journal*, 46(2):441–446, 1979.
- [LP20] Thibaud Lemanissier and Jérôme Poineau. Espaces de Berkovich sur \mathbf{Z} : catégorie, topologie, cohomologie. arXiv, 2020. <https://arxiv.org/abs/2010.08858>.
- [Mar74a] Albert Marden. The geometry of finitely generated kleinian groups. *Ann. of Math.*, 99:383–462, 1974.
- [Mar74b] Albert Marden. Schottky groups and circles. In *Contributions to Analysis*, pages 273–278. Academic Press, New York and London, 1974.
- [Mas67] Bernard Maskit. A characterization of Schottky groups. *Journal d’Analyse Mathématique*, 19(1):227–230, dec 1967.
- [MSW15] David Mumford, Caroline Series, and David Wright. *Indra’s pearls. The vision of Felix Klein*. Cambridge University Press, Cambridge, 2015.
- [Mum72] David Mumford. An analytic construction of degenerating curves over complete local rings. *Compositio Math.*, 24:129–174, 1972.
- [Poi10a] Jérôme Poineau. La droite de Berkovich sur \mathbf{Z} . *Astérisque*, (334):xii+284, 2010.
- [Poi10b] Jérôme Poineau. Raccord sur les espaces de Berkovich. *Algebra Number Theory*, 4(3):297–334, 2010.
- [Poi13] Jérôme Poineau. Espaces de Berkovich sur \mathbf{Z} : étude locale. *Invent. Math.*, 194(3):535–590, 2013.
- [PT21] Jérôme Poineau and Daniele Turchetti. Berkovich curves and Schottky uniformization. II: Analytic uniformization of Mumford curves. In *Arithmetic and geometry over local fields. VIASM 2018. Based on lectures given during the program “Arithmetic and geometry of local and global fields”, summer 2018, Hanoi, Vietnam*, pages 225–279. Cham: Springer, 2021.
- [Ser77] Jean-Pierre Serre. *Arbres, amalgames, SL_2* . Astérisque, No. 46. Société Mathématique de France, Paris, 1977. Avec un sommaire anglais, Rédigé avec la collaboration de Hyman Bass.
- [Uli21] Martin Ulirsch. A non-Archimedean analogue of Teichmüller space and its tropicalization. *Selecta Mathematica*, 27(3):39, 2021.
- [Vog14] Karen Vogtmann. On the geometry of outer space. *Bulletin of the American Mathematical Society*, 52:27–46, 2014.
- [Wer16] Annette Werner. Analytification and tropicalization over non-Archimedean fields. In Matt Baker and Sam Payne, editors, *Nonarchimedean and Tropical Geometry*, pages 145–171. Springer, 2016.
- [Yam91] Hiro-o Yamamoto. An example of a nonclassical Schottky group. *Duke Mathematical Journal*, 63(1):193 – 197, 1991.

UNIVERSITÉ DE CAEN - NORMANDIE
E-mail address: jerome.poineau@unicaen.fr
URL: <https://poineau.users.lmno.cnrs.fr/>

UNIVERSITY OF WARWICK
E-mail address: daniele.turchetti.math@gmail.com
URL: <https://danieleturchetti.github.io/>