

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/178842>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Phase Transitions in Biased Opinion Dynamics with 2-choices Rule

Arpan Mukhopadhyay

Department of Computer Science
University of Warwick

Abstract

We consider a model of binary opinion dynamics where one opinion is inherently ‘superior’ than the other and social agents exhibit a ‘bias’ towards the superior alternative. Specifically, it is assumed that an agent updates its choice to the superior alternative with probability $\alpha > 0$ irrespective of its current opinion and the opinions of the other agents. With probability $1 - \alpha$ it adopts the majority opinion among two randomly sampled neighbours and itself. We are interested in the time it takes for the network to converge to a consensus on the superior alternative. In a complete graph of size n , we show that irrespective of the initial configuration of the network, the average time to reach consensus scales as $\Theta(n \log n)$ when the bias parameter α is sufficiently high, i.e., $\alpha > \alpha_c$ where α_c is a threshold parameter that is uniquely characterised. When the bias is low, i.e., when $\alpha \in (0, \alpha_c]$, we show that the same rate of convergence can only be achieved if the initial proportion of agents with the superior opinion is above certain threshold $p_c(\alpha)$. If this is not the case, then we show that the network takes $\Omega(\exp(\Theta(n)))$ time on average to reach consensus.

Keywords— Biased opinion dynamics, 2-choices rule, phase transitions

1 Introduction

Opinion dynamical models are used extensively in statistical physics and computer science to study the effects of different local interaction rules on the adoption of new technologies and products. One key question in this context is how fast can a new/superior technology replace an old/outdated technology in a network of connected agents? Classical opinion dynamical models with two competing opinions (or technologies) assume the opinions to be indistinguishable; indeed under both voter and majority rule models, agents update their opinions purely based on the opinions of other agents in their neighbourhoods without exhibiting any preference for any opinion. However, to capture the inherent superiority of one opinion over another, we need to incorporate some form of “bias” towards one of the two opinions.

Opinion dynamical models with bias have been studied recently in [1, 2, 3, 4]. A strong form of bias is considered in [3, 4]; a bias parameter $\alpha \in (0, 1)$ is introduced; with probability α an agent is assumed to perform a *biased update* in which it adopts the superior opinion independently of its current opinion and the opinions of all other agents; with probability $1 - \alpha$ the agent performs a *regular update* in which it adopts the majority opinion among *all* agents in its neighbourhood. The later update rule is often referred to as the *majority rule* in the literature. Clearly, in this model, any network will eventually reach consensus on the superior opinion. It is shown in [3] that the expected time taken for the network to reach consensus on the superior opinion is exponential in the minimum degree of the underlying graph. Thus, in graphs where the neighbourhood size of each agent is proportional to the network size, the consensus time grows exponentially

with the network size. This naturally prompts the question if simpler rules exist under which consensus can be achieved faster.

To address this question, we consider a simpler rule called the *2-choices rule* [5] for *regular updates*. Under the 2-choices rule, an agent samples two other agents from its neighbourhood and updates to the majority opinion among the sampled agents and the agent itself. For graphs where the neighbourhood sizes are large, this modified rule greatly reduces the communication overhead associated with computing the majority opinion since an agent no longer needs to know the opinions of all other agents in its neighbourhood; only knowing the opinions of the two sampled agents suffices. The 2-choices rule also reduces the chance that an updating agent adopts the worse alternative when a majority of its neighbours have chosen this alternative. Hence, this modification should not only reduce the communication overhead, but also ‘facilitate’ consensus on the superior alternative. In this paper, we analytically characterise the improvement to the speed of consensus brought about by this modified rule.

Specifically, we show that if the network is a complete graph and the bias parameter α is sufficiently high ($\alpha > 1/9$), then consensus is achieved on the superior opinion in $\Theta(n \log n)$ time, where n denotes the number of agents in the network. When bias is small (i.e., when $\alpha < 1/9$) we show that the consensus time depends on the initial configuration of the network. More specifically, when the bias parameter α is small, fast consensus (i.e., $\Theta(n \log n)$ time) is achieved only if the initial proportion p of agents with the superior opinion is above a certain threshold value $p_c(\alpha)$ (explicitly characterised). If $p < p_c(\alpha)$ we show that the network takes exponentially long time to reach consensus. Thus, the speed with which the network reaches consensus on the superior opinion undergoes a sharp *phase transition* depending on the values of α and p . Through simulations, we observe similar behaviour on other classes of graphs, e.g., on random d -regular graphs both with $d = \Theta(\log n)$ and Erdős-Rényi graphs with edge probability $\Theta(\log n/n)$. To establish our theoretical results, we use a novel characterisation of the expected number of visits of a random walk to a given state using a branching process. We expect this technique to be useful in the analysis of other interacting particle system models. Thus, in summary, our contributions are the following:

- (Fast consensus) For complete graphs, we show the existence of a sharp threshold such that if the bias parameter α is above the threshold, then consensus is achieved on the superior opinion starting from any initial configuration in $\Theta(n \log n)$ time on average. If the bias parameter α is below the threshold, we show the existence of another sharp threshold such that if the initial proportion p of agents with the superior opinion is above this threshold, then consensus can be achieved in $\Theta(n \log n)$ time.
- (Slow consensus) We show that when both the bias parameter α and the initial proportion p of agents with the superior opinion are below their corresponding thresholds, the average consensus time on complete graphs is $\Omega(\exp(\Theta(n)))$, i.e., grows exponentially with the network size.
- (Other classes of graphs) Through extensive simulations we study consensus on other classes of graphs. Specifically, we observe similar behaviour on random d -regular graphs with $d = \Theta(\log n)$ and on Erdős-Rényi graphs with edge probability $\log n/n$. For d -regular graphs with constant degrees $d = O(1)$ we do observe a phase transition but the behaviour below criticality is different from that in complete graphs or other dense graphs studied in this paper.

1.1 Related Literature

The simplest model of opinion dynamics studied in the literature is the *voter model* where an agent simply copies the opinion of an agent sampled randomly from its neighbourhood. Thus, in the voter model, the probability that an agent adopts a specific opinion is equal to the proportion of agents having the same opinion in its neighbourhood. Due to the linearity of the resulting dynamics and its duality with coalescing random walks, the voter model has been extensively studied in the literature. The duality between the voter model and coalescing random walks was first observed independently in [6] and [7]. Using this duality, the model has been analysed on different classes of graphs such as regular lattices [8], random d -regular graphs [9], and ER graphs [10]. It is known that for connected graphs the probability of reaching consensus on a specific opinion is proportional to the initial volume (sum of degrees) of nodes having that opinion. It is also known that the mean consensus time for the asynchronous version of the voter model (where in each round only one randomly sampled agent updates its opinion) is $\Omega(n^2)$ for most graphs.

Another important model of opinion dynamics is the majority rule model wherein an agent adopts the majority opinion among all agents in its neighbourhood. It was shown in [11] that with high probability for a family of expander graphs with sufficiently large spectral gap, the majority dynamics leads to a consensus on the opinion having the initial majority, provided that the imbalance between the initial majority opinion and the alternate opinion is sufficiently high. Bounds on the consensus time for the majority rule model was obtained in [12] for expanders and Erdos-Renyi random graphs. For expanders with sufficiently large spectral gaps, it was shown that consensus can be achieved on the initial majority opinion in $O(\log n)$ steps in the synchronous model where all agents update in each round. For ER graphs, it was shown that if the edge probability is above the connectivity threshold of $\log n/n$, then consensus can be achieved on the initial majority opinion in constant number of rounds. The majority rule model has also been studied for other classes of graphs such as finite lattices [13], random regular graphs [14], and infinite trees [15].

Although the majority rule leads to faster consensus on many classes of graphs, it requires an agent to know the states of all other agents in its neighbourhood. This may be too computationally expensive when neighbourhood sizes are large. A simpler alternative is to consider the 2-choices rule where an agent only samples two random neighbours and changes to the majority opinion among the sampled agents and the agent itself. Rules similar to the 2-choices rule, where groups of agents are formed at each instant and all agents in the group update to the majority opinion within the group, were analysed in the physics literature [16, 17, 18]. A generalisation of 2-choices rule, where the updating agent samples m agents from its neighbourhood and only changes its opinion if d or more of the sampled agent differ from the updating agent, was analysed in the continuous time in [19]. The 2-choices rule that we consider in this paper was first analysed for random d -regular graphs and expanders in [5]. It was shown that consensus can be achieved in $O(\log n)$ time with high probability on the initial majority opinion provided that the initial imbalance is sufficiently high.

Opinion dynamical models with bias have been considered in the recent literature. These models are designed to capture the superiority of one alternative over another. Accordingly, in these models, the agents exhibit some form of bias towards the superior opinion. In [1, 2] a weak form of bias is considered. Here, agents with the superior opinion update with a lesser frequency than agents with the alternative opinion. It has been shown in [1] that, under the voter rule and the 2-choices rule, consensus is achieved in $O(\log n)$ time for complete graphs. For the voter model, the probability of achieving consensus on the superior opinion approaches to one as the network size grows. For the 2-choices rule, consensus is achieved on the superior opinion with high probability only when the initial proportion of agents with the superior opinion is above a certain threshold.

The form bias studied in this paper is introduced recently in [3, 20]. These papers show that on dense graphs the speed of consensus can be slow (depending on the minimum degree of a node) if the agents follow the majority rule during a regular update. The papers also analyse the voter rule under this form bias and show that consensus with the voter rule can be achieved in $O(n \log n)$ time. Our model is different from these models as we consider the 2-choices rule as the main update rule. Furthermore, we study the dynamics under the 2-choices rule as a function of the bias parameter α as well as the initial proportion p of agents with the superior opinion. This is unlike the previous papers [3, 20], where the dynamics is studied as a function of α only with a fixed value of p (specifically, $p = 0$). We show that phase transitions can occur with respect to both α and p . Similar phase transitions have been recently reported in [21, 22, 23] for noisy versions of the k -majority dynamics. However, the focus of these papers is a state of ‘near consensus’ where both opinions co-exist but the proportion of agents with one of the opinions is arbitrarily small. In contrast, our results focus on the state of full consensus in which the inferior opinion is completely eliminated. Furthermore, we obtain bounds on the mean of the consensus time that are stronger than high probability bounds obtained in previous papers. The technique used here is also quite different from those used in the earlier works; while earlier works use concentration around the mean drift, we use suitably constructed branching processes to obtain bounds on the number of visits to different states.

1.2 Organisation

The rest of the paper is organised as follows. In Section 2, we introduce the model studied in this paper. Next, in Section 3 we present the theoretical analysis of the model for complete graphs. Section 4 provides numerical results to support our theoretical findings and also demonstrates similar behaviour for other classes of graphs. Finally, we conclude the paper in Section 5.

2 Model

In this section, we describe the model studied in this paper. The model consists of a network of n agents described by an undirected graph $G_n = (V_n, E_n)$, where the nodes in V_n (with $|V_n| = n$) represent the agents and the edges in E_n represent the connections between the agents. For each agent (node) $u \in V_n$, we denote by $N_u = \{v : (u, v) \in E_n\}$, the set of neighbours of u .

Time is assumed to be discrete and at each discrete time instant $t \in \mathbb{Z}_+$ each agent is assumed to have an opinion in the set $\{0, 1\}$. Without loss of generality, we assume that 1 is the superior opinion. Let $X_u(t) \in \{0, 1\}$ denote the opinion of agent u at time t . At $t = 0$, the opinions of the agents are initialised such that $|\{u \in V_n : X_u(0) = 1\}| = \lceil pn \rceil$ for some $p \in [0, 1)$. Hence, p fraction of agents initially have the superior opinion 1.

At each instant $t \geq 0$, an agent, sampled uniformly at random, updates its opinion: with probability α , it performs a *biased update* in which it adopts the superior opinion irrespective of its current opinion and the opinions of all other agents; with probability $1 - \alpha$ it performs a *regular update* following the *2-choices rule* in which the updating agent samples two neighbours uniformly at random (with replacement¹) and adopts the majority opinion among the sampled agents and the agent itself. Therefore, if $U(t)$ denotes the randomly sampled agent at time t , then at time $t + 1$ the opinion of the agent is given by

$$X_{U(t)}(t + 1) = \begin{cases} 1 & \text{w.p. } \alpha, \\ M(t) & \text{w.p. } 1 - \alpha, \end{cases} \quad (1)$$

where $M(t) = \mathbb{1}(X_{U(t)}(t) + X_{N_1(t)}(t) + X_{N_2(t)}(t) \geq 2)$ denotes the majority opinion among two randomly sampled neighbours $N_1(t)$ and $N_2(t)$ of $U(t)$ and the agent $U(t)$ itself. The parameter $\alpha \in (0, 1]$ represents the bias towards the superior opinion and is referred to as the *bias parameter* of the model.

The state of the network at any time $t \geq 0$ can be represented by the vector $\mathbf{X}(t) = (X_u(t), u \in V_n)$ of opinions of all the agents. The process $\mathbf{X}(\cdot)$ is Markov on the state space $\{0, 1\}^n$ with a single absorbing state $\mathbf{1}$ where all agents have opinion 1. We refer to this absorbing state as the *consensus state*. Since it is possible to reach the consensus state from any other state in a finite number of steps, with probability one the chain is absorbed in the consensus state in a finite time. We refer to this time as the *consensus time*. The objective of the rest of the paper is to analyse the mean consensus time for different values of the parameters n, α, p and for different classes of graphs.

3 Analysis for complete graphs

In this section, we assume that G_n is a complete graph on n nodes. For complete graphs, the process $\bar{X}^n = (\bar{X}^n(t) = |\{u \in V_n : X_u(t) = 1\}|, t \geq 0)$ counting the number of agents with opinion 1 is a Markov chain on \mathbb{Z}_+ with absorbing state n . For the chain \bar{X}^n , the transition probability $\tilde{p}_{i,j}$ from state i to j is given by $\tilde{p}_{i,j} = p_{i,j} + o(1/n)$ where

¹Note that sampling with or without replacement does not make any difference when the neighbourhood size is proportional to n since the probability of choosing the same neighbour twice tends to zero as $n \rightarrow \infty$.

$$p_{i,j} = \begin{cases} \left(1 - \frac{i}{n}\right) \left(\alpha + (1-\alpha) \left(\frac{i}{n}\right)^2\right), & \text{if } j = i + 1, \\ \frac{i}{n} (1-\alpha) \left(1 - \frac{i}{n}\right)^2, & \text{if } j = i - 1, \\ 1 - p_{i,i+1} - p_{i,i-1}, & \text{if } j = i, \\ 0, & \text{otherwise.} \end{cases} \quad (2)$$

Note that $p_{i,j}$ denotes the transition probability of a slightly modified chain \bar{Y}^n (with the same absorbing state n) where an agent during a regular update can sample itself in addition to its neighbours. Below we analyse this modified chain as it is simpler to do so and the asymptotic (in n) results we obtain for this modified chain also hold for the original chain (see Remark 2 for more explanation).

Below we characterise the scaling law of $\bar{T}_n(p) = \mathbb{E}_{\lceil pn \rceil} [T_n]$, where $\mathbb{E}_x [\cdot]$ denotes expectation conditioned on $\bar{Y}^n(0) = x$ and $T_k = \inf \{t \geq 0 : \bar{Y}^n(t) = k\}$ denotes the first time the network reaches state k . Our main result is described in Theorem 1 below.

Theorem 1. *For complete graphs we have the following*

1. (Fast consensus) *For each $\alpha \in (1/9, 1)$, we have $\bar{T}_n(p) = \Theta(n \log n)$ for all $p \in [0, 1)$.*
2. (Fast consensus) *For each $\alpha \in (0, 1/9)$, there exists $p_c(\alpha) \in (0, 1)$ such that if $p \in [p_c(\alpha), 1)$, then $\bar{T}_n(p) = \Theta(n \log n)$. Furthermore, the threshold $p_c(\alpha)$ is the unique solution in the range $(\bar{x}_\alpha, 1)$ of the equation*

$$\int_{\bar{x}_\alpha}^{p_c(\alpha)} \log(f_\alpha(x)) dx = 0, \quad (3)$$

where $f_\alpha(x) = \frac{(1-\alpha)x(1-x)}{\alpha+(1-\alpha)x^2}$, $\bar{x}_\alpha = \frac{1}{4} \left(1 - \sqrt{1 - \frac{8\alpha}{1-\alpha}}\right)$ and $\bar{x}_\alpha = \frac{1}{4} \left(1 + \sqrt{1 - \frac{8\alpha}{1-\alpha}}\right)$.

3. (Slow consensus) *For each $\alpha \in (0, 1/9)$ and $p \in [0, p_c(\alpha))$, we have $\bar{T}_n(p) = \Omega(\exp(\Theta(n)))$.*

The above theorem implies that when the bias parameter is sufficiently high ($\alpha > 1/9$), the network quickly (in $\Theta(n \log n)$ time) reaches consensus on the superior opinion irrespective of its initial state. This is in sharp contrast to the result of [20] where the consensus time is exponential in n for all values of the bias parameter α . The theorem further implies that even with low value of the bias parameter α (for $\alpha < 1/9$) fast consensus can be achieved as long as the initial proportion p of agents with the superior opinion is above a threshold value denoted by $p_c(\alpha)$. We explicitly characterise this threshold $p_c(\alpha)$ required to ensure fast consensus when the bias is low. If the bias parameter is low ($\alpha < 1/9$) and the initial proportion of agents with the superior opinion is below the threshold, i.e., $p < p_c(\alpha)$, then the mean consensus time is exponential in n which corresponds to slow speed of convergence. Thus, our model exhibits rich behaviour in terms of the parameters α and p .

We breakdown the proof of the above theorem into several simpler steps. The first step is to make the following simple observation which expresses the mean consensus time as a function of the number of visits of the chain \bar{Y}^n to different states in its state-space $\{0, 1, \dots, n\}$.

Lemma 1. *Let Z_k denote the number of visits of the chain \bar{Y}^n to the state $k \in \{0, 1, \dots, n\}$ before absorption. Then, for any starting state $x = \bar{X}^n(0) \in \{0, 1, \dots, n\}$, the average consensus time is given by*

$$\mathbb{E}_x [T_n] = \sum_{k=0}^{n-1} \frac{\mathbb{E}_x [Z_k]}{(1 - p_{k,k})}, \quad (4)$$

where $p_{k,k}$ is the transition probability from state k to itself given by (2).

Proof. Observe that the consensus time T_n can be written as

$$T_n = \sum_{k=0}^{n-1} \sum_{j=1}^{Z_k} M_{k,j}, \quad (5)$$

where Z_k denotes the number of visits to state k before absorption and $M_{k,j}$ denotes the time spent in state k in the j^{th} visit. Clearly, the random variables Z_k and $(M_{k,j})_{j \geq 1}$ are independent of each other. Furthermore, $(M_{k,j})_{j \geq 1}$ is a sequence of i.i.d. random variables with geometric distribution given by $\mathbb{P}_x(M_{k,j} = i) = p_{k,k}^{i-1}(1 - p_{k,k})$ for all j . Hence, applying Wald's identity to (5) we have

$$\begin{aligned}
\bar{T}_n(p) &= \mathbb{E}_x [T_n] \\
&= \sum_{k=0}^{n-1} \mathbb{E}_x \left[\sum_{j=1}^{Z_k} M_{k,j} \right] \\
&= \sum_{k=0}^{n-1} \mathbb{E}_x [Z_k] \mathbb{E}_x [M_{k,j}] \\
&= \sum_{k=0}^{n-1} \frac{\mathbb{E}_x [Z_k]}{(1 - p_{k,k})}, \tag{6}
\end{aligned}$$

where the last step follows from the fact that $\mathbb{E}_x [M_{k,j}] = 1/(1 - p_{k,k})$ for each $j \in [Z_k]$. \square

From the above lemma, it is evident that in order to obtain bounds on $\mathbb{E}_x [T_n]$ we need to obtain bounds on the expected number of visits $\mathbb{E}_x [Z_k]$ to different states and the transition probabilities $p_{k,k}$. Using this approach, we first obtain a lower bound on $\bar{T}_n(p)$.

Lemma 2. *For any $\alpha \in (0, 1)$ and $p \in (0, 1)$ we have $\bar{T}_n(p) = \Omega(n \log n)$.*

Proof. From (2) we obtain

$$\begin{aligned}
1 - p_{k,k} &= \left(1 - \frac{k}{n}\right) \left(\alpha + (1 - \alpha) \frac{k}{n}\right) \\
&\leq \max(\alpha, 1 - \alpha) \left(1 - \frac{k}{n}\right) \left(1 + \frac{k}{n}\right)
\end{aligned}$$

Furthermore, we have $\mathbb{E}_x [Z_k] \geq \mathbb{1}(k \geq x)$ since states $k \geq x$ are visited at least once. Hence, using the above two inequalities in (4) we obtain

$$\begin{aligned}
\bar{T}_n(p) &\geq \sum_{k=0}^{n-1} \frac{\mathbb{1}(k \geq \lceil np \rceil)}{\max(\alpha, 1 - \alpha) \left(1 - \frac{k}{n}\right) \left(1 + \frac{k}{n}\right)} \\
&= \frac{n}{2 \max(\alpha, 1 - \alpha)} \sum_{k=\lceil np \rceil}^{n-1} \left(\frac{1}{n - k} + \frac{1}{n + k}\right) \\
&> \frac{n}{2 \max(\alpha, 1 - \alpha)} \sum_{k=\lceil np \rceil}^{n-1} \left(\frac{1}{n - k}\right) \\
&\geq \frac{n}{2 \max(\alpha, 1 - \alpha)} \log(n - \lceil np \rceil + 1),
\end{aligned}$$

which completes the proof. \square

To obtain an upper bound on the mean consensus time $\bar{T}_n(p)$ similarly, we need an upper bound on the expected number of visits $\mathbb{E}_x [Z_k]$ to state k for each $k \in \{0, 1, 2, \dots, n\}$. We obtain such an upper bound using a technique developed in [1] where the number of visits to each state is expressed as a function of a branching process. Specifically, we define ζ_k to be the number of jumps from state k to state $k - 1$ for

any $k \in \{1, 2, \dots, n\}$. Clearly, $\zeta_n = 0$ and, if the starting state is $x = \bar{Y}^n(0) \in [1, n)$, then ζ_k satisfies the following recursion:

$$\zeta_k = \begin{cases} \sum_{l=0}^{\zeta_{k+1}} \xi_{l,k}, & \text{if } k \in \{x, x+1, \dots, n-1\}, \\ \sum_{l=1}^{\zeta_{k+1}} \xi_{l,k}, & \text{if } k \in \{0, 1, \dots, x-1\}, \end{cases} \quad (7)$$

where $\xi_{l,k}$ denotes the number of jumps from state k to state $k-1$ between the l^{th} and $(l+1)^{\text{th}}$ visit to state $k+1$. Note that the first sum in (7) (for $k \geq x$) starts from $l=0$ whereas the second one starts from $l=1$. This is because for $k \geq x$, jumps from k to $k-1$ can occur even before the first jump from $k+1$ to k . This is not the case for states $k < x$. Furthermore, $\{\xi_{l,k}\}_{l \geq 0}$ is a sequence of i.i.d. random variables independent of ζ_{k+1} . Hence, the sequence $\{\xi_{n-k}\}_k$ defines a branching process. Moreover, the number of visits to state k can be written as

$$Z_k = \begin{cases} 1 + \zeta_k + \zeta_{k+1}, & \text{if } k \in \{x, x+1, \dots, n-1\}, \\ \zeta_k + \zeta_{k+1}, & \text{if } k \in \{0, 1, \dots, x-1\}, \end{cases} \quad (8)$$

where the additional one appears in the first case (for $k \geq x$) because the states $k \geq x$ are visited at least once before the chain is absorbed in state n . It is the above characterisation of the Z_k that we shall use to obtain upper bounds on $\mathbb{E}_x [Z_k]$. Precisely, we obtain bounds on $\mathbb{E}_x [\zeta_k]$ to bound $\mathbb{E}_x [Z_k]$. In the lemma below, we express $\mathbb{E}_x [\zeta_k]$ in terms of the transition probabilities of the chain \bar{Y}^n .

Lemma 3. *For any $k \in \{1, 2, \dots, n\}$, let ζ_k denote the number of jumps of the chain \bar{Y}^n from state k to state $k-1$. We have $\zeta_n = 0$ and*

$$\mathbb{E}_x [\zeta_k] = \begin{cases} \sum_{t=k}^{n-1} \prod_{i=k}^t \frac{p_{i,i-1}}{p_{i,i+1}}, & \text{for } x \leq k \leq n-1, \\ \left(\prod_{i=k}^{x-1} \frac{p_{i,i-1}}{p_{i,i+1}} \right) \mathbb{E}_x [\zeta_x], & \text{for } 0 \leq k < x, \end{cases} \quad (9)$$

where for each $i, j \in \{0, 1, \dots, n\}$, $p_{i,j}$ denotes the transition probability of the Markov chain \bar{Y}^n from state i to state j and is given by (2).

Proof. We observe that ζ_{k+1} and $\{\xi_{l,k}\}_{l \geq 0}$ are independent of each other and $\{\xi_{l,k}\}_{l \geq 0}$ is a sequence of i.i.d. random variables with

$$\mathbb{P}_x (\xi_{l,k} = i) = \left(\frac{p_{k,k-1}}{1 - p_{k,k}} \right)^i \left(\frac{p_{k,k+1}}{1 - p_{k,k}} \right)$$

for all $i \in \mathbb{Z}_+$ and all $l \geq 0$. Hence, we have

$$\mathbb{E}_x [\xi_{l,k}] = \frac{p_{k,k-1}}{p_{k,k+1}},$$

for all $l \geq 0$. Applying Wald's identity to (7) we obtain the following recursions

$$\mathbb{E}_x [\zeta_k] = \begin{cases} (1 + \mathbb{E}_x [\zeta_{k+1}]) \frac{p_{k,k-1}}{p_{k,k+1}}, & \text{if } k \geq x, \\ \mathbb{E}_x [\zeta_{k+1}] \frac{p_{k,k-1}}{p_{k,k+1}}, & \text{if } k < x, \end{cases} \quad (10)$$

Upon solving the above recursions with boundary condition $\zeta_n = 0$, we obtain desired result. \square

From (9), we observe that the ratio $p_{i,i-1}/p_{i,i+1}$ plays a crucial role in the expression of $\mathbb{E}_x [\zeta_k]$. Hence, by characterising this ratio, we can characterise $\mathbb{E}_x [\zeta_k]$. We note from (2) that

$$\frac{p_{i,i-1}}{p_{i,i+1}} = f_\alpha(i/n), \quad (11)$$

where $f_\alpha : [0, 1] \rightarrow \mathbb{R}_+$, for each $\alpha \in (0, 1)$, is as defined in Theorem 1. In the lemma below, we obtain a list some important properties of the function f_α .

Lemma 4. For $\alpha \in (0, 1)$, define $f_\alpha : [0, 1] \rightarrow \mathbb{R}_+$

$$f_\alpha(x) \triangleq \frac{(1-\alpha)x(1-x)}{\alpha + (1-\alpha)x^2} \quad (12)$$

Then f_α satisfies the following properties

1. For all $\alpha \in (0, 1)$, f_α is strictly increasing in $[0, x_\alpha)$, strictly decreasing in $(x_\alpha, 1]$, and, in the domain $[0, 1]$, attains its maximum value at x_α , where

$$x_\alpha \triangleq \sqrt{\left(\frac{\alpha}{1-\alpha}\right)^2 + \frac{\alpha}{1-\alpha}} - \frac{\alpha}{1-\alpha} \in [0, 1). \quad (13)$$

2. Let $r_\alpha \triangleq f_\alpha(x_\alpha) = \max_{x \in [0, 1]} f_\alpha(x)$. Then, for $\alpha \in (1/9, 1)$, we have $r_\alpha < 1$, and, for $\alpha \in (0, 1/9)$, we have $r_\alpha > 1$.
3. For $\alpha \in (0, 1/9]$, we have $f_\alpha(x) \geq 1$ iff $x \in [\underline{x}_\alpha, \bar{x}_\alpha]$, where

$$\underline{x}_\alpha = \frac{1}{4} \left(1 - \sqrt{1 - \frac{8\alpha}{1-\alpha}} \right) \quad (14)$$

$$\bar{x}_\alpha = \frac{1}{4} \left(1 + \sqrt{1 - \frac{8\alpha}{1-\alpha}} \right) \quad (15)$$

Furthermore, $f_\alpha(\underline{x}_\alpha) = f_\alpha(\bar{x}_\alpha) = 1$ and $x_\alpha \in [\underline{x}_\alpha, \bar{x}_\alpha]$.

4. For $\alpha \in (0, 1/9)$, define $g_\alpha(x) \triangleq \int_{\underline{x}_\alpha}^x \log(f_\alpha(x)) dx$. Then g_α has a unique root $p_c(\alpha) \in (\bar{x}_\alpha, 1)$. Furthermore, $g_\alpha(p) > 0$ if $p \in [\bar{x}_\alpha, p_c(\alpha))$ and $g_\alpha(p) < 0$ if $p \in (p_c(\alpha), 1)$

Proof. Taking the derivative of (12) with respect x we obtain

$$f'_\alpha(x) = \frac{-(1-\alpha)^2 x^2 - 2\alpha(1-\alpha)x + \alpha(1-\alpha)}{(\alpha + (1-\alpha)x^2)^2}.$$

We note that the numerator of the above expression is zero when $x = x_\alpha$, positive when $x \in [0, x_\alpha)$ and negative when $x \in (x_\alpha, 1]$, where x_α is as defined in (13). This proves the first statement of the lemma.

Next, we note from (12) that the condition $f_\alpha(x) \lesseqgtr 1$ is equivalent to $2x^2 - x + \frac{\alpha}{1-\alpha} \gtrless 0$. Hence, $f_\alpha(x) < 1$ for all $x \in [0, 1]$ if and only if $1 - 8\frac{\alpha}{1-\alpha} < 0$ or equivalently iff $\alpha > 1/9$. Furthermore, for $\alpha \in (0, 1/9]$, we have $2x^2 - x + \frac{\alpha}{1-\alpha} = (x - \underline{x}_\alpha)(x - \bar{x}_\alpha)$ where \underline{x}_α and \bar{x}_α are as defined in (14) and (15), respectively, and for $\alpha < 1/9$ we have $\underline{x}_\alpha < \bar{x}_\alpha$ and $x_\alpha \in (\underline{x}_\alpha, \bar{x}_\alpha)$. Combining the above facts we have the second and the third statements of the lemma.

We now turn to the last statement of the lemma. Note that for $\alpha \in (0, 1/9)$, we have

$$g_\alpha(\bar{x}_\alpha) = \int_{\underline{x}_\alpha}^{\bar{x}_\alpha} \log(f_\alpha(x)) dx \geq \log(f_\alpha(x_\alpha)) = \log(r_\alpha) > 0,$$

where the last inequality follows from the fact that $r_\alpha = f_\alpha(x_\alpha) > 1$ for $\alpha \in (0, 1/9)$ as established in the previous paragraph. Using (12), we can compute g_α in closed form. This is given as follows

$$\begin{aligned} g_\alpha(x) &= x \log\left(\frac{(1-\alpha)x}{\alpha + (1-\alpha)x^2}\right) - (1-x) \log(1-x) + \log(1-\underline{x}_\alpha) \\ &\quad - 2\sqrt{\frac{\alpha}{1-\alpha}} \left(\arctan\left(\sqrt{\frac{1-\alpha}{\alpha}} x\right) - \arctan\left(\sqrt{\frac{1-\alpha}{\alpha}} \underline{x}_\alpha\right) \right) \end{aligned} \quad (16)$$

From the above it follows that $\lim_{x \rightarrow 1^-} g(x) < 0$ since $\underline{x}_\alpha < 1$ and $\arctan(\cdot)$ is an increasing function. Furthermore, we have $g'_\alpha(x) = \log(f_\alpha(x)) < 0$ for $x \in (\bar{x}_\alpha, 1)$. Hence, there must exist a unique root $p_c(\alpha)$ of g_α in $(\bar{x}_\alpha, 1)$ and $g_\alpha(p)$ must be strictly negative for $p \in (p_c(\alpha), 1)$ and strictly positive for $p \in [\bar{x}_\alpha, p_c(\alpha))$. \square

We are now in a position to complete the proof of Theorem 1. We use Lemma 3 and the properties of f_α proved in Lemma 4 to obtain upper bounds of $\mathbb{E}_x [\zeta_k]$. These upper bounds, in turn, provide upper bounds on $\mathbb{E}_x [Z_k]$ using (8). Finally, we use Lemma 1 and the upper bounds on $\mathbb{E}_x [Z_k]$ to obtain an upper bound on the mean consensus time $\bar{T}_n(p)$. The complete proof is given below.

Proof of Theorem 1: From Lemma 1 it follows that

$$\begin{aligned}\bar{T}_n(p) &= \sum_{k=0}^{n-1} \frac{\mathbb{E}_{[pn]} [Z_k]}{(1 - p_{k,k})} \\ &= \sum_{k=0}^{n-1} \frac{\mathbb{E}_{[pn]} [Z_k]}{(1 - \frac{k}{n})(\alpha + (1 - \alpha)\frac{k}{n})} \\ &= n \sum_{k=0}^{n-1} \mathbb{E}_{[pn]} [Z_k] \left(\frac{1}{n-k} + \frac{1}{\frac{\alpha}{1-\alpha}n+k} \right)\end{aligned}\quad (17)$$

Hence, to prove $T_n(p) = O(n \log n)$, it is sufficient to establish that $\mathbb{E}_{[pn]} [Z_k] \leq C$ for all $k \in \{0, 1, \dots, n\}$, where $C > 0$ is a constant independent of k . Indeed, if $\mathbb{E}_{[pn]} [Z_k] \leq C$ for each state $k \in \{0, 1, \dots, n\}$, then for $n \geq \frac{1-\alpha}{\alpha}$ we have

$$\begin{aligned}\bar{T}_n(p) &\leq nC \sum_{k=0}^{n-1} \left(\frac{1}{n-k} + \frac{1}{k+1} \right) \\ &= 2nC \sum_{k=1}^{n-1} \frac{1}{k} \\ &= O(n \log n)\end{aligned}\quad (18)$$

Hence, to prove the first two statements of the theorem, it is sufficient to show that, under the conditions stated in these statements, $\mathbb{E}_{[pn]} [Z_k]$ is uniformly bounded by some constant for all $k \in \{0, 1, \dots, n-1\}$. This what we prove next.

Using the first two properties of f_α proved in Lemma 4 and (9), we see that for $\alpha \in (1/9, 1)$ we have

$$\mathbb{E}_x [\zeta_k] \leq \begin{cases} \sum_{t=k}^{n-1} r_\alpha^{t-k+1} < \frac{r_\alpha}{1-r_\alpha}, & \text{for } x \leq k \leq n-1 \\ \mathbb{E}_x [\zeta_x] < \frac{r_\alpha}{1-r_\alpha}, & \text{for } k < x, \end{cases}\quad (19)$$

where $r_\alpha < 1$ is as defined in Lemma 4. Hence, from (8) we have that for all $\alpha \in (1/9, 1)$ and all $k \in \{0, 1, \dots, n-1\}$,

$$\mathbb{E}_x [Z_k] \leq \frac{1+r_\alpha}{1-r_\alpha}.$$

This establishes the first statement of the theorem.

We now prove the second statement of the theorem. For $\alpha \in (0, 1/9)$, the third statement of Lemma 4 implies $p_c(\alpha) > \bar{x}_\alpha$. Furthermore, by the first and third statements of Lemma 4, it follows that f_α is strictly decreasing in the range $[\bar{x}_\alpha, 1)$. Hence, for $p \geq p_c(\alpha) > \bar{x}_\alpha$ we have $f_\alpha(p) < f_\alpha(\bar{x}_\alpha) = 1$. Furthermore, for all $k \geq x \geq pn$ we have

$$\frac{p_{k,k-1}}{p_{k,k+1}} = f_\alpha(k/n) \leq f_\alpha(p) < 1.$$

Let $r_p \triangleq f_\alpha(p) < 1$. Then, from (9), we have that for $k \geq x$

$$\mathbb{E}_x [\zeta_k] \leq \sum_{t=k}^{n-1} r_p^{t-k+1} < \frac{r_p}{1-r_p}.$$

Hence, from (8) we have that

$$\mathbb{E}_x [Z_k] \leq \frac{1+r_p}{1-r_p}$$

for $k \geq x$. Moreover, for $k < n\bar{x}_\alpha \leq x = \lceil pn \rceil$ we have

$$\begin{aligned}
\prod_{i=k}^{x-1} \frac{p_{i,i-1}}{p_{i,i+1}} &= \prod_{i=k}^{\lceil pn \rceil - 1} f_\alpha(i/n) \\
&\stackrel{(a)}{\leq} \prod_{i=\lceil n\bar{x}_\alpha \rceil}^{\lceil pn \rceil - 1} f(i/n) \\
&= \exp \left(\sum_{i=\lceil n\bar{x}_\alpha \rceil}^{\lceil pn \rceil - 1} \log(f_\alpha(i/n)) \right) \\
&\stackrel{(b)}{=} \exp \left(n \left(\int_{\bar{x}_\alpha}^p \log(f_\alpha(x)) dx + O(1/n) \right) \right) \\
&\stackrel{(c)}{=} \exp(n g_\alpha(p) + O(1)) \\
&\stackrel{(d)}{=} O(1),
\end{aligned}$$

where (a) follows from the facts that $f_\alpha(i/n) \leq 1$ for $i \leq n\bar{x}_\alpha$ and $f_\alpha(i/n) \geq 1$ for $n\bar{x}_\alpha \leq i \leq n\bar{x}_\alpha$; (b) follows from the fact that the Riemannian sum $(1/n) \sum_{i=\lceil n\bar{x}_\alpha \rceil}^{\lceil pn \rceil - 1} \log(f_\alpha(i/n))$ converges to the integral $\int_{\bar{x}_\alpha}^p \log(f_\alpha(x)) dx$ as $n \rightarrow \infty$ with an error of $O(1/n)$; (c) follows from the definition of g_α in Lemma 4; (d) follows from the fact that for $p \geq p_c(\alpha) > \bar{x}_\alpha$ we have $g_\alpha(p) \leq 0$ by the last statement of Lemma 4. For $x > k \geq n\bar{x}_\alpha$ we have

$$\prod_{i=k}^{x-1} \frac{p_{i,i-1}}{p_{i,i+1}} = \prod_{i=k}^{\lceil pn \rceil - 1} f(i/n) \leq 1,$$

because for $i \geq n\bar{x}_\alpha$ we have $f_\alpha(i/n) \leq 1$. Hence, for all $k < x$, it follows from (9) that $\mathbb{E}_x[\zeta_k] = O(1)\mathbb{E}_x[\zeta_x] = O(1)$. Finally, from (8) we obtain that $\mathbb{E}_x[Z_k] = O(1)$ for all $k < x$. This establishes the second statement of the theorem.

We now turn to the third statement of the theorem. Note from (17) that to prove this statement, it suffices to show that $\mathbb{E}_{\lceil pn \rceil}[Z_k] = \Omega(\exp(\Theta(n)))$ for $k = \lceil n\bar{x}_\alpha \rceil$ when $\alpha \in (0, 1/9)$ and $p \in (0, p_c(\alpha))$. First, note that for $\alpha \in (0, 1/9)$, $p \in (\bar{x}_\alpha, p_c(\alpha))$, and $k = \lceil n\bar{x}_\alpha \rceil$ we have

$$\begin{aligned}
\prod_{i=\lceil n\bar{x}_\alpha \rceil}^{x-1} \frac{p_{i,i-1}}{p_{i,i+1}} &= \prod_{i=\lceil n\bar{x}_\alpha \rceil}^{\lceil np \rceil - 1} f_\alpha(i/n) \\
&= \exp \left(n \int_{\bar{x}_\alpha}^p \log(f_\alpha(x)) dx + O(1) \right) \\
&= \Omega(\exp(\Theta(n))),
\end{aligned}$$

where the last line follows from the fact that $\int_{\bar{x}_\alpha}^p \log(f_\alpha(x)) dx = g_\alpha(p) > 0$ for $p \in (\bar{x}_\alpha, p_c(\alpha))$. Hence, $\mathbb{E}_x[\zeta_{\lceil \bar{x}_\alpha n \rceil}] = \Omega(\exp(\Theta(n)))\mathbb{E}_x[\zeta_x] = f_\alpha(p)\Omega(\exp(\Theta(n)))$ since $\mathbb{E}_x[\zeta_x] > f_\alpha(x/n) = f_\alpha(p) + o(1)$. Hence,

$\mathbb{E}_x [Z_{\lceil n\underline{x}_\alpha \rceil}] = \Omega(\exp(\Theta(n)))$. For $p \in [0, \underline{x}_\alpha]$ we have

$$\begin{aligned} \mathbb{E}_x [\zeta_{\lceil n\underline{x}_\alpha \rceil}] &> \prod_{i=\lceil n\underline{x}_\alpha \rceil}^{\lceil n\bar{x}_\alpha \rceil-1} \frac{p_{i,i-1}}{p_{i,i+1}} = \prod_{i=\lceil n\underline{x}_\alpha \rceil}^{\lceil n\bar{x}_\alpha \rceil-1} f_\alpha(i/n) \\ &= \exp\left(n \int_{\underline{x}_\alpha}^{\bar{x}_\alpha} \log(f_\alpha(x)) dx + O(1)\right) \\ &= \Omega(\exp(\Theta(n))), \end{aligned}$$

This proves the third part of the theorem. \square

Remark 1. From the proof above, it is clear that the absorption time of the chain \bar{Y}^n crucially depends on the ratio of the down-transition probability $p_{i,i-1}$ to the up-transition probability $p_{i,i+1}$ at any given state i . If this ratio is smaller than one at a given state i , then the chain has a tendency to drift quickly towards the absorbing state n . Similarly, if the ratio is larger than one, then the chain has a tendency to move towards state 0 and hence it takes longer to reach the absorbing state n . Since the value of this ratio depends on the bias parameter α as well as on the initial proportion p of agents with the superior opinion, we observe phase transitions with respect to both the parameters.

Remark 2. Although we have proved the theorem for the modified chain \bar{Y}^n , the proof can be extended to the original chain \bar{X}^n . This is because the results of Lemma 1 and Lemma 3 are also applicable to the original chain if the transition probabilities are replaced by those of the original chain. Furthermore, the transition probabilities of the two chains satisfy the following relations

$$1 - \tilde{p}_{k,k} = 1 - p_{k,k} + (1 - \alpha) \frac{k}{n} \left(1 - \frac{k}{n}\right) \left(\frac{n^2}{(n-1)^2} - 1\right) \leq 1 - p_{k,k},$$

and $\tilde{p}_{k,k-1}/\tilde{p}_{k,k+1} = p_{k,k-1}/p_{k,k+1} + o(1)$, where we recall that $\tilde{p}_{k,k}$ denotes the transition probability from state k to itself for the original chain \bar{X}^n . We note that the same lower bound as in Lemma 2 can be obtained for the chain \bar{X}^n since we have $1 - \tilde{p}_{k,k} \leq (\max(\alpha, 1 - \alpha) + o(1))(1 - k/n)(1 + k/n)$. Similarly, the upper and lower bounds on the expected number of visits to each state derived for the modified chain also hold for the original chain for large enough n . Combining the above, we obtain the same asymptotic bounds for the original chain \bar{X}^n as we have for the modified chain \bar{Y}^n .

4 Numerical Results

In this section, we present numerical results for the model presented in this paper. We first present results to support our theoretical findings on complete graphs. We then present simulation results for other classes of graphs. Error-bars in the plots represent 95% confidence intervals. Also, to understand the growth rates better, we overlay the plots obtained from simulations on theoretical growth rates obtained by plotting appropriately scaled functions.

4.1 Complete graphs

For complete graphs, the mean consensus time $\bar{T}_n(p)$ can be computed numerically using the first step analysis of the Markov chain \bar{Y}^n . This method is more exact and computationally less expensive than simulating the chain a large number of times to obtain the average absorption time. Hence, we adopt this method for complete graphs. With a slight abuse of notation let $\bar{T}_n(k)$ denote the average absorption time of the chain starting from state $k \in \{0, 1, \dots, n\}$. Then, from first step analysis we have for each $k \in \{0, 1, \dots, n\}$

$$\bar{T}_n(k) = 1 + p_{k,k-1}\bar{T}_n(k-1) + p_{k,k+1}\bar{T}_n(k+1) + p_{k,k}\bar{T}_n(k), \quad (20)$$


where $p_{i,j}$ is as defined in (2). Simplifying the above and using the boundary condition $\bar{T}_n(n) = 0$ we obtain

$$\bar{T}_n(p) := \bar{T}_n(\lceil np \rceil) = \sum_{k=\lceil np \rceil}^{n-1} S_n(k), \quad (21)$$


where $S_n(k)$ satisfies the following recursion:

$$S_n(k) = \frac{1}{p_{k,k+1}} + \frac{p_{k,k-1}}{p_{k,k+1}} S_n(k-1), \quad (22)$$

with $S_n(0) = 1/\alpha$. We find the expected consensus time numerically by solving the above recursion for different values of n, α , and p . First, we choose $\alpha = 0.1 < 1/9$. For this value of α , we can numerically compute the value of $p_c(\alpha)$ by solving (3). This value turns out to be 0.431 (accurate to the third decimal place). In Figures 1a and 1b we plot the normalised (by the network size) average consensus time as a function of the network size for $p = 0.4 < p_c(\alpha) = 0.431$ and $p = 0.5 > p_c(\alpha) = 0.431$, respectively. As expected from Theorem 1 we observe that the mean consensus time grows exponentially for $p < p_c(\alpha)$ and as $\Theta(n \log n)$ for $p > p_c(\alpha)$.



(a) $\alpha = 0.1 < 1/9, p = 0.4 < p_c(\alpha) = 0.431$



(b) $\alpha = 0.1 < 1/9, p = 0.5 > p_c(\alpha) = 0.431$


Figure 1: Mean consensus time per node $\bar{T}_n(p)/n$ as a function of the network size for $\alpha = 0.1 < 1/9$ for complete graphs

Next, we choose $\alpha = 0.125 > 1/9$. For this choice of α , we expect (from Theorem 1) the mean consensus time to grow as $\Theta(n \log n)$ for all values of p . This is verified in Figures 2a and 2b, where we choose $p = 0$ and $p = 0.5$, respectively. We observe that in both cases $\bar{T}_n(p) = \Theta(n \log n)$.


4.2 Random d -regular graphs with $d = \Theta(\log n)$

We now present simulation results for random d -regular graphs where the degree d for each node is chosen to be $d = \lceil \log n \rceil$. We use the `networkx` package to generate random d -regular graphs. For each randomly generated graph, we run the protocol until the network reaches consensus. The above procedure is repeated 500 times and mean consensus time is computed over these 500 runs. We further repeat this for different values of n, α , and p . In Figures 3a and 3b we fix α to be 0.05 and plot the normalised mean consensus time as a function of the network size for $p = 0.05$ and $p = 0.8$, respectively. Similar to complete graphs, we observe that the mean consensus time grows exponentially with n when both α and p are low. In contrast, when the initial proportion of agents having the superior opinion is sufficiently large, the mean consensus time grows as $\Theta(n \log n)$ even when the bias parameter α is small.

In Figures 4a and 4b we choose $\alpha = 0.8$ plot the normalised mean consensus time as a function of the network size for $p = 0.005$ and $p = 0.8$, respectively. We observe that in both cases the mean consensus time grows as $\Theta(n \log n)$. This indicates that if the bias parameter α is sufficiently large then the network reaches consensus to the superior opinion in $\Theta(n \log n)$ time irrespective of the initial proportion of agents having the superior opinion.




(a) $\alpha = 0.125 > 1/9, p = 0$




(b) $\alpha = 0.125 > 1/9, p = 0.5$

Figure 2: Mean consensus time per node $\bar{T}_n(p)/n$ as a function of the network size for $\alpha = 0.125 > 1/9$ for complete graphs



(a) $\alpha = 0.05, p = 0.05$




(b) $\alpha = 0.05, p = 0.8$


Figure 3: Mean consensus time per node $\bar{T}_n(p)/n$ as a function of the network size for $\alpha = 0.05$ for random d -regular graphs with $d = \lceil \log n \rceil$.

4.3 Random d -regular graphs with $d = O(1)$

We now investigate the biased opinion dynamics on random d -regular with constant degree, i.e., $d = O(1)$. Specifically, we set $d = 5$ and simulate the network for different values of n, α , and p . As before, we generate a random 5-regular graph using the `networkx` package and run the protocol on this randomly generated instance until consensus is reached. The above procedure is repeated multiple times (each with a newly generated random graph) to obtain the mean consensus time within the desired confidence bounds. In Figures 5a and 5b, we fix $\alpha = 0.05$ and choose $p = 0.05$ and $p = 0.8$, respectively. We observe that the normalised mean consensus time grows approximately linearly in n (i.e., $\bar{T}_n(p) = O(n^2)$) when both α and p are small. This is unlike the previous results for dense graphs, where, for small values of α and p , the mean consensus time is exponential in n . This reduction in absorption time can be intuitively explained by the fact that a smaller neighbourhood size reduces the chance of an agent updating to the worse opinion when a fixed proportion of all the agents have the worse opinion. Based on this intuition we conjecture that for small values of p and α the mean consensus time grows polynomially in n when the neighbourhood size d is constant higher than 2. The case of cycles (where $d = 2$) has already been considered in [20]. But in the case of cycles there is no phase transition; indeed, it has been shown in [20] that for cycles consensus is achieved on the superior opinion in $O(n \log n)$ time for all values of α even when $p = 0$.




(a) $\alpha = 0.8, p = 0.05$




(b) $\alpha = 0.8, p = 0.8$

Figure 4: Mean consensus time per node $\bar{T}_n(p)/n$ as a function of the network size for $\alpha = 0.8$ for random d -regular graphs with $d = \lceil \log n \rceil$.



(a) $\alpha = 0.05, p = 0.05$




(b) $\alpha = 0.05, p = 0.8$

Figure 5: Mean consensus time per node $\bar{T}_n(p)/n$ as a function of the network size for $\alpha = 0.05$ for random d -regular graphs with $d = 5$.


In Figures 6a and 6b, we plot the mean consensus time as a function of the network size for $p = 0.05$ and $p = 0.8$, respectively, keeping $\alpha = 0.8$. In both cases, we observe that the mean consensus time grows as $\Theta(n \log n)$. Thus, in these regimes, the behaviour of random d -regular graphs with $d = O(1)$ is similar to that of random d -regular graphs with $d = \Theta(\log n)$.

4.4 Erdős-Rényi graphs with edge probability $\log n/n$

Finally, we present results for Erdős-Rényi graphs where the edge probability is fixed at the connectivity threshold $\log n/n$. The experiments are designed in the same way as described for other classes of graphs. The normalised mean consensus time as a function of the network size n is plotted for different values of α and p : in Figures 7a and 7b for $\alpha = 0.05$ and in Figures 8a and 8b for $\alpha = 0.8$. We observe similar phase transitions as in the case of complete graphs, i.e., the mean consensus time grows as $\Theta(n \log n)$ in all cases except when both α and p are low.




(a) $\alpha = 0.8, p = 0.05$




(b) $\alpha = 0.8, p = 0.8$

Figure 6: Mean consensus time per node $\bar{T}_n(p)/n$ as a function of the network size for $\alpha = 0.8$ for random d -regular graphs with $d = 5$.



(a) $\alpha = 0.05, p = 0.05$



(b) $\alpha = 0.05, p = 0.8$

Figure 7: Mean consensus time per node $\bar{T}_n(p)/n$ as a function of the network size for $\alpha = 0.05$ for Erdős-Rényi graphs with edge probability $\log n/n$.

5 Conclusion and future work

In this paper we have studied a model of binary opinion dynamics where the agents show a strong form of bias towards one of the opinions, called the superior opinion. We showed that for complete graphs the model exhibits rich phase transitions based on the values of the bias parameter and the initial proportion of agents with the superior opinion. Specifically, we proved that fast consensus can be achieved on the superior opinion irrespective of the initial configuration of the network when bias is sufficiently high. When bias is low, we show that fast consensus can only be achieved when the initial proportion of agents with the superior opinion is above a certain threshold. If this is not the case, then we show that consensus takes exponentially long time. Through simulations, we observed similar behaviour for several classes of dense graphs where the average degree scales at least logarithmically with the network size. For sparse graphs, where the average degree is constant, we observed that phase transitions do occur but the behaviour below criticality is different to that of dense graphs. Specifically, we observed that when both bias and the initial proportion of agents with the superior opinion are low, the average consensus time is still polynomial in the network size.

Several directions remain open for theoretical investigation. One immediate problem is to theoretically establish the observed phase transitions for dense graphs. Here, it will be interesting to find out how ‘dense’




Figure 8: Mean consensus time per node $\bar{T}_n(p)/n$ as a function of the network size for $\alpha = 0.8$ for Erdős-Rényi graphs with edge probability $\log n/n$.

a graph must be in order for it to exhibit the same phase transitions as in complete graphs. Similar questions remain open for sparse graphs such as d -regular expanders with constant d . Here, it will be of interest to find out how the threshold value of the parameters depend on the average degree d or the spectral properties of the expander. It is also worth analysing the exact behaviour below criticality. The numerical experiments conducted in this paper suggests that the consensus time is still polynomial below criticality (unlike in complete graphs where it is exponential), but it would be challenging to obtain exact bounds in this case.

References

- [1] A. Mukhopadhyay, R. R. Mazumdar, and R. Roy, “Voter and majority dynamics with biased and stubborn agents,” *Journal of Statistical Physics*, vol. 181, no. 4, pp. 1239–1265, 2020.
- [2] A. Mukhopadhyay, R. R. Mazumdar, and R. Roy, “Binary opinion dynamics with biased agents and agents with different degrees of stubbornness,” in *2016 28th International Teletraffic Congress (ITC 28)*, vol. 1, pp. 261–269, IEEE, 2016.
- [3] A. Anagnostopoulos, L. Becchetti, E. Cruciani, F. Pasquale, and S. Rizzo, “Biased opinion dynamics: When the devil is in the details,” in *Proceedings of the Twenty-Ninth International Joint Conference on Artificial Intelligence, IJCAI-20* (C. Bessiere, ed.), pp. 53–59, International Joint Conferences on Artificial Intelligence Organization, 7 2020. Main track.
- [4] H. Lesfari, F. Giroire, and S. Pérennes, “Biased majority opinion dynamics: Exploiting graph k -domination,” in *IJCAI 2022-International Joint Conference on Artificial Intelligence*, 2022.
- [5] C. Cooper, R. Elsässer, and T. Radzik, “The power of two choices in distributed voting,” in *International Colloquium on Automata, Languages, and Programming*, pp. 435–446, Springer, 2014.
- [6] R. A. Holley and T. M. Liggett, “Ergodic theorems for weakly interacting infinite systems and the voter model,” *The Annals of Probability*, pp. 643–663, 1975.
- [7] P. Clifford and A. Sudbury, “A model for spatial conflict,” *Biometrika*, vol. 60, no. 3, pp. 581–588, 1973.
- [8] J. T. Cox, “Coalescing random walks and voter model consensus times on the torus in zd ,” *The Annals of Probability*, pp. 1333–1366, 1989.
- [9] C. Cooper, R. Elsässer, H. Ono, and T. Radzik, “Coalescing random walks and voting on connected graphs,” *SIAM Journal on Discrete Mathematics*, vol. 27, no. 4, pp. 1748–1758, 2013.

- [10] T. Nakata, H. Imahayashi, and M. Yamashita, “Probabilistic local majority voting for the agreement problem on finite graphs,” in *International Computing and Combinatorics Conference*, pp. 330–338, Springer, 1999.
- [11] E. Mossel, J. Neeman, and O. Tamuz, “Majority dynamics and aggregation of information in social networks,” *Autonomous Agents and Multi-Agent Systems*, vol. 28, no. 3, pp. 408–429, 2014.
- [12] A. N. Zehmakan, “Opinion forming in erdős-rényi random graph and expanders,” *Discrete Applied Mathematics*, vol. 277, pp. 280–290, 2020.
- [13] R. H. Schonmann, “Finite size scaling behavior of a biased majority rule cellular automaton,” *Physica A: Statistical Mechanics and its Applications*, vol. 167, no. 3, pp. 619–627, 1990.
- [14] B. Gärtner and A. N. Zehmakan, “Majority model on random regular graphs,” in *Latin American Symposium on Theoretical Informatics*, pp. 572–583, Springer, 2018.
- [15] Y. Kanoria and A. Montanari, “Majority dynamics on trees and the dynamic cavity method,” *The Annals of Applied Probability*, vol. 21, no. 5, pp. 1694–1748, 2011.
- [16] P. L. Krapivsky and S. Redner, “Dynamics of majority rule in two-state interacting spin systems,” *Physical Review Letters*, vol. 90, no. 23, p. 238701, 2003.
- [17] S. Galam, “Minority opinion spreading in random geometry,” *The European Physical Journal B-Condensed Matter and Complex Systems*, vol. 25, no. 4, pp. 403–406, 2002.
- [18] P. Chen and S. Redner, “Consensus formation in multi-state majority and plurality models,” *Journal of Physics A: Mathematical and General*, vol. 38, no. 33, p. 7239, 2005.
- [19] J. Cruise and A. Ganesh, “Probabilistic consensus via polling and majority rules,” *Queueing Systems*, vol. 78, no. 2, pp. 99–120, 2014.
- [20] A. Anagnostopoulos, L. Becchetti, E. Cruciani, F. Pasquale, and S. Rizzo, “Biased opinion dynamics: when the devil is in the details,” *Information Sciences*, vol. 593, pp. 49–63, 2022.
- [21] E. Cruciani, E. Natale, A. Nusser, and G. Scornavacca, “Phase transition of the 2-choices dynamics on core-periphery networks,” *Distributed Computing*, vol. 34, no. 3, pp. 207–225, 2021.
- [22] F. d’Amore and I. Ziccardi, “Phase transition of the 3-majority dynamics with uniform communication noise,” in *International Colloquium on Structural Information and Communication Complexity*, pp. 98–115, Springer, 2022.
- [23] E. Cruciani, H. A. Mimun, M. Quattropani, and S. Rizzo, “Phase transitions of the k-majority dynamics in a biased communication model,” in *International Conference on Distributed Computing and Networking 2021*, pp. 146–155, 2021.