

**Original citation:**

Carstensen, Carsten and Ortner, Christoph. (2010) Analysis of a class of penalty methods for computing singular minimizers. Computational Methods in Applied Mathematics, Volume 10 (Number 2). pp. 137-163.

**Permanent WRAP url:**

<http://wrap.warwick.ac.uk/43755>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**


2010 First published in Computational Methods in Applied Mathematics

<http://www.cmam.info/index.php>

**A note on versions:**

The version presented in WRAP is the published version or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: [publications@warwick.ac.uk](mailto:publications@warwick.ac.uk)



<http://wrap.warwick.ac.uk/>

# ANALYSIS OF A CLASS OF PENALTY METHODS FOR COMPUTING SINGULAR MINIMIZERS

C. CARSTENSEN<sup>1</sup> AND C. ORTNER<sup>2</sup>

**Abstract** — Amongst the more exciting phenomena in the field of nonlinear partial differential equations is the Lavrentiev phenomenon which occurs in the calculus of variations. We prove that a conforming finite element method fails if and only if the Lavrentiev phenomenon is present. Consequently, nonstandard finite element methods have to be designed for the detection of the Lavrentiev phenomenon in the computational calculus of variations.

We formulate and analyze a general strategy for solving variational problems in the presence of the Lavrentiev phenomenon based on a splitting and penalization strategy. We establish convergence results under mild conditions on the stored energy function. Moreover, we present practical strategies for the solution of the discretized problems and for the choice of the penalty parameter.

**2000 Mathematics Subject Classification:** 65N12; 65N25; 65N30; 65N50.

**Keywords:** Lavrentiev phenomenon, finite element method, computational calculus of variations.

## 1. Introduction

The calculus of variations is concerned with the minimisation problem

$$\inf E(\mathcal{A}_1) := \inf_{v \in \mathcal{A}_1} E(v), \quad (1.1)$$

where  $E : \mathcal{A}_1 \rightarrow \mathbb{R} \cup \{+\infty\}$  and where  $\mathcal{A}_1$  (or more generally  $\mathcal{A}_p$ ) is the first-order Sobolev space

$$\mathcal{A}_p := W_0^{1,p}(\Omega; \mathbb{R}^m) = \{v \in W^{1,p}(\Omega)^m : v|_{\partial\Omega} = 0\},$$

based on a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^n$  with piecewise hyperplanar boundary  $\partial\Omega$ .

We shall assume throughout that  $E$  is *proper* on  $\mathcal{A}_\infty$ , i.e., there exists  $v \in \mathcal{A}_\infty$  so that  $E(v) < +\infty$ . In particular,  $\mathcal{A}_\infty \subset \mathcal{A}_1$  always implies

$$-\infty \leq \inf E(\mathcal{A}_1) \leq \inf E(\mathcal{A}_\infty) < +\infty.$$

The *Lavrentiev phenomenon*, named after its first occurrence in the literature [18], is the surprising property that, in some some variational problems,

$$\inf E(\mathcal{A}_1) < \inf E(\mathcal{A}_\infty). \quad (\text{L})$$

---

<sup>1</sup>*Humboldt-Universität zu Berlin, Unter den Linden 6, 10099 Berlin, Germany; Department of Computational Science and Engineering, Yonsei University, 120-749 Seoul, Korea. E-mail: cc@mathematik.hu-berlin.de*

<sup>2</sup>*Mathematical Institute, University of Oxford, 24-29 St Giles', Oxford OX1 3LB, UK. E-mail: ortner@maths.ox.ac.uk*

Other well-known examples are the one-dimensional examples of Mania [23] and of Ball and Mizel [7, 6], or the convex example of Foss, Hrusa and Mizel [16]. In nonlinear elasticity, the Lavrentiev phenomenon is closely related to the occurrence of cavitation [4].

For the conforming finite element discretization of (1.1) assume we are given a family of finite element spaces

$$V_0, V_1, V_2, \dots \subset \overline{\bigcup_{\ell=0}^{\infty} V_{\ell}} \subseteq \mathcal{A}_{\infty},$$

to solve the discrete minimization problem

$$\inf E(V_{\ell}) := \inf_{v_{\ell} \in V_{\ell}} E(v_{\ell}). \quad (1.2)$$

The respective infimal energies are possibly convergent towards some limit

$$\inf E(\mathcal{A}_{\infty}) \leq \liminf_{\ell \rightarrow \infty} \inf E(V_{\ell}).$$

We say that the finite element method (FEM) is convergent if  $E$  and the sequence of discrete subspaces  $V_0, V_1, V_2, \dots$  allow for

$$\inf E(\mathcal{A}_1) = \liminf_{\ell \rightarrow \infty} \inf E(V_{\ell}). \quad (\text{C})$$

Therein, the convergence of the entire sequence of energy minima (not merely of *some* subsequence but for all subsequences) is part of the statement as well as the equality of that limit to  $\inf E(\mathcal{A}_1)$ .

However, since conforming finite element functions are always Lipschitz continuous any finite element space  $V_{\ell}$  is contained in  $\mathcal{A}_{\infty}$  and hence standard finite element methods cannot compute singular minimisers, that is, if (L) holds then

$$\inf E(\mathcal{A}_1) < \inf E(\mathcal{A}_{\infty}) \leq \inf E(V_{\ell}).$$

In particular, it follows that (C) implies that (L) is false. Section 2 below provides a general framework that allows for the converse and establishes

$$(\text{C}) \iff \text{NOT } (\text{L}),$$

under natural assumptions on the energy density.

A consequence of this equivalence is that conforming finite element methods are inappropriate tools for detecting the singular minimisers associated to the Lavrentiev phenomenon (L).

Several classes of numerical schemes have been introduced in the literature to allow for a numerical detection of (L), including the penalty method of Ball and Knowles [5, 17] and its extension to polyconvex integrands by Negron–Marrero [24], the element-removal method of Li [19, 20], and the truncation method of Li, and Bai and Li [1, 2, 21].

Section 3 introduces a general concept for the construction of a new class of splitting and penalty methods. We establish general convergence results in Sections 4 and 5. In Section 6 we discuss some connections of our results with the theory of  $\Gamma$ -convergence.

Similarly as in the methods of Ball & Knowles [5] and of Negron–Marrero [24] we decouple a problematic variable, for example the gradient  $\nabla u$ , by introducing a new variable  $\eta$  in its place and then penalizing the difference  $\nabla u - \eta$ . The main difference between the methods

[5, 24] and our approach is how this penalization is achieved. While [5, 24] use a constraint of the form

$$\|\nabla u - \eta\|_{L^p} \leq \varepsilon,$$

we add a penalization term

$$\varepsilon^{-1}\Psi(\nabla u, \eta),$$

to the total energy functional. Moreover, we design this penalization term with practical implementation issues in mind. For example, by choosing a non-differentiable penalty functional (similar to an  $L^1$ -norm), we obtain the desirable property that the difference  $\nabla u - \eta$  is non-zero only in a small subregion of the computational domain.

As a result of our careful design of the penalty functional our method is potentially easier to use and more efficient in practise. In particular, we also include a detailed description of a practical implementation and various computational examples in the final section of the paper.

In [25, 26] non-conforming finite element methods were analyzed as an alternative to the penalty methods discussed in the present paper. The main advantage of non-conforming methods is that they require no penalty parameter. However, even though this is a promising new direction, it is at present entirely unclear how to generalize the results in [25, 26] to the vectorial non-convex case. By contrast, our convergence results in the present paper hold under far less restrictive conditions on the stored energy functions.

## 2. Finite Element Failure is Equivalent to the Lavrentiev Phenomenon

This section is devoted to the proof of the equivalence of (C) and NOT (L), in a general setting which is entirely free of growth conditions and notions of convexity. However, we assume uniform convergence of the mesh-size to zero in the finite element methods as well as global continuity of the energy density.

Suppose that  $\mathcal{T}_1, \mathcal{T}_2, \mathcal{T}_3, \dots$  is a sequence of regular triangulations into simplices of a Lipschitz domain  $\Omega \subset \mathbb{R}^n$  with piecewise flat boundary  $\partial\Omega$  that is perfectly matched by the triangulations. Suppose that the triangulation is shape regular in the sense that the largest  $n$  dimensional ball inside each simplex  $T$  and the smallest ball outside have uniformly bounded ratios: There exists a universal positive constant  $C_{\text{shaperegular}}$ , which does not depend on  $T$  or  $\ell$ , such that one finds midpoints  $m_T$  and  $M_T$ , and radii  $r_T$  and  $R_T$ , satisfying

$$B(m_T, r_T) \subset T \subset B(M_T, R_T) \quad \text{and} \quad R_T/r_T \leq C_{\text{shaperegular}}.$$

We assume throughout that the mesh-size tends to zero, written  $h_\ell \rightarrow 0$ , by which we mean that

$$\lim_{\ell \rightarrow \infty} \max_{T \in \mathcal{T}_\ell} R_T = 0.$$

The finite-dimensional space  $V_\ell$  of piecewise affine finite element functions (piecewise with respect to the triangulation  $\mathcal{T}_\ell$ ),

$$V_\ell := \{v_\ell \in C_0(\Omega; \mathbb{R}^m) : \forall T \in \mathcal{T}_\ell, v_\ell|_T \text{ affine} \},$$

belongs to  $\mathcal{A}_\infty$ . For future reference we also define

$$\begin{aligned} P^0(\mathcal{T}_\ell) &:= \{v_\ell \in L^1(\Omega) : \forall T \in \mathcal{T}_\ell, v_\ell|_T \text{ constant} \}, \\ P^1(\mathcal{T}_\ell) &:= \{v_\ell \in C(\Omega; \mathbb{R}^m) : \forall T \in \mathcal{T}_\ell, v_\ell|_T \text{ affine} \}, \quad \text{and} \\ P_0^1(\mathcal{T}_\ell) &:= \{v_\ell \in C_0(\Omega; \mathbb{R}^m) : \forall T \in \mathcal{T}_\ell, v_\ell|_T \text{ affine} \}. \end{aligned}$$

Note that with this notation,  $V_\ell = P^1(\mathcal{T}_\ell) \cap \mathcal{A}_\infty = P_0^1(\mathcal{T}_\ell)$ . In the following sections we will redefine  $V_\ell$  in order to take into account nonhomogeneous boundary conditions.

Let the energy density  $W : \bar{\Omega} \times \mathbb{R}^m \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$  be continuous and define the energy

$$E(v) := \int_{\Omega} W(x, v(x), Dv(x)) dx,$$

for all  $v \in \mathcal{A}_\infty$ . In fact, if  $v$  is Lipschitz continuous, then the set of triples  $\{(x, v(x), Dv(x)) : x \in \bar{\Omega}\}$  as well as the set  $\{W(x, v(x), Dv(x)) : x \in \bar{\Omega}\}$  are contained in compact sets. Consequently,  $E(v) \in \mathbb{R}$  and  $E : \mathcal{A}_\infty \rightarrow \mathbb{R}$  is well defined. For an arbitrary function  $v \in \mathcal{A}_1$  this is no longer clear. Throughout this section we simply assume that

$$E : \mathcal{A}_1 \rightarrow \mathbb{R} \cup \{+\infty\},$$

is some extension of  $E|_{\mathcal{A}_\infty}$ . In applications, this may be guaranteed by growth control from below and we refer to the literature (e.g., [12]) for this well-understood argument in the direct method of the calculus of variations. The question of attainment of a global or discrete minimum is irrelevant here and bypassed by a consequent discussion of infima instead of minima, e.g., for any  $\ell = 0, 1, 2, \dots$ ,

$$E_\ell := \inf E(V_\ell) := \inf_{v_\ell \in V_\ell} E(v_\ell) \in \mathbb{R} \cup \{\pm\infty\}.$$

We emphasize that there is no nestedness assumption on the finite element spaces and so the convergence of the infimal energies  $E_\ell$  does *not* follow automatically. In fact, it is stated in the following theorem as a conclusion. We remark that an extension of the following result to non-homogeneous Dirichlet conditions is not straightforward since, by approximating the boundary condition, the discrete admissible set would not be contained in  $\mathcal{A}_\infty$  any more.

**Theorem 2.1 Finite Element Failure  $\Leftrightarrow$  Lavrentiev Phenomenon.** *If  $W : \bar{\Omega} \times \mathbb{R}^m \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$  is continuous then  $\lim_{\ell \rightarrow \infty} E_\ell = \inf E(\mathcal{A}_\infty)$  and, in particular,*

$$\lim_{\ell \rightarrow \infty} E_\ell = \inf E(\mathcal{A}_1) \quad \Leftrightarrow \quad \inf E(\mathcal{A}_1) = \inf E(\mathcal{A}_\infty).$$

The direction  $\Rightarrow$  in the theorem's assertion is obvious from the introduction and  $V_\ell \subset \mathcal{A}_\infty$ :

$$\inf E(\mathcal{A}_\infty) \leq \liminf_{\ell \rightarrow \infty} E_\ell = \inf E(\mathcal{A}_1) \leq \inf E(\mathcal{A}_\infty).$$

The converse  $\Leftarrow$  requires a density argument stated in terms of the nodal interpolation operator. Given a continuous function  $v : \bar{\Omega} \rightarrow \mathbb{R}^m$  and a triangulation  $\mathcal{T}_\ell$  the nodal interpolation  $v_\ell := I_\ell v$  of  $v$  is defined on each simplex  $T \in \mathcal{T}_\ell$  with vertices  $z_1, \dots, z_{n+1}$  through linear interpolation of the values  $v(z_j)$  at the  $n + 1$  vertices  $z_j$ .

**Lemma 2.1.** *There exists a constant  $C$ , which depends only on  $C_{\text{shaperegular}}$ , such that, for any  $v \in W^{1,\infty}(\Omega; \mathbb{R}^m)$ , the piecewise affine function  $v_\ell = I_\ell v$  satisfies*

$$\|v_\ell\|_{W^{1,\infty}(\Omega)} \leq C \|v\|_{W^{1,\infty}(\Omega)} \quad \text{for all } \ell = 0, 1, 2, \dots$$

Moreover,  $v_\ell \rightarrow v$  in  $L^\infty(\Omega; \mathbb{R}^m)$ , and  $Dv_\ell \rightarrow Dv$  pointwise a.e. in  $\Omega$ , as  $\ell \rightarrow \infty$ .

*Proof.* The stability of the nodal interpolation operator as well as the convergence in the  $L^\infty$ -norm are standard results and can, for example, be found in [10].

The theorem of Rademacher implies that, for almost all  $x$  in some simplex  $T$ ,  $Dv(x)$  exists in the sense of a Fréchet derivative, i.e.,

$$Dv(x)(y - x) = v(y) - v(x) + o(|x - y|),$$

for some function  $y \mapsto o(|x - y|)$  with

$$\lim_{y \rightarrow x} o(|x - y|)/|x - y| = 0.$$

Fix some  $x \in \Omega$  so that, for any  $\ell \in \mathbb{N}_0$ ,  $x$  lies in the interior of an element  $T \in \mathcal{T}_\ell$  then

$$\begin{aligned} Dv(x)(z_j - z_k) &= v_\ell(z_j) - v_\ell(z_k) + o(|x - z_j|) + o(|x - z_k|) \\ &= Dv_\ell(x)(z_j - z_k) + o(|x - z_j|) + o(|x - z_k|) \quad \text{for all } j, k = 1, \dots, n + 1, \end{aligned}$$

where  $|\cdot|$  denotes the  $\ell^2$ -norm of a vector, or as below, the Frobenius norm of a matrix. Since the tangential vectors are linearly independent and the interior angles do not deteriorate we have

$$\sup_{j,k=1,\dots,n+1} (Dv(x) - Dv_\ell(x))(z_j - z_k) \geq c |Dv(x) - Dv_\ell(x)| r_T,$$

where  $c$  depends only on  $C_{\text{shaperegular}}$ . It now follows easily that

$$\lim_{\ell \rightarrow \infty} |Dv(x) - Dv_\ell(x)| = 0.$$

□

*Proof of Theorem 2.1.* Given  $v \in \mathcal{A}_\infty$  and its nodal interpolant  $v_\ell := I_\ell v$  for all  $\ell \in \mathbb{N}_0$ , the previous lemma shows that

$$\lim_{\ell \rightarrow \infty} (v_\ell(x), Dv_\ell(x)) = (v(x), Dv(x)) \in \mathbb{R}^m \times \mathbb{R}^{m \times n} \quad \text{for a.e. } x \in \Omega.$$

Since  $W$  is continuous this yields pointwise convergence of the energy density

$$\lim_{\ell \rightarrow \infty} W(x, v_\ell(x), Dv_\ell(x)) = W(x, v(x), Dv(x)) \quad \text{for a.e. } x \in \Omega.$$

Furthermore, the boundedness of  $v_\ell$  in  $W^{1,\infty}(\Omega)$  and the assumption that  $W$  is continuous implies that  $W(x, v_\ell(x), Dv_\ell(x))$  is bounded uniformly in  $x$  and  $\ell$ . Consequently, Lebesgue's dominated convergence theorem shows

$$\lim_{\ell \rightarrow \infty} \int_{\Omega} W(x, v_\ell(x), Dv_\ell(x)) dx = \int_{\Omega} W(x, v(x), Dv(x)) dx = E(v).$$

Therefore,

$$\inf E(\mathcal{A}_\infty) \leq \liminf_{\ell \rightarrow \infty} E_\ell \leq \limsup_{\ell \rightarrow \infty} E_\ell \leq \lim_{\ell \rightarrow \infty} E(v_\ell) = E(v).$$

Since  $v$  was an arbitrary element in  $\mathcal{A}_\infty$ , we deduce

$$\liminf_{\ell \rightarrow \infty} E_\ell = \limsup_{\ell \rightarrow \infty} E_\ell = \inf E(\mathcal{A}_\infty).$$

In particular, we can conclude that  $\lim_{\ell \rightarrow \infty} E_\ell = \inf E(\mathcal{A}_\infty)$  exists. From this, the assertion of Theorem 2.1 follows immediately.  $\square$

### 3. Penalisation and Discrete Scheme

In many examples there exists a *coupling function*

$$\gamma : \Omega \times \mathbb{R}^m \times \mathbb{R}^{m \times n} \rightarrow \mathbb{M},$$

where  $\mathbb{M} \equiv \mathbb{R}^\mu$  is a space of matrices, and an *extended energy density*

$$\phi : \Omega \times \mathbb{R}^m \times \mathbb{R}^{m \times n} \times \mathbb{M} \rightarrow \mathbb{R},$$

such that the energy density  $W$  is given by

$$W(x, v, F) := \phi(x, v, F, \gamma(x, v, F)),$$

for all  $x \in \Omega$ ,  $v \in \mathbb{R}^m$ ,  $F \in \mathbb{R}^{m \times n}$ . In this case, we also define

$$\Phi(v, \eta) := \int_{\Omega} \phi(x, v(x), Dv(x), \eta(x)) dx \quad \text{for } (v, \eta) \in \mathcal{A}_1 \times L^1(\Omega; \mathbb{M}),$$

and, with the abbreviation  $\gamma(\cdot, v, Dv)(x) := \gamma(x, v(x), Dv(x))$  for  $x \in \Omega$ , we observe that

$$E(v) = \Phi(v, \gamma(\cdot, v, Dv)). \tag{3.1}$$

**Example 3.1 Polyconvex Materials.** By definition, at almost all material points  $x \in \Omega$  and all  $v \in \mathbb{R}^m$ , a polyconvex energy density  $W(x, v, \cdot) : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$  can be written in the form

$$W(x, v, F) = \phi(x, v, \gamma(F)),$$

where  $\phi$  is convex in its third component (with  $x, v$  fixed), and  $\gamma : \mathbb{R}^{m \times n} \rightarrow \mathbb{M}$  maps a deformation gradient  $F$  to the vector of minors (sub-determinants) of  $F$  and  $\mathbb{M}$  is the space of all those minors (e.g.  $\mathbb{M} = \mathbb{R}^{19}$  for  $m = n = 3$  and  $\mathbb{M} = \mathbb{R}^5$  for  $m = n = 2$ ).

**Example 3.2 Decoupling the Gradient.** For stored energy functions  $W : \Omega \times \mathbb{R}^m \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$  where no obvious coupling mechanism is present, it is sometimes useful to let  $\mathbb{M} = \mathbb{R}^{m \times n}$  and consider

$$\phi(x, v, F, \eta) := W(x, v, \eta) \quad \text{and} \quad \gamma(x, v, F) := F.$$

This decoupling of the gradient variable will help us to overcome the Lavrentiev gap phenomenon.

On the continuous level this looks as a trivial complication of the formulation but the point is that the discretisation relaxes the condition

$$\eta = \gamma(x, v, F) \quad \text{in} \quad W(x, v, F) = \phi(x, v, F, \eta).$$

Since the immediate substitution cannot detect singular minimisers with a Lavrentiev phenomenon the ‘coupling’  $\eta = \gamma(x, v(x), Dv(x))$  will be weakened by introducing a penalty functional,

$$\Psi_\ell : L^1(\Omega; \mathbb{M}) \times L^1(\Omega; \mathbb{M}) \rightarrow \mathbb{R} \cup \{+\infty\},$$

which is written, via some density  $\psi_\ell : \Omega \times \mathbb{M} \times \mathbb{M} \rightarrow [0, \infty]$ , as

$$\Psi_\ell(\eta, \zeta) := \int_{\Omega} \psi_\ell(x, \eta(x), \zeta(x)) dx \quad \text{for } \eta, \zeta \in L^1(\Omega; \mathbb{M}).$$

The proposed discrete minimisation problem reads: Minimise the discrete energy

$$E_\ell(v, \eta) := \Phi(v, \eta) + \Psi_\ell(\eta, \gamma(\cdot, v, Dv)),$$

over  $(v, \eta) \in V_\ell \times Y_\ell$  where  $V_\ell$  and  $Y_\ell$  are suitable finite element spaces.

**Example 3.3 Penalisation.** A typical class of distance functionals is given for  $1 \leq p < \infty$  and positive parameters  $\varepsilon_\ell$  which possibly depend on the position  $x$  in the spatial domain (e.g., piecewise constant with respect to the triangulation  $\mathcal{T}_\ell$ ) and

$$\psi_\ell(x, \eta, \zeta) := \varepsilon_\ell^{-1} |\eta - \zeta|^p,$$

for all  $x \in \Omega$  and  $\eta, \zeta \in \mathbb{M}$ .

## 4. Polyconvex Energy Densities

An important class of energy functionals, especially in the field of nonlinear elasticity, are those where the stored energy density is polyconvex. As a prototypical model problem, we consider the stored energy density

$$W(x, u, F) = \phi(x, F, \det F) - f(x) \cdot u, \quad (4.1)$$

where  $f \in L^q(\Omega)^n$  for some  $q > 1$ , and  $\phi : \Omega \times \mathbb{R}^{n \times n} \times \mathbb{R} \rightarrow [0, +\infty]$ ,  $n \geq 2$ . We assume throughout this section that  $\phi$  satisfies

$$\begin{aligned} |F|^n + \Gamma(\eta) &\lesssim \phi(x, F, \eta) \lesssim 1 + |F|^n + \Gamma(\eta), \quad \text{and} \\ \phi(x, \cdot, \cdot) &\text{ is convex and l.s.c. in } \mathbb{R}^{n \times n} \times \mathbb{R} \quad \text{for a.a. } x \in \Omega, \end{aligned} \quad (4.2)$$

where  $\Gamma : \mathbb{R} \rightarrow [0, +\infty]$  is convex and has superlinear growth, i.e.,  $\liminf_{|s| \rightarrow \infty} \Gamma(s)/s = +\infty$  [3, 11]. We remark that the growth condition  $|F|^n + \Gamma(\eta)$  may be replaced by  $|F|^p$  for some  $p > n$ . In fact, the latter implies the former.

The space of admissible functions is defined as

$$V = u_D + W_0^{1,n}(\Omega)^n,$$



where  $u_D \in W^{1,n}(\Omega)^n$  and  $E(u_D) < +\infty$ . Under these conditions the minimization problem

$$u \in \operatorname{argmin} E(V), \tag{4.3}$$

has at least one solution [12, Theorem 2.10].

To discretize the problem we fix a sequence  $u_{D,\ell} \in P^1(\mathcal{T}_\ell)^n$  such that  $u_{D,\ell} \rightarrow u_D$  strongly in  $W^{1,n}(\Omega)^n$ , and we discretize  $V$  and  $L^1(\Omega)$ , respectively, by

$$V_\ell = u_{D,\ell} + P_0^1(\mathcal{T}_\ell)^n, \quad \text{and} \quad Y_\ell = P^0(\mathcal{T}_\ell).$$

We remark that, throughout,  $V$  denotes the admissible set,  $V_\ell$  the discrete admissible set, and  $Y_\ell$  the discrete admissible set for the penalty variable.

Further, we assume that we have a *penalty functional*  $\Psi : L^1(\Omega)^2 \rightarrow [0, +\infty]$  such that, for all sequences  $(\eta_\ell)$  and  $(\zeta_\ell) \subset L^1(\Omega)$ ,

$$\Psi(\eta_\ell, \zeta_\ell) \rightarrow 0 \quad \Leftrightarrow \quad \|\eta_\ell - \zeta_\ell\|_{L^1} \rightarrow 0. \tag{4.4}$$

Given a sequence  $\varepsilon_\ell \searrow 0$ , we discretize (4.3) by

$$(u_\ell, \xi_\ell) \in \operatorname{argmin} E_\ell(V_\ell, Y_\ell),$$

where

$$\begin{aligned} E_\ell(v_\ell, \eta_\ell) &= \Phi(v_\ell, \eta_\ell) + \varepsilon_\ell^{-1} \Psi(\det Dv_\ell, \eta_\ell) \\ &= \int_{\Omega} (\phi(x, Dv_\ell, \eta_\ell) - f \cdot v_\ell) dx + \varepsilon_\ell^{-1} \Psi(\det Dv_\ell, \eta_\ell). \end{aligned}$$

**Theorem 4.1.** *Assume that (4.1), (4.2), and (4.4) hold. Then there exists a sequence  $\varepsilon_\ell \searrow 0$  such that, for any sequence  $(u_\ell, \xi_\ell) \in V_\ell \times X_\ell$  of approximate minimizers, that is,*

$$|E_\ell(u_\ell, \xi_\ell) - \inf E_\ell(V_\ell, Y_\ell)| \rightarrow 0 \quad \text{as } \ell \rightarrow \infty,$$

we have

$$\Phi(u_\ell, \xi_\ell) \rightarrow \inf E(V) \quad \text{and} \quad \varepsilon_\ell^{-1} \Psi(\det Du_\ell, \xi_\ell) \rightarrow 0.$$

Moreover, the family  $\{u_\ell; \ell \in \mathbb{N}\}$  is precompact in the weak topology of  $W^{1,n}(\Omega)^n$  and each accumulation point  $u$  is a minimizer of  $E$  in  $V$ . In particular, there exists a subsequence  $\ell_k \nearrow \infty$  such that

$$\begin{aligned} u_{\ell_k} &\rightharpoonup u && \text{weakly in } W^{1,n}(\Omega)^n, \\ \xi_{\ell_k} &\rightharpoonup \det Du && \text{weakly in } L^1(\Omega), \end{aligned}$$

where  $u$  solves (4.3).

The proof of Theorem 4.1 is contained in the following three lemmas.

**Lemma 4.1.** *Assume that (4.1), (4.2) and (4.4) hold. For every  $v \in V$  there exists a sequence  $(v_\ell, \eta_\ell) \in V_\ell \times Y_\ell$  such that*

$$v_\ell \rightarrow v \quad \text{strongly in } W^{1,n}(\Omega)^n, \tag{4.5}$$

$$\lim_{\ell \rightarrow \infty} \Psi(\det Dv_\ell, \eta_\ell) = 0, \quad \text{and} \tag{4.6}$$

$$\lim_{\ell \rightarrow \infty} \Phi(v_\ell, \eta_\ell) = \Phi(v, \det Dv) = E(v). \tag{4.7}$$

*Proof.* Let  $v \in V$ . If  $E(v) = +\infty$ , then we take an arbitrary sequence  $v_\ell \in V_\ell$  converging strongly in  $W^{1,n}(\Omega)^n$  to  $v$ , and  $\eta_\ell = \det Dv_\ell$ . From the lower semicontinuity of  $E$  we obtain that  $E(v_\ell) = \Phi(v_\ell, \eta_\ell) \rightarrow \infty$  as  $\ell \rightarrow +\infty$ , since, otherwise,  $E(v)$  would be finite. Moreover, we have  $\Psi(\det Dv_\ell, \eta_\ell) = 0$ .

We may now assume that  $E(v) < \infty$ . We take an arbitrary sequence  $v_\ell \in V_\ell$  such that  $v_\ell \rightarrow v$  strongly in  $W^{1,n}(\Omega)^n$  which also implies  $\det Dv_\ell \rightarrow \det Dv$  strongly in  $L^1(\Omega)$ . The variable  $\eta_\ell \in Y_\ell$  is defined as

$$\eta_\ell(x) = |T|^{-1} \int_T \det Dv \, dx \quad x \in T \in \mathcal{T}_\ell.$$

It follows that  $\eta_\ell \rightarrow \det Dv$  strongly in  $L^1(\Omega)$  and in particular that  $\Psi(\det Dv_\ell, \eta_\ell) \rightarrow 0$ . Thus, we have shown (4.5) and (4.6).

To prove (4.7) we first use Jensen's inequality to estimate, for  $x \in T \in \mathcal{T}_\ell$ ,

$$\Gamma(\eta_\ell(x)) = \Gamma\left(|T|^{-1} \int_T \det Dv \, dx\right) \leq |T|^{-1} \int_T \Gamma(\det Dv) \, dx =: \Gamma_\ell(x),$$

i.e.,  $\Gamma_\ell$  is a majorant for  $\Gamma(\eta_\ell)$ . From its definition, and since  $\Gamma(\det Dv) \in L^1(\Omega)$  (which follows from the fact that  $E(v)$  is finite), it follows immediately that  $\Gamma_\ell \rightarrow \Gamma(\det Dv)$  strongly in  $L^1(\Omega)$ .

Hence, we obtain that

$$\phi(x, Dv_\ell, \eta_\ell) \lesssim 1 + |Dv_\ell|^n + \Gamma_\ell =: a_\ell,$$

where  $a_\ell$  is strongly convergent in  $L^1(\Omega)$ . For any subsequence we can extract a further subsequence such that  $(Dv_\ell, \eta_\ell) \rightarrow (Dv, \eta)$  pointwise, and hence we can use a variant of Lebesgue's dominated convergence theorem [15, Sec. 1.3, Th. 4] to deduce (4.7).  $\square$

**Lemma 4.2.** *Assume that (4.1), (4.2), and (4.4) hold. There exists a sequence  $\varepsilon_\ell \searrow 0$  such that*

$$\limsup_{\ell \rightarrow \infty} \min E_\ell(V_\ell, Y_\ell) \leq \min E(V). \tag{4.8}$$

*Proof.* Let  $u \in \operatorname{argmin} E(V)$  and let  $(u_\ell, \xi_\ell)$  be the sequence constructed in Lemma 4.1 (for  $v = u$ ). Then

$$\Psi(\det Du_\ell, \xi_\ell) \rightarrow 0,$$

and choosing  $\varepsilon_\ell = \Psi(\det Du_\ell, \xi_\ell)^{1/2}$  we obtain

$$\limsup_{\ell \rightarrow \infty} \inf E_\ell(V_\ell, Y_\ell) \leq \limsup_{\ell \rightarrow \infty} E_\ell(u_\ell, \xi_\ell) = E(u).$$

$\square$

In the previous lemma, we showed that it is possible to choose a sequence  $\varepsilon_\ell$  such that the upper bound (4.8) holds. It remains to show that the limit is in fact equal.

**Lemma 4.3.** *Assume that (4.1), (4.2), and (4.4) hold. Suppose that a sequence  $\varepsilon_\ell \searrow 0$  is fixed. Suppose furthermore that  $u_\ell \in V_\ell, \xi_\ell \in Y_\ell$  such that*

$$\limsup_{\ell \rightarrow \infty} E_\ell(u_\ell, \xi_\ell) \leq \inf E(V), \tag{4.9}$$

then there exists a subsequence  $\ell_k \uparrow \infty$  and  $u \in \operatorname{argmin} E(V)$  such that

$$\begin{aligned} u_{\ell_k} &\rightharpoonup u && \text{weakly in } W^{1,n}(\Omega)^n \\ \xi_{\ell_k} &\rightharpoonup \det Du && \text{weakly in } L^1(\Omega), \end{aligned}$$

and moreover, we have separate convergence of the entire sequences of energy contributions:

$$\Phi(u_\ell, \xi_\ell) \rightarrow E(u), \quad \text{and} \quad \varepsilon_\ell^{-1} \Psi(\det Du_\ell, \xi_\ell) \rightarrow 0.$$

*Proof.* It follows from (4.9) that  $E_\ell(u_\ell, \xi_\ell)$  is bounded by some constant  $M$ . Using (4.2) and the assumption that  $u_D$  has finite energy, we obtain

$$M \geq E_\ell(u_\ell, \xi_\ell) \gtrsim \|\nabla u_\ell\|_{L^n}^n - C\|u_\ell\|_{L^{q'}} + \int_{\Omega} \Gamma(\xi_\ell) dx + \varepsilon_\ell^{-1} \Psi(\det Du_\ell, \xi_\ell),$$

and since  $W^{1,n}(\Omega)^n$  is continuously embedded in  $L^{q'}(\Omega)^n$ , there exists  $M' \in \mathbb{R}$  such that

$$\|u_\ell\|_{W^{1,n}}^n + \int_{\Omega} \Gamma(\xi_\ell) dx + \varepsilon_\ell^{-1} \Psi(\det Du_\ell, \xi_\ell) \leq M'.$$

We can therefore deduce the existence of a subsequence  $\ell_k \nearrow \infty$ , and of functions  $u \in W^{1,n}(\Omega)^n$  and  $\xi \in L^1(\Omega)$  such that

$$u_\ell \rightharpoonup u \text{ weakly in } W^{1,n}(\Omega)^n \quad \text{and} \quad \xi_\ell \rightharpoonup \xi \text{ weakly in } L^1(\Omega).$$

(We note that the superlinear bound implies equi-integrability of the sequence  $(\xi_\ell)$  which implies its precompactness in the weak topology of  $L^1(\Omega)$  [14, Cor. IV.8.11].)

Since  $\det Du_\ell \rightharpoonup' \det Du$  in the sense of distributions [12, Sec. 4.2, Th. 2.6, (5)], and using (4.4), it follows that  $\xi = \det Du$ . Using sequential weak lower semi-continuity of energies with convex integrands [12, Sec. 3.3, Th. 3.4] we can estimate

$$\begin{aligned} E(u) &\leq \liminf_{k \rightarrow \infty} \int_{\Omega} \left( \phi(x, u_{\ell_k}, \xi_{\ell_k}) - f \cdot u_{\ell_k} \right) dx \\ &\leq \liminf_{k \rightarrow \infty} \int_{\Omega} \left( \phi(x, u_{\ell_k}, \xi_{\ell_k}) - f \cdot u_{\ell_k} \right) dx \\ &\quad + \limsup_{k \rightarrow \infty} \varepsilon_{\ell_k}^{-1} \Psi(\det Du_{\ell_k}, \xi_{\ell_k}) \\ &\leq \limsup_{\ell \rightarrow \infty} E_\ell(u_\ell, \xi_\ell) \leq \inf E(V). \end{aligned}$$

It follows therefore that  $E(u) = \inf E(V)$ . Moreover, this implies that all inequalities in the above chain of estimates must be equalities, and hence,

$$\limsup_{k \rightarrow \infty} \varepsilon_{\ell_k}^{-1} \Psi(\det Du_{\ell_k}, \xi_{\ell_k}) = 0.$$

Since the proof applies also if we begin with an arbitrary subsequence, it follows that the energy of the entire sequence converges in this sense.  $\square$

*Proof of Theorem 4.1.* Lemma 4.2 guarantees the existence of a sequence  $\varepsilon_\ell \searrow 0$  such that the conditions of Lemma 4.3 are satisfied. Hence, Lemma 4.3 guarantees the existence of a weakly convergent subsequence of approximate minimizers  $E_\ell$  and establishes the various convergence statements in the theorem.  $\square$

**Remark 4.1.** 1. In practise, the condition that  $\Psi$  is continuous in the strong topology of  $L^1(\Omega; \mathbb{R}^n)$  requires that  $\Psi$  takes the form

$$\Psi(\eta, \zeta) = \int_{\Omega} \psi(|\eta - \zeta|) dx,$$

where  $\psi$  has 1-growth at infinity. Typical penalty densities  $\psi$  are  $\psi(t) = |t|$ , or, if one prefers a smooth functional,  $\psi(t) = (t^2 + 1)^{1/2} - 1$ . The condition (4.4) can be obtained, for example, by requiring that  $\psi \geq 0$  and  $\psi(t) = 0$  if and only if  $t = 0$ .

2. If  $\phi$  satisfies a stronger growth condition, for example  $\phi(x, F, g) \gtrsim |F|^p$  for some  $p > n$  then this additional integrability allows us to use a penalty functional which is only continuous in  $L^{p/n}(\Omega; \mathbb{R}^n)$ .

3. We have only shown the existence of some sequence  $\varepsilon_\ell$  for which we obtain convergence of the penalty method. We will show in Section 7 below how this sequence can be constructed in practise.

4. More general polyconvex material models where  $\phi$  depends on all minors of the gradient can be easily incorporated in our analysis. One would then have to decouple all minors which appear in the definition of the functional. Similar convergence can then be obtained whenever the growth conditions from above and below are the same and are sufficiently strong so that the direct method can be applied.

## 5. Examples with Lavrentiev Phenomenon

In many problems decoupling the gradient is sufficient, and it is the goal of this section to make this precise. This is possible whenever  $W$  is convex in the third component, but it is also a useful approach if it is unclear which variable should be relaxed. We begin again with a more general discussion which we then make precise at two classes of problems, general one-dimensional functionals with continuous integrands, and higher-dimensional examples with mild  $v$ -dependence of the integrand.

We assume throughout that  $W = \phi : \Omega \times \mathbb{R}^m \times \mathbb{R}^{m \times n} \rightarrow (-\infty, +\infty]$  is lower semi-continuous in all three variables, continuous at every point  $(x, v, \eta)$  where  $\phi(x, v, \eta) < \infty$ , and that it satisfies the lower bound

$$\phi(x, v, \eta) \gtrsim -1 - |v|^q, \tag{5.1}$$

where  $1 \leq q < n/(n - 1)$  if  $n \geq 2$  and  $1 \leq q < \infty$  if  $n = 1$ . This implies in particular that, for  $v \in W^{1,1}(\Omega)^m$  and  $\eta \in L^1(\Omega)^{m \times n}$ , the functionals

$$\Phi(v, \eta) = \int_{\Omega} W(x, v, \eta) dx, \quad \text{and} \quad E(v) = \Phi(v, Dv),$$

are well-defined in  $(-\infty, +\infty]$ . Let  $u_D \in W^{1,1}(\Omega)^m$  such that  $E(u_D) < \infty$  and define  $V = u_D + W_0^{1,1}(\Omega)^m$ .

We will assume that the penalty has 1-growth, namely, that there exists a continuous penalty density  $\psi : \mathbb{R}^{m \times n} \rightarrow [0, \infty)$  satisfying

$$\begin{aligned} |\eta| - 1 &\lesssim \psi(\eta) \lesssim |\eta| + 1 && \text{for all } \eta \in \mathbb{R}^{m \times n}, \quad \text{and} \\ \psi(\eta) &= 0 && \text{if and only if } \eta = 0, \end{aligned} \tag{5.2}$$

such that the functional  $\Psi$  is of the form

$$\Psi(\eta, \zeta) = \int_{\Omega} \psi(\eta - \zeta) dx \quad \text{for all } \eta, \zeta \in L^1(\Omega)^{m \times n}. \tag{5.3}$$

To discretize the problem of minimizing  $E$  over  $V$  we take  $u_{D,\ell} \in P^1(\mathcal{T}_\ell)$  such that  $u_{D,\ell} \rightarrow u_D$  strongly in  $W^{1,1}(\Omega)^m$ , and define

$$V_\ell = u_{D,\ell} + P_0^1(\mathcal{T}_\ell)^m \quad \text{and} \quad Y_\ell = P^0(\mathcal{T}_\ell)^{m \times n},$$

to discretize, respectively, the variables  $u$  and  $\eta$ . We approximate  $\Phi$  using the midpoint rule: For  $v_\ell \in P^1(\mathcal{T}_\ell)^m$ , we set  $\bar{v}_\ell(x) = (v_\ell)_T := |T|^{-1} \int_T v_\ell dx$  for  $x \in T \in \mathcal{T}_\ell$ , and for  $v_\ell \in V_\ell$  and  $\eta_\ell \in Y_\ell$ , we define

$$\Phi_\ell(v_\ell, \eta_\ell) = \int_{\Omega} \phi(\bar{x}_\ell, \bar{v}_\ell, \eta_\ell) dx = \sum_{T \in \mathcal{T}_\ell} |T| \phi((x)_T, (v_\ell)_T, \eta_\ell|_T).$$

The functional  $\Phi_\ell$  is extended in an obvious way to  $V_\ell \times L^1(\Omega)^{m \times n}$ .

**Remark 5.1.** We could have included a quadrature approximation in our analysis in Section 4 as well. For the sake of simplicity, we decided not to do so. In the present case, we are in fact unable to prove convergence of the penalty method *without* the quadrature approximation. The reason for this is essentially that we have chosen  $\eta_\ell \in P^0(\mathcal{T}_\ell)^{m \times n}$  and hence we can only adjust its value to a single point within each element. Since we assume no control on  $\phi$  from above we cannot control an integral over an element from information at a single quadrature point.

Our first aim is an approximation result akin to Lemma 4.1. In Lemma 5.1 below we reduce this task to the following general condition which can be quite easily checked for different problems: *for all  $v \in V$  there exists a function  $\zeta \in L^1(\Omega)^{m \times n}$  and a sequence  $v_\ell \in V_\ell$  such that the following conditions are satisfied:*

$$\begin{aligned} (i) \quad &\phi(x, v, \zeta) \in L^1(\Omega), \\ (ii) \quad &v_\ell \rightarrow v \text{ strongly in } W^{1,1}(\Omega)^m, \text{ and} \\ (iii) \quad &\limsup_{\ell \rightarrow \infty} \Phi_\ell(v_\ell, \zeta) \leq \Phi(v, \zeta). \end{aligned} \tag{5.4}$$

**Example 5.1 1D Examples.** Suppose that  $n = 1$ , that  $\phi : \Omega \times \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}$  is globally continuous, and assume that  $u_{D,\ell} = u_D$  for all  $\ell$ . This class includes in particular problems of Maniá type [7, 23].

We now prove that (5.4) holds under this assumption. Let  $v \in V$  and let  $v_\ell$  be its piecewise affine nodal interpolant. Then  $v_\ell \rightarrow v$  strongly in  $W^{1,1}(\Omega)^m$ ,  $(\bar{x}_\ell, \bar{v}_\ell(x)) \rightarrow (x, v(x))$  uniformly in  $\Omega$  and, since  $\phi$  is globally continuous,

$$\Phi_\ell(v_\ell, \zeta) \rightarrow \Phi(v, \zeta),$$

for any fixed  $\zeta \in \mathbb{R}^m$ . □

**Example 5.2 Weak Coupling of  $u$  and  $Du$ .** Suppose that, in addition to (5.1),

$$\phi(x, v, \eta) \lesssim |v|^q + \Gamma(\eta), \quad (5.5)$$

where  $\Gamma : \mathbb{R}^{m \times n} \rightarrow [0, +\infty]$  is proper. We note that this class includes in particular the example of Foss, Hrusa, and Mizel [16] and Ball's example of cavitation [4].

We now prove that (5.4) holds under this assumption. Let  $v \in V$  and take  $v_\ell \in V_\ell$  converging strongly in  $W^{1,1}(\Omega)^m \cap L^q(\Omega)^m$  to  $v$ . In particular, we also have  $\bar{v}_\ell \rightarrow v$  strongly in  $L^q(\Omega)^m$  by Lebesgue's differentiation theorem. Further, let  $\zeta \in \mathbb{R}^{m \times n}$  such that  $\Gamma(\zeta) < +\infty$ . In view of the growth condition imposed in (5.5) we obtain  $\phi(x, v(x), \zeta) \in L^1(\Omega)$ . Let  $\ell_j \nearrow \infty$  be a subsequence such that

$$\limsup_{\ell \rightarrow \infty} \Phi_\ell(v_\ell, \zeta) = \lim_{j \rightarrow \infty} \Phi_{\ell_j}(v_{\ell_j}, \zeta).$$

Upon extracting a further subsequence we may assume that  $(\bar{x}_{\ell_j}, \bar{v}_{\ell_j}) \rightarrow (x, v)$  pointwise a.e. in  $\Omega$ . Since  $\phi(x, v(x), \zeta) \in L^1(\Omega)$  it is finite for a.a.  $x \in \Omega$  and hence continuous at those points. We therefore obtain

$$\lim_{j \rightarrow \infty} \phi(\bar{x}_j, \bar{v}_j, \zeta) = \phi(x, v, \zeta) \quad \text{pointwise a.e. in } \Omega.$$

The majorant

$$\phi(\bar{x}_{\ell_j}, \bar{v}_{\ell_j}, \zeta) \leq |\bar{v}_{\ell_j}|^q + \Gamma(\zeta),$$

is strongly convergent in  $L^1(\Omega)$  and hence we can use Fatou's Lemma to obtain (5.4) (iii).  $\square$

Having shown that (5.4) indeed holds for several interesting problem classes we establish the basic approximation result, which it implies.

**Lemma 5.1.** *Fix  $\varepsilon > 0$  and suppose that (5.2), (5.3) and (5.4) hold; then, for every  $v \in V$  there exists a sequence  $(v_\ell, \eta_\ell) \in V_\ell \times Y_\ell$  such that*

$$\limsup_{\ell \rightarrow \infty} [\Phi(v_\ell, \eta_\ell) + \varepsilon^{-1} \Psi(Dv_\ell, \eta_\ell)] \leq E(v).$$

Moreover, the sequence  $v_\ell$  can be chosen independent of the value of  $\varepsilon$ .

*Proof.* We take the sequence  $v_\ell$  specified in (5.4). For every  $T \in \mathcal{T}_\ell$  and  $x \in T$  we define

$$\bar{\phi}_\ell(x) = \inf_{\xi \in \mathbb{R}^{m \times n}} [\phi(\bar{x}_\ell(x), \bar{v}_\ell(x), \xi) + \varepsilon^{-1} \psi(\xi - Dv_\ell(x))].$$

Since  $\bar{v}_\ell$  and  $\bar{x}_\ell$  are piecewise constant  $\bar{\phi}_\ell$  may also be chosen as a piecewise constant function and it follows from the growth condition on  $\phi$  from below that it is finite. In particular, it is measurable and its integral is well-defined with a value in  $(-\infty, +\infty]$ .

There exists a subsequence  $\ell_j \nearrow \infty$  such that

$$\begin{aligned} \limsup_{\ell \rightarrow \infty} \int_{\Omega} \bar{\phi}_\ell dx &= \lim_{j \rightarrow \infty} \int_{\Omega} \bar{\phi}_{\ell_j}, \quad \text{and} \\ (\bar{v}_{\ell_j}, Dv_{\ell_j}) &\rightarrow (v, Dv) \quad \text{pointwise a.e. in } \Omega. \end{aligned}$$

From the definition of  $\bar{\phi}_\ell$ , we have

$$\bar{\phi}_\ell \leq \phi(\bar{x}_\ell, \bar{v}_\ell, Dv) + \varepsilon^{-1}\psi(Dv - Dv_\ell) \quad \text{for a.a. } x \in \Omega,$$

and since we assumed that  $\phi$  is continuous at every point where it is finite, and that  $\psi$  is globally continuous, we obtain

$$\limsup_{j \rightarrow \infty} \bar{\phi}_{\ell_j}(x) \leq \phi(x, v(x), Dv(x)) \quad \text{for a.a. } x \in \Omega. \quad (5.6)$$

Again using the definition of  $\bar{\phi}_\ell$  we obtain the majorant

$$\bar{\phi}_\ell \leq \phi(\bar{x}_\ell, \bar{v}_\ell, \zeta) + \varepsilon^{-1}\psi(\zeta - Dv_\ell) =: m_\ell,$$

where  $\zeta \in L^1(\Omega)^{m \times n}$  is taken from (5.4). Since  $\phi$  is continuous at  $(x, v(x), \zeta(x))$ , for a.a.  $x \in \Omega$ , it follows from (5.4) (ii) that

$$m_{\ell_j}(x) \rightarrow m(x) := \phi(x, v(x), \zeta(x)) + \varepsilon^{-1}\psi(\zeta(x) - Dv(x)) \quad \text{for a.a. } x \in \Omega.$$

Condition (5.4) (iii) translates as

$$\liminf_{j \rightarrow \infty} \int_{\Omega} m_{\ell_j} dx \leq \int_{\Omega} m dx.$$

Applying Fatou's lemma to the sequence  $m_\ell - \bar{\phi}_\ell$  gives

$$\int_{\Omega} \liminf_{j \rightarrow \infty} (m_{\ell_j} - \bar{\phi}_{\ell_j}) dx \leq \liminf_{j \rightarrow \infty} \int_{\Omega} (m_{\ell_j} - \bar{\phi}_{\ell_j}) dx,$$

which can, equivalently, be written as

$$\int_{\Omega} (m - \limsup_{j \rightarrow \infty} \bar{\phi}_{\ell_j}) dx \leq \int_{\Omega} m dx - \limsup_{j \rightarrow \infty} \int_{\Omega} \bar{\phi}_{\ell_j} dx,$$

and hence we obtain, using (5.6) in the last inequality,

$$\limsup_{\ell \rightarrow \infty} \int_{\Omega} \bar{\phi}_\ell dx = \limsup_{j \rightarrow \infty} \int_{\Omega} \bar{\phi}_{\ell_j} dx \leq \int_{\Omega} \limsup_{j \rightarrow \infty} \bar{\phi}_{\ell_j} dx \leq E(v).$$

It remains to show that there exists a sequence  $\eta_\ell \in Y_\ell$  such that

$$\limsup_{\ell \rightarrow \infty} \int_{\Omega} \bar{\phi}_\ell dx = \limsup_{\ell \rightarrow \infty} \Phi_\ell(v_\ell, \eta_\ell).$$

To this end we choose  $\eta_\ell \in Y_\ell$ , such that

$$\phi(\bar{x}_\ell, \bar{v}_\ell(x), \eta_\ell(x)) \leq \bar{\phi}_\ell(x) + 1/\ell \quad \text{for a.e. } x \in \Omega.$$

The existence of such functions follows from the definition of  $\bar{\phi}_\ell$ . □

Next, we will deduce from Lemma 5.1 the existence of a sequence  $\varepsilon_\ell \searrow 0$  for which the same upper bound still holds.

**Lemma 5.2.** *Suppose that (5.2), (5.3) and (5.4) hold; then there exists a sequence  $\varepsilon_\ell \searrow 0$  such that*

$$\limsup_{\ell \rightarrow \infty} \inf E_\ell(V_\ell, Y_\ell) \leq \inf E(V).$$

*Proof.* Let  $v_k \in V$  such that  $E(v_k) \leq \inf E(V) + 1/k$ . According to Lemma 5.1, for every  $k \in \mathbb{N}$ , there exists  $\ell_k \in \mathbb{N}$  such that, for all  $\ell \geq \ell_k$ ,

$$\inf_{(u_\ell, \xi_\ell) \in V_\ell \times Y_\ell} [\Phi_\ell(u_\ell, \xi_\ell) + k\Psi(\xi_\ell, Du_\ell)] \leq E(v_k) + 1/k \leq \inf E(V) + 2/k.$$

We may assume that  $\ell_k \leq \ell_{k+1}$  for all  $k$ . If we define

$$\varepsilon_\ell = 1/k \quad \text{for } \ell_k \leq \ell < \ell_{k+1}, \quad k = 1, 2, \dots,$$

and  $\varepsilon_\ell = 1$  for  $1 \leq \ell < \ell_1$ , then  $\varepsilon_\ell \searrow 0$  and

$$\inf E_\ell(V_\ell, Y_\ell) \leq \inf E(V) + 2\varepsilon_\ell \quad \text{for all } \ell \geq \ell_1.$$

□

We only need to prove a lower bound now. Here, we distinguish two cases: whether  $\phi$  is convex in the third component or only quasiconvex.

We adopt assumption (ii) in the following theorem as an abstract compactness assumption that we found difficult to verify for examples where we observe it in practise, such as the Foss/Hrusa/Mizel example in Section 7.5. Failure of this assumption will normally be displayed as an instability in the numerical calculation.

**Theorem 5.1 Convex Energies.** *Suppose that (5.2), (5.3) and (5.4) hold, and assume in addition that  $\phi$  is convex in its third component. Let  $\varepsilon_\ell \searrow 0$  be the sequence established in Lemma 5.2, and let  $(u_\ell, \xi_\ell) \in V_\ell \times Y_\ell$  be a sequence satisfying the following conditions:*

(i)  $(u_\ell, \xi_\ell)$  are approximate minimizers, i.e.,

$$|E_\ell(u_\ell, \xi_\ell) - \inf E_\ell(V_\ell, Y_\ell)| \rightarrow 0 \quad \text{as } \ell \rightarrow \infty. \quad (5.7)$$

(ii) There exists  $u \in V$  such that

$$u_\ell \rightharpoonup u \quad \text{weakly in } W^{1,1}(\Omega)^m. \quad (5.8)$$

Then  $u \in \operatorname{argmin} E(V)$ ,

$$\begin{aligned} \lim_{\ell \rightarrow \infty} \Phi_\ell(u_\ell, \xi_\ell) &= E(u), \\ \lim_{\ell \rightarrow \infty} \varepsilon_\ell^{-1} \Psi(\xi_\ell, Du_\ell) &= 0, \quad \text{and} \\ \xi_\ell &\rightharpoonup Du \quad \text{weakly in } L^1(\Omega)^{m \times n}. \end{aligned} \quad (5.9)$$



*Proof.* By the construction of  $\varepsilon_\ell$  and assumption (5.7) we have

$$\limsup_{\ell \rightarrow \infty} E_\ell(u_\ell, \xi_\ell) \leq \inf E(V).$$

In particular,  $\Psi(\xi_\ell, Du_\ell) \lesssim \varepsilon_\ell \rightarrow 0$  which implies  $\xi_\ell \rightharpoonup Du$  weakly in  $L^1(\Omega)^{m \times n}$ . We can therefore deduce that

$$E(u) \leq \liminf_{\ell \rightarrow \infty} \Phi_\ell(u_\ell, \xi_\ell).$$

Using the same arguments as in the proof of Lemma 4.3 we can conclude the proof of the theorem.  $\square$

In addition to assumption (ii) in Theorem 5.1 we require another stability assumption in the quasiconvex case. Assumption (iii) in the following theorem will be satisfied whenever singularities occur only in localized regions. This is again observed in typical numerical experiments but would be very difficult to prove rigorously.

**Theorem 5.2 Quasiconvex Energies.** *Suppose that (5.2), (5.3) and (5.4) hold, and assume in addition that  $\phi$  is quasiconvex in its third component. Let  $\varepsilon_\ell \searrow 0$  be the sequence established in Lemma 5.2 and let  $(u_\ell, \xi_\ell) \in V_\ell \times Y_\ell$  be a sequence satisfying (i) and (ii) in Theorem 5.1, as well as:*

(iii) *There exists a monotone family of subsets  $\Omega_k \nearrow \Omega$  such that*

$$\lim_{\ell \rightarrow \infty} \|\phi(\bar{x}_\ell, \bar{u}_\ell, Du_\ell) - \phi(\bar{x}_\ell, \bar{u}_\ell, \xi_\ell)\|_{L^1(\Omega_k)} = 0 \quad \text{and} \quad (5.10)$$

$$\forall k \in \mathbb{N} \quad \sup_{\ell \geq k} \|u_\ell\|_{W^{1,\infty}(\Omega_k)} < \infty. \quad (5.11)$$

Then  $u \in \operatorname{argmin} E(V)$  and the conclusion (5.9) remains true as well.

*Proof.* In view of the bound (5.11), for fixed  $k \in \mathbb{N}$ , we have

$$u_\ell \overset{*}{\rightharpoonup} u \quad \text{weakly-}^* \text{ in } W^{1,\infty}(\Omega_k)^m.$$

Since  $\phi$  is quasiconvex in its third component it follows from (5.10) that

$$\begin{aligned} \int_{\Omega_k} \phi(x, u, Du) dx &\leq \liminf_{\ell \rightarrow \infty} \int_{\Omega_k} \phi(\bar{x}_\ell, \bar{u}_\ell, Du_\ell) dx \\ &= \liminf_{\ell \rightarrow \infty} \int_{\Omega_k} \phi(\bar{x}_\ell, \bar{u}_\ell, \xi_\ell) dx. \end{aligned}$$

Using the lower bound (5.1), the compactness of the embedding  $W^{1,1}(\Omega)^m \subset L^q(\Omega)^m$ , and setting  $\Omega'_k = \Omega \setminus \Omega_k$ , we obtain

$$\begin{aligned} \int_{\Omega_k} \phi(x, u, Du) dx &\leq \liminf_{\ell \rightarrow \infty} \left( \Phi_\ell(u_\ell, \xi_\ell) - \int_{\Omega'_k} \phi(\bar{x}_\ell, \bar{u}_\ell, \xi_\ell) dx \right) \\ &\leq \liminf_{\ell \rightarrow \infty} \Phi_\ell(u_\ell, \xi_\ell) + \limsup_{\ell \rightarrow \infty} C(|\Omega'_k| + \|u_\ell\|_{L^q(\Omega'_k)}^q) \\ &= \liminf_{\ell \rightarrow \infty} \Phi_\ell(u_\ell, \xi_\ell) + C(|\Omega'_k| + \|u\|_{L^q(\Omega'_k)}^q). \end{aligned}$$

Setting  $\delta_k = C(|\Omega'_k| + \|u\|_{L^q(\Omega'_k)}^q)$  we can further estimate

$$\begin{aligned}
 \int_{\Omega_k} \phi(x, u, Du) dx &\leq \liminf_{\ell \rightarrow \infty} \Phi_\ell(u_\ell, \xi_\ell) + \delta_k \\
 &\leq \liminf_{\ell \rightarrow \infty} \Phi_\ell(u_\ell, \xi_\ell) + \limsup_{\ell \rightarrow \infty} \varepsilon_\ell^{-1} \Psi(Du_\ell, \xi_\ell) + \delta_k \\
 &\leq \limsup_{\ell \rightarrow \infty} E_\ell(u_\ell, \xi_\ell) + \delta_k \\
 &\leq \inf E(V) + \delta_k \quad \text{for all } k \in \mathbb{N}.
 \end{aligned} \tag{5.12}$$

Adding the term  $C(1 + |u|^q)$  to the integral on the left-hand side the integrand becomes non-negative and the bound becomes

$$\int_{\Omega_k} [\phi(x, u, Du) + C(1 + |u|^q)] dx \leq \inf E(V) + \int_{\Omega} C(1 + |u|^q) dx.$$

Taking the supremum over  $k$  on the left-hand side (employing, for example, the Beppo-Levi theorem), it follows that  $\phi(x, u, Du)$  is integrable and that  $u \in \operatorname{argmin} E(V)$ . Furthermore, we can let  $k \rightarrow \infty$  and thus  $\delta_k \rightarrow 0$  in (5.12) from which we can deduce the separate convergence of the energy contributions (compare also with the proof of Lemma 4.3).  $\square$

## 6. Connection with $\Gamma$ -Convergence

Our main results, Theorems 4.1, 5.1 and 5.2, can be understood as  $\Gamma$ -convergence (also known as epi-convergence) results. The purpose of the present section is to briefly explain this connection. We refer to the monographs of Braides [9] and Dal Maso [13] for an introduction to  $\Gamma$ -convergence.

We will demonstrate this point of view at the example of the polyconvex case. To this end, suppose that (4.1), (4.2), (4.4) and (5.2) hold, and define, for  $v \in W^{1,n}(\Omega)^n$ ,  $\eta \in L^1(\Omega)$  and  $\varepsilon \in [0, \infty)$ ,

$$\begin{aligned}
 F(v, \eta, \varepsilon) &= \begin{cases} E(v) & \text{if } v \in V, \eta = \det Dv, \varepsilon = 0, \\ \Phi(v, \eta) + \varepsilon^{-1} \Psi(\det Dv, \eta) & \text{if } v \in V, \varepsilon \in (0, \infty), \\ +\infty & \text{otherwise;} \end{cases} \\
 F_\ell(v, \eta, \varepsilon) &= \begin{cases} \Phi(v, \eta) + \varepsilon^{-1} \Psi(\det Dv, \eta) & \text{if } v \in V_\ell, \eta \in Y_\ell, \varepsilon \in (0, \infty), \\ +\infty & \text{otherwise.} \end{cases}
 \end{aligned}$$

A minor modification of Lemma 4.2 shows that, for each  $u \in V$ ,  $\xi = \det Du$ , there exists a sequence  $u_\ell \rightarrow u$  strongly in  $W^{1,n}(\Omega)^n$ ,  $\xi_\ell \rightarrow \xi$  strongly in  $L^1(\Omega)$ , and  $\varepsilon_\ell \rightarrow 0$  such that

$$\limsup_{\ell \rightarrow \infty} F_\ell(u_\ell, \xi_\ell, \varepsilon_\ell) \leq F(u, \xi, 0). \tag{6.1}$$

If  $\xi \neq \det Du$  then  $F(u, \xi, 0) = +\infty$  and hence (6.1) is trivially satisfied.

On the other hand, in Lemma 4.3, we have proven that, whenever  $u_\ell \rightharpoonup u$  weakly in  $W^{1,n}(\Omega)^n$ ,  $\xi_\ell \rightharpoonup \xi$  weakly in  $L^1(\Omega)$ , and  $\varepsilon_\ell \rightarrow 0$ , then

$$F(u, \xi, 0) \leq \liminf_{\ell \rightarrow \infty} F_\ell(u_\ell, \xi_\ell, \varepsilon_\ell). \tag{6.2}$$

Strictly speaking we have shown this for the case  $\xi = \det Du$ , but we have also shown that all accumulation points of families with bounded energy satisfy this. Hence, (6.2) is indeed correct.

In the language of  $\Gamma$ -convergence (6.1) and (6.2) are, respectively, called the *limsup* and *liminf conditions* (here only for  $\varepsilon = 0$ ), and together they can be written as

$$\Gamma\text{-}\lim_{\ell \rightarrow \infty} F_\ell(v, \eta, 0) = F(v, \eta, 0) \quad \text{for all } v \in V, \eta \in L^1(\Omega), \quad (6.3)$$

where  $\Gamma$ -convergence is understood with respect to the weak  $W^{1,n}(\Omega)^n \times L^1(\Omega) \times [0, \infty)$ -topology. In fact, it is straightforward to verify that

$$\Gamma\text{-}\lim_{\ell \rightarrow \infty} F_\ell = F,$$

holds in the entire space  $W^{1,n}(\Omega)^n \times L^1(\Omega) \times [0, \infty)$ , however, this is less relevant for our purposes.

Thus, Theorem 4.1 can be interpreted as a  $\Gamma$ -convergence result in the sense of (6.3). In an obvious way, Theorems 5.1 and 5.2 can also be written in this way. We note however, that our original statements are slightly stronger in that we obtain separate convergence of the different contributions to the energy.

To conclude, we note that the statement

$$\Gamma\text{-}\lim_{\ell \rightarrow \infty} F_\ell(\cdot, \cdot, \varepsilon_\ell) = F(\cdot, \cdot, 0),$$

for a fixed sequence  $\varepsilon_\ell \rightarrow 0$ , is in general *false*. To see this, observe that to obtain (6.1), the choice of the sequence  $(\varepsilon_\ell)$  may strongly depend on the limit point  $u$  which we are aiming to approximate.

## 7. Algorithms and Numerical Examples

In the preceding sections we have formulated a general class of numerical methods for the solution of problems of the calculus of variations. The purpose of the present section is to demonstrate how they can be efficiently implemented and to demonstrate their practicality at several examples. We aim to give as much detail as possible so that our numerical results may be easily reproduced.

### 7.1. Optimization of non-differentiable energies

We begin by describing the implementation of the non-differentiable functionals which arise in our penalization procedure. Recall that we are aiming to minimize an energy which can be written in the form

$$\begin{aligned} E(v) &= \int_{\Omega} W(x, v, Dv) dx \\ &= \int_{\Omega} \phi(x, v, Dv, \gamma(Dv)) dx, \end{aligned}$$

over a convex and closed subset  $V \subset W^{1,1}(\Omega)^m$ , where  $\phi(x, v, F, \eta)$  and  $\gamma(F)$  are assumed to be smooth (at least twice differentiable) in  $v$ ,  $F$ , and  $\eta$ . For the sake of simplicity we do not consider  $\gamma = \gamma(x, v, Dv)$ , but this is not a true restriction.

We shall consider general penalty functionals of the type

$$E_\varepsilon(v, \eta) = \int_{\Omega} \phi(x, v, Dv, \eta) dx + \varepsilon^{-1} \int_{\Omega} |\gamma(Dv) - \eta|_1 dx, \quad (7.1)$$

defined for  $v \in V_\ell = u_{D,\ell} + P^1(\mathcal{T}_\ell)^m$ ,  $\eta \in Y_\ell = P^0(\mathcal{T}_\ell)^\mu$ , and where  $|\cdot|_1$  denotes the  $\ell^1$ -norm. We will see in numerical experiments that the  $L^1$ -type penalty functional guarantees a compact support of the difference  $\gamma(Dv) - \eta$ . This gives us information about the location of the singularities and also significantly reduces the complexity of the optimization (the optimization software TRON [22] automatically removes the unnecessary degrees of freedom).

By a simple variable transformation, we can replace  $\eta$  by  $\eta + \gamma(F)$  to obtain a new functional

$$\int_{\Omega} \phi(x, v, Dv, \gamma(Dv) + \eta) dx + \int_{\Omega} |\eta|_1 dx.$$

Next, we split the variable  $\eta$  into  $\eta = \eta^+ - \eta^-$  where  $\eta_j^+ = \max(\eta_j, 0)$  and  $\eta_j^- = -\min(\eta_j, 0)$ ,  $j = 1, \dots, \mu$ , and hide  $\gamma(F)$  within a newly defined energy density

$$\tilde{\phi}(x, u, F, \eta) = \phi(x, u, F, \gamma(F) + \eta),$$

to rewrite the functional as

$$\tilde{E}_\varepsilon(v, \eta^+, \eta^-) = \int_{\Omega} \tilde{\phi}(x, v, Dv, \eta^+ - \eta^-) dx + \varepsilon^{-1} \int_{\Omega} |\eta^+|_1 + |\eta^-|_1 dx. \quad (7.2)$$

Upon making  $\eta^+$  and  $\eta^-$  independent variables but imposing the bound constraints  $\eta^+ \geq 0$  and  $\eta^- \geq 0$  we have thus turned the original non-differentiable problem to minimize (7.1) into a smooth but constrained optimization problem. In particular, we define (7.2) for all  $v \in V_\ell$  and for all  $\eta^+, \eta^- \in Y_\ell^+$ , where

$$Y_\ell^+ = \{\eta \in Y_\ell : \eta_j \geq 0 \text{ in } \Omega, j = 1, \dots, \mu\}.$$

The functionals in (7.2) can be easily implemented with its gradient and hessian provided exactly. Our own implementation uses the trust region software TRON [22] to solve the *local* minimization problem

$$\min_{\substack{u \in V_\ell \\ \xi^\pm \in Y_\ell^+}} \tilde{E}_\varepsilon(u, \xi^+, \xi^-). \quad (7.3)$$

## 7.2. Adaptive mesh refinement for the penalty method

At several points in the continuation algorithm for the penalty method, described in the following section, we have to refine the mesh based on one of two principles: (i) either to reduce the overall energy or (ii) to reduce the contribution from the penalty term.

(i) To reduce the overall energy we use a DWR-type idea [8]. Let  $(u_\ell, \xi_\ell^+, \xi_\ell^-)$  be a local minimum of  $\tilde{E}_\varepsilon$ , computed using the method described above. We then define the error indicators

$$\eta_e = \sum_{T \in \mathcal{T}_\ell} \eta_T, \quad \text{where}$$

$$\eta_T = \left| \int_T \partial_F \tilde{\phi}(x, u_\ell, Du_\ell, \xi_\ell^+ - \xi_\ell^-) : (Du_\ell - G_\ell) dx \right|,$$

where  $G_\ell \in P^1(\mathcal{T}_\ell)^{m \times n}$  is a gradient recovery defined at each node  $z$  of the mesh  $\mathcal{T}_\ell$  by

$$G_\ell(z) = - \int_{\cup\{T \in \mathcal{T}_\ell: z \in T\}} Du_\ell dx.$$

The value  $\eta_e$  gives an indication how much the “elastic” energy may be lowered by local mesh refinement. On the other hand, the value of the penalty integral

$$\eta_p = \varepsilon^{-1} \int_{\Omega} |\xi_\varepsilon^+|_1 + |\xi_\varepsilon^-|_1 dx,$$

indicates how much the “penalty” energy can be lowered. If  $\eta_e > C_{e,p} \eta_p$  then the mesh is refined by marking a fraction of all elements which have the largest indicators  $\eta_T$  for refinement. Otherwise all those elements are marked where  $\xi^+ + \xi^-$  is non-zero (up to a threshold which takes round-off errors and premature termination of the optimization into account).

(ii) To reduce the penalty energy we use the very same procedure. All those elements are marked for refinement where  $\xi^+ + \xi^-$  is non-zero.

### 7.3. Continuation algorithm

A major difficulty one encounters when solving problems involving the Lavrentiev phenomenon is the so-called repulsion property. For example, if  $u_j \rightarrow x^{1/3}$  strongly in  $L^1(0, 1)$ , but  $u_j \in W^{1,\infty}(0, 1)$  for all  $j$ , then

$$\int_0^1 |u_{j,x}|^6 (u_j^3 - x)^2 dx \rightarrow +\infty.$$

We can imagine this effect as a huge energy barrier that needs to be overcome (or a complicated path to be found) when moving from a Lipschitz function to the global minimum. In our computations, we see this effect in that even for sufficiently small meshes it is often difficult to find the correct minimizers and that the penalty method converges to the Galerkin solution instead. (By “*Galerkin solution*”, we mean any  $P^1$ -minimizer of the original non-penalized functional.) In particular, we observed that a local minimum when  $\varepsilon$  is chosen too small in relation to the current mesh since in that case the penalty method becomes in effect a Galerkin method again.

Thus the problem may be overcome by, either increasing  $\varepsilon$ , or decreasing the mesh size. The former is clearly not desirable while the latter may be prohibitively expensive. Our solution therefore was to consider a continuation with respect to the parameter  $\varepsilon$ . By initially choosing  $\varepsilon$  very large the Galerkin solution is automatically discarded even for coarse meshes. We then gradually decrease  $\varepsilon$  and adapt the mesh whenever there is a danger that we may “fall out” of the basin of attraction of the exact minimizer because  $\varepsilon$  has become too small for the current mesh. This may be controlled by requiring that at all times the total energy  $\tilde{E}_\varepsilon$  must be below a critical value which should be less than the energy of the Galerkin solution.

- (1) Choose  $\varepsilon_{\text{dec}} \in (0, 1)$ ,  $E_{\text{goal}} \in \mathbb{R}$ ,  $\varepsilon_0$ , an initial mesh  $\mathcal{T}_0$ , and two bounds  $N_{\text{opt}}^1, N_{\text{opt}}^2$  (see remarks below how to choose them) for number of iterations of the optimization. Set  $\ell = 0$  and  $N_{\text{opt}} = N_{\text{opt}}^2$ .

- (2) Minimize  $\tilde{E}_{\varepsilon_\ell}$ , allowing at most  $N_{\text{opt}}$  iterations.
- (3) Determine next action:
  - (3.1) If the optimization converged and  $\tilde{E}_{\varepsilon_\ell} \leq E_{\text{goal}}$  accept the step, set  $\ell \leftarrow \ell + 1$ ,  $\varepsilon_\ell = \varepsilon_{\ell-1} \cdot \varepsilon_{\text{dec}}$ ,  $\mathcal{T}_\ell = \mathcal{T}_{\ell-1}$ ,  $N_{\text{opt}} = N_{\text{opt}}^1$  and continue at (2).
  - (3.2) If the optimization converged but  $\tilde{E}_{\varepsilon_\ell} > E_{\text{goal}}$  use refinement strategy (i) of the previous section to obtain a new mesh  $\mathcal{T}_\ell$ , set  $N_{\text{opt}} = N_{\text{opt}}^2$ , and redo step (2).
  - (3.3) If the optimization did not converge use refinement strategy (ii) of the previous section to obtain a new mesh  $\mathcal{T}_\ell$ , set  $\varepsilon_\ell = \varepsilon_{\ell-1}$ ,  $N_{\text{opt}} = N_{\text{opt}}^2$ , and redo step (2).

Some further comments to refine the continuation algorithm are required.

- The initial parameters for step (1) have to be chosen in such a way that the first step is always succesful.
- The algorithm terminates unsuccessfully when a maximum number of elements is reached, and succesfully when a prescribed goal  $\varepsilon_{\text{goal}}$  for  $\varepsilon_\ell$  is achieved.
- If the algorithm has terminated succesfully we usually “postprocess” the solution by performing a few additional mesh refinements (but fixing  $\varepsilon$ ) using strategy (i) to confirm that the penalty energy and support of  $\xi_\ell^+ + \xi_\ell^-$  tend to zero.
- After  $\varepsilon_\ell$  is decreased in step (3.1) we only expect a small change in the solution. Therefore the optimization should essentially behave like Newton’s method and terminate in few steps. We therefore set the maximum number of iterations to a relatively small number (say  $N_{\text{opt}}^1 = 20$ ). This setting prevents us from spending many iterations on finding an entirely new equilibrium when  $\varepsilon_\ell$  becomes too small for the current mesh and the penalty solution ceases to be a local minimizer.
- On the other hand, after the mesh is refined in either step (3.2) or (3.3) we expect a large change in the solution because the support of  $\xi^+ + \xi^-$  may shrink and we therefore allow a larger number of iterations (say  $N_{\text{opt}}^2 = 10^6$ , but we usually observe termination in far fewer iterations).

We have not addressed the question under which the algorithm is considered to have failed. When no Lavrentiev phenomenon occurs, we observe, in general, that for large  $\varepsilon$  a state satisfying the requirement  $\tilde{E}_{\varepsilon_\ell} \leq E_{\text{goal}}$  is found but that eventually, the algorithm will keep refining the mesh without being able to uphold this bound. We have therefore implemented a safety check which terminates the algorithm when a prescribed number of elements is reached.

As a warning, we also note that for sufficiently large  $\varepsilon$  it is sometimes possible to find *reasonably looking* solutions which indicate a Lavrentiev gap, but which may disappear as  $\varepsilon$  becomes small. It is therefore crucial to be able to drive  $\varepsilon$  as close to zero as possible.

### 7.4. Maniá-type examples

In this section we present numerical results for one-dimensional problems of the type

$$E(v) = \int_0^1 \left( |v_x|^n (v^m - x^k)^2 + \nu |v_x|^2 \right) dx \tag{7.4}$$

$$V = \{v \in W^{1,1}(0,1) : v(0) = 0, v(1) = 1\} = \text{id} + W_0^{1,1}(0,1),$$

where  $k, m, n \in \mathbb{N}$  and  $\nu \geq 0$ . This class includes in particular Maniá’s original example [23] ( $n = 6, m = 3, k = 1, \nu = 0$ ), and the *regular* example of Ball and Mizel [6, 7] ( $n = 14, m = 3, k = 2, 0 < \nu < 2.4 \times 10^{-3}$ ). The idea behind these examples is that, for  $\nu = 0$  the infimum of the energy is always zero with exact solution  $u^*(x) = x^{k/m}$ , but that the power  $n$  can be chosen large to make approximation *difficult*. Moreover, if  $m$  and  $k$  are chosen such that  $u^* \in H^1(0,1)$  then a perturbation of the functional with sufficiently small positive  $\nu$  does not change whether  $E$  exhibits a Lavrentiev phenomenon or not [6, 7].

The  $x^{1/3}$  singularity for the original Maniá example is expensive (though not impossible) to resolve and so we have chosen to compute the solution for  $n = 8, m = 2, k = 1, \nu = 0$  instead. We have plotted an accurate Galerkin solution, the solution of the penalty method for  $\varepsilon = 10^{-1}$  and  $\#T = X$  in Fig. 7.1, and the iterations of the contributions to the energy of the penalty method as well as the support of  $\xi_\ell^+ + \xi_\ell^-$  in Fig. 7.2.

In addition, we also computed the solution for the regular example of Ball and Mizel with  $n = 14, m = 3, k = 2, \nu = 10^{-3}$ , and we have plotted the solution in Fig. 7.3. The evolution of the energy and of the support of the penalty variable is similar as in the previous example.




Fig. 7.1. Final solutions of the Galerkin and the Penalty methods for the Maniá problem (7.4) with parameters  $n = 8, m = 2, k = 1, \nu = 0$  before the reduction step. The error of the penalty solution in the  $L^\infty$ -norm is  $\|u_\ell - u^*\|_{L^\infty} \approx 7.83 \times 10^{-5}$




Fig. 7.2. Evolution of the contributions to the penalty energy  $\tilde{E}_{\epsilon_\ell}$  and of the support of the penalty variables at each step of the continuation algorithm outlined in Section 7.3. The clear convergence of  $|\text{supp}(\xi^+ + \xi^-)|$  to zero is a strong indicator for the convergence of the method




Fig. 7.3. Final solution of the Galerkin and Penalty methods for Ball and Foss' [7] version of the Maniá problem (7.4) with parameters  $n = 17, m = 3, k = 2, \nu = 10^{-3}$  before the reduction step. The different orders of the singularity at the origin are a clear indication for a Lavrentiev gap

## 7.5. A convex example in 2D

In this section, we present numerical results for a modification of the example provided by Foss, Hrusa and Mizel [16]. In their original example a semi-circle  $\Omega$  is transformed into a quarter-circle  $y(\Omega)$  with stored energy

$$E(y) = \int_{\Omega} \left[ (|Dy|^2 - 2 \det Dy)^4 + \nu \left( \frac{\kappa}{\det Dy} + 3^{2-\kappa} (1 + |Dy|^2)^{\kappa/2} \right) \right] dx,$$

where  $\kappa$  and  $\nu$  are parameters, creating a singularity at the origin. The idea of the example is similar as in the regular examples of Ball and Mizel. For  $\nu = 0$  the map  $y^*(x) =$



$r^{1/2}(\cos(\theta/2), \sin(\theta/2))$  gives zero energy but the large power makes approximation difficult and it can be shown that the problem exhibits the Lavrentiev phenomenon. Further, the deformation  $y^*$  has finite energy for  $\nu > 0$  and hence, for  $\nu$  sufficiently small the Lavrentiev effect remains [16].

We note that the map  $F \mapsto (|F|^2 - 2 \det F)$  is a non-negative quadratic form and hence the stored energy density

$$W_0(F) = (|F|^2 - 2 \det F)^4,$$

is convex. The polyconvex terms are fairly unimportant for the Lavrentiev effect and hence we decided to ignore them completely (though we should mention that we also performed succesful computations with the full Foss/Hrusa/Mizel example). Instead, upon noting that  $y^* \in H^1(\Omega)$  we regularize  $W_0$  by a quadratic and define

$$E(v) = \int_{\Omega} [W_0(Dy) + \nu |Dy|^2] dx, \tag{7.5}$$

$$V = \{v \in W^{1,1}(\Omega) : v(x) = r(\cos(\theta/2), \sin(\theta/2)) \text{ if } |x| = 1, \\ v_1(\{x_2 = 0, x_1 < 0\}) = \{0\} \text{ and } v_2(\{x_2 = 0, x_1 > 0\}) = \{0\}\}.$$

The solution and the evolution of the energy during optimization for the case  $\nu = 0$  are plotted in Fig. 7.4, 7.5 and 7.6. For the case  $\nu = 10^{-3}$ , we have only plotted the radial component of the solution in Fig. 7.7. The evolutions of energy and support of the penalty variables during the optimization is similar as in Fig. 7.6.




Fig. 7.4. Radial components of the solution of the Galerkin and the Penalty methods for the modified Foss/Hrusa/Mizel problem (7.5) with  $\nu = 0$  before the reduction step. The different orders in the singularities at the origin are a clear indicator for a Lavrentiev gap. The error of the penalty solution before reduction is  $\|u_\ell - u^*\|_{L^\infty} \approx 1.07 \times 10^{-1}$




Fig. 7.6. Evolution of the contributions to the penalty energy  $\tilde{E}_{\epsilon_\ell}$  and of the support of the penalty variables at each step of the continuation algorithm outlined in Section 7.3. The apparent convergence of  $|\text{supp}(\xi^+ + \xi^-)|$  to zero is a strong indicator for the convergence of the method

### Deformation and Penalty Variable




Fig. 7.5. Plot of the deformation given by the solution of the Penalty method for the Foss/Hrusa/Mizel problem (7.5) with  $\nu = 0$ . The shade of the elements represents the size of the penalty variables  $\xi^+ + \xi^-$




Fig. 7.7. Radial components of the solution of the Galerkin and the Penalty methods for the modified Foss/Hrusa/Mizel problem (7.5) with  $\nu = 0$  before the reduction step. The different orders in the singularities at the origin are a clear indicator for a Lavrentiev gap. For comparison, the exact solution for the case  $\nu = 0$  is plotted as well

To conclude, we briefly outline the result of an experiment that does not exhibit a Lavrentiev gap. We modify (7.5) as follows:

$$E(v) = \int_{\Omega} [W_0(Dy) + \nu |Dy|^p] dx,$$

keeping the same admissible set  $V$ . We choose  $\nu = 1/60$  and  $p = 6$  a case for which numerical experiments in [25, 26] indicate the absence of a Lavrentiev gap.

An adaptive Galerkin solution suggests that the infimum of the energy in the space of Lipschitz functions is approximately  $\inf E(V \cap W^{1,\infty}(\Omega; \mathbb{R}^2)) \approx 0.0093 + O(3 \times 10^4)$ . Hence, we try to minimize the penalty functional with target energy  $E_{\text{goal}} = 0.0085$ . We observe that up to  $\varepsilon \approx 2$  the algorithm behaves similar as in the case  $p = 2$  above. However, at this point it stagnates and is unable to lower the penalty parameter further without increasing the energy above  $E_{\text{goal}}$ . This is strong indication that no Lavrentiev gap exists or, more precisely, that no gap larger than  $10^{-3}$  exists, which is consistent with [25, 26].

Next, we considered the case  $\nu = 1/40$  and  $p = 4$ . This is a borderline case that is particularly difficult to resolve. In this case the adaptive Galerkin solution suggests that  $\inf E(V \cap W^{1,\infty}(\Omega; \mathbb{R}^2)) \approx 0.0212 + O(3 \times 10^4)$ . We tried to minimize the penalty functional with  $E_{\text{goal}} = 0.02$ . Our algorithm once again managed to decrease the penalty parameter to approximately  $\varepsilon \approx 1.8$  but not further, thus indicating the absence of a Lavrentiev gap. However, this is in contradiction with the numerical experiments shown in [26]. Due to the relative simplicity of the method used in [26] it is conceivable that their results are correct, and thus shows that in particularly difficult borderline cases our method may still require some improvements.

**Acknowledgments.** This work was supported by the DFG Research Center MATH-EON at Berlin, Germany, the World Class University (WCU) program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology R31-2008-000-10049-0, the German Research Group Microplast (Forschergruppe 797 der Deutschen Forschungsgemeinschaft), the Hausdorff Institute of Mathematics at Bonn, Germany, and the EPSRC Critical Mass programme ‘New Frontiers in the Mathematics of Solids’ (OxMOS), Oxford, UK.

## References

1. Y. Bai and Z. Li, *Numerical solution of nonlinear elasticity problems with Lavrentiev phenomenon*, Math. Models Methods Appl. Sci., **17** (2007), no. 10, pp. 1619–1640.
2. Y. Bai and Z.-P. Li, *A truncation method for detecting singular minimizers involving the Lavrentiev phenomenon*, Math. Models Methods Appl. Sci., **16** (2006), no. 6, pp. 847–867.
3. J. M. Ball, *Convexity conditions and existence theorems in nonlinear elasticity*, Arch. Rational Mech. Anal., **63** (1976/77), no. 4, pp. 337–403.
4. J. M. Ball, *Discontinuous equilibrium solutions and cavitation in nonlinear elasticity*, Philos. Trans. Roy. Soc. London Ser. A, **306** (1982), no. 1496, pp. 557–611.
5. J. M. Ball and G. Knowles, *A numerical method for detecting singular minimizers*, Numer. Math., **51** (1987), no. 2, pp. 181–197.
6. J. M. Ball and V. J. Mizel, *Singular minimizers for regular one-dimensional problems in the calculus of variations*, Bull. Amer. Math. Soc. (N.S.), **11** (1984), no. 1, pp. 143–146.
7. J. M. Ball and V. J. Mizel, *One-dimensional variational problems whose minimizers do not satisfy the Euler-Lagrange equation*, Arch. Rational Mech. Anal., **90** (1985), no. 4, pp. 325–388.
8. W. Bangerth and R. Rannacher, *Adaptive finite element methods for differential equations*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 2003.
9. A. Braides,  *$\Gamma$ -convergence for beginners*, vol. 22 of *Oxford Lecture Series in Mathematics and its Applications*, Oxford University Press, Oxford, 2002.
10. S. C. Brenner and L. R. Scott, *The mathematical theory of finite element methods*, vol. 15 of *Texts in Applied Mathematics*, 3rd edn., Springer, New York, 2008.
11. P. G. Ciarlet, *Mathematical elasticity. Vol. I*, vol. 20 of *Studies in Mathematics and its Applications*, North-Holland Publishing Co., Amsterdam, 1988, three-dimensional elasticity.
12. B. Dacorogna, *Direct methods in the calculus of variations*, vol. 78 of *Applied Mathematical Sciences*, Springer-Verlag, Berlin, 1989.
13. G. Dal Maso, *An introduction to  $\Gamma$ -convergence*, Progress in Nonlinear Differential Equations and their Applications, 8, Birkhäuser Boston Inc., Boston, MA, 1993.
14. N. Dunford and J. T. Schwartz, *Linear operators. Part I*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1988, general theory, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1958 original, A Wiley-Interscience Publication.
15. L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1992.
16. M. Foss, W. J. Hrusa, and V. J. Mizel, *The Lavrentiev gap phenomenon in nonlinear elasticity*, Arch. Ration. Mech. Anal., **167** (2003), no. 4, pp. 337–365.
17. G. Knowles, *Finite element approximation to singular minimizers, and applications to cavitation in nonlinear elasticity*, in: *Differential equations and mathematical physics (Birmingham, Ala., 1986)*, vol. 1285 of *Lecture Notes in Math.*, Springer, Berlin, 1987, pp. 236–247.
18. A. Lavrentiev, *Sur quelques problèmes du calcul des variations*, Ann. Mat. Pura Appl., **41** (1926), pp. 107–124.
19. Z.-P. Li, *Element removal method for singular minimizers in variational problems involving Lavrentiev phenomenon*, Proc. Roy. Soc. London Ser. A, **439** (1992), no. 1905, pp. 131–137.
20. Z.-P. Li, *Element removal method for singular minimizers in problems of hyperelasticity*, Math. Models Methods Appl. Sci., **5** (1995), no. 3, pp. 387–399.
21. Z.-P. Li, *A numerical method for computing singular minimizers*, Numer. Math., **71** (1995), no. 3, pp. 317–330.
22. C.-J. Lin and J. J. Moré, *Newton's method for large bound-constrained optimization problems*, SIAM J. Optim., **9** (1999), no. 4, pp. 1100–1127 (electronic), dedicated to John E. Dennis, Jr., on his 60th birthday.
23. B. Maniá, *Sopra un esempio di Lavrentieff*, Boll. Un. Mat. Ital., **13** (1934), pp. 147–153.
24. P. V. Negrón-Marrero, *A numerical method for detecting singular minimizers of multidimensional problems in nonlinear elasticity*, Numer. Math., **58** (1990), no. 2, pp. 135–144.
25. C. Ortner, *Non-conforming finite element discretisation of convex variational problems*, to appear in IMA J. Numer. Anal.
26. C. Ortner and D. Praetorius, *On the convergence of adaptive non-conforming finite element methods*, Tech. rep., Mathematical Institute, University of Oxford, 2008, oxMOS Preprint 14.