

Original citation:

Atwan, Zeenah, Wright, Jordan L., Leppard, Keith and Woodman, Andrew. (2016)
Promyelocytic leukemia protein isoform II inhibits infection by human adenovirus type 5
through effects on HSP70 and the interferon response. Journal of General Virology .

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/79401>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Link to published version: <http://wrap.warwick.ac.uk/79401/>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

Promyelocytic leukemia protein isoform II inhibits infection by human adenovirus type 5 through effects on HSP70 and the interferon response

Zeenah Atwan, Jordan Wright¹, Andrew Woodman² and Keith N Leppard*

School of Life Sciences, University of Warwick, Coventry CV4 7AL, U.K.

* Corresponding author

Phone: +44 (0)24 7652 3579

Fax: +44 (0)24 7652 3701

Email: Keith.Leppard@warwick.ac.uk

¹ Present address: Oxford Biomedica PLC, Medawar Building, Oxford Science Park, Oxford, UK

² Present address: Dept of Biochemistry and Molecular Biology, Penn State University, State College, PA 16802, U.S.A.

Running title: PML-II is inhibitory to Ad5 infection

Subject category: animal small DNA viruses

Keywords: adenovirus; promyelocytic leukaemia protein; heat-shock protein 70; interferon response

Word counts:

Abstract: 234

Main text: 6089

Abstract

Promyelocytic leukemia (PML) proteins have been implicated in antiviral responses but PML and associated proteins are also suggested to support virus replication. One isoform, PML-II, is required for efficient transcription of interferon and interferon-responsive genes. . We therefore investigated the PML-II contribution to human adenovirus 5 (Ad5) infection, using shRNA-mediated knock-down. HeLaΔII cells showed a 2 - 3 fold elevation in Ad5 yield, reflecting an increase in late gene expression. This increase was found to be due in part to the reduced innate immune response consequent upon PML-II depletion. However the effect was minor because the viral E4 Orf3 protein targets and inactivates this PML-II function. The major benefit to Ad5 in HeLaΔII cells was exerted via an increase in HSP70; depletion of HSP70 completely reversed this replicative advantage. Increased Ad5 late gene expression was not due either to the previously described inhibition of inflammatory responses by HSP70 or to effects of HSP70 on major late promoter or L4 promoter activity but might be linked to an observed increase in E1B 55K, as this protein is known to be required for efficient late gene expression. The induction of HSP70 by PML-II removal was specific for the HSPA1B gene among the HSP70 gene family and thus was not the consequence of a general stress response. Taken together, these data show that PML-II, through its various actions, has an overall negative effect on the Ad5 life-cycle.

Introduction

The promyelocytic leukemia (PML) gene encodes a series of protein isoforms via alternative splicing (Jensen *et al.*, 2001). Most of these contribute to the formation of PML nuclear bodies (PML-NB) that also transiently or permanently include many other proteins (Van Damme *et al.*, 2010). PML proteins and/or PML-NB are implicated in a wide range of cellular functions, including innate and intrinsic immune responses (Bernardi & Pandolfi, 2007; Geoffroy & Chelbi-Alix, 2011). The PML gene itself is an interferon-stimulated gene (ISG) (Chelbi-Alix *et al.*, 1995; Stadler *et al.*, 1995), which suggests PML might be an effector of IFN responses. PML isoform II (PML-II) in particular is also required for the effective induction of IFN β and ISG expression, which it achieves by promoting the assembly of functional transcription complexes at target promoters (Chen *et al.*, 2015; Kim & Ahn, 2015). Thus this PML isoform can act upstream of IFN production to create a feed-forward loop that potentiates type I IFN responses.

PML-NBs are intimately associated with the replication cycles of nucleus-replicating DNA viruses. Incoming viral genomes are found located in close proximity to PML-NBs (Ishov & Maul, 1996) and, for herpes simplex virus type 1 (HSV-1), PML-NBs have been shown to disassemble and reform close to the site of virus entry into the nucleus (Everett & Murray, 2005), suggestive of an early response by the cell to infection. During infection, PML-NBs are then targeted by proteins encoded by a wide variety of viruses (Leppard & Dimmock, 2006; Leppard & Wright, 2012). These findings fit a model in which PML-NBs or their components are broadly antiviral and hence viruses have evolved functions to disrupt these activities in order to favour virus replication. Alternatively, and not mutually exclusively, the interaction of viruses with PML-NB may have been selected to favour the virus, with disruption of PML-NBs liberating proteins that act to increase virus production (Berscheminski *et al.*, 2014).

One well-characterized example of PML-NB disruption is the action of herpes simplex virus type 1 (HSV1) ICP0 protein (Maul & Everett, 1994). Supporting an involvement of PML in IFN responses, HSV1 ICP0 mutants have a significant growth defect and attenuated pathogenicity in mice and both of these properties are substantially recovered in animals that are deficient in IFN responses (Halford *et al.*, 2006; Leib *et al.*, 1999); the ability of IFN to inhibit growth of ICP0 mutants in cell culture is also greatly reduced when PML-null or knock-down cells are used (Chee *et al.*, 2003; Everett *et al.*, 2008b). However, PML and other PML-NB components are also directly inhibitory to HSV1 independent of IFN (Everett *et al.*, 2008a; Everett *et al.*, 2008b), with PML-I and PML-II being particularly implicated (Cuchet *et al.*, 2011). More recently, PML-II was also found to be the most

potent inhibitor, among the six nuclear PML isoforms, of transduction by a recombinant parvovirus AAV-2 vector (Mitchell *et al.*, 2014).

Human adenovirus type 5 (HAdV-C5, Ad5) infection also targets PML, rearranging it from PML-NB into track-like structures (Carvalho *et al.*, 1995; Doucas *et al.*, 1996); other PML-NB components are also redistributed, including some into virus replication centres (Berscheminski *et al.*, 2014; Doucas *et al.*, 1996). The Ad5 E4 Orf3 protein, which forms nuclear tracks by self-association (Ou *et al.*, 2012; Patsalo *et al.*, 2012), acts directly on PML-II, binding to its unique C-terminal domain to cause the redistribution of all PML isoforms (Hoppe *et al.*, 2006; Leppard *et al.*, 2009). Functionally, E4 Orf3 is necessary for Ad5 to replicate in the face of a pre-established IFN response (Ullman *et al.*, 2007) and E4 Orf3 also disrupts the intrinsic antiviral effects of PML and another PML-NB component, Daxx (Ullman & Hearing, 2008). Taken together, these observations support the idea that the E4 Orf3 interaction with PML-II opposes antiviral responses so as to favour productive viral infection, which fits well with the more recent finding that PML-II is needed for a robust type 1 IFN response (Chen *et al.*, 2015). However, it has also been reported that PML-II serves a positive function during Ad5 infection (Berscheminski *et al.*, 2013).

In light of these findings, we investigated the circumstances under which PML-II could provide a positive or negative influence on Ad5 infection, and the mechanisms underlying these influences. Viral gene expression and replication were increased by the removal of PML-II within a background of ongoing expression of other PML isoforms, leading to an increase in virus yield. One factor in this increase was the reduction in the interferon type I response in PML-II depleted cells. The other more significant factor was the increased level of HSP 70 protein in PML-II depleted cells, which was found to support elevated Ad5 gene expression.

Results

Stable knockdown of PML-II in Hela cells. To investigate the effect of PML-II on the well-characterised Ad5 productive infection of Hela cells, we first generated PML-II knockdown Hela cells (HelaΔII) by lentiviral shRNA transduction, along with matched empty vector cells (HelaEV). HelaΔII cells were fully viable in long-term culture, showed similar morphology to both parental Hela cells and HelaEV cells (Fig. 1a) and grew at only a slightly slower rate than HelaEV cells under puromycin selection. HelaΔII cells showed significant reductions in PML-II mRNA (Fig. 1b) and protein (Fig. 1c). These cells also displayed functional knockdown of PML-II based on their reduced ability to express IL-6 and ISG56 (Fig. 1d), which was shown previously to depend on the presence of PML-II (Chen *et al.*, 2015).

Depletion of PML-II increases the productivity of Ad5 infection. To establish the effect of PML-II on Ad5 infection, HeLa Δ II and HeLaEV cells were infected in parallel with wild-type (wt) Ad5. Looking at protein expression over a time course, there was a strikingly higher level of late protein expression in HeLa Δ II cells (Fig. 2a); with an exposure selected to avoid grossly overexposing the HeLa Δ II lane, late proteins in the HeLaEV cells were barely detectable. In contrast, expression of the early protein E2A 72K DNA binding protein (DBP) was much less affected by the removal of PML-II though the E1B 55K protein was, by the late stage of infection (24 h p.i.), significantly increased (Fig. 2a). The expression of late proteins in HeLaEV cells was similar to that in untransduced standard HeLa cells (Fig 2b), confirming that the difference between HeLa Δ II and HeLaEV infections was not due to any unexpected negative effect of introducing the retroviral vector alone in HeLaEV cells. The effect of PML-II depletion on viral gene expression was confirmed and quantified by flow cytometry (Fig. 2c); both the proportion of cells positive for late proteins and their mean fluorescence intensity were increased in HeLa Δ II cells. The increased late protein expression in HeLa Δ II cells was reflected in a 2-3 - fold higher virus yield/cell as compared with HeLaEV cells, measured at 24 h and 48 h post-infection (Fig. 2d).

Lack of IFN response partially explains the beneficial effect of PML-II depletion. Infection by Ad5 is intrinsically an IFN response-inducing event, with both virus entry itself and later gene expression events triggering interferon and inflammatory signalling (Hartman *et al.*, 2007; Hendrickx *et al.*, 2014; Zhu *et al.*, 2007). Since PML-II plays an important role in the activation of an IFN response (Chen *et al.*, 2015), we considered the possibility that, even though Ad5 encodes functions that inhibit the IFN response in various ways, the beneficial effect on Ad5 infection of PML-II removal might nonetheless arise because of the consequent further defect in the IFN response. To test this, we directly disabled the IFN response by knockdown of IRF3, which is a key transcription factor in the induction of type 1 IFN responses (Au *et al.*, 1995). Physical and functional depletion of IRF3 from HeLaEV cells (Fig 3a, b) increased Ad5 wt300 late gene expression by a modest amount (Fig. 3c), indicating that the Ad5 functions deployed to inhibit IFN responses are not 100% effective. However hexon expression under IRF3 knockdown in HeLaEV cells was still very substantially lower than seen in HeLa Δ II cells, in which IRF3 knockdown had little additional effect. Thus, while some part of the benefit to Ad5 of PML-II removal reflects the loss of the IFN response, there is a significant additional component to be accounted for; this is considered further below.

Ad5 E4 Orf3 inhibits PML-II function in the IFN response

Ad5 E4 Orf3 binds PML-II directly (Hoppe *et al.*, 2006) and is necessary for Ad5 replication in IFN-treated cells, dependent on the presence of PML (Ullman & Hearing, 2008; Ullman *et al.*, 2007), so

we asked whether this interaction also inhibited the natural IFN response to Ad5 infection. To determine whether viral expression of E4 Orf3 had any measurable effect on induction of type 1 IFN during infection, culture media from Ad5-infected HEK293 cells (Fig. 4a) or MRC5 normal human fibroblasts (Fig. 4b) were tested in plaque reduction assays using Semliki Forest virus, an IFN-sensitive alphavirus. In both cell types, IFN activity was detected from an Orf3-deficient virus infection (*inOrf3*) while none was detected from *wt300* or mock infections. Based on a calibration of the assay with recombinant IFN α , which showed inhibition from 0% to 100% by IFN in the range 0.1 – 100 U/ml, *inOrf3* medium contained ~50 U/ml IFN. In separate experiments, IFN levels in infected HEK293 cell culture media were determined using an IFN-responsive reporter assay (Chen *et al.*, 2015). Again, IFN accumulation was detected only in *inOrf3*-infected cultures (Fig. 4c): amounts were equivalent to about 60 U/ml, in line with the estimate from the plaque-reduction assay. Thus, the presence of E4 Orf3 causes a measurable reduction in IFN production and secretion stimulated by Ad5 infection.

To test whether E4 Orf3 protein alone was sufficient to inhibit IFN responses, we employed transient expression IFN β promoter reporter assays. PML-II Δ RBCC is an artificially truncated form of PML-II that does not associate with PML-NB but retains E4 Orf3 binding (Leppard *et al.*, 2009) and has increased ability to potentiate IFN β promoter activation by inducers such as poly(I:C) (Chen *et al.*, 2015). E4 Orf3 fully reversed the increased response of the IFN β promoter to poly(I:C) due to PML-II Δ RBCC and further reduced reporter activity to levels below that of poly(I:C) stimulation in the absence of exogenous PML (Fig. 4d). This reduction below baseline reflected the contribution of endogenous PML-II, also an E4 Orf3 target, to the observed IFN β promoter activation as, in the presence only of endogenous PML, added Orf3 also gave a dose-dependent inhibition of poly(I:C)-stimulated reporter activity (Fig. 4e).

To correlate the activity of E4 Orf3 in regulating IFN β expression with its ability to bind PML-II, we compared the inhibitory effect of wild-type E4 Orf3 with that of selected Orf3 mutants (Hoppe *et al.*, 2006). Those mutants unable to bind PML-II (N82A, L103A) also failed to inhibit activation of the IFN β promoter while mutants that retained PML-II binding (R100A, D105-L106A) had inhibitory activity similar to wild-type (Fig. 4e-i). Importantly, mutant D105-L106A uniquely retains PML-NB rearrangement activity whilst lacking the ability to disrupt the location of the MRN protein complex involved in DNA damage repair (Evans & Hearing, 2005). The retention of activity by this mutant thus clearly links the inhibitory effect of E4 Orf3 on IFN induction to its interaction with PML-II.

The inhibitory effect of E4 Orf3 on PML-II function suggested that an Orf3-deficient virus should benefit more from the lack of PML-II in Hela Δ II cells than a virus that was able to make Orf3. When

wt300 and *inOrf3* late protein expression was compared in HeLaEV cells (Fig. 5, lanes 2, 4), amounts were very similar, as expected (Huang & Hearing, 1989). As also shown previously, in Vero cells and human fibroblasts (Ullman *et al.*, 2007), IFN α pre-treatment more severely inhibited *inOrf3* than *wt300* late protein synthesis in HeLaEV cells (Fig. 5, lanes 3, 5). Importantly, removal of PML-II in HeLa Δ II cells largely abolished this difference in viral gene expression (Fig. 5, lanes 8, 10), confirming that PML-II is a significant functional target of E4 Orf3 during infection. However, contrary to expectation, *wt300* gene expression benefitted more than that of mutant virus *inOrf3* from PML-II removal (Fig. 5, lanes 2, 7 and 4, 9), see below. Collectively, our results show that PML-II is inhibitory to Ad5 infection in part through its role in the development of an IFN response and that E4 Orf3 inhibits this function of PML-II.

Enhanced growth of Ad5 in HeLa Δ II cells reflects overexpression of HSP70. A significant part of the benefit to Ad5 of PML-II depletion was independent of IRF3 and hence was not directly related to the IFN response (Fig. 3). Hence, late protein expression by either *wt300* or *inOrf3* was greater in HeLa Δ II cells than in equivalently treated HeLaEV cells (Fig. 5). When studying stress responses in HeLa Δ II cells, we fortuitously observed that they displayed elevated levels of HSP70 mRNA and protein under normal growth conditions in comparison with HeLaEV cells (Fig. 6a). HSP70 was also induced to a lower level by transient knock-down of PML-II in standard HeLa cells (Fig. 6b), suggesting a direct link between loss of PML-II and HSP70 expression. Ad5 infection also induces HSP70 expression (Nevins, 1982) and, since it inhibits many other host genes whose activity is detrimental to infection (Zhao *et al.*, 2003) and pre-existing HSP70 levels correlate with permissivity to Ad infection (Imperiale *et al.*, 1984), we inferred that HSP70 might be the relevant positive factor for Ad5 growth in HeLa Δ II cells. To test this, hexon expression was compared in cells infected with or without HSP70 knock-down (Fig. 6c, d). Whilst HeLa Δ II cells showed substantially more hexon mRNA and protein than HeLaEV cells when treated with a control siRNA, this difference was abolished by HSP70 siRNA treatment. Moreover, HSP70 siRNA also reduced hexon expression further from its lower base level in HeLaEV cells. Thus, HSP70 contributes positively to Ad5 gene expression and the elevated expression of HSP70 in HeLa Δ II cells is a major factor in the increased efficiency of infection in these cells. The fact that *wt300* gene expression benefitted more than that of mutant virus *inOrf3* from the high HSP70 environment in HeLa Δ II cells suggests that E4 Orf3 might be involved in the beneficial action of HSP70.

The assay of HSP70 mRNA shown in Fig. 6a detects transcripts only from the two major heat-inducible loci, HSPA1A and HSPA1B. However HSP70 encompasses a number of related proteins encoded by the HSPA gene family, only some of which are heat-inducible (Brocchieri *et al.*, 2008). To

determine the specificity of HSP70 induction in HeLaΔII cells, mRNA levels from several HSPA genes were assessed alongside HSP60 (HSPD gene family). As before, the HSP70 assay detected elevated mRNA levels in HeLaΔII cells (Fig. 6e). Interestingly, despite the high level of similarity between the HSPA1A and HSPA1B genes (they encode identical 641 amino acid proteins), elevated HSP70 expression was accounted for almost entirely by HSPA1B mRNA; there was little difference in expression of HSPA1A between the two cell types. In contrast to HSPA1, expression of HSPA5 mRNA, which encodes the endoplasmic reticulum chaperone GRP78 also known as BiP, was if anything slightly reduced by removing PML-II. Another HSP70 family member, HSPA6, which shows no basal expression but is induced by heat stress (Brocchieri *et al.*, 2008), was detected only at low levels and was not induced by removal of PML-II; expression of HSP60 was also unaltered. Thus, the loss of PML-II leads to highly specific induction of the HSPA1B gene, providing HSP70 protein that supports enhanced Ad5 gene expression.

Possible roles of HSP70 during Ad5 infection

HSP70 has been shown previously to inhibit pro-inflammatory NF-κB signalling and hence both the production and the effect of TNF-α (Meng & Harken, 2002). Confirming that HSP70 had this effect in our system, we found that HSP70 knock-down significantly increased the expression of ISG56 in wt300-infected HeLaEV cells (Fig. 7a). This suggested that HSP70 might favour Ad5 replication by limiting the induction of innate and inflammatory responses through NF-κB. TNFα is a known activator of NF-κB signalling and is considered to be inhibitory to virus infection (McFadden *et al.*, 2009). Ad5 infection can stimulate NF-κB signalling in several ways (Higginbotham *et al.*, 2002; Pahl *et al.*, 1996; Schmitz *et al.*, 1996) while several viral E3 gene products counteract TNFα activity (Gooding *et al.*, 1988), suggesting NF-κB activation might be inhibitory to Ad5 infection. We therefore tested whether elevated HSP70 in HeLaΔII cells enhanced Ad5 infection by inhibiting NF-κB. Reasoning that exogenous TNFα would oppose such an effect and so reduce the benefit of PML-II removal, we analysed Ad5 late gene expression in HeLaΔII cells with or without TNFα treatment (Fig. 7b). However, TNFα actually modestly enhanced Ad5 late protein expression in both HeLaΔII cells and HeLaEV cells. The same effect was seen on hexon mRNA in HeLaEV cells and this was potentiated by HSP70 knock-down (Fig. 7c), as expected if HSP70 limits pro-inflammatory signalling that is beneficial to the virus. We also tested the effect of QNZ, an inhibitor of NF-κB activation (Tobe *et al.*, 2003) and found that, consistent with the effect of TNFα treatment, QNZ reduced Ad5 late gene expression in both cell types (Fig. 7d). These data indicate that NF-κB signalling increases rather than inhibits Ad5 gene expression in our system and that HSP70 limits rather than increases this effect.

The beneficial effect on Ad5 infection of the high levels of HSP70 in HelaΔII cells must therefore be due to some other function of HSP70.

HSP70's principal role is as a chaperone: during heat-stress it stabilises partially denatured proteins to prevent aggregation and facilitate re-folding (Clerico *et al.*, 2015). The Ad5 replication cycle involves both the disassembly and assembly of protein complexes, processes which might be facilitated by HSP70. Indeed, HSP70 interacts both with the hexon shell of Ad2 particles shortly after infection (Niewiarowska *et al.*, 1992) and with fibre protein during the late phase of Ad5 infection (Macejak & Luftig, 1991), and has been implicated in uncoating and import of the genome into the nucleus (Saphire *et al.*, 2000). We therefore examined whether increased HSP70 present in HelaΔII cells altered the subcellular location of hexon protein, as an indicator of possible effects on particle assembly. Prior depletion of HSP70 from these cells, as well as decreasing the overall level of hexon protein as already described, increased more than two-fold the cytoplasmic / nuclear ratio of hexon (Fig. 8a; quantitation under right panels) whereas it had little effect on the distribution of E1A or E2A DBP. This result suggests that HSP70 overexpression consequent on PML-II depletion may positively affect the assembly of progeny particles in the nuclear compartment and hence could contribute to the increased yield of virus.

Any impact of HSP70 level on assembly cannot explain the effect of HSP70 on hexon mRNA levels (Fig. 6). This mRNA is produced by processing of transcripts from the major late promoter (MLP), which itself is positively influenced by L4-22K protein expressed from L4P which is activated at the onset of the late phase (Morris *et al.*, 2010). We therefore tested the effect of HSP70 depletion on the activity of MLP and L4P luciferase reporters in Hela cells (Fig. 8b), but found that neither was significantly affected. Thus HSP70 does not increase directly the intrinsic activity of either promoter when taken out of the context of viral infection and must therefore affect late gene expression post-transcriptionally or dependent on the infected cell environment. In this regard, we noted the increase in E1B 55K protein upon PML-II depletion in HelaΔII cells (Fig. 8c), a protein which is known to positively regulate Ad5 late mRNA nucleo-cytoplasmic transport and accumulation (Leppard, 1998). HSP70 depletion in HelaΔII cells reversed this increase in E1B 55K (Fig. 8a), further suggesting that it could be significant in the elevation of late gene expression.

Discussion

Depletion of PML-II from Hela cells, a standard permissive cell line for Ad5, led to a substantial 2-3 fold enhancement in virus yield. This was attributed largely to a general increase in late gene expression, our experiments focusing mainly on the major capsid protein hexon and its mRNA as an

example. Thus, PML-II has an overall inhibitory effect on Ad5 infection. Two factors were identified that contributed to the increased infectious productivity upon PML-II depletion: a reduction in IFN response and an increase in HSP70 expression, the latter being the predominant factor. Depletion of HSP70 from HelaΔII cells completely eliminated the advantage to hexon expression of PML-II removal whilst blocking the IFN response in HelaEV cells by a means other than PML-II depletion only somewhat increased the levels of viral late gene expression.

Previously, Berscheminski and colleagues reported that PML-II was beneficial to Ad5 infection, in apparent contradiction to our findings (Berscheminski *et al.*, 2013). They showed that PML-II potentiated transcriptional activation of a viral early promoter by the E1A 13S protein and that, in a cellular context where all PML proteins were depleted, the addition of exogenous PML-II enhanced virus yield about 3-fold. Comparing these findings with our own, it is important to note the differences in cell environment employed. PML-II function will be a composite of its free and PML-NB associated activities. PML-V is the stable base of PML-NBs (Weidtkamp-Peters *et al.*, 2008), with other isoforms and many other proteins associating with these bodies by protein - protein interactions including SUMO-SIM interactions (Bernardi & Pandolfi, 2007). In a PML-null cell, functions observed for added PML-II will be essentially those of its soluble nuclear form. Indeed, a mutated form of PML-II with reduced ability to associate with PML-NBs was more active than the wild-type in cooperating with E1A in the presence of endogenous PML (Berscheminski *et al.*, 2013), suggesting that interactions with other PML isoforms limit this activity. The overexpressed PML-II will also potentially exceed the capacity of E4 Orf3 expressed upon infection to bind and inactivate it. In contrast, specific depletion of PML-II in an otherwise normal PML background demonstrates the combined net contribution of this protein to the Ad5 life-cycle in all its cellular contexts. Our finding that PML-II removal exerts an overall positive effect on Ad5 growth is therefore not in disagreement with this prior study but instead reveals a new aspect of the functional interaction of PML-II with the virus. Our findings are also consistent with an earlier study showing the importance of PML proteins generally in the inhibition of Ad5 by an established IFN response (Ullman & Hearing, 2008).

Although PML-II is necessary for an efficient IFN response (Chen *et al.*, 2015), Ad5 gained only modest benefit from the loss of this response in HelaΔII cells. This finding is expected since the virus possesses several functions that collectively oppose IFN responses, so allowing infection to succeed even when the cell is capable of launching a response. First, E1A proteins inhibit both the expression of ISGs and the activation of IFN β transcription (Ackrill *et al.*, 1991; Reich *et al.*, 1988). Second, E1B 55K protein blocks the induction of a number of IFN-inducible genes and is required for efficient

replication in normal fibroblasts (Chahal *et al.*, 2012). Third, E4 Orf3 protein is necessary for replication to proceed in permanent cell lines in the face of an established IFN response (Ullman *et al.*, 2007), implying it negatively regulates that response. Fourth, VA RNA I inhibits the induction of an IFN-induced antiviral state by inhibiting protein kinase R (Kitajewski *et al.*, 1986). Finally, activated STAT1 is sequestered in Ad replication centres (Sohn & Hearing, 2011). The further advantage to Ad5 of an inherent lack of IFN response may arise because of the time it otherwise takes for virus-encoded anti-IFN functions to become active. We directly tested the idea that E4 Orf3 would inhibit the IFN response via its targeting of PML-II, and showed that this was the case: wild-type E4 Orf3 inhibited type 1 IFN induction whilst mutant forms unable to bind PML-II could not; E4 Orf3 mutant virus infection elicited more IFN than wild-type; and E4 Orf3 mutant virus late protein expression was more strongly inhibited by prior IFN treatment in cells with functional PML-II. E4 Orf3 is expressed in the early phase but takes time to accumulate. In the period prior to this, our data suggest that inhibition of IFN induction is incomplete.

The principal factor in the enhanced growth of Ad5 in HelaΔII cells was the elevated level of HSP70. Investigating this, we observed a modest effect of HSP70 on Ad5 late protein nuclear accumulation that would favour progeny virus formation but this could not account for the significant enhancement of late gene expression. HSP70 expression/depletion in HelaΔII cells did however also affect the accumulation of the viral E1B 55K protein, which is known to regulate late mRNA accumulation and hence to increase late gene expression (Leppard, 1998). This increased amount of E1B 55K in the presence of elevated HSP70 may contribute to the observed elevation in late gene expression and virus yield under these conditions. HSP70 also opposes inflammatory responses (Meng & Harken, 2002); given the role of PML-II in regulating inflammatory gene expression we considered this to be a plausible basis for the positive effect of HSP70 on Ad5 growth. However, NF-κB activation was actually modestly beneficial to late gene expression. Since, as reported, HSP70 opposed this activation, HSP70 elevation cannot be benefitting Ad5 via effects on NF-κB. The positive effect of NF-κB on Ad5 was unexpected given that the virus encodes functions in its E3 region that inhibit TNFα signalling and hence NF-κB activation (Burgert *et al.*, 2002). However, although these functions will be important *in vivo* they are known to be dispensable for growth in culture. Thus the small increase in Ad5 gene expression when NF-κB is activated in cell culture should not imply that this response benefits the virus *in vivo*.

The elevated level of HSP70 in PML-II depleted cells reflected a highly specific increase in mRNA derived from the HSPA1B gene, one of two intronless genes that are strongly heat-inducible members of the HSPA gene family. HSPA1A and HSPA1B are very similar even in their promoter

sequences (Broccieri *et al.*, 2008); their products are not normally distinguished in analyses of heat-induced HSP70 expression. The specific upregulation of HSPA1B by PML-II depletion cannot be due to a general cell stress response, and in particular cannot be attributed to activation of the heat shock transcription factor, HSF, which regulates transcription of HSPA1A and HSPA1B as well as other classes of HSP (Singh *et al.*, 2010). Thus, these results indicate a novel mechanism whereby the HSPA1B promoter is selectively activated. Interestingly, HSPA1A and HSPA1B are located within the MHC III region, between the gene clusters encoding MHC class I and II antigen where specific depletion of individual PML isoforms has been shown to have effects on chromatin architecture and gene expression (Kumar *et al.*, 2007). Further work is needed to test whether HSPA1B induction by PML-II removal reflects a similar mechanism.

HSP70 is also induced during Ad5 infection (Nevins, 1982) and, whilst virus infection might be considered a stress that would lead to generalized activation of HSP expression, this induction is actually specific to HSP70 (Phillips *et al.*, 1991). These studies did not distinguish between HSPA1A and HSPA1B, which were not separately recognised at the time. HSP70 transcription is induced by the virus-coded transactivator, E1A 13S (Wu *et al.*, 1986), which acts via the cellular CCAAT-box factor (CBF) and its binding site in the context of a specific TATA box (Lum *et al.*, 1992; Simon *et al.*, 1988). In this way, induction is independent of HSF. The CBF site is also a target for p53-mediated inhibition of the HSP70 genes (Agoff *et al.*, 1993), suggesting that E1A might disrupt this inhibition. HSP70 expression is also favoured by the viral E1B 55K / E4 Orf6 complex promoting HSP70 mRNA export to the cytoplasm (Moore *et al.*, 1987). Since, as discussed, our study shows that HSP70 levels are also positively linked to E1B 55K accumulation, a feed-forward loop may be established that promotes efficient late gene expression.

The induction of HSP70 during Ad infection may be linked with positive roles for this protein in the virus life cycle. The best documented of these, in viral uncoating (Saphire *et al.*, 2000), may account for the modest increase in early gene expression seen in HelaΔII cells. However, this action must precede E1A-induced activation of HSP70 synthesis, suggesting that other roles may exist to justify this mechanism. This role also cannot account for the predominant effect on late rather than early gene expression that we observed. A study by White *et al.* suggested an involvement of HSP70 in nuclear events linked with PML during Ad2 infection (White *et al.*, 1988). HSP70 was recruited from the cytoplasm into discrete nuclear structures that co-localized with E1A and which appeared similar to the reorganized PML tracks that are formed by E4 Orf3 (Carvalho *et al.*, 1995; Doucas *et al.*, 1996). Indeed, Carvalho *et al.* found a small fraction of E1A and, in a few infected cells, HSP70 located in these Orf3 / PML structures, evidence of a physical and/or functional link between HSP70 and PML

that might be related to our observations. Interestingly, our work suggests that the presence of E4 Orf3 is required in order for Ad5 to benefit from the elevation of HSP70 that occurs in Hela Δ II cells.

Many other viruses induce and/or functionally interact with HSP70 (Santoro *et al.*, 2010) suggesting a general importance of this protein to infection. The avian adenovirus CELO Gam1 protein causes an increase in both HSP70 and HSP40 that is needed for replication, and loss of Gam1 can be complemented by heat-shock (Glutzer *et al.*, 2000). Gam1 is also responsible for the loss of PML from infected cells through an inhibition of sumoylation (Colombo *et al.*, 2002), raising the possibility that HSP induction and PML loss are also linked in this system. Human cytomegalovirus, HSV1, vaccinia virus and some paramyxoviruses all induce HSP70 expression (Santoro *et al.*, 2010). For HSV1, heat shock can complement deficiency in ICP0, the protein responsible for PML body disruption and PML degradation, (Bringhurst & Schaffer, 2006) while the same is true for E1A deficiency in Ad (Imperiale *et al.*, 1984; Madara *et al.*, 2005).

In conclusion, we have shown that PML-II opposes productive Ad5 infection, in part by supporting innate immune responses but mainly due to a suppressive effect on HSP70 expression. Our study reveals a previously undefined activity for HSP70 in supporting Ad5 late gene expression and demonstrates an inhibitory effect of PML-II on HSP70 expression.

Materials and Methods

Generation of Hela Δ II and HelaEV cell lines

Hela cells were transduced with either lentiviral particles encoding an shRNA specific for PML-II or equivalent particles with no shRNA insert. The PML-II shRNA incorporated the active siRNA sequence described by (Kumar *et al.*, 2007) which was used previously by our laboratory to achieve functional knock-down of PML-II (Chen *et al.*, 2015). Lentiviral particles were generated using pLKO.1 (Moffat *et al.*, 2006) following protocols supplied by the RNAi consortium (Addgene). Briefly, a double-stranded synthetic oligonucleotide corresponding to the shRNA was cloned into pLKO.1. Specific plasmid clones were verified by sequencing, then transfected with psPAX2 and pMD2.G packaging plasmids into HEK-293T cells using Transit LT-1 (Mirus) to produce VSV-G-pseudotyped particles. Particle stocks were then used to infect Hela cells and transduced cells were selected with 3 μ g/ml puromycin.

Antibodies and reagents

Specific primary antibodies were: AdJLB1 rabbit antiserum to Ad5 late proteins (Farley *et al.*, 2004); mouse monoclonal antibodies 2HX-2 to Ad5 hexon (Cepko *et al.*, 1983), B6-8 to Ad5 E2A DNA binding protein (DBP) (Reich *et al.*, 1983), and 2A6 to Ad5 E1B 55K (Sarnow *et al.*, 1982);

monospecific anti-peptide sera reactive against PML-II (Xu *et al.*, 2005), kindly provided by Prof K.-S. Chang, M.D. Anderson Cancer Center, University of Texas; FL-425 rabbit anti-IRF3 (SantaCruz); rabbit anti-HSP70 (StressMarq SPC-103C/D); and GA1R mouse anti-GAPDH (Thermo Scientific). Secondary antibodies were: Alexa488 - conjugated goat anti-mouse Ig (Life Technologies); horseradish peroxidase (HRP) - conjugated goat anti-mouse Ig (Sigma); and HRP-conjugated goat anti-rabbit Ig (SantaCruz). IFN α was from PBL Assay Science, TNF α from Invitrogen, poly(I:C) from Sigma and 6-amino-4-(phenoxyphenylethylamino)quinazolin (QNZ) from Santa Cruz. siRNAs were: IRF3 (ID 3661, Qiagen); HSP70 (targets HSPA1A and HSPA1B, Ambion); and control B (Chen *et al.*, 2015).

Cell culture and virus infection

HEK293, HEK293T, Hela and knock-down cell lines were maintained at 37 °C, 5% CO₂ in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal bovine serum (FBS); for maintenance purposes, HelaEV and Hela Δ II cells were alternated between media containing or not containing 3 μ g/ml puromycin. VERO cells were maintained in DMEM supplemented with 5% FBS and MRC5 cells in 10% Eagle's minimal essential medium supplemented with 10% FBS, 2mM L-glutamine and 1% non-essential amino acids. Cells were seeded at the appropriate density 24 h prior to the respective procedure. Light microscope images were recorded on an inverted microscope using a 5x objective. Virus stocks and experimental samples were titred in a fluorescent focus assay. Hela cell monolayers were infected in duplicate with serial dilutions of each stock, incubated at 37 °C, 5% CO₂ for 16 h, then fixed and stained with antibody to E2A DBP to visualize fluorescent cells for counting. Experimental infections were carried out with wild-type Ad5 *wt*300 or E4 Orf3 mutant *in*Orf3 (Huang & Hearing, 1989) at a multiplicity of 5 fluorescence focus units (ffu) per cell unless otherwise indicated. siRNA transfections were performed with Lipofectamine 2000 (Invitrogen), using a ratio of 1 μ l reagent per 25 pmol siRNA.

Protein and RNA analysis

For total protein analysis, cells were lysed directly in SDS gel sample buffer. Cytoplasmic and nuclear fractions were generated by lysing cells in 0.67% (v/v) NP40, 10 mM NaCl, 1.5 mM MgCl₂, 10 mM Tris.HCl pH7.5 for 10 min on ice, then nuclei were pelleted by low speed centrifugation; an equal volume of 2x SDS gel sample buffer was added to the supernatant (cytoplasmic fraction). Crude nuclei were washed once in PBS, pelleted as before and then lysed in SDS gel sample buffer (nuclear fraction). Proteins were separated by electrophoresis on 10% SDS polyacrylamide gels and detected by western blotting as previously described (Lethbridge *et al.*, 2003). For flow cytometry analysis, single cell suspensions produced by trypsinization were fixed on ice with 10% (v/v) formalin in PBS for 20 min, permeabilized with 0.5% (v/v) NP40 in PBS for 10 min and then incubated with 1% (w/v)

bovine serum albumin in PBS for 45 min to block nonspecific protein binding. Cells were resuspended in FACS buffer (PBS containing 3% v/v FBS, 0.07% w/v NaN₃), then incubated with specific primary antibodies to hexon or E2A DBP followed by Alexa488-conjugated secondary antibody. Washed cells in FACS buffer were analysed using a FACSCAN (Becton Dickinson) and WinMDI software. Immunofluorescence analysis was performed as previously described (Leppard & Everett, 1999); images were collected with a Leica SP5 confocal microscope system and processed using Leica confocal software. Total RNA was isolated and mRNA quantified by RT-qPCR as previously described (Chen *et al.*, 2015) using the following primers and amplicons: ISG56 and IL-6 (Chen *et al.*, 2015); E1A (113 bp, Ad genome 1422-1534) and hexon (137 bp, 21540-21576) (Schreiner *et al.*, 2013); PML-II 5'AGGCAGAGGAACGCGTTGT and 5'GGCTCCATGCACGAGTTTTTC (70 bp); HSP70 (Tanaka *et al.*, 2007); HSPA1A, HSPA1B, HSPA5, HSPA6, and HSP60 (www.rtpimerdb.org).

Interferon activity and luciferase reporter assays

MLP and L4P activity was determined in luciferase reporter assays as described (Morris & Leppard, 2009; Wright *et al.*, 2015). IFN β promoter activity was measured by transfecting IFN β -Luc (King & Goodbourn, 1994) in the presence of either wild-type or mutant E4 Orf3 expression plasmids (Hoppe *et al.*, 2006) and pcDNAHisLacZ as an internal control for 24 h and stimulating by transfection with poly(I:C) for a further 8 h, otherwise as previously described (Chen *et al.*, 2015; Morris & Leppard, 2009). IFN activity in cell culture fluids was measured by plaque-reduction assay using infection of VERO cells by Semliki Forest virus (SFV). Subconfluent 12 well cultures were incubated for 24 h with either standard IFN α or with an unknown sample at 1 in 10 dilution, both in normal growth medium. After 24 h, cells were infected with 25 plaque-forming units of SFV, overlaid with agar-solidified medium and then fixed after 48 h incubation and plaques detected with crystal violet. All determinations were made in triplicate. Alternatively, IFN activity was determined by measuring the stimulation of pISRE-Luc by IFN-containing samples for 20 h in a luciferase reporter assay (Chen *et al.*, 2015).

Acknowledgement

Zeenah Atwan was supported by a studentship from The Higher Committee for Education Development in Iraq. Jordan Wright was supported by a studentship from the Biotechnology and Biological Sciences Research Council. Andrew Woodman was supported by a University of Warwick Undergraduate Research Student Scholarship.

References

- Ackrill, A. M., Foster, G. R., Laxton, C. D., Flavell, D. M., Stark, G. R. & Kerr, I. M. (1991). Inhibition of the cellular response to interferons by products of the adenovirus type 5 E1a oncogene. *Nucleic Acids Res* **19**, 4387-4393.
- Agoff, S. N., Hou, J. K., Linzer, D. I. & Wu, B. (1993). Regulation of the human hsp70 promoter by p53. *Science* **259**, 84-87.
- Au, W. C., Moore, P. A., Lowther, W., Juang, Y. T. & Pitha, P. M. (1995). Identification of a member of the interferon regulatory factor family that binds to the interferon-stimulated response element and activates expression of interferon-induced genes. *Proc Natl Acad Sci USA* **92**, 11657-11661.
- Bernardi, R. & Pandolfi, P. P. (2007). Structure, dynamics and functions of promyelocytic leukaemia nuclear bodies. *Nat Revs Mol Cell Biol* **8**, 1006-1016.
- Berscheminski, J., Groitl, P., Dobner, T., Wimmer, P. & Schreiner, S. (2013). The adenoviral oncogene E1A-13S interacts with a specific isoform of the tumor suppressor PML to enhance viral transcription. *J Virol* **87**, 965-977.
- Berscheminski, J., Wimmer, P., Brun, J., Ip, W. H., Groitl, P., Horlacher, T., Jaffray, E., Hay, R. T., Dobner, T. & Schreiner, S. (2014). Sp100 isoform-specific regulation of human adenovirus 5 gene expression. *J Virol* **88**, 6076-6092.
- Bringhurst, R. M. & Schaffer, P. A. (2006). Cellular stress rather than stage of the cell cycle enhances the replication and plating efficiencies of herpes simplex virus type 1 ICP0- viruses. *J Virol* **80**, 4528-4537.
- Brocchieri, L., Conway de Macario, E. & Macario, A. J. L. (2008). hsp70 genes in the human genome: Conservation and differentiation patterns predict a wide array of overlapping and specialized functions. *BMC Evolutionary Biology* **8**, 19-19.
- Burgert, H. G., Ruzsics, Z., Obermeier, S., Hilgendorf, A., Windheim, M. & Elsing, A. (2002). Subversion of host defense mechanisms by adenoviruses. In *Viral Proteins Counteracting Host Defenses*, pp. 273-318.
- Carvalho, T., Seeler, J. S., Ohman, K., Jordan, P., Pettersson, U., Akusjarvi, G., Carmo-Fonseca, M. & Dejean, A. (1995). Targeting of adenovirus E1A and E4-ORF3 proteins to nuclear matrix-associated PML bodies. *J Cell Biol* **131**, 45-56.
- Cepko, C. L., Whetsone, C. A. & Sharp, P. A. (1983). Adenovirus hexon monoclonal antibody that is group specific and potentially useful as a diagnostic reagent. *J Clin Microbiol* **17**, 360-364.
- Chahal, J. S., Qi, J. & Flint, S. J. (2012). The Human Adenovirus Type 5 E1B 55 kDa Protein Obstructs Inhibition of Viral Replication by Type I Interferon in Normal Human Cells. *Plos Path* **8**.
- Chee, A. V., Lopez, P., Pandolfi, P. P. & Roizman, B. (2003). Promyelocytic leukemia protein mediates interferon-based anti- herpes simplex virus 1 effects. *J Virol* **77**, 7101-7105.
- Chelbi-Alix, M. K., Pelicano, L., Quignon, F., Koken, M. H., Venturini, L., Stadler, M., Pavlovic, J., Degos, L. & de Thé, H. (1995). Induction of the PML protein by interferons in normal and APL cells. *Leukemia* **9**, 2027-2033.
- Chen, Y., Wright, J., Meng, X. & Leppard, K. N. (2015). Promyelocytic leukemia protein isoform II promotes transcription factor recruitment to activate interferon β and interferon-responsive gene expression. *Mol Cell Biol* **35**, 1660-1672.
- Clerico, E. M., Tilitsky, J. M., Meng, W. & Gierasch, L. M. (2015). How Hsp70 Molecular Machines Interact with Their Substrates to Mediate Diverse Physiological Functions. *J Mol Biol* **427**, 1575-1588.
- Colombo, R., Boggio, R., Seiser, C., Draetta, G. F. & Chiocca, S. (2002). The adenovirus protein Gam1 interferes with sumoylation of histone deacetylase 1. *EMBO Rep* **3**, 1062-1068.
- Cuchet, D., Sykes, A., Nicolas, A., Orr, A., Murray, J., Sirma, H., Heeren, J., Bartelt, A. & Everett, R. D. (2011). PML isoforms I and II participate in PML-dependent restriction of HSV-1 replication. *J Cell Sci* **124**, 280-291.

- Doucas, V., Ishov, A. M., Romo, A., Juguilon, H., Weitzman, M. D., Evans, R. M. & Maul, G. G. (1996). Adenovirus replication is coupled with the dynamic properties of the PML nuclear structure. *Genes & Development* **10**, 196-207.
- Evans, J. D. & Hearing, P. (2005). Relocalization of the Mre11-Rad50-Nbs1 complex by the adenovirus E4 ORF3 protein is required for viral replication. *J Virol* **79**, 6207-6215.
- Everett, R. D. & Murray, J. (2005). ND10 components relocate to sites associated with herpes simplex virus type 1 nucleoprotein complexes during virus infection. *J Virol* **79**, 5078-5089.
- Everett, R. D., Parada, C., Gripon, P., Sirma, H. & Orr, A. (2008a). Replication of ICP0-Null mutant herpes simplex virus type 1 is restricted by both PML and Sp100. *J Virol* **82**, 2661-2672.
- Everett, R. D., Young, D. F., Randall, R. E. & Orr, A. (2008b). STAT-1- and IRF-3-dependent pathways are not essential for repression of ICP0-null mutant herpes simplex virus type 1 in human fibroblasts. *J Virol* **82**, 8871-8881.
- Farley, D. C., Brown, J. L. & Leppard, K. N. (2004). Activation of the early-late switch in adenovirus type 5 major late transcription unit expression by L4 gene products. *J Virol* **78**, 1782-1791.
- Geoffroy, M. C. & Chelbi-Alix, M. K. (2011). Role of Promyelocytic Leukemia Protein in Host Antiviral Defense. *J Interferon Cytokine Res* **31**, 145-158.
- Glotzer, J. B., Saltik, M., Chiocca, S., Michou, A.-I., Moseley, P. & Cotten, M. (2000). Activation of heat-shock response by an adenovirus is essential for virus replication. *Nature* **407**, 207-211.
- Gooding, L. R., Elmore, L. W., Tollefson, A. E., Brady, H. A. & Wold, W. S. M. (1988). A 14,700 MW protein from the E3 region of adenovirus inhibits cytolysis by tumor necrosis factor. *Cell* **53**, 341-346.
- Halford, W. P., Weisend, C., Grace, J., Soboleski, M., Carr, D. J. J., Balliet, J. W., Imai, Y., Margolis, T. & Gebhardt, B. M. (2006). ICP0 antagonizes Stat 1-dependent repression of herpes simplex virus: implications for the regulation of viral latency. *Virology Journal* **3**, 44.
- Hartman, Z. C., Black, E. P. & Amalfitano, A. (2007). Adenoviral infection induces a multi-faceted innate cellular immune response that is mediated by the toll-like receptor pathway in A549 cells. *Virology* **358**, 357-372.
- Hendrickx, R., Stichling, N., Koelen, J., Kuryk, L., Lipiec, A. & Greber, U. F. (2014). Innate Immunity to Adenovirus. *Hum Gene Ther* **25**, 265-284.
- Higginbotham, J. N., Seth, P., Blaese, R. M. & Ramsey, W. J. (2002). The release of inflammatory cytokines from human peripheral blood mononuclear cells in vitro following exposure to adenovirus variants and capsid. *Hum Gene Ther* **13**, 129-141.
- Hoppe, A., Beech, S. J., Dimmock, J. & Leppard, K. N. (2006). Interaction of the adenovirus type 5 E4 Orf3 protein with promyelocytic leukemia protein isoform II is required for ND10 disruption. *J Virol* **80**, 3042-3049.
- Huang, M.-M. & Hearing, P. (1989). Adenovirus early region 4 encodes 2 gene products with redundant effects in lytic infection. *J Virol* **63**, 2605-2615.
- Imperiale, M. J., Kao, H. T., Feldman, L. T., Nevins, J. R. & Strickland, S. (1984). Common control of the heat shock gene and early adenovirus genes - evidence for a cellular E1a-like activity. *Mol Cell Biol* **4**, 867-874.
- Ishov, A. M. & Maul, G. G. (1996). The periphery of nuclear domain 10 (ND10) as site of DNA virus deposition. *J Cell Biol* **134**, 815-826.
- Jensen, K., Shiels, C. & Freemont, P. S. (2001). PML protein isoforms and the RBCC/TRIM motif. *Oncogene* **20**, 7223-7233.
- Kim, Y.-E. & Ahn, J.-H. (2015). Positive Role of Promyelocytic Leukemia Protein in Type I Interferon Response and Its Regulation by Human Cytomegalovirus. *PLoS Pathog* **11**, e1004785.
- King, P. & Goodbourn, S. (1994). The beta-interferon promoter responds to priming through multiple independent regulatory elements. *J Biol Chem* **269**, 30609-30615.
- Kitajewski, J., Schneider, R. J., Safer, B., Munemitsu, S. M., Samuel, C. E., Thimmappaya, B. & Shenk, T. (1986). Adenovirus VAI RNA antagonizes the antiviral action of interferon by preventing activation of the interferon-induced eIF-2-alpha kinase. *Cell* **45**, 195-200.

- Kumar, P. P., Bischof, O., Purbey, P. K., Notani, D., Urlaub, H., Dejean, A. & Galande, S. (2007).** Functional interaction between PML and SATB1 regulates chromatin-loop architecture and transcription of the MHC class I locus. *Nat Cell Biol* **9**, 45-56.
- Leib, D. A., Harrison, T. E., Laslo, K. M., Machalek, M. A., Moorman, N. J. & Virgin, H. W. (1999).** Interferons regulate the phenotype of wild-type and mutant herpes simplex viruses in vivo. *J Exp Med* **189**, 663-672.
- Leppard, K. N. (1998).** Regulated RNA processing and RNA transport during adenovirus infection. *Semin Virol* **8**, 301-307.
- Leppard, K. N. & Dimmock, J. (2006).** Virus interactions with PML nuclear bodies. In *Viruses and the nucleus*, pp. 213-245. Edited by J. Hiscox: J Wiley.
- Leppard, K. N., Emmott, E., Cortese, M. S. & Rich, T. (2009).** Adenovirus type 5 E4 Orf3 protein targets promyelocytic leukaemia (PML) protein nuclear domains for disruption via a sequence in PML isoform II that is predicted as a protein interaction site by bioinformatic analysis. *J Gen Virol* **90**, 95-104.
- Leppard, K. N. & Everett, R. D. (1999).** The adenovirus type 5 E1b 55K and E4 Orf3 proteins associate in infected cells and affect ND10 components. *J Gen Virol* **80**, 997-1008.
- Leppard, K. N. & Wright, J. (2012).** Targeting of promyelocytic leukaemia proteins and promyelocytic leukaemia nuclear bodies by DNA tumour viruses. In *Small DNA tumour viruses*, pp. 255 - 280. Edited by K. Gaston. Norfolk, UK: Caister Academic Press.
- Lethbridge, K. J., Scott, G. E. & Leppard, K. N. (2003).** Nuclear matrix localization and SUMO-1 modification of adenovirus type 5 E1b 55K protein are controlled by E4 Orf6 protein. *J Gen Virol* **84**, 259-268.
- Lum, L. S., Hsu, S., Vaewhongs, M. & Wu, B. (1992).** The hsp70 gene CCAAT-binding factor mediates transcriptional activation by the adenovirus E1a protein. *Mol Cell Biol* **12**, 2599-2605.
- Macejak, D. G. & Luftig, R. B. (1991).** Association of HSP70 with the adenovirus type 5 fiber protein in infected HEp-2 cells. *Virology* **180**, 120-125.
- Madara, J., Krewet, J. A. & Shah, M. (2005).** Heat shock protein 72 expression allows permissive replication of oncolytic adenovirus dl1520 (ONYX-015) in rat glioblastoma cells. *Molecular Cancer* **4**, 12.
- Maul, G. G. & Everett, R. D. (1994).** The nuclear location of PML, a cellular member of the C3HC4 zinc binding domain protein family, is rearranged during herpes simplex virus infection by the C3HC4 viral protein ICP0. *J Gen Virol* **75**, 1223-1233.
- McFadden, G., Mohamed, M. R., Rahman, M. M. & Bartee, E. (2009).** Cytokine determinants of viral tropism. *Nat Revs Immunol* **9**, 645-655.
- Meng, X. Z. & Harken, A. H. (2002).** The interaction between HSP70 and TNF-alpha expression: A novel mechanism for protection of the myocardium against post-injury depression. *Shock* **17**, 345-353.
- Mitchell, A. M., Hirsch, M. L., Li, C. W. & Samulski, R. J. (2014).** Promyelocytic Leukemia Protein Is a Cell-Intrinsic Factor Inhibiting Parvovirus DNA Replication. *J Virol* **88**, 925-936.
- Moffat, J., Grueneberg, D. A., Yang, X. P., Kim, S. Y., Kloepper, A. M., Hinkle, G., Piquani, B., Eisenhaure, T. M., Luo, B., Grenier, J. K., Carpenter, A. E., Foo, S. Y., Stewart, S. A., Stockwell, B. R., Hacohen, N., Hahn, W. C., Lander, E. S., Sabatini, D. M. & Root, D. E. (2006).** A lentiviral RNAi library for human and mouse genes applied to an arrayed viral high-content screen. *Cell* **124**, 1283-1298.
- Moore, M., Schaack, J., Baim, S. B., Morimoto, R. I. & Shenk, T. (1987).** Induced heat-shock messenger-RNAs escape the nucleocytoplasmic transport block in adenovirus-infected Hela cells. *Mol Cell Biol* **7**, 4505-4512.
- Morris, S. J. & Leppard, K. N. (2009).** Adenovirus serotype 5 L4-22K and L4-33K proteins have distinct functions in regulating late gene expression. *J Virol* **83**, 3049-3058.
- Morris, S. J., Scott, G. E. & Leppard, K. N. (2010).** Adenovirus late phase infection is controlled by a novel L4 promoter. *J Virol* **84**, 7096-7104.

- Nevins, J. R. (1982).** Induction of the synthesis of a 70,000 dalton mammalian heat shock protein by the adenovirus E1a gene-product. *Cell* **29**, 913-919.
- Niewiarowska, J., D'Halluin, J. C. & Belin, M.-T. (1992).** Adenovirus capsid proteins interact with HSP70 proteins after penetration in human or rodent cells. *Exp Cell Res* **201**, 408-416.
- Ou, H. D., Kwiatkowski, W., Deerinck, T. J., Noske, A., Blain, K. Y., Land, H. S., Soria, C., Powers, C. J., May, A. P., Shu, X. K., Tsien, R. Y., Fitzpatrick, J. A. J., Long, J. A., Ellisman, M. H., Choe, S. & O'Shea, C. C. (2012).** A Structural Basis for the Assembly and Functions of a Viral Polymer that Inactivates Multiple Tumor Suppressors. *Cell* **151**, 304-319.
- Pahl, H. L., Sester, M., Burgert, H. G. & Baeuerle, P. A. (1996).** Activation of transcription factor NF-kappa B by the adenovirus E3/19K protein requires its ER retention. *J Cell Biol* **132**, 511-522.
- Patsalo, V., Yondola, M. A., Luan, B. W., Shoshani, I., Kisker, C., Green, D. F., Raleigh, D. P. & Hearing, P. (2012).** Biophysical and functional analyses suggest that adenovirus E4-ORF3 protein requires higher-order multimerization to function against promyelocytic leukemia protein nuclear bodies. *J Biol Chem* **287**, 22573-22583.
- Phillips, B., Abravaya, K. & Morimoto, R. I. (1991).** Analysis of the specificity and mechanism of transcriptional activation of the human hsp70 gene during infection by DNA viruses. *J Virol* **65**, 5680-5692.
- Reich, N., Pine, R., Levy, D. & Darnell, J. E. (1988).** Transcription of interferon-stimulated genes is induced by adenovirus particles but is suppressed by E1a gene products. *J Virol* **62**, 114-119.
- Reich, N. C., Sarnow, P., Duprey, E. & Levine, A. J. (1983).** Monoclonal antibodies which recognize native and denatured forms of the adenovirus DNA-binding protein. *Virology* **128**, 480-484.
- Santoro, M. G., Amici, C. & Rossi, A. (2010).** Role of heat shock proteins in viral infection. In *Prokaryotic and eukaryotic heat shock proteins in infectious disease*, pp. 51-84. Edited by A. G. Pockley, S. K. Calderwood & M. G. Santoro: Springer.
- Saphire, A. C., Guan, T., Schirmer, E. C., Nemerow, G. R. & Gerace, L. (2000).** Nuclear import of adenovirus DNA in vitro involves the nuclear protein import pathway and hsc70. *J Biol Chem* **275**.
- Sarnow, P., Sullivan, C. A. & Levine, A. J. (1982).** A monoclonal antibody detecting the Ad5 E1b 58K tumor antigen: characterization of the E1b 58K tumor antigen in adenovirus infected and transformed cells. *Virology* **120**, 510-517.
- Schmitz, M. L., Indorf, A., Limbourg, F. P., Städtler, H., Traenckner, E. B. & Baeuerle, P. A. (1996).** The dual effect of adenovirus type 5 E1A 13S protein on NF-kappaB activation is antagonized by E1B 19K. *Mol Cell Biol* **16**, 4052-4063.
- Schreiner, S., Bürck, C., Glass, M., Groitl, P., Wimmer, P., Kinkley, S., Mund, A., Everett, R. D. & Dobner, T. (2013).** Control of human adenovirus type 5 gene expression by cellular Daxx/ATRX chromatin-associated complexes. *Nucleic Acids Res* **41**, 3532-3550.
- Simon, M. C., Fisch, T. M., Benecke, B. J., Nevins, J. R. & Heintz, N. (1988).** Definition of multiple, functionally distinct tata elements, one of which is a target in the hsp70 promoter for e1a regulation. *Cell* **52**, 723-729.
- Singh, I. S., Shah, N. G., Almutairy, E. & Hasday, J. D. (2010).** Role of HSF1 in infectious disease. In *Prokaryotic and eukaryotic heat shock proteins in infectious disease*, pp. 1-31. Edited by A. G. Pockley, S. K. Calderwood & M. G. Santoro: Springer.
- Sohn, S. Y. & Hearing, P. (2011).** Adenovirus sequesters phosphorylated STAT1 at viral replication centers and inhibits STAT dephosphorylation. *J Virol* **85**, 7555-7562.
- Stadler, M., Chelbi-Alix, M. K., Koken, M. H., Venturini, L., Lee, C., Saïb, A., Quignon, F., Pelicano, L., Guillemin, M. C. & Schindler, C. (1995).** Transcriptional induction of the PML growth suppressor gene by interferons is mediated through an ISRE and a GAS element. *Oncogene* **11**, 2565-2573.
- Tanaka, K.-I., Namba, T., Arai, Y., Fujimoto, M., Adachi, H., Sobue, G., Takeuchi, K., Nakai, A. & Mizushima, T. (2007).** Genetic Evidence for a Protective Role for Heat Shock Factor 1 and Heat Shock Protein 70 against Colitis. *J Biol Chem* **282**, 23240-23252.

- Tobe, M., Isobe, Y., Tomizawa, H., Nagasaki, T., Takahashi, H. & Hayashi, H. (2003).** A novel structural class of potent inhibitors of NF-kappa B activation: structure-activity relationships and biological effects of 6-aminoquinazoline derivatives. *Bioorg Med Chem* **11**, 3869-3878.
- Ullman, A. J. & Hearing, P. (2008).** Cellular proteins PML and Daxx mediate an innate antiviral defense antagonized by the adenovirus E4 ORF3 protein. *J Virol* **82**, 7325-7335.
- Ullman, A. J., Reich, N. C. & Hearing, P. (2007).** Adenovirus E4 ORF3 protein inhibits the interferon-mediated antiviral response. *J Virol* **81**, 4744-4752.
- Van Damme, E., Laukens, K., Dang, T. H. & Van Ostade, X. (2010).** A manually curated network of the PML nuclear body interactome reveals an important role for PML-NBs in SUMOylation dynamics. *Int J Biol Sci* **6**, 51-67.
- Weidtkamp-Peters, S., Lenser, T., Negorev, D., Gerstner, N., Hofmann, T. G., Schwanitz, G., Hoischen, C., Maul, G., Dittrich, P. & Hemmerich, P. (2008).** Dynamics of component exchange at PML nuclear bodies. *J Cell Sci* **121**, 2731-2743.
- White, E., Spector, D. & Welch, W. (1988).** Differential distribution of the adenovirus E1A proteins and colocalization of E1A with the 70-kilodalton cellular heat shock protein in infected cells. *J Virol* **62**, 4153-4166.
- Wright, J., Atwan, Z., Morris, S. J. & Leppard, K. N. (2015).** The Human Adenovirus Type 5 L4 Promoter Is Negatively Regulated by TFII-I and L4-33K. *J Virol* **89**, 7053-7063.
- Wu, B. J., Hurst, H. C., Jones, N. C. & Morimoto, R. I. (1986).** The E1A 13S product of adenovirus 5 activates transcription of the cellular human HSP70 gene. *Mol Cell Biol* **6**, 2994-2999.
- Xu, Z. X., Zou, W. X., Lin, P. & Chang, K. S. (2005).** A role for PML3 in centrosome duplication and genome stability. *Mol Cell* **17**, 721-732.
- Zhao, H. X., Granberg, F., Elfineh, L., Pettersson, U. & Svensson, C. (2003).** Strategic attack on host cell gene expression during adenovirus infection. *J Virol* **77**, 11006-11015.
- Zhu, J. G., Huang, X. P. & Yang, Y. P. (2007).** Innate immune response to adenoviral vectors is mediated by both Toll-like receptor-dependent and -independent pathways. *J Virol* **81**, 3170-3180.

Figures




Fig. 1. HelixDeltaII cells show physical and functional knockdown of PML-II. (a) Phase -contrast microscopic images of control Helix and shRNA Helix cells; scale bar 100 μ m. (b,c,d) HelixEV and HelixDeltaII cells were plated for 24 h, then RNA or protein samples were harvested. (b) PML-II mRNA was detected by RT-qPCR; results are normalized to the level detected in HelixEV cells and are the means and standard deviation of 3 technical replicates. (c) PML-II protein was detected by western blotting. (d) IL-6 and ISG56 mRNAs were analysed as in (b).




Fig. 2. Removal of PML-II increases Ad5 protein expression and virus yield. (a) HeLaEV and HeLaΔII cells were infected with wild type Ad5 at moi of 5, and total protein extracts at various times post-infection analysed by western blotting. Upper panel: Ad5 late protein; middle panels: Ad5 E1B 55K and E2A DNA binding protein (DBP) ; lower panel: GAPDH. The 24 h samples were loaded at a 1:100 dilution compared to the earlier time points. (b) Total protein extracts of HeLaEV and standard HeLa cells, infected for 20 h as in (a), were analysed for hexon expression. (c) Adenovirus gene expression by FACS analysis. Upper panel: late gene expression; lower panel: DBP expression; grey curves are the background (mock infected cells) while the black curves represent infected cells; the % of fluorescence-positive cells and their mean fluorescence intensity (MFI) are indicated on each panel. (d) Total virus in infected culture lysates was determined by fluorescent focus assay. Error bars show the standard deviation of replicates within an experiment; the experiment shown is representative of multiple experiments.




Fig. 3 PML-II inhibits Ad5 infection by both IFN-dependent and independent mechanisms. HeLaEV and HeLaΔII cells were plated for 24 h, transfected with 62.5 pmol/ml siRNA as indicated for 48 h and then either stimulated with poly I:C for 16 h (a, b) or infected with Ad5 wt300 at moi of 5 for 20 h (c), after which samples were prepared for analysis. (a) IRF3 protein or GAPDH (loading control) was detected by western blot. (b) RT-qPCR analysis detecting ISG56 mRNA; results are the means and standard deviation of 3 technical replicates. (c) As panel (a), but detecting hexon protein using anti-late protein polyclonal antibodies.





Fig. 4 Adenovirus E4 Orf3 inhibits IFN production and IFN β promoter activation. (a) HEK293 cells were infected at a multiplicity of 10 p.f.u./cell with wild type Ad5 (wt300) or mutant *inOrf3*, or mock-infected. Media was harvested at 8 h p.i. and IFN activity measured by plaque-reduction assay. (b) As panel (a) but using media from Ad5-infected MRC5 fibroblasts harvested at 16 h p.i. (c) Media from HEK293 cell cultures infected as in (a) was harvested at 6 h p.i. and IFN activity measured using an ISRE-luciferase reporter construct in HEK293 cells. Known amounts of recombinant IFN α were analyzed in parallel to provide a standard curve. (d) HEK293 cells were transfected with IFN β promoter luciferase reporter and β -galactosidase control plasmids together with PML-II Δ RBCC (125 ng) and from 125 – 625 ng E4 Orf3 plasmid as appropriate and then stimulated with poly(I:C) and reporter activities assayed 8 h later. (e-i) HEK293 cells were transfected with reporter plasmids as in (d) plus 150 – 600 ng (150 – 750 ng, panels e, g) of either wild-type E4 Orf3 plasmid (e), or mutant E4 Orf3 R100A (f), N82A (g), L103A (h) or D105A-L106A (i), and then stimulated or not with poly(I:C) as indicated and assayed as in panel (d). Data are the means and standard deviation of three biological replicates.



747

748 **Fig. 5. Role of E4 Orf3 in the response of Ad5 to PML-II depletion and IFN- α .** HeLaEV and HeLa Δ II
749 cells were either mock-treated (lanes 1, 2, 4, 6, 7, 9) or treated with 1000 U/ml of IFN- α for 24 h (+I;
750 lanes 3, 5, 8, 10), then mock-infected or infected with Ad5 wt300 or inOrf3 as indicated for 20 h.
751 Total protein lysates were collected and analysed by western blotting for Ad5 late proteins (upper)
752 or GAPDH (lower). Hexon protein bands were quantified using QuantityOne software, normalized to
753 GAPDH and expressed relative to the value for wt300 in HeLaEV cells.




Fig. 6. Elevated Hsp70 enhances the expression of Ad5 proteins when PML-II is reduced. (a, e) Samples were harvested from HeLaEV and HeLaΔII cells and analysed for HSP70 mRNA (a) or a selection of HSP mRNAs (e) by RT-qPCR, or (a, lower) for HSP70 protein by western blot. (b) HeLa cells were transfected or not with 125 pmol/ml siRNA as indicated and RNA harvested after 48 h for analysis of HSP70 mRNA by RT-qPCR. (c) HeLaEV and HeLaΔII cells were transfected with 125 pmol/ml HSP70 or control siRNA for 48 h, then infected with Ad5 wt300 for 20 h before RNA was harvested and analysed for hexon mRNA by RT-qPCR. (d) HeLaΔII cells were treated with siRNA and infected as in (c), then lysed and analysed by western blotting as in Fig. 5. In (a - c), data were standardized to an internal control and then normalized to values from: (a) HeLaEV; (b) siControl-treated HeLa; (c) siControl-treated HeLaEV. Panel (e) shows mRNA amounts measured separately for each amplicon, standardized in each case to an internal control. Graphs show the means and standard deviation of 3 technical replicates.





Fig. 7. Effects of elevated Hsp70 on NF-κB signalling do not cause enhanced Ad5 gene expression.

(a) HeLaEV cells were treated with HSP70 or control siRNA as in Fig. 6(c), then infected with Ad5 wt300 at moi of 5 for 20 h and ISG56 mRNA quantified by RT-qPCR. (b, d) Cells as indicated were treated or not with 50 ng TNFα for 1 h (b) or with 100 nM of the NF-κB inhibitor QNZ for 45 min (d), then infected with Ad5 wt300 at moi of 5 for 20 h. Protein samples were harvested and analysed for late protein expression by western blot. (c) HeLaEV cells, treated with siRNA as in (a) were treated with TNFα as in (b), infected with Ad5 wt300 as in (a) and hexon mRNA quantified by RT-qPCR. Results in panels a, c are the means and standard deviation of 3 technical replicates.

781



782

783 **Fig. 8. Hsp70 promotes nuclear accumulation of Ad5 hexon but has no effect on late promoter**
784 **activity.** (a) HeLaΔII cells were treated with HSP70 or control siRNA as in Fig. 6(c), then infected with
785 Ad5wt300 at moi of 5 for 16 hours. Cytoplasmic and nuclear fractions were analysed by SDS-PAGE
786 and western blotting. Replicate blots were probed with antibodies to the proteins indicated and
787 bands quantified as in Fig 5, normalized to GAPDH (cytoplasmic) or SP1 (nuclear). (b) HeLa cells were
788 treated with siRNA as in (a) and then transfected with MLP or L4P luciferase reporter. Luciferase
789 activity was measured after 20 h and normalized to a β -galactosidase transfection control as
790 described (Wright *et al.*, 2015). (c) HeLaEV and HeLaΔII cells were infected with wild type Ad5 at moi
791 of 5, and total protein extracts analysed by western blotting at 20 h p.i. for the proteins indicated.