

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/86064>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

INDUCED RANDOM β -TRANSFORMATION

SIMON BAKER AND KARMA DAJANI

ABSTRACT. In this article we study the first return map defined on the switch region induced by the greedy and lazy maps. In particular we study the allowable sequences of return times, and when the first return map is a generalised Lüroth series transformation. We show that there exists a countable collection of disjoint intervals $(\mathcal{I}_n)_{n=1}^{\infty}$, such that all sequences of return times are permissible if and only if $\beta \in \mathcal{I}_n$ for some n . Moreover, we show that there exists a set $M \subseteq (1, 2)$ of Hausdorff dimension 1 and Lebesgue measure zero, for which the first return map is a generalised Lüroth series transformation if and only if $\beta \in M$.

1. INTRODUCTION

Let $\beta \in (1, 2)$ and $I_\beta := [0, \frac{1}{\beta-1}]$. Given $x \in I_\beta$ we call a sequence $(b_n)_{n=1}^{\infty} \in \{0, 1\}^{\mathbb{N}}$ a β -expansion for x if

$$x = \sum_{n=1}^{\infty} \frac{b_n}{\beta^n}.$$

Non-integer representations of real numbers were pioneered in the papers of Rényi [13] and Parry [12]. Since then they have been studied by many authors and have connections with ergodic theory, fractal geometry, and number theory (see the survey articles [10] and [15]). Perhaps one of the most interesting objects to study within expansions in non-integer bases is the set of expansions, i.e.,

$$\Sigma_\beta(x) := \left\{ (b_n)_{n=1}^{\infty} \in \{0, 1\}^{\mathbb{N}} : \sum_{n=1}^{\infty} \frac{b_n}{\beta^n} = x \right\}.$$

A result of Sidorov states that given $\beta \in (1, 2)$ then Lebesgue almost every $x \in I_\beta$ satisfies $\text{card } \Sigma_\beta(x) = 2^{\aleph_0}$ [14]. Moreover, for any $k \in \mathbb{N} \cup \{\aleph_0\}$ there exists $\beta \in (1, 2)$ and $x \in I_\beta$ such that $\text{card } \Sigma_\beta(x) = k$, see [6, 8, 9]. The situation described above is completely different to the case of integer base expansions where every number has a unique expansion except for a countable set of exceptions which have precisely two.

A useful observation when studying expansions in non-integer bases is that a β -expansion has a natural dynamical interpretation. Namely, let $T_0(x) = \beta x$, $T_1(x) = \beta x - 1$, and

$$\Gamma_\beta(x) := \left\{ (a_n)_{n=1}^{\infty} \in \{T_0, T_1\}^{\mathbb{N}} : (a_n \circ \dots \circ a_1)(x) \in I_\beta \text{ for all } n \in \mathbb{N} \right\}.$$

It was shown in [1] that $\text{card } \Sigma_\beta(x) = \text{card } \Gamma_\beta(x)$ and the map sending (b_n) to (T_{b_n}) is a bijection between these two sets. As such, performing the map T_0 corresponds to taking the digit 0, and T_1 corresponds to taking the digit 1. An all encompassing method by which we can use the maps T_0 and T_1 to generate β -expansions is the *random β -transformation*. This map is defined as follows. Set $\Omega = \{0, 1\}^{\mathbb{N}}$ and denote by σ the left shift on Ω . Consider the transformation

2010 *Mathematics Subject Classification.* 11A63, 37A45.

Key words and phrases. β -expansions, First return maps, Lüroth transformations.

$K_\beta : \Omega \times [0, \frac{1}{\beta-1}] \rightarrow \Omega \times [0, \frac{1}{\beta-1}]$ defined by

$$K_\beta(\omega, x) = \begin{cases} (\omega, T_0x), & \text{if } 0 \leq x < \frac{1}{\beta}, \\ (\sigma\omega, T_{\omega_1}x), & \text{if } \frac{1}{\beta} \leq x \leq \frac{1}{\beta(\beta-1)}, \\ (\omega, T_1x), & \text{if } \frac{1}{\beta(\beta-1)} < x \leq \frac{1}{\beta-1}. \end{cases}$$

The random β -transformation K_β was introduced and studied in [3, 4, 5]. Given $x \in I_\beta$, the map K_β generates all possible β -expansions of x . Furthermore, it is a random mix of the classical greedy and lazy maps defined by

$$G_\beta(x) = \begin{cases} T_0(x), & \text{if } 0 \leq x < \frac{1}{\beta}, \\ T_1(x), & \text{if } \frac{1}{\beta} \leq x \leq \frac{1}{\beta-1}, \end{cases}$$

and

$$L_\beta(x) = \begin{cases} T_0(x), & \text{if } 0 \leq x < \frac{1}{\beta(\beta-1)}, \\ T_1(x), & \text{if } \frac{1}{\beta(\beta-1)} \leq x \leq \frac{1}{\beta-1} \end{cases}$$

respectively. Let $S := [\frac{1}{\beta}, \frac{1}{\beta(\beta-1)}]$, we refer to S as the *switch region*. This is the region where the greedy map G_β and lazy map L_β differ, and is the region where the coordinates of ω are used to decide which map to use. Understanding the dynamics of the maps T_0 and T_1 on the switch region provides valuable insight into the possible $\Gamma_\beta(x)$, and thus the possible $\Sigma_\beta(x)$.

This paper is concerned with the dynamics of the first return map defined on the switch region. We consider the induced transformation U_β of K_β on the set $\Omega \times S$. More precisely, $U_\beta : \Omega \times S \rightarrow \Omega \times S$ is defined as follows:

$$U_\beta(\omega, x) := K_\beta^{r_1(\omega, x)}(\omega, x), \text{ where } r_1(\omega, x) = \inf\{m \geq 1 : K_\beta^m(\omega, x) \in \Omega \times S\}.$$

Similarly we set

$$U_{\beta,0}(x) := U_\beta((0)^\infty, x) \text{ and } U_{\beta,1}(x) := U_\beta((1)^\infty, x).$$

Note that when we have fixed the sequence ω to equal $(0)^\infty$ or $(1)^\infty$ the maps $U_{\beta,0}$ and $U_{\beta,1}$ are well defined maps from S to S .

Remark 1.1. *The map U_β is defined on $\Omega \times S$, and both $U_{\beta,0}$ and $U_{\beta,1}$ are defined on S . However, there exists ω and x for which $K_\beta(\omega, x)$ is never mapped back into $\Omega \times S$, thus for this choice of ω and x the map U_β is not well defined. Similarly, there exists x for which U_{β,ω_i} is not well defined. However, it is a consequence of the work of Sidorov [14] that the set of x for which $U_\beta^n(\omega, x)$ is well defined for all $n \in \mathbb{N}$ and $\omega \in \Omega$ is of full Lebesgue within S . Similarly, the set of x for which $U_{\beta,\omega_i}^n(x)$ is well defined for every $n \in \mathbb{N}$ is of full Lebesgue measure within S . Throughout this article we will abuse notation and let S denote both the switch region and the full measure subset of S for which U_β and U_{β,ω_i} are well defined. It should be clear which interpretation of S we mean from the context.*

For $i \geq 1$ let $r_i(\omega, x) := r_1(U_\beta^{i-1}(\omega, x))$ be the i th return time to the switch region $\Omega \times S$. Note that for any β and ω , the set $\{r_1(\omega, x)\}_{x \in S}$ equals $\mathcal{R}_\beta := \{m, m+1, \dots\}$ where m is some natural number that only depends upon β . We emphasise that \mathcal{R}_β has no dependence on ω . One of the goals of this paper is to understand the sequences $(r_i(\omega, x))_{i=1}^\infty$ and to answer the following question: given $\omega \in \Omega$ and a sequence of integers $(j_i)_{i=1}^\infty \in \mathcal{R}_\beta^\mathbb{N}$, when is it possible to find $x \in S$ such that $r_i(\omega, x) = j_i$, for $i = 1, 2, \dots$? The following theorem provides an answer to this question. Before we state this theorem we have to introduce two classes of algebraic integers. Let α_k denote the unique solution in $(1, 2)$ of the equation

$$x^{k+1} - 2x^k + x - 1 = 0,$$

k	α_k	γ_k	η_k
1	$\frac{1+\sqrt{5}}{2}$	$\frac{1+\sqrt{5}}{2}$	$1 + 2^{-1/2}$
2	1.7549 ...	1.8393 ...	1.8546 ...
3	1.8668 ...	1.9276 ...	1.9305 ...
4	1.9332 ...	1.9660 ...	1.9666 ...
5	1.9672 ...	1.9836 ...	1.9837 ...

FIGURE 1. Tables of values for α_k , γ_k and η_k

and let γ_k denote the k -th multinacci number. Recall that the k -th multinacci number is the unique root of

$$x^{k+1} - x^k - x^{k-1} - \dots - x - 1 = 0$$

contained in (1, 2).

Theorem 1.1. *Let $\beta \in (\alpha_k, \gamma_k]$ for some $k \geq 2$, then for any $\omega \in \Omega$ and $(j_i) \in \mathcal{R}_\beta^{\mathbb{N}}$ there exists $x \in S$ such that $r_i(\omega, x) = j_i$, for $i = 1, 2, \dots$. Moreover, if $\beta \notin (\alpha_k, \gamma_k]$ for all $k \geq 2$, then there exists $\omega \in \Omega$ and $(j_i) \in \mathcal{R}_\beta^{\mathbb{N}}$ such that no $x \in S$ satisfies $r_i(\omega, x) = j_i$, for $i = 1, 2, \dots$.*

As we will see, the algebraic properties of α_k and γ_k correspond naturally to conditions on the orbit of 1 and its reflection $\frac{1}{\beta-1} - 1$. These points determine completely the dynamics of the greedy map G_β and lazy map L_β respectively, and hence it is not surprising that these points play a crucial role in our situation as well. For values of β lying outside of the intervals $(\alpha_k, \gamma_k]$ it is natural to ask whether the following weaker condition is satisfied: given $(j_i) \in \mathcal{R}_\beta^{\mathbb{N}}$ does there exist $\omega \in \Omega$ and $x \in S$ such that $r_i(\omega, x) = j_i$ for $i = 1, 2, \dots$. Let η_k denote the unique root of the equation

$$2x^{k+1} - 4x^k + 1 = 0$$

contained in (1, 2).

Theorem 1.2. *Let $\beta \in (\alpha_k, \eta_k]$ for some $k \geq 1$, then for any sequence $(j_i) \in \mathcal{R}_\beta^{\mathbb{N}}$ there exists $\omega \in \Omega$ and $x \in S$ such that $r_i(\omega, x) = j_i$ for $i = 1, 2, \dots$.*

If β satisfies the hypothesis of Theorem 1.2 then the orbit of 1 and $\frac{1}{\beta-1} - 1$ satisfy a *cross over property*. This cross over property is sufficient to prove Theorem 1.2. Note that $\alpha_k \leq \gamma_k \leq \eta_k$ for each $k \geq 1$. We include a tables of values for α_k , γ_k and η_k in Figure 1.

The second half of this paper is concerned with the maps $U_{\beta,0}$ and $U_{\beta,1}$. Before we state our results it is necessary to make a definition. Given a closed interval $[a, b]$, we call a map $T : [a, b] \rightarrow [a, b]$ a generalized Lüroth series transformation (abbreviated to GLST) if there exists a countable set of bounded subintervals $\{I_n\}_{n=1}^\infty$ ($I_n = (l_n, r_n), [l_n, r_n], (l_n, r_n], [l_n, r_n)$) for which the following criteria are satisfied:

- (1) $I_n \cap I_m = \emptyset$ for $n \neq m$.
- (2) $\sum_{n=1}^\infty (r_n - l_n) = b - a$.
- (3)

$$T(x) = a + \frac{(x - l_n)(b - a)}{r_n - l_n}$$

for $x \in I_n$.

Property (3) is equivalent to the map T restricted to the interval I_n being the unique surjective linear orientation preserving map from the interval I_n into S .

The traditional Lüroth expansion of a number $x \in (0, 1]$ is a sequence of natural numbers $(a_n)_{n=1}^\infty$ where each $a_n \geq 2$ and

$$x = \frac{1}{a_1} + \frac{1}{a_1(a_1 - 1)a_2} + \dots + \frac{1}{a_1(a_1 - 1)a_2(a_2 - 1) \dots a_n} + \dots$$

This Lüroth expansion (a_n) can be seen to be generated by the map $T : [0, 1] \rightarrow [0, 1]$ where

$$T(x) = \begin{cases} n(n+1)x - n, & \text{if } x \in (\frac{1}{n+1}, \frac{1}{n}] \\ 0, & \text{if } x = 0 \end{cases}$$

GLST's were introduced in [2]. Our definition is slightly different to that appearing in this paper but all of the main results translate over into our context. Namely if $T : [a, b] \rightarrow [a, b]$ is a GLST then the normalised Lebesgue measure on $[a, b]$ is a T -invariant ergodic measure. Our main result for the maps $U_{\beta,0}$ and $U_{\beta,1}$ is the following theorem.

Theorem 1.3. *There exists a set $M \subseteq (1, 2)$ of Hausdorff dimension 1 and Lebesgue measure zero such that:*

- (1) *If $\beta \in M$ then both $U_{\beta,0}$ and $U_{\beta,1}$ are GLSTs.*
- (2) *If $\beta \notin M$ then both $U_{\beta,0}$ and $U_{\beta,1}$ are not GLSTs.*

What is more we can describe the set M explicitly.

Before we move on to our proofs of Theorems 1.1, 1.2 and 1.3 we provide a worked example. Namely we consider the case where $\beta = \frac{1+\sqrt{5}}{2}$. This case exhibits some of the important features of our later proofs.

Example 1.1. *When $\beta = \frac{1+\sqrt{5}}{2}$ then $S = [\frac{1}{\beta}, 1]$. Let $C_j = \{\omega \in \Omega : \omega_1 = j\}$, $j = 0, 1$, then for any $\omega \in C_0$, $r_1(\omega, 1) = \infty$, and $r_1(\omega, \frac{1}{\beta}) = 1$, while for any $\omega \in C_1$, we have $r_1(\omega, 1) = 1$ and $r_1(\omega, \frac{1}{\beta}) = \infty$. If $x \in (\frac{1}{\beta}, 1)$, then $r_1(\omega, x) \geq 2$ for all $\omega \in \Omega$.*

Let

$$(1) \quad B_i^0 := \{x \in S : U_{\beta,0}(x) = (T_1^{i-1} \circ T_0)(x)\}$$

and

$$(2) \quad B_i^1 := \{x \in S : U_{\beta,1}(x) = (T_0^{i-1} \circ T_1)(x)\}$$

where $i \geq 2$. A simple calculation shows that

$$(3) \quad B_i^0 = \left(\sum_{n=2}^{i+1} \frac{1}{\beta^n}, \sum_{n=2}^{i+2} \frac{1}{\beta^n} \right] = (T_1^{i-1} \circ T_0)^{-1} \left(\frac{1}{\beta}, 1 \right]$$

and


$$(4) \quad B_i^1 = \left[\frac{1}{\beta} + \frac{1}{\beta^{i+1}}, \frac{1}{\beta} + \frac{1}{\beta^i} \right) = (T_0^{i-1} \circ T_1)^{-1} \left[\frac{1}{\beta}, 1 \right),$$

where $i \geq 2$.

The collection $\{B_i^0 : i \geq 2\}$ is a partition of $(\frac{1}{\beta}, 1)$, and $\{B_i^1 : i \geq 2\}$ is a partition of $(\frac{1}{\beta}, 1)$. Equation (3) demonstrates that $U_{\beta,0}$ restricted to B_i^0 is a full branch, thus $U_{\beta,0}$ is a GLST. Similarly equation (4) implies $U_{\beta,1}$ is a GLSTs. We include a diagram of the graph of $U_{\beta,0}$ in Figure 2.

By the aforementioned results of [2] we know that a GLST is ergodic with respect to the normalised Lebesgue measure μ . As such we can state the average return time. For $\beta = \frac{1+\sqrt{5}}{2}$ Lebesgue almost every $x \in S$ satisfies

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} r_j((0)^\infty, x) &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \sum_{i=2}^{\infty} i \chi_{B_i^0}((U_{\beta,0})^j(x)) \\ &= \int \sum_{i=2}^{\infty} i \chi_{B_i^0} d\mu \\ &= 2\beta^2 - \beta \\ &= 3.6178 \dots \end{aligned}$$

FIGURE 2. The graph of $U_{\beta,0}$ when $\beta = \frac{1+\sqrt{5}}{2}$

Where in the above $\chi_{B_i^0}$ denotes the characteristic function on B_i^0 . Note that the result stated above holds with $r_j((0)^\infty, x)$ replaced with $r_j((1)^\infty, x)$.

By Theorem 1.1 we know that there exists $\omega \in \Omega$ and $(j_i)_{i=1}^\infty \in R_{\frac{1+\sqrt{5}}{2}}^\mathbb{N}$ for which no x satisfies $r_i(\omega, x) = j_i$ for $i = 1, 2, \dots$. This is essentially a consequence of the fact mentioned above that if $\omega \in C_0$ then $r_1(\omega, 1) = \infty$, and $r_1(\omega, \frac{1}{\beta}) = 1$, while for any $\omega \in C_1$, we have $r_1(\omega, 1) = 1$ and $r_1(\omega, \frac{1}{\beta}) = \infty$. This statement implies that we cannot have a return time 1 followed by any other natural number.

2. SEQUENCES OF RETURN TIMES

2.1. Proof of Theorem 1.1. In this section we prove Theorem 1.1. The proofs of Theorems 1.1 and 1.2 both make use of a nested interval construction. We begin by examining the condition $\beta \in (\alpha_k, \gamma_k]$. It is easy to show that the following statements hold:

$$(5) \quad \beta \in (\alpha_k, 2) \iff (T_1^{k-1} \circ T_0)\left(\frac{1}{\beta}\right) > \frac{1}{\beta(\beta-1)} \iff (T_0^{k-1} \circ T_1)\left(\frac{1}{\beta(\beta-1)}\right) < \frac{1}{\beta}$$

and

$$(6) \quad \beta \in (1, \gamma_k] \iff (T_1^k \circ T_0)\left(\frac{1}{\beta}\right) \leq \frac{1}{\beta} \iff (T_0^k \circ T_1)\left(\frac{1}{\beta(\beta-1)}\right) \geq \frac{1}{\beta(\beta-1)}.$$

Thus $\beta \in (\alpha_k, \gamma_k]$ is equivalent to the orbit of $\frac{1}{\beta}$ either jumping over the switch region, or satisfying $U_{\beta,0}(\frac{1}{\beta}) = (T_1^k \circ T_0)(\frac{1}{\beta}) = \frac{1}{\beta}$. Similarly, $\beta \in (\alpha_k, \gamma_k]$ is equivalent to the orbit of $\frac{1}{\beta(\beta-1)}$ either jumping over the switch region, or satisfying $U_{\beta,1}(\frac{1}{\beta(\beta-1)}) = (T_0^k \circ T_1)(\frac{1}{\beta(\beta-1)}) = \frac{1}{\beta(\beta-1)}$. The

following properties are important consequences of the above. First of all it is straightforward to see that for $\beta \in (\alpha_k, \gamma_k]$ we have $R_\beta = \{k+1, k+2, \dots\}$. Secondly we have

$$(7) \quad B_i^0 = (T_1^{i-1} \circ T_0)^{-1}(S)$$

and

$$(8) \quad B_i^1 = (T_0^{i-1} \circ T_1)^{-1}(S)$$

for $i \geq k+1$ where B_i^0 and B_i^1 are as in Example 1.1, but in this case they do not form a partition of S . We now prove Theorem 1.1, we separate our proof into the following propositions..

Proposition 2.1. *Let $\beta \in (\alpha_k, \gamma_k]$ for some $k \geq 2$, then for any $\omega \in \Omega$ and $(j_i) \in \mathcal{R}_\beta^\mathbb{N}$ there exists $x \in S$ such that $r_i(\omega, x) = j_i$, for $i = 1, 2, \dots$*

Proof. Let $\beta \in (\alpha_k, \gamma_k]$ and let us fix $(\omega_i) \in \Omega$ and $(j_i) \in \{k+1, k+2, \dots\}^\mathbb{N}$. We let $\mathcal{I}_1 = B_{j_1}^{\omega_1}$ and

$$(9) \quad \mathcal{I}_i := B_{j_1}^{\omega_1} \cap (T_{\bar{\omega}_1}^{j_1-1} \circ T_{\omega_1})^{-1}(B_{j_2}^{\omega_2}) \cap \dots \cap \left((T_{\bar{\omega}_{i-1}}^{j_{i-1}-1} \circ T_{\omega_{i-1}}) \circ \dots \circ (T_{\bar{\omega}_1}^{j_1-1} \circ T_{\omega_1}) \right)^{-1}(B_{j_i}^{\omega_i})$$

for $i \geq 2$. In the above and throughout we let $\bar{\omega}_i = 1 - \omega_i$. Any element of \mathcal{I}_i satisfies $r_l(\omega, x) = j_l$ for $1 \leq l \leq i$. Note that by Equations (7) and (8) we have $(T_{\bar{\omega}_1}^{j_1-1} \circ T_{\omega_1})(\mathcal{I}_1) = S$, by an induction argument it can be shown that

$$(10) \quad \left((T_{\bar{\omega}_i}^{j_i-1} \circ T_{\omega_i}) \circ \dots \circ (T_{\bar{\omega}_1}^{j_1-1} \circ T_{\omega_1}) \right)(\mathcal{I}_i) = S$$

for all $i \in \mathbb{N}$. Equation (10) guarantees that \mathcal{I}_i is nonempty and well defined for each $i \in \mathbb{N}$. Moreover, $\mathcal{I}_{i+1} \subseteq \mathcal{I}_i$ by equation (9). Thus (\mathcal{I}_i) is a decreasing sequence of compact intervals and

$$E = \bigcap_{i=1}^{\infty} \mathcal{I}_i$$

is nonempty. Finally, any $x \in E$ satisfies $r_i(\omega, x) = j_i$ for all $i \in \mathbb{N}$. \square

Proposition 2.2. *Let $\beta \in (1, \frac{1+\sqrt{5}}{2}]$, then there exists $\omega \in \Omega$ and $(j_i) \in \mathcal{R}_\beta^\mathbb{N}$ such that no $x \in S$ satisfies $r_i(\omega, x) = j_i$, for $i = 1, 2, \dots$*

Proof. Any $\beta \in (1, \frac{1+\sqrt{5}}{2}]$ satisfies $R_\beta = \{1, 2, \dots\}$. We now fix the sequence $\omega = (0)^\infty$ and $(j_i) = (1)^\infty$. There exists no $x \in S$ satisfying $r_i((0)^\infty, x) = 1$ for all $i \geq 1$, as this would imply there exists $x \in S$ satisfying $T_0^i(x) \in S$ for all $i \geq 1$. This is not possible as repeated iteration of T_0 eventually maps any element of S outside of S . \square

Proposition 2.3. *Let $\beta \in (\frac{1+\sqrt{5}}{2}, 2) \setminus \bigcup_{k=2}^{\infty} (\alpha_k, \gamma_k]$, then there exists $\omega \in \Omega$ and $(j_i) \in \mathcal{R}_\beta^\mathbb{N}$ such that no $x \in S$ satisfies $r_i(\omega, x) = j_i$, for $i = 1, 2, \dots$*

Proof. For $\beta \in (\frac{1+\sqrt{5}}{2}, 2)$ we have $r_1(\omega, 1/\beta) \geq 2$ for any $\omega \in C_0$. Moreover, by our assumption that $\beta \notin (\alpha_k, \gamma_k]$ for any $k \geq 2$ we must have

$$(T_1^k \circ T_0)\left(\frac{1}{\beta}\right) \in \left(\frac{1}{\beta}, \frac{1}{\beta(\beta-1)}\right]$$

for some $k \geq 1$. For such a β we have $R_\beta := \{k+1, k+2, \dots\}$. Let $\omega = (0)^\infty$ and $(j_i) = (k+1)^\infty$, we now show that there exists no $x \in S$ satisfying $r_i((0)^\infty, x) = k+1$ for all i . Since $k+1$ is the earliest return times there exists a single interval \mathcal{I} for which $\mathcal{I} := \{x \in S : r_1((0)^\infty, x) = k+1\}$, moreover for any $x \in \mathcal{I}$ we have $U_{\beta,0}(x) = (T_1^k \circ T_0)(x)$. Thus, any x satisfying $r_i((0)^\infty, x) = k+1$ for all $i \in \mathbb{N}$ must satisfy

$$(11) \quad (T_1^k \circ T_0)^i(x) \in S \text{ for all } i \in \mathbb{N}.$$

We now explain why this is not possible.

The map $T_1^k \circ T_0$ scales distances by a factor β^{k+1} and satisfies $(T_1^k \circ T_0)(x) > x$ for x to the right of the fixed point of $T_1^k \circ T_0$. We previously observed that $(T_1^k \circ T_0)(1/\beta) \in (\frac{1}{\beta}, \frac{1}{\beta(\beta-1)}]$ thus the fixed point of $T_1^k \circ T_0$ is to the left of S . Therefore under repeated iteration of the map $T_1^k \circ T_0$

every $x \in S$ is eventually mapped outside of S . This implies that equation (11) cannot hold and we have proved our result. \square

Combining Propositions 2.1, 2.2 and 2.3 we conclude Theorem 1.1.

2.2. Proof of Theorem 1.2. We now prove Theorem 1.2 our proof is similar to Theorem 1.1 in that we make use of a nested interval construction. However, with our proof we do not explicitly construct the desired ω , we can only show existence, as such our proof takes on an added degree of abstraction.

Let us start by examining the consequences of $\beta \in (\alpha_k, \eta_k]$ for some $k \geq 1$. For $k \geq 2$ we ignore the intervals $(\alpha_k, \gamma_k]$ as their proof is covered by Theorem 1.1. For β in the remaining parameter space the following inclusions hold

$$(12) \quad \begin{aligned} (T_1^k \circ T_0)\left(\frac{1}{\beta}\right) &\in \left(\frac{1}{\beta}, \frac{1}{2(\beta-1)}\right] \\ (T_0^k \circ T_1)\left(\frac{1}{\beta(\beta-1)}\right) &\in \left[\frac{1}{2(\beta-1)}, \frac{1}{\beta(\beta-1)}\right). \end{aligned}$$

We emphasise that for any $\beta \in (1, 2)$ the point $\frac{1}{2(\beta-1)}$ is the midpoint of the interval S and is thus always in the interior of S . Equation (12) is equivalent to $U_{\beta,0}(\frac{1}{\beta})$ being contained in the left hand side of S , and $U_{\beta,1}(\frac{1}{\beta(\beta-1)})$ being contained in the right hand side of S . As such the two orbits cross over when they return to S .

The cross over property described by equation (12) implies

$$(13) \quad (T_1^k \circ T_0)(B_{k+1}^0) \cup (T_0^k \circ T_1)(B_{k+1}^1) = S.$$

Moreover, for any $i \geq k+1$ we have

$$(14) \quad (T_1^i \circ T_0)(B_{i+1}^0) = S$$

With the identities (13) and (14) we may now prove Theorem 1.2.

Proof of Theorem 1.2. Let $\beta \in (\gamma_k, \eta_k]$ and let us fix a sequence of return times $(j_i) \in R_\beta^\mathbb{N} = \{k+1, k+2, \dots\}^\mathbb{N}$. We will construct a set J , such that for any $x \in J$ there exists a sequence ω satisfying $r_i(\omega, x) = j_i$ for all $i \in \mathbb{N}$. We construct J by building a sequence of levels J_1, J_2, \dots . Each J_i will denote a finite collection of compact intervals $\{\mathcal{I}_l^i\}_{l=1}^{2^i}$. Moreover,

$$(15) \quad \bigcup_{l=1}^{2^{i+1}} \mathcal{I}_l^{i+1} \subseteq \bigcup_{l=1}^{2^i} \mathcal{I}_l^i$$

for each $i = 1, 2, \dots$. Thus

$$J = \bigcap_{i=1}^{\infty} \bigcup_{l=1}^{2^i} \mathcal{I}_l^i$$

is nonempty, and as we will see, for each $x \in J$ there exists an $\omega \in \Omega$ such that $r_i(\omega, x) = j_i$ for $i = 1, 2, \dots$. We emphasise that in our construction not every \mathcal{I}_l^i will necessarily be nonempty.

For each level J_i it is useful to define a collection of maps $M_i = \{f_l^i\}_{l=1}^{2^i}$. Each f_l^i will be a map from \mathcal{I}_l^i into S . These maps will also have the property that

$$(16) \quad \bigcup_{l=1}^{2^i} f_l^i(\mathcal{I}_l^i) = S.$$

We start by letting

$$J_1 = \{B_{j_1}^0, B_{j_1}^1\} \text{ and } M_1 = \{T_1^{j_1-1} \circ T_0, T_0^{j_1-1} \circ T_1\}.$$

By Equations (13) and (14) we have

$$(T_1^{j_1-1} \circ T_0)(B_{j_1}^0) \cup (T_0^{j_1-1} \circ T_1)(B_{j_1}^1) = S$$

So we satisfy (16) when $i = 1$. Assume we have constructed J_i and M_i for $1 \leq i \leq N$, and (15) holds for $1 \leq i \leq N-1$, and (16) holds for $1 \leq i \leq N$. We now construct J_{N+1} and M_{N+1} . To each $f_i^N \in M_N$ we associate the compact intervals $(f_i^N)^{-1}(B_{j_{N+1}}^0)$ and $(f_i^N)^{-1}(B_{j_{N+1}}^1)$, the set of these new intervals is our J_{N+1} . By (16) this collection of intervals $\{(f_i^N)^{-1}(B_{j_{N+1}}^0), (f_i^N)^{-1}(B_{j_{N+1}}^1)\}$ is nonempty. Each f_i^N is a map from \mathcal{I}_i^N into S , thus $(f_i^N)^{-1}(B_{j_{N+1}}^1) \subseteq \mathcal{I}_i^N$ and we have that equation (15) holds for $i = N$.

To each $(f_i^N)^{-1}(B_{j_{N+1}}^0)$ we associate the map $(T_1^{j_{N+1}-1} \circ T_0) \circ f_i^N$, and to each $(f_i^N)^{-1}(B_{j_{N+1}}^1)$ we associate the map $(T_0^{j_{N+1}-1} \circ T_1) \circ f_i^N$ respectively. This collection of maps is our new M_{N+1} . Moreover

$$\begin{aligned} & \left(\bigcup_{l=1}^{2^N} ((T_1^{j_{N+1}-1} \circ T_0) \circ f_l^N) \circ (f_l^N)^{-1}(B_{j_{N+1}}^0) \right) \cup \left(\bigcup_{l=1}^{2^N} ((T_0^{j_{N+1}-1} \circ T_1) \circ f_l^N) \circ (f_l^N)^{-1}(B_{j_{N+1}}^1) \right) \\ &= (T_1^{j_{N+1}-1} \circ T_0) \left(\bigcup_{l=1}^{2^N} f_l^N (f_l^N)^{-1}(B_{j_{N+1}}^0) \right) \cup (T_0^{j_{N+1}-1} \circ T_1) \left(\bigcup_{l=1}^{2^N} f_l^N (f_l^N)^{-1}(B_{j_{N+1}}^1) \right) \\ &= (T_1^{j_{N+1}-1} \circ T_0)(B_{j_{N+1}}^0) \cup (T_0^{j_{N+1}-1} \circ T_1)(B_{j_{N+1}}^1) \quad (\text{By (16) for } i = N) \\ &= S \quad (\text{By (13) and (14)}). \end{aligned}$$

Therefore we satisfy (16) for $i = N+1$. As such we can repeat the above steps indefinitely and J_i and M_i are well defined for all $i \in \mathbb{N}$ and satisfy equations (15) and (16). This implies that the set J is well defined and nonempty.

It is not immediately obvious why an $x \in J$ admits an $\omega \in \Omega$ such that $r_i(\omega, x) = j_i$ for all $i \geq 1$. We now explain why. If $x \in J$, then by our construction for each $n \in \mathbb{N}$ there exists $(\omega_i^n)_{i=1}^n \in \{0, 1\}^n$ such that

$$(17) \quad (T_{\omega_i^n}^{j_i} \circ T_{\omega_i^n}) \circ \dots \circ (T_{\omega_1^n}^{j_1} \circ T_{\omega_1^n})(x) \in S$$

for all $1 \leq i \leq n$. We identify the finite sequence (ω_i^n) with the infinite sequence $v_n = (\omega_1^n, \dots, \omega_n^n, (0)^\infty)$. We equip Ω with the usual metric $d(\cdot, \cdot)$ where $d((\epsilon_i), (\delta_i)) = 2^{-n((\epsilon_i), (\delta_i))}$ where $n(x, y) = \inf\{i : \epsilon_i \neq \delta_i\}$. With respect to this metric Ω is a compact metric space, thus there exists $v \in \Omega$ and a subsequence of the (v_n) such that $v_{n_k} \rightarrow v$. This v has the property that

$$(18) \quad (T_{v_i}^{j_i} \circ T_{v_i}) \circ \dots \circ (T_{v_1}^{j_1} \circ T_{v_1})(x) \in S$$

for all $i \in \mathbb{N}$. (18) is a consequence of v being the limit of sequences satisfying (17). Clearly (18) implies that $r_i(v, x) = j_i$ for all $i \in \mathbb{N}$. \square

Remark 2.1. We end this section by pointing out that there are non trivial examples of $\beta \in (1, 2)$ for which there exists $(j_i) \in \mathbb{R}_\beta^\mathbb{N}$ and no $x \in S$ and $\omega \in \Omega$ for which $r_i(\omega, x) = j_i$ for all $i \in \mathbb{N}$. For example take $\beta = 1.754$. We chose β to be this value because it is slightly less than α_2 . Thus $T_1 \circ T_0(\frac{1}{\beta}) \in S$, but it is only slightly less than the right end point of the switch region. Clearly $R_\beta := \{2, 3, \dots\}$. However, any point that can have a return time two gets mapped close to the endpoints of S under the corresponding map. Being close to the endpoints of the switch suggests either a large return time or a small return time. This is the case for $\beta = 1.754$, and a simple calculation shows that it is not possible for $r_1(\omega, x) = 2$ and $r_2(\omega, x) = 3$.

3. PROOF OF THEOREM 1.3

Let us begin our proof of Theorem 1.3 by defining the set M that appear in its statement. Let

$$M := \left\{ \beta \in (1, 2) : \text{card } \Sigma_\beta(1) = 1 \right\} \cup \left\{ \beta \in (1, 2) : U_{\beta, 0}\left(\frac{1}{\beta}\right) \in \left\{ \frac{1}{\beta}, \frac{1}{\beta(\beta-1)} \right\} \right\}.$$

The first set in this union is the set of univoque bases, the study of this set is classical within expansions in noninteger bases, we refer the reader to the following papers for more on this subject [7, 8, 9, 11]. In [9] Erdős and Joó showed that the set of univoque bases has Hausdorff dimension 1 and Lebesgue measure zero. The second set in the above union is a countable set of algebraic

numbers, thus M has Hausdorff dimension 1 and Lebesgue measure zero. It is worth noting that if $\beta \in \{\beta \in (1, 2) : U_{\beta,0}(\frac{1}{\beta}) \in \{\frac{1}{\beta}, \frac{1}{\beta(\beta-1)}\}\}$ then $\text{card } \Sigma_\beta(1) = \aleph_0$. The important observation to make from the definition of M is that the following statement holds

$$\beta \in M \iff \frac{1}{\beta} \text{ and } \frac{1}{\beta(\beta-1)} \text{ are never mapped into the interior of } S.$$

This property will be sufficient to prove that both $U_{\beta,0}$ and $U_{\beta,1}$ are GLSTs. Our proof of Theorem 1.3 is split over the following propositions.

Proposition 3.1. *If $\beta \notin M$ then $U_{\beta,0}$ and $U_{\beta,1}$ are not GLSTs.*

Proof. If $\beta \notin M$ then $U_{\beta,0}(\frac{1}{\beta}) \in S^0$. In which case at the left endpoint of S the graph of $U_{\beta,0}$ has an incomplete branch. Thus it is not possible that $U_{\beta,0}$ is a GLST as all of the branches are full for this class of transformation. The proof that $U_{\beta,1}$ is not a GLST is similar and appeals to the fact that $U_{\beta,1}(\frac{1}{\beta(\beta-1)}) \in S^0$. \square

Proposition 3.2. *If $\beta \in M$ then $U_{\beta,0}$ and $U_{\beta,1}$ are GLSTs.*

We will only show that if $\beta \in M$ then $U_{\beta,0}$ is a GLST, the proof for $U_{\beta,1}$ being analogous. Moreover, as we previously demonstrated in Example 1.1 that the maps $U_{\beta,0}$ and $U_{\beta,1}$ were GLSTs for $\beta = \frac{1+\sqrt{5}}{2}$ we restrict our attention to the interval $(\frac{1+\sqrt{5}}{2}, 2)$, where the rest of the set M exists.

Before proceeding with our proof that $U_{\beta,0}$ is a GLST we make several observations. Let $\beta \in (\frac{1+\sqrt{5}}{2}, 2)$ and $x \in S$ be such that $U_{\beta,0}$ is well defined, then

$$(19) \quad U_{\beta,0}(x) = (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)(x)$$

for some $\omega_i \in \{0, 1\}$ that alternate digits with $\omega_1 = 1$. Equation (19) holds because the map T_0 maps every element of S outside of S . The quantity $i - 1$ is the number of times x jumps over S before eventually being mapped inside. Note that if i is even then $\omega_i = 0$ and if i is odd then $\omega_i = 1$.

Let

$$C_n := T_0^{-n}(S) \text{ and } D_n := T_1^{-n}(S)$$

where $n \in \mathbb{N}$. Equation (19) demonstrates that if $U_{\beta,0}(x)$ is well defined then x must eventually map into a C_n or a D_n . Note that for $\beta \in (\frac{1+\sqrt{5}}{2}, 2)$ the C_n are all disjoint and contained in the interval $(0, \frac{1}{\beta})$, and similarly the D_n are all disjoint and contained in $(\frac{1}{\beta(\beta-1)}, \frac{1}{\beta-1})$

It is instructive here to make a final notational remark before we give our proof. As we will see, the proof of Proposition 3.2 relies heavily on understanding the trajectories of certain intervals under certain maps and where they lie relative to C_n, D_n and S . Often we will be in a situation where a relation $(I \cap J = \emptyset, I \subseteq J)$ is true only if we ignore the endpoints of these intervals. For ease of exposition instead of repeatedly emphasising the fact that this relation holds modulo the endpoints we will simply state that the equation holds. This is technically not correct, but our proof still holds and is far more succinct by adopting this convention.

Proof of Proposition 3.2. To prove $U_{\beta,0}$ is a GLST it suffices to show that for any $x \in S$ such that $U_{\beta,0}(x)$ is well defined then we have

$$(20) \quad \{y \in S : U_{\beta,0}(y) = (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)(y)\} = (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)^{-1}(S).$$

Where we have assumed $U_{\beta,0}(x) = (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)(x)$. We now explain why Equation (20) implies $U_{\beta,0}$ is a GLST. The intervals on the left hand side of equation (20) are all disjoint, thus we satisfy part (1) of the definition of a GLST. By Sidorov's result we know that for Lebesgue almost every $x \in S$ the map $U_{\beta,0}(x)$ is well defined, thus the lengths of the intervals on the left hand side of equation (20) sum up to equal the length of S and we satisfy part (2) of the definition of a GLST. Lastly, the right hand side of equation (20) demonstrates that $U_{\beta,0}$ restricted to this interval is surjective onto S , since there is a unique surjective linear orientation preserving map from this interval onto S we also satisfy part (3) of the definition of a GLST.

We begin with the most simple case, we assume that $U_{\beta,0}(x) = (T_1^{n_1} \circ T_0)(x)$, i.e. $T_0(x) \in D_{n_1}$. Importantly, since $\beta \in M$ we know that $1 \notin D_{n_1}^0$. Thus $T_0(S) \cap D_{n_1} = [1, \frac{1}{\beta-1}] \cap D_{n_1} = D_{n_1}$. Therefore $T_0^{-1}(D_{n_1}) \subseteq S$ and any y in this interval satisfies $U_{\beta,0}(y) = (T_1^{n_1} \circ T_0)(y)$. This implies that

$$(21) \quad \{y \in S : U_{\beta,0}(x) = (T_1^{n_1} \circ T_0)(y)\} = (T_1^{n_1} \circ T_0)^{-1}(S).$$

It remains to show that equation (20) holds in the general case. Obviously

$$(22) \quad \{y \in S : U_{\beta,0}(y) = (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)(y)\} \subseteq (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)^{-1}(S).$$

So we have to show that the opposite inclusion holds, for this we examine the formula for $U_{\beta,0}$ more closely. We assume $U_{\beta,0}(x) = (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)(x)$ for some $i \geq 2$. Since $i \geq 2$ we have $T_0(x)$ is contained in a connected component of $[1, \frac{1}{\beta-1}] \setminus \cup_{n=1}^{\infty} D_n$. Let us denote this interval by \mathcal{I}_1 . We also let

$$E := \left\{ T_0^{-n} \left(\frac{1}{\beta} \right), T_0^{-n} \left(\frac{1}{\beta(\beta-1)} \right), T_1^{-n} \left(\frac{1}{\beta} \right), T_1^{-n} \left(\frac{1}{\beta(\beta-1)} \right), G_{\beta}^n(1), G_{\beta}^n \left(\frac{1}{\beta-1} - 1 \right) : n \geq 0 \right\}.$$

Here G_{β} is the greedy map defined earlier. Since $\beta \in M$ no element of E is contained in the interior of a C_n , a D_n , or S .

Importantly $\mathcal{I}_1 = (a_1, b_1)$ where $a_1, b_1 \in E$. In this case either

$$(a_1, b_1) = \left(1, T_1^{-n_1} \left(\frac{1}{\beta} \right) \right) \text{ or } (a_1, b_1) = \left(T_1^{-(n_1-1)} \left(\frac{1}{\beta(\beta-1)} \right), T_1^{-n_1} \left(\frac{1}{\beta} \right) \right).$$

Therefore

$$T_1^k(\mathcal{I}_1) \cap S = \emptyset \text{ for } 1 \leq k \leq n_1 - 1 \text{ and } T_1^{n_1}(\mathcal{I}_1) \subseteq \left(\frac{2-\beta}{\beta-1}, \frac{1}{\beta} \right).$$

The endpoints of $T_1^{n_1}(\mathcal{I}_1)$ are elements of E and are therefore not contained in the interior of any C_n . Either $(T_1^{n_1} \circ T_0)(x) \in C_n$ for some n or maybe $(T_1^{n_1} \circ T_0)(x) \in T_1^{n_1}(\mathcal{I}_1) \setminus \cup_{n=1}^{\infty} C_n$. If $(T_1^{n_1} \circ T_0)(x) \in T_1^{n_1}(\mathcal{I}_1) \setminus \cup_{n=1}^{\infty} C_n$ then let the connected component it is contained in be denoted by \mathcal{I}_2 . Let $\mathcal{I}_2 = (a_2, b_2)$ then again $a_2, b_2 \in E$. In which case

$$(23) \quad T_0^k(\mathcal{I}_2) \cap S = \emptyset \text{ for } 1 \leq k \leq n_2 - 1 \text{ and } T_0^{n_2}(\mathcal{I}_2) \subseteq \left(\frac{1}{\beta(\beta-1)}, 1 \right)$$

The endpoints of $T_0^{n_2}(\mathcal{I}_2)$ are again contained in E and therefore do not intersect the interior of any D_n . The point x has either been mapped into a D_n or is contained in a connected component of $T_0^{n_2}(\mathcal{I}_2) \setminus \cup_{n=1}^{\infty} D_n$. If it is contained in a connected component of $T_0^{n_2}(\mathcal{I}_2) \setminus \cup_{n=1}^{\infty} D_n$ then we repeat the previous steps. Eventually x is mapped into either C_{n_i} or D_{n_i} and our algorithm terminates. Without loss of generality we assume x is eventually mapped into D_{n_i} . The above algorithm yields a finite sequence of intervals $(\mathcal{I}_j)_{j=1}^{i-1}$ which satisfy the following properties:

- (1) $\mathcal{I}_1 \subseteq T_0(S)$.
- (2) For $1 \leq j \leq i-1$

$$T_{\omega_j}^k(\mathcal{I}_{n_j}) \cap S = \emptyset \text{ for } 1 \leq k \leq n_j$$

- (3) For $1 \leq j \leq i-2$ we have $\mathcal{I}_{j+1} \subseteq T_{\omega_j}^{n_j}(\mathcal{I}_j)$
- (4)

$$D_{n_i} \subseteq T_{\omega_{i-1}}^{n_{i-1}}(\mathcal{I}_{i-1}).$$

Where in the above $\omega_j = 0$ if j is even and $\omega_j = 1$ if j is odd. These properties have the following consequences:

- (5) $(T_{\omega_i}^{n_i})^{-1}(S) \subseteq \mathcal{I}_{i-1}$
- (6) For $1 \leq j \leq i-1$

$$(T_{\omega_i}^{n_i} \circ \dots \circ T_{\omega_j}^{n_j})^{-1}(S) \cap S = \emptyset \text{ for } 1 \leq k \leq n_j.$$

- (7) For $1 \leq j \leq i-1$

$$(T_{\omega_i}^{n_i} \circ \dots \circ T_{\omega_j}^{n_j})^{-1}(S) \subseteq \mathcal{I}_{n_j}$$

- (8)

$$(T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)^{-1}(S) \subseteq S.$$

Property (8) states that $(T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)^{-1}(S) \subseteq S$. Moreover, properties (5), (6) and (7) imply that every $y \in (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)^{-1}(S)$ satisfies $U_{\beta,0}(y) = (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)(y)$. Thus

$$(T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)^{-1}(S) \subseteq \{y \in S : U_{\beta,0}(y) = (T_{\omega_i}^{n_i} \circ \dots \circ T_1^{n_1} \circ T_0)(y)\},$$

which when combined with equation (22) yields (20). □

REFERENCES

- [1] S. Baker, *Generalised golden ratios over integer alphabets*, Integers 14(2014), Paper No A15.
- [2] J. Barrionuevo, R. Burton, K. Dajani and C. Kraaikamp, *Ergodic properties of generalized Lüroth series*, Acta Arith. 74 (1996), no. 4, 311-327.
- [3] K. Dajani and C. Kraaikamp, *Random β -expansions*, Ergodic Theory Dynam. Systems 23 (2003), no. 2, 461-479.
- [4] K. Dajani and M. de Vries, *Measures of maximal entropy for random β -expansions*, J. Eur. Math. Soc. (JEMS) 7 (2005), no. 1, 51-68.
- [5] K. Dajani and M. de Vries, *Invariant densities for random β -expansions*, J. Eur. Math. Soc. (JEMS) 9 (2007), no. 1, 157-176.
- [6] Z. Daróczy and I. Katai, *Univoque sequences*, Publ. Math. Debrecen **42** (1993), 397-407.
- [7] M. de Vries and V. Komornik, *Unique expansions of real numbers*, Adv. Math. 221 (2009), 390-427.
- [8] P. Erdős, M. Horváth, I. Joó, *On the uniqueness of the expansions $1 = \sum_{i=1}^{\infty} q^{-n_i}$* , Acta Math. Hungar. 58 (1991), no. 3-4, 333-342.
- [9] P. Erdős and I. Joó, *On the number of expansions $1 = \sum q^{-n_i}$* , Ann. Univ. Sci. Budapest **35** (1992), 129-132.
- [10] V. Komornik, *Expansions in non-integer bases*, Integers 11B (2011), Paper No. A9, 30 pp. 11A63.
- [11] V. Komornik and P. Loreti, *Unique developments in noninteger bases*, Amer. Math. Monthly, 105 (1998), 636-639
- [12] W. Parry, *On the β -expansions of real numbers*, Acta Math. Acad. Sci. Hung. **11** (1960) 401-416.
- [13] A. Rényi, *Representations for real numbers and their ergodic properties*, Acta Math. Acad. Sci. Hung. **8** (1957) 477-493.
- [14] N. Sidorov, *Almost every number has a continuum of β -expansions*, Amer. Math. Monthly 110 (2003), no. 9, 838-842.
- [15] N. Sidorov, *Arithmetic dynamics*, Topics in dynamics and ergodic theory, 145-189, London Math. Soc. Lecture Note Ser., 310, Cambridge Univ. Press, Cambridge, 2003.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF READING, READING, RG6 6AX, UK. E-MAIL: SIMONBAKER412@GMAIL.COM

DEPARTMENT OF MATHEMATICS, UTRECHT UNIVERSITY, 3508 TA UTRECHT, THE NETHERLANDS. E-MAIL: K.DAJANI1@UU.NL