

Original citation:

Hyseni, L., Elliot-Green, A., Lloyd-Williams, F., Kypridemos, C., O'Flaherty, M., McGill, R., Orton, L., Bromley, H., Cappuccio, Francesco and Capewell, S.. (2017) Systematic review of dietary salt reduction policies : evidence for an effectiveness hierarchy? PLoS One, 12 (5). e0177535.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/88526>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 4.0 International license (CC BY 4.0) and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/4.0/>

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk


RESEARCH ARTICLE

Systematic review of dietary salt reduction policies: Evidence for an effectiveness hierarchy?

Lirije Hyseni^{1*}, Alex Elliot-Green¹, Ffion Lloyd-Williams¹, Chris Kypridemos¹, Martin O'Flaherty¹, Rory McGill¹, Lois Orton¹, Helen Bromley¹, Francesco P. Cappuccio², Simon Capewell¹

1 Department of Public Health and Policy, Institute of Psychology, Health and Society, University of Liverpool, Liverpool, United Kingdom, **2** University of Warwick, WHO Collaborating Centre, Warwick Medical School, Coventry, United Kingdom

* L.hyseni@liv.ac.uk



Abstract

OPEN ACCESS

Citation: Hyseni L, Elliot-Green A, Lloyd-Williams F, Kypridemos C, O'Flaherty M, McGill R, et al. (2017) Systematic review of dietary salt reduction policies: Evidence for an effectiveness hierarchy? PLoS ONE 12(5): e0177535. <https://doi.org/10.1371/journal.pone.0177535>

Editor: Bhavani Shankar, SOAS, University of London, UNITED KINGDOM

Received: July 18, 2016

Accepted: April 29, 2017

Published: May 18, 2017

Copyright: © 2017 Hyseni et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: Lirije Hyseni, Rory McGill and Lois Orton were funded by LiLaC/NIHR. This article presents independent research funded by the National Institute for Health Research's School for Public Health Research (NIHR SPHR). The School for Public Health Research (SPHR) is funded by the National Institute for Health Research (NIHR). SPHR is a partnership between the Universities of

Background

Non-communicable disease (NCD) prevention strategies now prioritise four major risk factors: food, tobacco, alcohol and physical activity. Dietary salt intake remains much higher than recommended, increasing blood pressure, cardiovascular disease and stomach cancer. Substantial reductions in salt intake are therefore urgently needed. However, the debate continues about the most effective approaches. To inform future prevention programmes, we systematically reviewed the evidence on the effectiveness of possible salt reduction interventions. We further compared “downstream, agentic” approaches targeting individuals with “upstream, structural” policy-based population strategies.

Methods

We searched six electronic databases (CDSR, CRD, MEDLINE, SCI, SCOPUS and the Campbell Library) using a pre-piloted search strategy focussing on the effectiveness of population interventions to reduce salt intake. Retrieved papers were independently screened, appraised and graded for quality by two researchers. To facilitate comparisons between the interventions, the extracted data were categorised using nine stages along the agentic/structural continuum, from “downstream”: dietary counselling (for individuals, worksites or communities), through media campaigns, nutrition labelling, voluntary and mandatory reformulation, to the most “upstream” regulatory and fiscal interventions, and comprehensive strategies involving multiple components.

Results

After screening 2,526 candidate papers, 70 were included in this systematic review (49 empirical studies and 21 modelling studies). Some papers described several interventions. Quality was variable. Multi-component strategies involving both upstream and downstream interventions, generally achieved the biggest reductions in salt consumption across an

Sheffield, Bristol, Cambridge, Exeter, UCL; The London School for Hygiene and Tropical Medicine; the LiLaC collaboration between the Universities of Liverpool and Lancaster and Fuse; The Centre for Translational Research in Public Health. The work was carried out under the remit of the WHO Collaborating Centre for Nutrition of the University of Warwick. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, the Department of Health or the World Health Organization. All other authors were funded by HEFCE. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

entire population, most notably 4g/day in Finland and Japan, 3g/day in Turkey and 1.3g/day recently in the UK. Mandatory reformulation alone could achieve a reduction of approximately 1.45g/day (three separate studies), followed by voluntary reformulation (-0.8g/day), school interventions (-0.7g/day), short term dietary advice (-0.6g/day) and nutrition labelling (-0.4g/day), but each with a wide range. Tax and community based counselling could, each typically reduce salt intake by 0.3g/day, whilst even smaller population benefits were derived from health education media campaigns (-0.1g/day). Worksite interventions achieved an increase in intake (+0.5g/day), however, with a very wide range. Long term dietary advice could achieve a -2g/day reduction under optimal research trial conditions; however, smaller reductions might be anticipated in unselected individuals.

Conclusions

Comprehensive strategies involving multiple components (reformulation, food labelling and media campaigns) and “upstream” population-wide policies such as mandatory reformulation generally appear to achieve larger reductions in population-wide salt consumption than “downstream”, individually focussed interventions. This ‘effectiveness hierarchy’ might deserve greater emphasis in future NCD prevention strategies.

Introduction

Non-communicable diseases (NCDs) kill over 35 million people annually. Common cancers, cardiovascular diseases, diabetes, respiratory diseases and dementia together now account for over two thirds of the entire global burden of disability and death.[1,2] These NCDs are mainly attributable to just four major risk factors. Furthermore, the contribution from poor diet exceeds the combined contribution from alcohol, tobacco and physical inactivity.[3] This poor diet mainly reflects a predominantly unhealthy global food environment, dominated by processed foods high in sugar, saturated fat, trans-fat and, crucially, salt.[3]

In the UK and other high income countries, over 70% of dietary salt is consumed in processed foods such as bread, breakfast cereals, processed meats, snack foods, soups and sauces.[4–6] This food environment contributes to excessive salt intake among adults, on average 10g/day or more,[7] far in excess of what the body actually needs.[8] High salt intake is a major risk factor for increasing blood pressure,[9–11] cardiovascular disease,[12–14] stroke,[15,16] and stomach cancer.[17–19] Moreover, a reduction in salt intake would substantially reduce this risk.[10]

WHO recommends a maximum adult salt intake of 5g/day.[20] Different strategies and policy options have been proposed to achieve this goal. Individual level interventions often involve behavioural approaches, for example dietary counselling, leaflets or medical advice. These are sometimes termed “downstream” or “agentic” interventions, and are dependent on the individual responding. [21,22] Conversely, “upstream” structural interventions take place at the population level and typically involve policies such as regulatory approaches, taxes or subsidies. Finally, intermediate interventions target subgroups in worksites, schools or communities.[23]

National salt reduction strategies were identified in 75 countries in 2015, a substantial increase from 32 in 2010.[24] However, the debate regarding the most effective and acceptable salt reduction strategy continues.

Notable policy approaches have been seen in Finland,[25] Japan,[26] and more recently, the United Kingdom.[27] In the UK, a combination of awareness campaigns, agreed target settings, voluntary reformulation from industry and population monitoring of salt consumption have led to a 1.4g per day reduction in population salt intake between 2001 and 2011 (the campaign started in 2003).[27] However, health inequalities in salt consumption have persisted.[28,29] Furthermore, the introduction of the UK Responsibility Deal in 2010 shifted emphasis to ‘downstream’ interventions, coupled with ineffective voluntary agreements and, controversially, the direct involvement of the industry in policy decisions.[30,31]

Geoffrey Rose famously advocated population wide approaches rather than targeting high-risk individuals.[32] Furthermore, there seems to be some evidence for a public health ‘effectiveness hierarchy’ whereby “upstream” structural interventions consistently achieve larger improvements in population health, are more equitable and often reduce health inequalities [33,34] compared to “downstream” agentic interventions targeting individuals, for instance in tobacco control and alcohol policies.[35,36] Emerging evidence suggests that a comparable effectiveness hierarchy might also exist for salt reduction strategies, whereby upstream interventions apparently achieve bigger reductions in salt intake.[37,38]. To test this hypothesis and hence inform future preventive health strategies, we have systematically reviewed the evidence for studies focusing on the effectiveness of salt interventions to reduce salt intake.

Methods

Study design

We conducted a systematic review of interventions intended to decrease population dietary salt intake. To ensure proper conduct, we adhered to the PRISMA checklist (Preferred Reporting Items for Systematic Reviews and Meta-Analyses)(S1 Table).[39] We used a narrative synthesis and formally investigated evidence to support or refute an effectiveness hierarchy. The research protocol can be found in S1 File.

Search strategy

We first identified exemplar studies to define and refine search terms needed for targeted searches. The search strategy consisted of a combination of four sets of key words:

1) salt, sodium; 2) health promotion, nutrition education, campaigns, dietary counselling, regulation, legislation, tax, self-regulation, reformulation, social marketing, promotion, provision, labelling, marketing control, primary care advice, food industry; 3) public policy, health policy, nutrition policy, policies, interventions, strategies, initiatives, programmes, policy option, actions; and 4) effectiveness, effect, intake, consumption, reduction, cost-benefit analysis, and cardiovascular diseases.

A pilot search was conducted to determine appropriate databases, identify relevant studies and highlight potential issues to be addressed. This process identified six databases which were then used for the targeted searches: Ovid MEDLINE, Science Citation Index, SCOPUS, Cochrane Database of Systematic Reviews, The Campbell Collaboration Library of Systematic Reviews and the CRD Wider Public Health database. We searched for all studies published in the last four decades (from 1975 onwards). The final searches were conducted on 30 October 2015. All papers identified by the searches were imported into the Zotero data management programme to identify duplicates and help screen titles, abstracts and full texts as appropriate. The reference lists of included studies were scanned for potential additional papers and topic experts (FPC and SC) were also consulted for additional data sources.[40,41]

Study selection and inclusion criteria

Studies were included if they investigated the effectiveness of specific interventions on population dietary salt intake and contained quantitative outcomes. Only studies in English were included. We included a wide range of study designs including meta-analyses, trials, observational studies and natural experiments. Empirical studies and modelling studies were analysed separately, in view of their profound differences. The retrieved studies were assessed using the PICOS approach (Participants, Interventions, Comparators, Outcomes and Study design), summarised in Table 1. The primary outcome was salt intake (g/day). Studies reporting urinary sodium excretion (mmol/day) or sodium mg/day were converted to g/day. Where necessary, we simultaneously considered studies reporting solely on salt intake data in a specific population with the corresponding studies describing the interventions during that same time period.

One reviewer (LH) conducted the searches; extracted potential papers and removed duplicates. Two reviewers (LH and AEG) then independently screened titles and abstracts for eligibility using the inclusion and exclusion criteria. Full text was retrieved for all papers deemed potentially eligible and these were also screened independently by the two reviewers. Any discrepancies were resolved by consensus or by involving the senior author (SC).

Data extraction and management

Pre-designed and pre-piloted tables were used to extract data from all included studies. To ensure that all relevant information was captured, extracted data included: first author; year of publication; funder(s); study aim(s); sample size; study design; methods; participants; policies analysed; geographical scope; length of follow-up; outcomes, effect and response;

Table 1. PICOS; Inclusion/exclusion criteria.

Participants	
Include	Exclude
Studies for all age groups from all populations, from high-, middle- and low-income countries	Studies on animals, cells and pregnant women
Interventions	
Systematic Reviews and primary studies evaluating the effects of actions to promote salt reduction by government policy or adopted in specific real or experimental settings	Studies evaluating the effect of a general or specific diet
Comparators	
Systematic and non-systematic reviews where actions to promote salt reduction were evaluated or compared	No comparisons of different actions to promote salt reduction presented
Outcomes	
Primary outcome of interest was dietary salt intake (g/day). Studies including urinary sodium excretion as an outcome were converted to g/day. Secondary outcomes included changes in clinical/physiological indicators related to NCDs and behaviours associated with a healthy diet	Process evaluations reporting on implementation of interventions/policies without any quantitative outcome data; feasibility or acceptability without an assessment or primary outcomes (intake); studies on individuals as opposed to populations; data on cost only and BMI
Study design	
Primary studies, RCTs, Systematic Reviews (SRs), empirical observational studies, natural experiments, and modelling studies, secondary analysis, and before vs. after interventions	Commentary/opinion articles and purely qualitative evaluations with no quantitative assessment

<https://doi.org/10.1371/journal.pone.0177535.t001>

authors' assessment of limitations and our own assessment of potential risk of bias. The sources referenced for the effect sizes used in each modelling study were also specified in the tables (recognising that some modelling studies are based on empirical studies, potentially some included in this review). This data extraction was done independently by two reviewers (LH and AEG).

Quality assessment of included studies

Two reviewers (LH and AEG) independently assessed the methodological quality of each study (poor, fair or good). We used the National Heart, Lung and Blood Institute (NHLBI) tools specific for each research design (i.e. RCTs, cross-sectional studies, before and after studies, and systematic reviews).[42] Several questions were asked for each study design (varying from 8 to 14) and depending on the points scored, the studies were labelled as good, fair or poor. However, we also took into consideration as to which questions points were allocated. For example, if an RCT scored 10 out of 14 points, but did not conduct an intention to treat analysis, it would be rated as fair rather than good. Modelling studies were independently assessed by two modelling experts (MOF & CK) using a different tool adapted from Fattore et al. (2014).[43] Discrepancies in quality assessment were reconciled by consensus or by involving a third, senior member of the team (SC or HB).

Data synthesis and effectiveness hierarchy continuum

The evidence was summarised as a narrative synthesis according to intervention type, ranging from downstream to upstream interventions, to facilitate comparisons between the interventions. Summary tables of the studies included in this review can be found in Tables 2–10 for empirical studies and Table 11 for modelling studies. A more detailed data extraction of these studies can be found in S2 Table. We defined UPSTREAM interventions as those targeting the entire population (not a subset, however large) and creating structural changes (effectively removing individual choice from the equation). This accorded with the Nuffield's ladder taxonomy,[44] and with McLaren's structural/agentive continuum.[21] Conversely, we defined DOWNSTREAM interventions as those where the principal mechanism of action is "agentive", being dependent on an individual altering their behaviour.

Interventions were then categorised according to their position in the McLaren et al. (2010) continuum from "upstream" to "downstream" (Fig 1).[21]

Multi-component interventions were considered separately.

Patient involvement

Individual patients were not involved in this research; this is a secondary analysis of published data.

Results

The literature search identified 3336 potentially relevant papers. An additional 26 papers were identified through other sources, including reference lists and key informants. After removing 836 duplicates, 2526 publications were left to be screened by title and abstract, after which 134 full-text papers were assessed for eligibility. A total of 70 papers were finally included (49 empirical studies and 21 modelling studies, Fig 2). The interventions and their effect sizes are presented in Fig 3 (empirical studies) and Fig 4 (modelling studies).

Table 2. Dietary counselling (individuals).

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Hooper et al. (2002) ⁴⁵	SR and meta-analysis of RCTs	US, Australia, New Zealand, UK	<i>Aim:</i> to assess the long term effects of advice to restrict dietary sodium in adults with and without hypertension. <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	Dietary advice	<i>Meta-analysis (11 studies included).</i> They found reductions in salt intake at both intermediate, <12 months (2.8g/day) and late follow up, 13–60 months (2.0g/day).	Good
Appel et al. (2003) ⁴⁶	Randomised trial	US	<i>Aim:</i> to determine the effect on BP of 2 multicomponent, behavioral interventions <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	Dietary advice	Only the reduction in the established group differed significantly from that of advice only group. 24-hour dietary recall data indicated both behavioral interventions significantly reduced sodium intake in comparison with advice only group (P value = 0.01). Advice group <ul style="list-style-type: none"> • Baseline = 10.0g/day • 6 months = 8.8g/day • Mean difference = -1.2g/day Intervention group <ul style="list-style-type: none"> • Established: mean difference = -1.82 g/day • Established + DASH: mean difference = -1.83 g/day 	Good
Brunner et al. (1997) ⁴⁷	Meta-analysis of RCTs	UK, US, Netherlands and Australia	<i>Aim:</i> to evaluate the effectiveness of dietary advice in primary prevention of chronic disease. <i>Outcomes:</i> salt intake	Dietary advice	Overall mean net reduction of 1.8g/day which is a 20% reduction in salt intake. The heterogeneity test was highly significant (P < .0005) for the 3- to 6-month trials, with a net reduction of 3.4 (95% CI = 45, 72) g/day. Summary effect of the two trials with SE was somewhat larger at 9–18 months than at 3–6 months.	Fair
Francis & Taylor (2009) ⁴⁸	Randomised control group study	US	<i>Aim:</i> to implement a health-healthy diet-education programme. <i>Outcomes:</i> salt intake	Dietary counselling	Intervention salt consumption decreased significantly (P0.020) from record 1 to record 3. The reduction in control group participants' sodium intake was not significant <i>Intervention:</i> (Mean ± SEM (g/day); P-value) <ul style="list-style-type: none"> • Record 1: 7.0 ± 0.5; 0.020e • Record 2: 5.9 ± 0.3; 0.067 • Record 3: 5.9 ± 0.4; 0.937 <i>Control</i> (Mean ± SEM (g/day), P-value) <ul style="list-style-type: none"> • Record 1: 6.2 ± 0.5; 0.323 • Record 2: 6.1 ± 0.4; 0.880 • Record 3: 5.7 ± 0.4; 0.284 Mean effect size:- 0.6g/day	Fair
Parekh et al. (2012) ⁴⁹	RCT	Australia	<i>Aim:</i> to evaluate the effectiveness of a minimal intervention on multiple lifestyle factors including diet using computer tailored feedback. <i>Outcomes:</i> salt intake (%)	Health promotion—computer tailored advice	<i>Salt (%)</i> Intervention +5.43 net change. Control +1.23 net change. Significant changes between groups were observed for reduced salt intake (OR 1.19, CI 1.05–1.38). The intervention group were 20% more likely to reduce salt intake	Fair

(Continued)

Table 2. (Continued)

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Petersen et al. (2013) ⁵⁰	RCT	Australia	<i>Aim:</i> to investigate whether urinary sodium excretion can be reduced by educating people with T2DM to read food labels and choose low sodium products. <i>Outcomes:</i> salt intake	Nutrition education	Baseline reported salt intake: 6.8 ± 3.2 g/day Intervention • Baseline: 10.0 ± 0.7 • 3 months: 10.1 ± 0.7 • Change: +0.06 ± 0.9 Control • Baseline: 9.6 ± 0.9 • 3 months: 9.3 ± 0.7 • Change: -0.3 ± 0.8 There was no between group difference (p > 0.05)	Fair
Kokanović et al. (2014) ⁵¹	Before and after study	Croatia	<i>Aim:</i> to assess eating habits of adolescent population diagnosed with one or more cardiovascular risks before and after two months of individual dietary intervention <i>Outcomes:</i> salt intake	Nutrition education	Difference in intake on initial and control examination statistically significant for intake of sodium p = 0.013. <i>Salt intake g/day.</i> Initial examination: 18.9g/day; Control examination: 15.4g/day; Difference: -3.5g/day (= -18.8%)	Fair
Heino et al. (2000) ⁵²	Prospective randomized trial	Finland	<i>Aim:</i> to examine sodium intake of 1-5-y-old children in a CHD prevention trial, focused on dietary fat modification. <i>Outcomes:</i> salt intake	Dietary counselling	Intervention children (+1.5g/day) • 13 months: 4.1 ± 1.2 • 3 years: 4.9 ± 1.2 • 5 years: 5.6 ± 1.3 Control children (+1.6g/day) • 13 months: 3.9 ± 1.4 • 3 years: 4.7 ± 1.3 • 5 years: 5.5 ± 1.4 No significant differences between the intervention and control group found	Poor
Wang et al. (2013) ⁵³	RCT	US	<i>Aim:</i> one year dietary intervention study to examine patterns and amount of daily sodium intake among participants with metabolic syndrome <i>Outcomes:</i> salt intake	Dietary counselling	Intervention arm at one year follow-up found participants who consumed sodium greater than 5.8g/day declined from 75% at baseline to 59%. Those consumed higher than 3.8g/day declined from 96% (at baseline) to 85%. Average salt intake decreased from 7.5 g/day at baseline to 6.4 g/day at one-year (P<0.001). At one-year visit, salt intake was consistently reduced; significant difference only observed between males (7.6±0.4 g/day) and females (6.0 ± 0.2 g/day; p < 0.001)	Poor

<https://doi.org/10.1371/journal.pone.0177535.t002>

Dietary counselling—individual level (Table 2)

Nine empirical studies (two of good quality;[45–46] five of fair quality;[47–51] and two of poor quality [52–53]), and three modelling studies (all of good quality [54–56]) investigated the effect on salt intake of dietary counselling targeted at consenting individuals.

Two separate meta-analyses investigated the effect of dietary advice on salt intake. The first included eleven randomised controlled trials (RCTs) and found a 1.8g/day salt reduction after up to 18 months of dietary advice.[47] The second meta-analysis included eight RCTs and

Table 3. Dietary counselling (worksite/schools).

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
He et al. (2015) ⁴⁰	Cluster RCT	China	<i>Aim:</i> to determine whether an education programme targeted at schoolchildren could lower salt intake in children and their families <i>Outcomes:</i> salt intake as measured by urinary excretion	Health education	At baseline, the mean salt intake in children was 7.3 (SE 0.3) g/day in the intervention group and 6.8 (SE 0.3) g/day in the control group. The mean effect on salt intake for intervention versus control group was -1.9 g/day (95% confidence interval -2.6 to -1.3 g/day; $P < 0.001$). In adult family members the salt intakes were 12.6 (SE 0.4) and 11.3 (SE 0.4) g/day, respectively. During the study there was a reduction in salt intake in the intervention group, whereas in the control group salt intake increased. The mean effect on salt intake for intervention versus control group was -2.9 g/day (-3.7 to -2.2 g/day; $P < 0.001$)	Good
Cotter et al. (2013) ⁵⁷	School based RCT	Portugal	<i>Aim:</i> to examine the influence on salt intake and blood pressure of three different educational interventions for 6 months <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	Nutrition education	<i>Baseline:</i> mean salt intake of 7.8 ± 2.5 g per day. Estimated salt intake (g/d): CRT • Baseline: 7.7 ± 2.0 • Final: 7.4 ± 3.0 • Change: 0.35 ± 2.42 THEOR • Baseline: 8.1 ± 3.0 • Final: 7.5 ± 3.0 • Change: 0.60 ± 3.24 PRACT • Baseline: 7.5 ± 2.4 • Final: 6.4 ± 2.2 • Change: $1.08 \pm 2.47^*$	Fair
Katz et al. (2011) ⁵⁸	School based RCT	US	<i>Aim:</i> to evaluate the effects of a nutrition education programme in distinguishing between healthful and less healthful choices in diverse food categories. <i>Outcomes:</i> salt intake	Nutrition education	There were no statistically significant improvements in dietary patterns from baseline between the intervention (-0.23g/day) and control groups (-0.04g/day) for salt intake ($p = .44$)	Poor
Aldana et al. (2005) ⁵⁹	RCT	US	<i>Aim:</i> to determine behavioral and clinical impact of a worksite chronic disease prevention program <i>Outcomes:</i> salt intake	Health education	<i>Intervention group (salt g/day)</i> • Baseline: 7.5 • $\Delta 6$ weeks: -0.5 • $\Delta 6$ months: -1.7 <i>Control group (salt g/day)</i> • Baseline: 6.3 • $\Delta 6$ weeks: -0.5 • $\Delta 6$ months: -0.5 Significant differences in mean change scores were not observed at 6 weeks ($P = 0.88$) but they were seen at 6 months ($P = 0.0097$)	Fair
Chen et al. (2008) ⁶⁰	Intervention control trial	China	<i>Aim:</i> to report the effects of these two programmes on blood pressure and changes in morbidity and mortality from CHD and stroke <i>Outcomes:</i> salt intake	Health education	Mean daily salt intake declined from 16.0 to 10.6 g d ⁻¹ in the intervention factory, compared with the control factory from 16.9 to 15.4 g d ⁻¹ , with the net reduction of 3.9 g d ⁻¹ , which was significantly different ($P < 0.05$).	Fair

(Continued)

Table 3. (Continued)

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Levin et al. (2009) ⁶¹	Worksite based dietary intervention	US	<i>Aim:</i> to examine whether a worksite nutrition programme using a low-fat vegan diet could significantly improve nutritional intake <i>Outcomes:</i> salt intake	Dietary counselling	Intervention group participants significantly increased the reported intake and mean intake ($P = 0.04$) of salt compared to the control group. <i>Salt (g/day)</i> Intervention group • Baseline: 4.1 ± 0.1 • 22 weeks: 5.0 ± 0.2 • Mean difference: 0.9 ± 0.2 Control group • Baseline: 4.5 ± 0.2 • 22 weeks: 4.9 ± 0.2 • Mean difference: 0.4 ± 0.2 Mean effect size: $+0.5$ (95% CI 9.2, 394.4; $P = 0.04$)	Fair

<https://doi.org/10.1371/journal.pone.0177535.t003>

reported an overall reduction in salt consumption of 2.8g/day at 12 months and 2g/day up to 60 months.[45] The two meta-analyses overlapped in respect of only three studies.

One additional RCT found a statistically significant net reduction of 0.6g/day between the groups,[48] whilst a second RCT found no effect between the control and intervention group.[50]

All three modelling studies predicted that dietary advice is less effective in reducing the disease burden of high salt intake, only gaining 180–2,600 quality-adjusted life years (QALYs) compared to other interventions (7,900–195,000 QALYs).[54–56]

Dietary counselling—school based and worksite interventions (Table 3)

Three school-based interventions (one of good quality;[40] one of fair quality;[57] one of poor quality [58]) and three worksite-based studies (all of fair quality) were included.[59–61] No modelling studies were identified for this section.

Schools. A nutrition programme in schools aimed at distinguishing between healthy and less healthy choices reported a non-significant reduction.[58] In the second school based RCT, the practical intervention group achieved a significant net reduction of 0.7g/day compared with the control group.[57] In a cluster RCT in China, education and training significantly reduced salt intake by a mean of -1.9 g/day in 279 school children (and -2.9 g/day in adult family members).[40]

Worksites. A randomised trial of a chronic disease prevention programme achieved a net reduction of 1.2g/day between the intervention and control group ($P = 0.01$).[59] A factory-based intervention study in China assessed health education aimed at altering diet, together with a high-risk strategy of hypertension control. Salt intake was reduced by 3.9g/day from a mean of 16g/day ($P < 0.05$).[60]

Dietary counselling—community level (Table 4)

Four empirical studies and one review, all of fair quality,[62–66] investigated community based dietary counselling. One study reported a statistically significant difference of -0.4 g/day in salt intake between the intervention and control groups.[62] Two intervention trials of nutrition education reported significant reductions of 0.7g/day and 2.2g/day reductions

Table 4. Dietary counselling (community).

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Yanek et al. (2001) ⁶²	RCT	US	<i>Aim:</i> to test the impact on cardiovascular risk profiles after one year of participation in one of three church-based nutrition and physical activity strategies <i>Outcomes:</i> salt intake	Health promotion–education	Salt (g/day) <i>Combined standard and spiritual intervention groups</i> • Baseline: 6.7 ±2.5 • Change: -0.4 ±0.06 <i>Self-help control group</i> • Baseline: 7.4 ±3.0 • Change: -0.02 ±0.09 Between group P value = 0.0167	Fair
Cappuccio et al. (2006) ⁶³	Community-based cluster randomised trial	Ghana	<i>Aim:</i> to establish the feasibility of salt reduction as a way of reducing BP <i>Outcomes:</i> salt intake	Health education	Sodium intake as measured by sodium excretion fell in four out of six villages in the intervention group and in 5 out of six villages in the control group. The net intervention effect was non-significant. <i>Control</i> Baseline: 6.0 g/day 3 months: 5.6 g/day 6 months: 5.2 g/day <i>Intervention</i> Baseline: 5.8 g/day 3 months: 5.4 g/day 6 months: 5.3 g/day	Fair
Takahashi et al. (2006) ⁶⁴	Community based open randomizer controlled cross-over trial	Japan	<i>Aim:</i> to assess whether dietary intervention in free-living healthy subjects is effective in improving blood pressure levels. <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	Dietary education	Salt intake as measured by sodium excretion, collected at two points, in the intervention group decreased by 2.8 (95% CI: -3.6, -2.1) and 0.6 g/day (-1.4, +0.2) in the control group. This difference in change between the two groups was statistically significant (P < 0.001). Dietary counselling for 1 year reduced salt intake by 2.2 g/day as measured by 24-h urinary sodium	Fair
Robare et al. (2010) ⁶⁵	Community based intervention trial	US	<i>Aim:</i> to evaluate a dietary Na reduction trial in a community setting <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	Nutrition education	Salt intake decreased by 0.3g/day (7.8 to 7.5g/day) from baseline to 6 months follow up which was not significant (p = 0.30). When comparing baseline with 12 months follow up, salt intake decreased by 0.7g/day (7.8 to 7.2g/day) which was significant (p = 0.03)	Fair
Van de Vijver et al. (2012) ⁶⁶	Review	Ghana and China	<i>Aim:</i> to evaluate the effectiveness of the community-based interventions for CVD prevention programmes in LMIC <i>Outcomes:</i> BP and salt intake (g/day and n, %)	Health education	Cappuccio et al. (2006) • BP: reduction SBP 2.5 mmHg (1.45 to 6.54), DBP 3.9 mmHg (0.78 7.11)* vs control • Salt: no significant reduction in salt intake vs control Chen, Wu, and Gu (2008) (urban) • BP: reduction SBP 1.9 mmHg, reduction DBP 2.2 mmHg* vs control • Salt: reduction in salt intake of 3.9 g/day* vs control Yu et al. (1999) • BP: reduction among men in prevalence in HT 2%,* SBP 0%, among women prevalence of HT 2%,* SBP 2 mmHg • Salt: reduction in salt intake 6.0% Huang et al. (2011) • BP: reduction prevalence HT 12.9%* pre vs post • Salt: reduction in salt intake 30%* (n, %)	Fair

<https://doi.org/10.1371/journal.pone.0177535.t004>

respectively in salt intake after 12 months.[63–64] One RCT reported a favourable trend; however, this was non-significant and could have been caused by contamination between the groups.[63]

Mass media campaigns (Table 5)

One empirical study of fair quality [67] and five modelling studies; four of good quality[56, 68–70] and one of fair quality[71] were included.

Table 5. Media campaigns.

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Shankar et al. (2012) ⁶⁷	Cross-sectional	UK	<i>Aim:</i> to examine the trend in salt intake over a set period and deduce the effects of the policy on the intake of socio-demographic groups <i>Outcomes:</i> salt intake as measured by spot urinary sodium readings	Salt campaign (and potential effect on reformulation and table salt use)	The results are consistent with a previous hypothesis that the campaign reduced salt intakes by approximately 10%. The impact is shown to be stronger among women than among men. Salt as measured by spot urinary sodium readings <ul style="list-style-type: none"> • 2003: 6.3 g/day • 2004: 6.4 g/day • 2005: 5.7 g/day • 2006: 5.6 g/day • 2007: 5.4 g/day Difference in g/day between 2003–2007 = 0.9 g/day = 13.5%	Fair

<https://doi.org/10.1371/journal.pone.0177535.t005>

The UK FSA salt reduction programme involved media campaigns to discourage table salt use, plus sustained pressure on industry to reformulate. Although salt consumption declined by 0.9g/day using spot urinary sodium readings from 2003–2007, the media contribution was unclear but likely modest.[67]

The modelling studies likewise suggested media campaigns were generally considered less effective than food labelling or reformulation.[56, 69–71] The Change4Life campaign in the UK was predicted to reduce salt intake by 0.16g/day, less than labelling or reformulation.[68] Gillespie et al. (2015) similarly estimated that social marketing might modestly reduce salt consumption by 0.03g/day to 0.13g/day.[69]

Nutrition labelling (Table 6)

Two empirical studies, both of poor quality, investigated the effect of nutrition labelling on salt intake [72–73]. Reduced salt intake was not observed in participants who reported frequent vs. non-frequent label use (7.7g/day vs. 7.6g/day).[73]

Ten modelling studies also examined labelling, four of good quality[56,68–70] and two of fair quality.[71, 74–77] These suggested that labelling might modestly reduce UK salt intake by

Table 6. Labelling.

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Babio et al. (2013) ⁷²	Randomised cross-over trial	Spain	<i>Aim:</i> to compare two models of front-of-pack guideline daily amounts (GDA) and the ability to choose a diet that follows the nutritional recommendations. <i>Outcomes:</i> salt intake based on choices	Labelling	Participants using the multiple-traffic-light GDA system chose significantly less salt (0.4g/day; P <0.001) than those using the monochrome GDA labels	Poor
Elfassy et al. (2015) ⁷³	Cross-sectional	US	<i>Aim:</i> to examine independent association between hypertension and frequency use of NF label for sodium information and whether this was associated with differences in intake <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	Labelling (use)	Daily sodium intake was not lower in those who reported frequent vs non-frequent use of the NF label for sodium information (7.7g/day vs 7.6g/day; P = 0.924)	Poor

<https://doi.org/10.1371/journal.pone.0177535.t006>

Table 7. Reformulation.

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Chang et al. (2006) ⁷⁸	Cluster RCT	Taiwan	<i>Aim:</i> to examine the effects of potassium-enriched salt on CVD mortality and medical expenditures in elderly veterans. <i>Outcomes:</i> incidence, CVD mortality, LYG	Reformulation—low sodium salt	The incidence of CVD-related deaths was 13.1 per 1000 persons (27 deaths in 2057 person-years) and 20.5 per 1000 (66 deaths in 3218 person years) for the experimental and control groups, respectively. A significant reduction in CVD mortality (age-adjusted hazard ratio: 0.59; 95% CI: 0.37, 0.95) was observed in the experimental group. Persons in the experimental group lived 0.3–0.90 y longer	Fair

<https://doi.org/10.1371/journal.pone.0177535.t007>

0.03g/day to 0.16g/day [68, 69]; much less than the 0.9g/day estimated by Roodenburg et al. (2013).[77] Another study suggested that salt intake might be lowered by 1.2g/day if the population were to choose products labelled as low-salt, or increased by 1.6g/day if they choose products labelled as high salt content.[74]

Reformulation (Table 7)

Very few studies which focused on reformulation included quantified results of salt intake. In one empirical Taiwanese study of fair quality,[78] salt was enriched with potassium in the intervention group and their outcomes were an apparent reduction in cardiovascular deaths by 41%, compared to the control group rather than salt intake. Furthermore, people in the intervention group lived 0.3–0.9 years longer.[78]

Fourteen modelling studies evaluated reformulation, eleven of good quality[41, 54–56, 68–70, 79–82] and three of fair quality[71, 83, 84]. Mandatory reformulation could consistently achieve bigger salt reductions than voluntary reformulation; 1.6g/day compared with 1.2g/day; [68] and 1.4g/day versus 0.5g/day.[69] Mandatory reformulation might also achieve higher reductions in disability-adjusted life years (DALYs) and QALYs compared to voluntary reformulation.[54, 56, 79]

Table 8. Taxes.

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Thow et al. (2014) ⁸⁵	Systematic Review	US (with UK data)	<i>Aim:</i> to assess the effect of food taxes on consumption <i>Outcomes:</i> sodium consumption	Sodium tax	A modelling study predicted that a sodium tax increasing the price of salty foods by 40% would reduce sodium consumption by 6%	Fair
Niebylski et al. (2015) ⁸⁶	Systematic Review	France and US	<i>Aim:</i> to evaluate the evidence base to assess the effect of unhealthy food taxation. <i>Outcomes:</i> energy intake	1) Tax on salty snacks 2) Tax on cheese/butter	1) Modelling study of tax on chips/salty snacks on energy intake in US. Predicted a 1% tax had no effect on consumption or body weight 2) Modelling study of effect of 1% VAT on cheese/butter, sugar, and fat products along with ready-made meals in France. Predicted proposed taxes reduced saturated fat, cholesterol, sodium, and energy intake but suggest 1% is insufficient to have positive health effect.	Fair

<https://doi.org/10.1371/journal.pone.0177535.t008>

Table 9. Multi-component interventions.

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
He et al. (2014) ⁸⁸	Comprehensive analysis	UK	<i>Aim:</i> to analyse the UK salt reduction programme <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	1) Reformulation 2) Labelling 3) Health promotion campaigns	15% decrease, there have been a steady fall in salt intake at a rate of ~2% per year since the introduction of the salt reduction strategy. The 0.9g/day reduction in salt intake achieved by 2008 led to E 6000 fewer CVD deaths per year. • 2000–2001: salt intake = 9.5g/day • 2005–2006: salt intake = 9.0g/day • 2008: salt intake = 8.6g/day • 2011: salt intake = 8.1g/day	Good
Mozaffarian et al. (2012) ⁸⁹	Systematic review	Finland and China	<i>Aim:</i> to systematically review and grade the current scientific evidence for effective population approaches to improve dietary habits. <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	1) Education 2) Combined effects of labelling, reformulation and campaigns	Tian et al. (1995) 1) <i>Education:</i> In the intervention neighborhoods, mean sodium intake decreased by 1.3 and 0.6 mmol/day in men and women, respectively, compared with increases of 1.0 and 0.2 mmol/day, respectively, in the control neighborhoods (P0.001 for men, P0.065 for women) Pekka et al. (2002) + Puska & Stahl (2010) 2) From the 1970s to the late 1990s, mean daily salt consumption in Finland declined from approximately 14.5 g in men (unknown in women) to approximately 11 g in men and 7 g in women; mean diastolic blood pressure declined by 5% in men and 13% in women	Good
Fattore et al. (2014) ⁴³	Systematic review	Australia, US and Vietnam	<i>Aim:</i> to summarize and critically assess economic evaluation studies conducted on direct (e.g., counseling) or indirect (e.g., food labeling) interventions aimed at promoting voluntary dietary improvements through reduction of fat intake <i>Outcomes:</i> DALYs	1) Voluntary reformulation, mandatory reformulation and dietary advice 2) Reduction in daily caloric intake of 100 to 500 kcal below current estimated energy requirements 3) A set of personal (e.g., individual treatment of SBP >160 mmHg) and non-personal (e.g., a mass media campaign for reducing consumption of salt) prevention strategies to reduce CVD 4) Voluntary reformulation and sodium tax	1) Cobiac et al. (2010) 610,000 DALYs averted (95%CI: 480,000–740,000) if everyone reduced their salt intake to recommended limits. Dietary advice: <0.5% disease burden (IHD & stroke cases) averted; Tick program: <1%; making Tick limits mandatory: 18% 2) Dall et al. (2009) 400 mg/d sodium intake reduction 3) Ha & Chisholm (2011) A health education program to reduce salt intake (VND 1,945,002 or USD 118 per DALY averted) & individual treatment of SBP >160 mmHg (VND 1,281,596 or USD 78 per DALY averted) are the most cost-effective measures 4) Smith-Spangler (2010) (1) vs. (2): 1.25-mm Hg vs. 0.93-mm Hg decrease in mean SBP; 513,885 vs. 327,892 strokes averted; 480,358 vs. 306,137 MIs averted; 1.3 million vs. 840,113 years LE increase. Collaboration with industry: 2.1 million QALYs gained; USD 32.1 billion medical cost savings. Tax on sodium: 1.3 million QALYs gained; USD 22.4 billion medical cost savings	Fair
He & MacGregor (2009) ⁹⁰	Review	Japan, Finland and UK	<i>Aim:</i> to provide an update on the current experience of worldwide salt reduction programmes. <i>Outcomes:</i> salt intake, blood pressure, stroke & CHD mortality and life expectancy	1) Reformulation to reduce the salt content of all foods 2) Health promotion campaigns 3) Labelling to highlight salt content	<i>Japan.</i> The Japanese Government initiated a campaign to reduce salt intake. Over the following decade salt intake was reduced from an average of 13.5 to 12.1 g/day. However, in the north of Japan salt intake fell from 18 to 14 g/day. Paralleling this reduction in salt intake, there was an 80% reduction in stroke mortality despite large increases in population fat intake, cigarette smoking, alcohol consumption and an increase in BMI. <i>Finland.</i> Since the 1970s, Finland aimed to reduce salt intake by reformulation and raising general awareness of the harmful effects of salt on health. This led to a significant reduction in salt intake of 3g/day from 1979 to 2002 (12 to 9g/day) as measured by urinary sodium. This was accompanied by a fall of over 10mmHg in both systolic and diastolic BP, a pronounced decrease of 75–80% in both stroke and CHD mortality, and a remarkable increase of 5–6 years in life expectancy. UK. Salt added to cooking or at the table: estimated that 15% of the total 9.5g/day consumed was added (1.4g/day). Naturally present in food: approximately 5% (0.6g/day). Reformulation: 80% (7.5g/day) was added by the food industry. The UK salt reduction strategy started in 2003/2004 and the adult daily salt intake has already fallen, as measured by urinary sodium, from an average of 9.5 g/day to 8.6 g/day by May 2008	Fair
Pietinen et al. (2010) ⁹¹	Before and after study	Finland	<i>Aim:</i> to describe the main actions in Finnish nutrition policy during the past decades. <i>Outcomes:</i> salt intake	1) Education 2) Voluntary reformulation 3) Labelling	1981; Eastern Finland: salt intake was about 13 g in men and 11 g in women. Salt intake has decreased continuously to a level of about 9 g in men and 7 g in women in 2007	Fair

(Continued)

Table 9. (Continued)

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Wang et al. (2011) ⁹²	Literature review	US	<i>Aim:</i> to summarize cost-effectiveness evidence on selected interventions to reduce sodium intake that would be intended as population-wide approaches to control hypertension <i>Outcomes:</i> stroke and MI averted	1) Reformulation 2) Sodium tax	Smith-Spangler et al. For US adults aged 40–85 years, collaboration with industry that decreased mean intake of sodium by 9.5% was estimated to avert 513 885 strokes and 480 358 myocardial infarctions over their lifetimes and to save US\$ 32.1 billion in annual medical costs. Over the same period, a tax on sodium that decreased the population's intake of sodium by 6% was projected to save US\$ 22.4 billion in such costs	Fair
Webster et al. (2011) ⁹³	Review	Finland, France, Japan and UK	<i>Aim:</i> to provide an overview of national salt reduction initiatives around the world and describe core characteristic. <i>Outcomes:</i> salt intake, LYG, CHD and stroke mortality	1) Reformulation 2) Labelling 3) Health promotion campaigns	<i>Finland:</i> started salt reduction strategy in 1978 (reformulation, labelling and mass media campaigns) and by 2002 had demonstrated a 3 g reduction in average population salt intake (from 12 to 9 g/person per day). During the same period there was a corresponding 60% fall in CHD and stroke mortality <i>UK:</i> the Food Standards Agency (FSA) started working with the food industry in 2003 and launched its consumer education campaign in 2005. By 2008 the UK had achieved an average 0.9 g/person per day reduction in daily salt consumption, which is predicted to be saving some 6000 lives a year. <i>France:</i> the Food Safety Authority recommended a reduction in population salt consumption in 2000 and has since reported a decline in intake provided by foods from 8.1 to 7.7 g/day in the overall adult population. Focus was on bread reformulation and nutrition campaigns <i>Japan:</i> 60s started a salt campaign through a sustained public education campaign. Over the following decade average salt intake was reduced from 13.5 to 12.1 g/day with a parallel fall in blood pressure in adults and children, and an 80% reduction in stroke mortality despite large adverse changes in a range of other cardiovascular risk factors.	Fair
Wang & Bowman (2013) ⁹⁴	Literature review	US, UK	<i>Aim:</i> to summarize recent economic analyses of interventions to reduce sodium intake. <i>Outcomes:</i> SBP, hypertension, cardiovascular events	1) reducing the sodium content of all foods 2) reducing sodium content by labelling foods and by promoting, subsidizing, and providing low sodium food options 3) Legislation	<i>US (1&2):</i> If the sodium-reduction strategies were implemented, adults in the county would reduce their intake of sodium by 233 mg per day, on average, in 2010. This would correspond to an average decrease of 0.71 mmHg in SBP among adults with hypertension, 388 fewer cases of uncontrolled hypertension, and a decrease per year of \$629,724 in direct health care costs <i>UK (3):</i> Legislation or other measures to reduce the intake of salt by 3 g per person per day (in a population where the current mean intake was about 8.5 g per person per day) would reduce the mean population SBP by approximately 2.5 mmHg, prevent about 30,000 cardiovascular events and approximately 4,450 deaths, and produce discounted savings overall of approximately £347 million (about \$684 million) over a decade, which would be equivalent to annual savings of approximately £40 million	Fair
He et al. (2014) ⁹⁵	Cross-sectional	England	<i>Aim:</i> to determine the relationship between the reduction in salt intake that occurred in England, and BP, as well as mortality from stroke and IHD <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	Combined 1) Reformulation 2) Health promotion campaigns 3) Labelling	From 2003 to 2011, salt intake decreased by 1.4 g/day (15%, $p<0.05$ for the downward trend). From 2003 to 2011, stroke mortality decreased from 128/1 000 000 to 82/1 000 000 (36% reduction, $p<0.001$) and IHD mortality decreased from 423/1 000 000 to 272/1 000 000 (36% reduction, $p<0.001$). • 2003: 9.5g/day • 2005/2006: 9.0g/day • 2008: 8.6g/day • 2011: 8.1g/day	Fair
Enkhungalag et al. (2015) ⁹⁶	Before and after study	Mongolia	<i>Aim:</i> to reduce salt intake of the employees of three of the main food producing factories. <i>Outcomes:</i> salt intake as measured by 24h urine excretion	Education on salt consumption and provision of reduced salt foods	Salt intake reduced from 11.5g/day in 2011 to 8.7g/day in 2013	Fair

(Continued)

Table 9. (Continued)

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment																		
Trieu et al. (2015) ²⁴	Systematic review	75 countries	<i>Aim:</i> to quantify progress with the initiation of salt reduction strategies around the world in the context of the global target to reduce population salt intake by 30% by 2025. <i>Outcomes:</i> salt (g/day)	Labelling, mass media campaigns, education, reformulation	<i>Denmark:</i> from 2006 to 2010 salt intake reduced from 10.7 to 9.9g/day in men and 7.5g to 7.0g/day in women (7%) <i>Japan:</i> salt intake reduced from 13.5in 1997 to 10.4g/day in 2012 (23%) <i>Korea:</i> salt intake reduced from 13.4g in 2005 to 11.6g/day in 2012 (13.6%) <i>Slovenia:</i> salt intake reduced from 12.4g in 2007 to 11.3g/day in 2012 (8.9%) Du et al. (2014) <i>China:</i> salt intake reduced from 16.8g in 1999 to 12g/day in 2009 (28%) Pietinen et al. (2010) & Laatikainen et al. (2006) <i>Finland:</i> from 1979 to 2007 salt intake reduced from 13g to 8.3g/day in men and 11g to 7.0g/day in women (36%) European commission (2008) <i>France:</i> salt intake reduced from 8.1g in 1999 to 7.7g/day in 2007(4.9%) WHO (2013) <i>Iceland:</i> salt intake reduced from 8.4g in 2002 to 7.9g/day in 2010 (6%) Walton (2013) <i>Ireland:</i> salt intake reduced from 8.1g in 2001 to 7g/day in 2011(13.6%) National Food and Veterinary Risk Assessment Institute <i>Lithuania:</i> salt intake reduced from 10.8g in 1997 to 8.8g/day in 2007(18.6%) WHO (2013) <i>Turkey:</i> salt intake reduced from 18.0g in 2008 to 15g/day in 2012(16.7%) Sadler et al. (2011) <i>UK:</i> Salt intake reduced from 9.5g in 2001 to 8.1g/day in 2011(14.7%)	Fair																		
Luft et al. (1997) ⁹⁷	Review	Finland and US	<i>Aim:</i> to discuss the approaches used in a community-wide salt-reduction project. <i>Outcomes:</i> salt intake as measured by urinary excretion	1) Nutrition education 2) Reformulation	Pietinen et al. (1984) — <i>Health education & reformulation.</i> After 3 y salt intake had not changed significantly. Hypertensive subjects <table><tr><td>Men</td><td>Women</td></tr><tr><td>1979: 13.8 ± 5.3</td><td>1979: 10.4 ± 4.7</td></tr><tr><td>1982: 13.7 ± 5.5</td><td>1982: 10.0 ± 4.1</td></tr></table> Normotensive subjects <table><tr><td>Men</td><td>Women</td></tr><tr><td>1979: 12.4 ± 4.8</td><td>1979: 9.8 ± 3.8</td></tr><tr><td>1982: 12.2 ± 4.8</td><td>1982: 9.1 ± 3.6</td></tr></table> Lang et al. (1985) — <i>Dietary counselling.</i> Women reduced their salt intake from 7.5 ± 0.4 to 3.6 ± 0.2 g/day and men reduced their salt intake from 10.3 ± 0.8 to 4.7 ± 0.3 g/day. Wassertheil-Smoller et a. (1992) — <i>Education.</i> Salt intake as measured by urinary sodium excretion was reduced from 7.9 to 1 6.4 g/day. Analysis of 3-d food records indicated that sodium intake decreased from 8.1 to 4.9 g/day. Hypertension prevention collaborative research group (1992) — <i>Nutrition education.</i> Salt intake as measured by urinary sodium excretion <table><tr><td><i>Intervention</i></td><td><i>Control</i></td></tr><tr><td>Baseline: 8.9 ± 3.4</td><td>Baseline: 9.0 ± 3.5</td></tr><tr><td>Change: -3.2 ± 4.4</td><td>Change: -0.6 ± 4.4</td></tr></table>	Men	Women	1979: 13.8 ± 5.3	1979: 10.4 ± 4.7	1982: 13.7 ± 5.5	1982: 10.0 ± 4.1	Men	Women	1979: 12.4 ± 4.8	1979: 9.8 ± 3.8	1982: 12.2 ± 4.8	1982: 9.1 ± 3.6	<i>Intervention</i>	<i>Control</i>	Baseline: 8.9 ± 3.4	Baseline: 9.0 ± 3.5	Change: -3.2 ± 4.4	Change: -0.6 ± 4.4	Poor
Men	Women																							
1979: 13.8 ± 5.3	1979: 10.4 ± 4.7																							
1982: 13.7 ± 5.5	1982: 10.0 ± 4.1																							
Men	Women																							
1979: 12.4 ± 4.8	1979: 9.8 ± 3.8																							
1982: 12.2 ± 4.8	1982: 9.1 ± 3.6																							
<i>Intervention</i>	<i>Control</i>																							
Baseline: 8.9 ± 3.4	Baseline: 9.0 ± 3.5																							
Change: -3.2 ± 4.4	Change: -0.6 ± 4.4																							
Mohan et al. (2009) ⁹⁸	Review	UK	<i>Aim:</i> to review the evidence related to dietary sodium and health in the context of the Ottawa Charter for health promotion. <i>Outcomes:</i> salt intake, stroke, CVD & coronary artery mortality	1) Reformulation 2) Labelling 3) Health promotion campaign	<i>UK:</i> Consumer-friendly labelling indicating sodium content in processed foods by use of a colour system implemented in several UK food chains. Together with other efforts population salt intake decreased from 9.5g/day in 2004 to 8.6g/day in 2008	Poor																		

(Continued)

Table 9. (Continued)

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
He & MacGregor et al. (2010) ⁹⁹	Comprehensive review	Japan, Finland and UK	<i>Aim:</i> to provide an update on the current salt reduction programmes that have been successfully carried out <i>Outcomes:</i> salt intake	1) Reformulation 2) Labelling 3) Health promotion campaigns	<i>Japan:</i> over a decade national salt intake fell from 13.5g/day to 12.1g/day. In the North, salt intake was reduced from 18 to 14g/day. There was also an 80% reduction in stroke mortality despite large increases in fat intake, cigarette smoking, alcohol consumption, and obesity <i>Finland:</i> reformulation, labelling and campaigns led to a significant reduction in salt from 12g/day in 1979 to 9g/day in 2002 <i>UK:</i> salt reduction strategy started in 2003/2004 and salt intake has already fallen from 9.5 to 8.6 g/d by May 2008	Poor
Wyness et al. (2012) ¹⁰⁰	Literature review	UK	<i>Aim:</i> to describe the UK Food Standards Agency's (FSA) salt reduction programme undertaken between 2003 and 2010 and to discuss its effectiveness <i>Outcomes:</i> salt intake	1) Health promotion campaigns 2) Voluntary reformulation 3) Labelling	<ul style="list-style-type: none"> • 2000–2001: salt intake = 9.5g/day • 2005–2006: salt intake = 9.0g/day • 2008: : salt intake = 8.6g/day 	Poor

<https://doi.org/10.1371/journal.pone.0177535.t009>

In the Netherlands, reformulation of processed foods was predicted to reduce median salt intake by 2.3g/day,[84] compared with a 0.9g/day from a two-year salt reformulation initiative in Argentina.[82]

Fiscal interventions (Table 8)

Two systematic reviews of fair quality [85, 86] included three modelling studies eligible for this review. Furthermore, three additional tax modelling studies were included, all of good quality. [56, 81, 87] Two studies included in Niebylski et al's. systematic review (2015) modelled a 1% tax on salty snacks or on cheese and butter; neither reduced salt consumption.[86] Another modelling study suggested that a very high (40%) tax might achieve a 6% reduction in salt consumption (0.6g/day).[81]

One modelling study predicted that a 20% tax on major dietary sodium sources might prevent or postpone 2000 deaths annually,[87] whilst Nghiem et al. (2015) predicted that a sodium tax might gain more QALYs than other interventions.[56]

Multi-component interventions (Table 9 and Table 10)

Fifteen papers were included under multi-component interventions. Most studies came from Japan, Finland and the UK. Two were of good quality;[88, 89] ten of fair quality;[24, 43, 89–96] and four of poor quality.[97–100]

Four studies were included which presented dietary salt intake and linked to papers describing the interventions; (one of good quality;[25]; two of fair quality;[101, 102] and one of poor quality.[103]

Japan. The Japanese government initiated a sustained campaign in the 1960s.[26] Over the following decade, mean salt intake fell from 13.5g/day to 12.1g/day overall (and from 18g/day to 14g/day in Northern Japan). Miura et al. (2000) reported that salt intake subsequently decreased from 14.5g/day in 1972 to 10.6g/day in 2010, a fall of almost 4g/day [103]. Stroke mortality was predicted to fall by 80%.[90, 93]

Finland. Starting in 1978, Finland pursued a comprehensive salt reduction strategy using mass media campaigns, mandatory labelling and voluntary reformulation by the food industry. Population salt consumption was monitored regularly by using 24h urinary assessment and dietary survey data.[72] By 2007, salt intake had reduced by approximately 4g/day, from

Table 10. Salt intake outcomes with interventions detailed in other publications.

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Laatikainen et al. (2006) ²⁵	Cross-sectional population surveys	Finland	<i>Aim:</i> to present trends in urinary sodium and potassium excretion from 1979 to 2002 <i>Outcomes:</i> salt intake as measured by urinary sodium excretion	1) Reformulation 2) Mass media campaigns 3) Labelling	Between 1979 and 2002 salt intake as measured by sodium excretion decreased from over 12.7g/day to less than 9.8g/day among men and from nearly 10.4 to less than 7.5g/day among women. In 1979 the most educated North Karelian men had lower salt intake compared to the least educated being 11.4 g in the highest education tertile and 13.1 g in the lowest tertile. Respectively, in 2002, the salt intake in southwestern Finland among women in the highest education tertile was 6.7g compared to 8.1g in the lowest tertile	Good
Otsuka et al. (2011) ¹⁰¹	Longitudinal study	Japan	<i>Aim:</i> to describe salt intake over 8 years according to age groups. Also to examine whether salt intake changes over time in middle-aged and elderly Japanese subjects <i>Outcomes:</i> salt intake		In stratified analyses by age, mean salt intake in men decreased 0.08 g/year among 40- to 49-year-olds, 0.09 g/year among 50- to 59-year-olds, 0.16 g/year among 60- to 69-year-olds, and 0.14 g/year among 70- to 79-year-olds. For women, mean salt intake decreased 0.08 g/year among 70- to 79-year-olds (P0.098).	Fair
Du et al. (2014) ¹⁰²	Ongoing open cohort study	China	<i>Aim:</i> to analyse the patterns and trends of dietary sodium intake, potassium intake and the Na/K ratio and their relations with incident hypertension. <i>Outcomes:</i> salt intake as measured per 24h dietary recalls	Labelling & media campaign	Salt intake decreased from 16.5g/day in 1991 to 11.8g/day in 2009	Fair
Miura et al. (2000) ¹⁰³	Report	Japan	<i>Aim:</i> to present the status of salt consumption, salt-reducing measures/guidance methods in individual and population strategies to reduce salt intake <i>Outcomes:</i> salt intake		The National Health and Nutrition Survey in 2010 reported that the mean salt intake in adults was 10.6 g/day. There was an ~4 g decrease in comparison with that in 1972 (14.5 g), when salt intake was investigated for the first time in the National Nutrition Survey	Poor

<https://doi.org/10.1371/journal.pone.0177535.t010>

13 to 8.3g/day in men, and from 11 to 7g/day in women.[24, 25] Stroke and coronary heart disease (CHD) mortality fell by over 75% during that period.[90]

United Kingdom. The UK salt reduction strategy included voluntary reformulation, a consumer awareness campaign, food labelling, target settings and population monitoring.[95] By 2011, population salt intake, measured by 24h urinary sodium excretion, had decreased by 1.4g/day (9.5g/day to 8.1g/day)[88]. He et al. (2014b) estimated that this might reduce stroke and coronary heart disease mortality by some 36%.[88]

Table 11. Modelling studies included in the systematic review.

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Salt Cobiac et al. (2010) ⁴⁴	Modelling study	Australia	Aim: to evaluate population health benefits and cost-effectiveness of interventions for reducing salt in the diet. Outcomes: DALYs and proportion of DALYs averted	1) Voluntary reformulation 2) Mandatory reformulation 3) Dietary advice	Mandatory reformulation: could avert 18% of the disease burden (110,000 DALYs). Dietary advice: might avert less than 0.5% of the disease burden (1,700–2,600 DALYs) Voluntary Reformulation: modelled for breads, margarine and cereals would avert less than 1% of the disease burden (5,300 DALYs) Voluntary reformulation and mandatory salt reduction had a 100% probability of being dominant (i.e., cost saving to the health sector) under all modelled scenarios. Dietary advice had zero probability of being cost-effective.	Good
Cobiac et al. (2012) ⁴⁵	Modelling study	Australia	Aim: to evaluate the optimal mix of lifestyle, pharmaceutical and population-wide interventions for primary prevention of cardiovascular disease Outcomes: DALYs	1) Mandatory reformulation 2) Community heart health programme 3) Dietary advice	Mandatory reformulation in breads, margarine and cereals is easily the most effective and cost-effective strategy for primary prevention of CVD; (80,000 DALYs) and cost saving (dominant). Community heart health program (3,000 DALYs; \$44,000) Dietary advice (180–370 DALYs) are least cost-effective of all the primary prevention strategies (\$1,000,000 to \$1,400,000)	Good
Nghiem et al. (2015) ⁴⁶	Modelling study	New Zealand	Aim: to compare the impact of eight sodium reduction interventions Outcomes: QALYs	1) Dietary counselling 2) Labelling 3) Mandatory 3G reformulation (reads, processed meats and sauces) 4) Mandatory 'All' reformulation 5) UK package (multiple policies) 6) Mass media campaign 7) Tax	QALYs gained in order of effectiveness: 1) Salt tax (195,000) 2) Mandatory 'all' reformulation (110,000) 3) UK package (85,100) 4) Mandatory 3G reformulation (61,700) 5) Mass media campaign as per the UK one (25,200) 6) Voluntary labelling (7,900) 7) Dietary counselling (200)	Good
Collins et al. (2014) ⁴⁸	Modelling study	UK	Aim: to evaluate the cost-effectiveness of four population health policies to reduce dietary salt intake on an English population to prevent coronary heart disease (CHD). Outcomes: life years gained and salt intake	1) Health promotion campaign 2) Labelling 3) Voluntary salt reformulation 4) Mandatory salt reformulation	Primary outcomes: Salt intake reductions: Campaign = 0.16g/d; Labelling = 0.16g/d; Voluntary reformulation = 1.21g/d; Mandatory reformulation = 1.62g/d Secondary outcomes: Gains: Changelife and labelling might each gain approximately = 1,960 life-years; Voluntary reformulation = 14,560 life-years; and Mandatory reformulation 19,320 life-years.	Good
Gillespie et al. (2015) ⁴⁹	Modelling study	England	Aim: to forecast the potential impact on English adults of policies implemented during the 2015 UK parliament, projecting the health consequences to 2025 Outcomes: salt intake, CHD deaths prevented, LYG	1) Mandatory reformulation 2) Voluntary reformulation 3) Social marketing 4) Nutrition labelling	Mandatory reformulation (30% reduction in salt content) • Salt intake = -1.45g/day • SBP = -0.81mmHg • CHD deaths = 4,500 prevented or postponed • LYG = 44,000 Mandatory reformulation (10% reduction in salt content) • Salt intake = -0.48g/day • SBP = -0.27mmHg • CHD deaths = 1,500 prevented or postponed • LYG = 15,000 Voluntary reformulation • Salt intake = -0.48g/day • SBP = -0.27mmHg • CHD deaths = 1,500 prevented or postponed • LYG = 14,000 Social marketing (50% impact) • Salt intake = -0.13g/day • SBP = -0.078mmHg • CHD deaths = 400–500 prevented or postponed • LYG = 5,000 Social marketing (10% impact) • Salt intake = -0.027g/day • SBP = -0.015mmHg • CHD deaths = 100 prevented or postponed • LYG = 780 Nutrition labelling (50% impact) • Salt intake = -0.16g/day • SBP = -0.091mmHg • CHD deaths = 500 prevented or postponed • LYG = 5,000 Nutrition labelling (10% impact) • Salt intake = -0.031g/day • SBP = -0.018mmHg • CHD deaths = 100 prevented or postponed • LYG = 1,000	Good
Wilcox et al. (2014) ⁷⁰	Modelling study	Syria	Aim: to present a cost-effectiveness analysis of salt reduction policies to lower coronary heart disease in Syria. Outcomes: salt intake, deaths prevented and life years gained	1) Health promotion 2) Labelling 3) Reformulation	Health promotion campaign – 5% reduction in salt intake. 252 deaths prevented; 5,679 LYG Labelling – 10% reduction in salt intake. 497 deaths prevented; 11,192 LYG Reformulation – 10% reduction in salt intake. 497 deaths prevented; 11,192 LYG Reformulation + HP & Reformulation + Labelling – 15% reduction in salt intake. 735 deaths prevented; 16,543 LYG All 3 policies – 30% reduction in salt intake. 1,413 deaths prevented; 31,674 LYG	Good

(Continued)

Table 11. (Continued)

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Mason et al. (2014) ⁷¹	Modelling study	Tunisia, Syria, Palestine and Turkey	Aim: to present an economic evaluation of population based salt reduction policies in Tunisia, Syria, Palestine and Turkey. <i>Outcomes</i> : life years gained	1) Health promotion campaign (HP) 2) Labelling (L) 3) Mandatory reformulation (R)	Tunisia: HP = 1.15 LY; L = 2.272 LY; R = 2.272 LY; All 3 policies = 6.455 LY Syria: HP = 5.679 LY; L = 11.192 LY; R = 11.192 LY; All 3 policies = 31.674 LY Palestine: HP = 479 LY; L = 945 LY; R = 945 LY; All 3 policies = 2.682 LY Turkey: HP = 68.816 LY; L = 135.221 LY; R = 135.221 LY; All 3 policies = 199.303 LY	Fair
Pietinen et al. (2008) ⁷⁴	Modelling study	Finland	Aim: to estimate the impact of choosing food products labelled either as low or high in salt. <i>Outcomes</i> : salt intake	Salt labelling	If the entire population were to choose low-salt breads, cheeses, processed meat and fish, fat spreads, and breakfast cereals, then salt intake could be lowered by 1.5 g in men and by 0.9 g in women. If everybody was to select high-salt products, then salt intake would go up by 1.9 g in men and by 1.2 g in women. Thus, the potential difference between the low and the high alternatives would be 3.4 g in men and 2.9 g in women. If all prepared foods had a reduced salt content, the mean salt intake would go further down by 2.3 g in men and by 1.7 g in women. Excluding under reporters, the mean salt intake was 11.1 g in men and would go down to 9.5 g if all men chose lightly salted products and further down to 6.8 g if also all prepared foods would have a lower salt content. In women, the respective numbers are 7.8, 6.7 and 4.9 g	Fair
Terme et al. (2010) ⁷⁵	Modelling study	Netherlands	Aim: to evaluate the effects of changed food compositions according to health logo criteria on the intake of saturated fat, sugar and sodium. <i>Outcomes</i> : salt intake	Labelling (health logos)	At baseline salt intake was 7.3 (95% CI 2.8, 2.9) g/day. For salt, in a 100% market share scenario (scenario II), salt reduction expected is 0.3g/day (4% reduction; non-significant). In scenario III, when all non-complying foods are replaced with foods complying with health logo criteria, sodium intake reduced by 23% to 5.5g/d (-1.8g/day).	Fair
De Menezes et al. (2013) ⁷⁶	Modelling study	Brazil	Aim: to evaluate the impact of introducing products that are in agreement with the Choices criteria in the usual diet. <i>Outcomes</i> : salt intake	Food Labelling	Salt would still be considered an important reduction, 36% in relation to the typical menus (TM), but it would be 7.6 g/day, which is above the recommended by the program (6.5 g/day). Salt (g/day) • Typical menus (TM): 11.9 ± 1.2 • Choices menus (CM): 6.3 ± 0.1 • Choices menus energy (CME)—same as CM, but adjusted for energy of TM: 7.6 ± 0.9	Fair
Roodenburg et al. (2013) ⁷⁷	Modelling study	The Netherlands	Aim: to evaluate these nutritional criteria by investigating the potential effect on nutrient intakes. <i>Outcomes</i> : salt intake	Labelling	A reduction of ~23% for sodium was seen for sodium compared to the 'actual scenario'. Salt (g/day) • Actual: 7.4 • Choices: 5.7 • Choices energy adjusted: 6.5 • Snacks (partially replaced): 5.8 • Snacks (not replaced): 5.9	Fair
Choi et al. (2015) ⁷¹	Modelling study	US	Aim: to estimate the cardio-vascular impact of the expanded NSRI among different segments of the US population and under varying possible producer and consumer responses to the initiative. <i>Outcomes</i> : sodium intake, MIs, stroke and hypertension	Reformulation (restaurants and manufacturers)	<i>Restaurants and manufacturers reaching agreed upon sodium targets</i> . Expansion of the initiative to ensure all restaurants and manufacturers reach agreed-upon sodium targets would be expected to avert from 0.9 to 3.0 MIs (a 1.6%–5.4% reduction) and 0.5 to 2.8 strokes (a 1.1%–6.2% reduction) per 10,000 Americans per year over the next decade, after incorporating consumption patterns and variations in the effect of sodium reduction on blood pressure among different demographic groups. The expanded NSRI covering both packaged and restaurant food items would be expected to reduce mean daily sodium intake by 447 mg per person per day on average, or 13.0%. If the NSRI included only restaurant food items, the program would lower MI and stroke mortality by an estimated 2.7% (95% CI, 1.4–4.0) and 2.1% (95% CI, 0.4–3.9), respectively. Hence, most of the benefit from the program would likely be due to sodium changes among packaged foods	Good
Murray et al. (2003) ⁷⁹	Modelling study	South East Asia (SEA), Latin America (LA), Europe (EU)	Aim: to report estimates of the population health effects, and costs of selected interventions to reduce the risks associated with high cholesterol and blood pressure in areas of the world with differing epidemiological profiles. <i>Outcomes</i> : DALYs	1) Voluntary reformulation 2) Mandatory reformulation	Measures to decrease salt intake appeared cost effective. Legislation appeared more cost effective than voluntary agreements with assumption it would lead to a larger reduction in dietary salt intake. A 15% reduction in mean population salt intake could avert 8.5 million cardiovascular deaths Voluntary reformulation: • EU 7 X10 ⁶ DALYs averted (\$44 per DALY) • SEA: 5X10 ⁶ DALYs averted (\$37 per DALY) • LA: 3 X10 ⁶ DALYs averted (\$24 per DALY) Mandatory reformulation: • EU: 13x10 ⁶ DALYs averted (\$23 per DALY) • SEA: 10x10 ⁶ DALYs averted (\$19 per DALY) • LA: 6x10 ⁶ DALYs averted (\$13 per DALY)	Good
Rubinstein et al. (2010) ⁸⁰	Modelling study	Argentina	Aim: to estimate the burden of acute CHD and stroke and the cost-effectiveness of preventative population-based and clinical interventions. <i>Outcomes</i> : DALYs	Salt reduction in bread	<i>Reducing salt in bread is cost-saving</i> • DALY averted: 672.80 • % of DALY saved: 0.11% • International Dollars per DALY saved: 1,406.93	Good

(Continued)

Table 11. (Continued)

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Smith-Spangler et al. (2010) ⁸¹	Modelling study	US	Aim: to assess the cost-effectiveness of two population strategies to reduce sodium intake. <i>Outcomes:</i> strokes and MIs averted, life years and QALYs gained	1) Voluntary reformulation 2) Sodium tax	<i>Collaboration with the industry:</i> a 9.5% reduction in sodium intake resulted in • Averted strokes = 513 885 s • Averted MIs = 480 358 • LYG = 1.3 million • QALYs = 2 million <i>Sodium tax:</i> would lead to a 6% decrease in sodium intake. • Averted Strokes = 327 892 • Averted MIs = 306 137 • LYG = 840 113 • QALYs = 1.3 million	Good
Konfino et al. (2013) ⁸²	Modelling study	Argentina	Aim: to use Argentina-specific data to project impact of Argentina's sodium reduction policies under two scenarios—the 2-year intervention currently being undertaken or a more persistent 10-year sodium reduction strategy. <i>Outcomes:</i> salt intake as measured by urinary sodium excretion, systolic blood pressure, deaths and cases averted, mortality	Reformulation	<i>Scenario 1: current initiative (2 year intervention)</i> • Projected to reduce mean salt consumption by 0.96 g/day in men and 0.79 g/day in women • SBP would reduce by 0.93 mmHg to 1.81 mmHg depending on population subgroup • 19,000 deaths, 6,000 CHD deaths and 2,000 stroke deaths, 13,000 MIs and 10,000 stroke cases averted • Overall mortality reduction of 0.6% in adults >35 years, 1.5% in total MIs, 1% in total stroke cases in the next decade <i>Scenario 2: current initiative maintained for 10 years</i> • Projected to reduce mean salt consumption by 4.83 g/day in men and 3.98 g/day in women • SBP would reduce by 4.68 mmHg to 9.04 mmHg depending on subgroup • 55,000 deaths, 16,000 CHD deaths and 5,000 stroke deaths, 38,000 MIs and 27,000 strokes averted • Overall mortality decreased by 2% in adults > 35 years, 4.3% MIs and 2.7% stroke cases in the next decade	Good
Rubinstein et al. (2009) ⁸³	Modelling study	Argentina	Aim: to use generalised cost-effectiveness analysis to identify the most efficient interventions to decrease CVD. <i>Outcomes:</i> cost-effectiveness, DALYs	Reformulation in bread	Lowering salt intake in the population through reducing salt in bread was found to be the most cost-effective (\$17 per DALY averted). Less salt in bread • Total Cost per year (AR\$S): \$9,644 • DALY Age weighted, 3% discounted per year: 579 • DALY No age-weight 3% discounted per year: 713 • DALY # age-weight, undiscounted per year: 1,107 • AR\$S (1)/DALY (2): \$17	Fair
Hendriksen et al. (2014) ⁸⁴	Modelling study	Netherlands	Aim: to evaluate the health benefits of salt-reduction strategies related to processed foods. <i>Outcomes:</i> AMI, CHF and CVA averted, life expectancy and DALYs gained, salt intake (g/day)	1) Reformulation 2) Substitution of high salt foods with low salt foods 3) Adherence to the recommended intake	If salt intake is reduced to the recommended maximum salt intake (6 g/d): Prevented cases: • 31,800 cases of AMI; • 15,300 cases of CHF; • 51,900 cases of CVA. • Mortality reduction: 0.77%. • LE increased by 0.15 years • 56,000 DALYs gained <i>Salt reduction processed foods scenario:</i> median salt intake would decrease by -2.3g/d (28%). Prevented Cases: • 29,200 AMI cases; • 16,600 CHF cases; • 53,400 CVA. • Mortality Reduction: 0.8%. • LE increased by 0.15 years • 56,400 DALYs gained <i>Substitution:</i> median salt intake would decrease by -3.0g/day (35%). Prevented Cases: • 35,500 cases of AMI; • 20,000 cases of CHF; • 64,300 cases of CVA • LE increased by 0.18 years • 67,900 DALYs gained	Fair
Ni Mhurchu et al. (2015) ⁸⁷	Modelling study	New Zealand	Aim: to estimate the effects of health-related food taxes and subsidies. <i>Outcomes:</i> deaths prevented or postponed	Tax on major dietary sodium products	A 20% tax on major dietary sources of sodium might result in 2,000 (1300 to 2,700) DPP (6.8%)	Good
Asaria et al. (2007) ¹⁰⁴	Modelling study	23 low and middle income countries	Aim: to investigate potential deaths averted over 10 years by implementation of selected population-based interventions. <i>Outcomes:</i> CVD deaths averted, salt reduction (g/day)	Combined: 1) Mass media campaign 2) Voluntary reformulation	8.5 million deaths would be averted by implementation of the salt-reduction strategy (15%) alone. Salt interventions: 15% reduction in mean salt intake • risk factor reduction of 1.69g/day • 8.4 million CVD deaths averted 30% reduction in mean intake • risk factor reduction of 3.38g/day • 16.0 million CVD deaths averted Reducing salt intake to 5g/d • risk factor reduction of 6.28g/d day • 28.3 million CVD deaths averted Reduction in deaths • CVD = 75.6% Respiratory disease = 15.4% Cancer = 8.7%	Good

(Continued)

Table 11. (Continued)

Study	Study type	Geographical scope	Aim and main outcomes	Policies analysed	Relevant results	Quality assessment
Dodhia et al. (2012) ¹⁰⁵	Modelling study	England	Aim: to assess the impact of cost-effective interventions in terms of the avoidable CVD burden and costs by comparing these strategies to the current situation Outcomes: IHD and stroke events and deaths avoided, DALYs	Combined 1) Health promotion 2) Reformulation	30% reformulation through agreement with the food industry. Interventions: Na–2mmHg • IHD events avoided: 56,116 • Stroke events avoided: 98,497 • IHD deaths avoided: 26,781 • Stroke deaths avoided: 39,557 • DALYs averted: 288,043 • Cost per DALY (\$): -4,228 Na–5mmHg • IHD events avoided: 120,138 • Stroke events avoided: 257,508 • IHD deaths avoided: 57,322 • Stroke deaths avoided: 103,492 • DALYs averted: 579,869 • Cost per DALY (\$): -5,021 Reducing salt intake to 6g/day through reformulation NA–MRC review • IHD events avoided: 80,366 • Stroke events avoided: 128,032 • IHD deaths avoided: 38,372 • Stroke deaths avoided: 51,419 Reducing salt intake in the population with a 5 mmHg reduction in SBP had the greatest population impact and cost-saving to the NHS.	Good
Gase et al. (2011) ¹⁰⁶	Modelling study	US	Aim: to examine approaches to reduce sodium content of food served in settings operated or funded by the government of the County of Los Angeles, California Outcomes: salt intake and BP	Combined: 1) Labelling 2) Promotion 3) Subsidy 4) Provide low sodium food options	Hospital cafeterias: Average sodium reduction of 1.8g/day (23%). Overall SBP: 1.59 County government cafeterias: Average sodium reduction of 0.7g/day (11%). Overall SBP: 0.63	Fair
Ha & Christholm (2011) ¹⁰⁷	Modelling study	Vietnam	Aim: to assess costs, health effects and cost-effectiveness prevention strategies to reduce CVD Outcomes: DALYs	Combined 1) Mass media campaign 2) Voluntary reformulation	Media salt campaign • Cost per year (USD, million): 4.1 • DALYs averted per year: 45,939 • VND per DALY saved: 89.2	Fair
Barton et al. (2011) ¹²⁰	Modelling study	England and Wales	Aim: to estimate the potential cost effectiveness of a population-wide risk factor reduction programme aimed at preventing cardiovascular disease. Outcomes: BP, CVD deaths averted	Salt legislation	Reducing salt intake by 3 g/day might reduce mean population systolic blood pressure by approximately 2.5 mm Hg preventing approximately 4450 deaths from cardiovascular disease	Good

<https://doi.org/10.1371/journal.pone.0177535.t011>




Fig 1. Interventions classified on the upstream / downstream continuum.

<https://doi.org/10.1371/journal.pone.0177535.g001>

Other countries have implemented several strategies including labelling, media campaigns and voluntary reformulation and effect sizes ranged from -0.4g/day in France [24, 93] to -4.8g/day in China [24, 102].

Modelling studies of combined interventions. Six modelling studies investigated the effect of multi-component interventions, three were of good quality; [70, 104, 105] whilst three others were of fair quality. [70, 106, 107]

Several modelling studies consistently suggested that multi-component salt reduction strategies (e.g. labelling, health promotion and reformulation) would be more effective than any single intervention. [70, 71] For instance, Gase et al. (2011) suggested that using labelling, promotion, subsidies and provision of low sodium options could lead to a 0.7–1.8g/day reduction. [106]

Discussion

Main results

This systematic review of salt reduction interventions suggests that comprehensive strategies could generally achieve the biggest reductions in salt consumption across an entire population, approximately 4g/day in Finland and Japan, 3g/day in Turkey and 1.3g/day recently in the UK. Mandatory reformulation alone could achieve a reduction of approximately 1.4g/day, followed by voluntary reformulation (median 0.7g/day) school interventions (0.7g/day) and worksite interventions (+0.5g/day). Smaller population benefits were generally achieved by short-term dietary advice (0.6g/day), community-based counselling (0.3g/day), nutrition labelling (0.4g/day), and health education media campaigns (-0.1g/day). Although dietary advice to




Fig 2. PRISMA flowchart.

<https://doi.org/10.1371/journal.pone.0177535.g002>

individuals achieved a -2g/day reduction, this required optimal research trial conditions (smaller reductions might be anticipated in unselected individuals).

Comparison with other research

Geoffrey Rose famously argued that a greater net benefit came from the population-wide approach, (achieving a small effect in a large number of people) when compared with targeting high risk individuals (a large effect but only achieved in a small number of people).[32]




Fig 3. Effectiveness of interventions to reduce salt intake (empirical studies). Forest plot of the empirical studies that were included in this systematic review. Negative values of salt reduction are interpreted as reported increase in salt consumption. For most combined interventions the sample size and confidence intervals were not reported. NA denotes not applicable or not reported.

<https://doi.org/10.1371/journal.pone.0177535.g003>




Fig 4. Effectiveness of interventions to reduce salt intake (modelling studies). Forest plot of the modelling studies that were included in this systematic review. Because of the different modelling approaches in these studies, their uncertainty measures are not comparable. Therefore we do not plot them in this graph. Different scenarios were considered for different studies. NA denotes not applicable or not reported.

<https://doi.org/10.1371/journal.pone.0177535.g004>

Multi-component interventions. Multi-component salt reduction strategies involving a series of structural initiatives together with campaigns to increase population awareness have been successful in Japan and Finland where they substantially reduced dietary salt consumption, and associated high stroke and cardiovascular disease mortality rates. In Finland, some credit should also go to other dietary changes e.g. fat quality.[108]

Between 2003 and 2010, a multi-component approach in the UK including voluntary reformulation and political pressure on industry to agree category-specific targets achieved some success (1.3g/day reduction in population salt consumption over 8 years to 8.1g/day in 2011). Interestingly, pre-existing health inequalities in salt consumption persisted.[29] However,

from 2010, the Responsibility Deal simply advocated a voluntary scheme. This was ineffective, and MacGregor therefore subsequently recommended mandatory reformulation.[31] Other useful reductions were demonstrated in other countries mostly using dietary surveys and some from grey literature. However, the -4.8g/day reduction reported in China appears extra-ordinarily large and perhaps merits some caution [24]. Multi-component interventions clearly have more potential than single interventions, and synergies might be anticipated. [13,93] Similarly powerful benefits have also been observed with comprehensive strategies for tobacco control and alcohol reduction.[35,36]

Reformulation. In high income countries, the majority of dietary salt intake comes in processed food (75%) and reformulation can be very effective in reducing salt consumption. [109] Though mandatory reformulation is more powerful, most countries currently use voluntary reformulation.[54,56,68,69,110] Success may then be very dependent on the degree of political pressure applied to the food industry and on regular, independent monitoring, as recently achieved in the UK. [111,112]

Food labelling. Nutrition labelling can be potentially effective, as demonstrated in Finland [72] and Brazil.[74] Nutrition labelling allows consumers to make informed choices whilst also putting pressure on the food industry to reformulate.[89] However, interpretation of labels depends on health literacy and different labelling systems may confuse consumers, [113] and reinforce inequalities.[29]. Consumers generally want simple (traffic light) labels which are easier to understand.[76,77,113,114]

Dietary interventions in diverse settings: communities, worksites, schools and homes.

Dietary interventions can be delivered at different levels, such as communities, worksites, schools or to individuals. However, effectiveness varies widely.[45,47,50] Furthermore, the benefits of dietary counselling decrease over time and are thus generally not sustainable; much smaller reductions might therefore be anticipated in unselected individuals in the general population.[44] Furthermore, for many individuals, issues such as competing priorities and financial constraints might reduce compliance and adherence,[8,13,21,22] and thus reduce net population benefits.

Mass media campaigns. Few empirical studies have examined salt media campaigns. However, benefits appear to be generally modest.[56, 67,68,69,115] or negligible.[111] Many individuals may not perceive any personal relevance and hence fail to engage in any behaviour change.[22,116,117]

Taxation. Price increases can powerfully reduce consumption of tobacco or alcohol. [35,36] However, salt is cheap, and a substantial tax of at least 40% might be needed to reduce consumption by just 6%.[81,118]

Public health benefits and cost-effectiveness

Most economic analyses have consistently predicted substantial reductions in cardiovascular mortality, and consequent gains in life-years, QALYs, DALYs and healthcare savings. This is consistent with the growing evidence that population-wide prevention policies can often be powerful, rapid, equitable and cost-saving.[38,119–122]

Several modelling studies also investigated the cost-effectiveness of the salt interventions described above. Mandatory and voluntary reformulation appeared far more cost-effective than labelling or [54,55,68] dietary advice targeting individuals.[122]

Strengths and limitations

This systematic review has multiple strengths. Firstly, two independent reviewers screened all papers and assessed quality using appropriate validated tools. Secondly, the inclusion of

modelling studies (presented separately) adds value by allowing the evaluation of certain interventions where empirical studies failed (e.g. labelling). In addition, we recorded the effect size used in each modelling paper together with the source reference. Furthermore, most of the better quality modelling studies confirmed the superiority of upstream approaches. Finally, the studies reviewed included a wide variety of interventions, thus providing a useful spread of estimates.

Our review also has limitations. We were unable to conduct a formal meta-analysis due to the profound heterogeneity of the diverse studies, many of which included multiple interventions. Furthermore, studies were only included if the full text was available in English (15 non-English papers were excluded). We also had to exclude two potentially relevant studies which lacked the full text.[123,124] Publication bias remains possible, potentially over-estimating the true effect of some interventions. The primary outcome of this study was dietary intake (consumption); we excluded studies considering other dietary behaviours such as awareness, knowledge, preferences or purchasing behaviour. Also, the positive benefits of policy changes may sometimes appear larger if favourable underlying secular trends have not been formally considered. Furthermore, we did not contact authors for missing data. However, all the key information was presented in all but two papers. [123,124] Finally, generalization of the results should be cautioned as countries may vary in baseline salt intake.

Socio-economic Inequalities

More deprived groups more often consume foods high in salt, (and sugar and fat); all are associated with poor health.[125–127] These inequalities persist in Britain [28,29] and Italy.[128]

Downstream interventions focused on individuals typically widen inequalities whereas upstream “structural” interventions may reduce inequalities.[33,129,130]

Future research

This review highlights the greater power of combined (multi-component) strategies, mandatory reformulation and traffic light labelling. Most were cost-effective and many were cost-saving. However, the feasibility of implementing policy changes also deserves further study. Many factors can facilitate or obstruct successful policy development, notably including political feasibility and stakeholder influence.[114,131,132]

Stoeckle and Zola’s “upstream”/“downstream” concept was disseminated by John McKinlay,[133] critiqued by Krieger,[134] and then refined as a structural/agentive continuum by McLaren et al 2010.[21] To test our effectiveness hierarchy hypothesis, one ideally needs to quantify the “average” effect of each category of salt reduction intervention. Yet, the limited number and heterogeneity of these studies precludes a formal meta-analysis. However, the consistency with the effectiveness hierarchies demonstrated by tobacco and alcohol control interventions is encouraging. The effectiveness hierarchy hypothesis now clearly needs to be tested in other fields.

Conclusions

There are clear implications for public health. The biggest population-wide reductions in salt consumption were consistently achieved by comprehensive multi-component strategies involving “upstream” population-wide policies (regulation, mandatory reformulation, and food labelling). “Downstream” individually-based interventions appear relatively weak (e.g. dietary counselling to individuals and school children, and media campaigns in isolation).

This ‘effectiveness hierarchy’ might deserve greater emphasis on the agendas of the WHO and other global health organizations reviewing action plans for NCD prevention.

Supporting information

S1 Table. PRISMA checklist.
(DOCX)

S2 Table. Full data extraction tables empirical and modelling studies.
(DOC)

S1 File. Research protocol.
(DOCX)

Acknowledgments

We thank Mark Petticrew and Cecile Knai for their very helpful comments. FPC contributed under the remit of the Terms of Reference of the World Health Organization Collaborating Centre for Nutrition of the University of Warwick.

Author Contributions

Conceptualization: SC.

Formal analysis: CK LH SC.

Funding acquisition: SC.

Investigation: LH AEG FLW RMG LO.

Methodology: LH LO RMG SC.

Project administration: LH.

Supervision: SC LH.

Validation: LH AEG.

Visualization: CK LH.

Writing – original draft: LH AEG FLW CK MOF RMG LO HB FPC SC.

Writing – review & editing: LH AEG FLW CK MOF RMG LO HB FPC SC.

References

1. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012; 380(9859): 2095–128. [https://doi.org/10.1016/S0140-6736\(12\)61728-0](https://doi.org/10.1016/S0140-6736(12)61728-0) PMID: 23245604
2. Vos T, Barber RM, Bell B, Bertozzi-Villa A, Biryukov S, Bolligeret I, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. In press, corrected proof.
3. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet*. 2012; 380: 2224–2260.
4. Anderson CA, Appel LJ, Okuda N, Brown IJ, Chan Q, Zhao L, et al. Dietary Sources of Sodium in China, Japan, the United Kingdom, and the United States, Women and Men Aged 40 to 59 Years: The INTERMAP Study. *J Am Diet Assoc*. 2010; 110(5): 736–45. <https://doi.org/10.1016/j.jada.2010.02.007> PMID: 20430135

5. Webster JL, Dunford EK, Neal BC. A systematic survey of the sodium contents of processed foods. *Am J Clin Nutr.* 2010; 91(2): 413–420. <https://doi.org/10.3945/ajcn.2009.28688> PMID: 19955402
6. Ni Mhurchu C, Capelin C, Dunford EK, Webster JL, Neal BC, Jebb SA. Sodium content of processed foods in the United Kingdom: analysis of 44,000 foods purchased by 21,000 households. *Am J Clin Nutr.* 2011; 93(3): 594–600. <https://doi.org/10.3945/ajcn.110.004481> PMID: 21191142
7. World Health Organization. Salt reduction [online], WHO 2010. Available at: <http://www.who.int/mediacentre/factsheets/fs393/en/>
8. Cappuccio FP & Capewell S. Facts, Issues, and Controversies in Salt Reduction for the Prevention of Cardiovascular Disease. *Functional Food Reviews.* 2015; 7: 41–61.
9. He FJ & MacGregor GA. Effect of modest salt reduction on blood pressure: a meta-analysis of randomized trials. Implications for public health. *J Hum Hypertens.* 2002; 16: 761–770. <https://doi.org/10.1038/sj.jhh.1001459> PMID: 12444537
10. Aburto NJ, Ziolkovska A, Hooper L, Elliott P, Cappuccio FP, Meerpohl JJ. Effect of lower sodium intake on health: systematic review and meta-analyses. *BMJ.* 2013; 346: f1326. <https://doi.org/10.1136/bmj.f1326> PMID: 23558163
11. He FJ, Li J, MacGregor GA. Effect of longer term modest salt reduction on blood pressure: Cochrane systematic review and meta-analysis of randomised trials. *BMJ.* 2013; 346: f1325. <https://doi.org/10.1136/bmj.f1325> PMID: 23558162
12. Cook NR, Cutler JA, Obarzanek E, Buring JE, Rexrode KM, Kumanyika SK, et al. Long term effects of dietary sodium reduction on cardiovascular disease outcomes: observational follow-up of the trials of hypertension prevention (TOHP). *BMJ.* 2007; 334: 885. <https://doi.org/10.1136/bmj.39147.604896.55> PMID: 17449506
13. He FJ & MacGregor GA. A comprehensive review on salt and health and current experience of world-wide salt reduction programmes. *J Hum Hypertens.* 2009; 23(6): 363–84. <https://doi.org/10.1038/jhh.2008.144> PMID: 19110538
14. Strazzullo P, D'Elia L, Kandala NB, Cappuccio FP. Salt intake, stroke, and cardiovascular disease: meta-analysis of prospective studies. *BMJ.* 2009; 339: b4567. <https://doi.org/10.1136/bmj.b4567> PMID: 19934192
15. Perry IJ & Beevers DG. Salt intake and stroke: a possible direct effect. *J Hum Hypertens.* 1992; 6(1): 23–5. PMID: 1583626
16. Karppanen H & Mervaala E. Sodium intake and hypertension. *Prog Cardiovasc Dis.* 2006; 49(2): 59–75. <https://doi.org/10.1016/j.pcad.2006.07.001> PMID: 17046432
17. Joossens JV, Hill MJ, Elliott P, Stamler R, Lesaffre E, Dyer A et al. Dietary salt, nitrate and stomach cancer mortality in 24 countries. European Cancer Prevention (ECP) and the INTERSALT Cooperative Research Group. *Int J Epidemiol.* 1996; 25: 494–504. PMID: 8671549
18. Wang X- Q, Terry PD, Yan H. Review of salt consumption and stomach cancer risk: Epidemiological and biological evidence. *World J Gastroenterol.* 2009; 15(18): 2204–13. <https://doi.org/10.3748/wjg.15.2204> PMID: 19437559
19. D'Elia L, Rossi G, Ippolito R, Cappuccio FP, Strazzullo P. Habitual salt intake and risk of gastric cancer: a meta-analysis of prospective studies. *Clin Nutr.* 2012; 31: 489–98. <https://doi.org/10.1016/j.clnu.2012.01.003> PMID: 22296873
20. WHO. Guideline: Sodium intake for adults and children. Geneva, World Health Organization (WHO), 2012
21. McLaren L, McIntyre L & Kirkpatrick S. Rose's population strategy of prevention need not increase social inequalities in health. *Int J Epidemiology.* 2010; 39: 372–377.
22. Adams J, Mytton O, White M, Monsivais P. Why Are Some Population Interventions for Diet and Obesity More Equitable and Effective Than Others? The Role of Individual Agency. *PLoS Med* (2016); 13(4): e1001990. <https://doi.org/10.1371/journal.pmed.1001990> PMID: 27046234
23. Brownson RC, Seiler R, Eyler AA. Measuring the impact of public health policy. *Prev Chronic Dis.* 2010; 7(4): A77. PMID: 20550835
24. Trieu K, Neal B, Hawkes C, Dunford E, Campbell N, Rodriguez-Fernandez R, et al. Salt initiatives around the world—A systematic review of progress towards the global target. *Plos One.* 2015; 10(7): e0130247 <https://doi.org/10.1371/journal.pone.0130247> PMID: 26201031
25. Laatikainen T, Pietinen P, Valsta L, Sundvall J, Reinivuo H, Tuomilehto J. Sodium in the Finnish diet: 20-year trends in urinary sodium excretion among the adult population. *Eur J Clin Nutr.* 2006; 60(8): 965–70. <https://doi.org/10.1038/sj.ejcn.1602406> PMID: 16482074
26. Sasaki N. The salt factor in apoplexy and hypertension: epidemiological studies in Japan. In: Yamori Y, editor. Prophylactic approach to hypertensive diseases. New York: Raven Press; 1979. p. 467–74.

27. He FJ, Pombo-Rodrigues S, Macgregor GA. Salt reduction in England from 2003 to 2011: its relationship to blood pressure, stroke and ischaemic heart disease mortality. *BMJ Open*. 2014a; 4(4): e004549.
28. Ji C, Kandala N- B & Cappuccio FP. Spatial variation of salt intake in Britain and association with socio-economic status. *BMJ Open*. 2013; 3: e002246. <https://doi.org/10.1136/bmjopen-2012-002246> PMID: 23295624
29. Ji C & Cappuccio FP. Socio-economic inequality in salt intake in Britain 10 years after a national salt reduction programme. *BMJ Open*. 2014; 4: e005683. <https://doi.org/10.1136/bmjopen-2014-005683> PMID: 25161292
30. Knai C, Petticrew M, Durand MA, Eastmure E, James L, Mehrotra A, et al. Has a public-private partnership resulted in action on healthier diets in England? An analysis of the Public Health Responsibility Deal food pledges. *Food Policy*. 2015; 54: 1–10.
31. MacGregor GA, He FJ, Pombo-Rodrigues S. Food and the responsibility deal: how the salt reduction strategy was derailed by Andrew Lansley and the coalition government. *Br Med J*. 2015; 350: h1936.
32. Rose G. Sick individuals and sick populations. *International Journal of Epidemiology*. 2001; 30:427–432 PMID: 11416056
33. Capewell S & Graham H. Will cardiovascular disease prevention widen health inequalities? *PLoS Med*. 2010; 7(8): e1000320. <https://doi.org/10.1371/journal.pmed.1000320> PMID: 20811492
34. Hogberg L, Cnattingius S, Lundholm C, Sparén P, Iliadou AN. Intergenerational social mobility and the risk of hypertension. *J Epidemiol Community Health*. 2012; 66(6): e9. <https://doi.org/10.1136/jech.2010.130567> PMID: 21747130
35. Joossens L & Raw M. The Tobacco Control Scale: a new scale to measure country activity. *Tob Control*. 2006; 15: 247–253. <https://doi.org/10.1136/tc.2005.015347> PMID: 16728757
36. Anderson P, Chisholm D, Fuhr DC. Alcohol and Global Health 2. Effectiveness and cost-effectiveness of policies and programmes to reduce the harm caused by alcohol. *Lancet*. 2009; 373: 2234–46. [https://doi.org/10.1016/S0140-6736\(09\)60744-3](https://doi.org/10.1016/S0140-6736(09)60744-3) PMID: 19560605
37. Cappuccio FP, Capewell S, Lincoln P, McPherson K. Population salt reduction to prevent cardiovascular disease: identifying policy options. *BMJ*. 2011; 343: d4995. <https://doi.org/10.1136/bmj.d4995> PMID: 21835876
38. NICE Public Health Guidance: Prevention of cardiovascular disease at population level. 2010. Reviewed and updated in 2014: <http://guidance.nice.org.uk/PH25/Review>
39. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for the systematic reviews and meta-analysis: the PRISMA statement. *BMJ*. 2009; 339: b2535. <https://doi.org/10.1136/bmj.b2535> PMID: 19622551
40. He FJ, Wu Y, Feng X- X, Ma J, Ma Y, Wang H, et al. School based education programme to reduce salt intake in children and their families (School-EduSalt): cluster randomised controlled trial. *BMJ*. 2015; 350: h770. <https://doi.org/10.1136/bmj.h770> PMID: 25788018
41. Choi SE, Brandeau ML, Basu S. Expansion of the National Salt Reduction Initiative: A Mathematical Model of Benefits and Risks of Population-Level Sodium Reduction. *Medical Decision Making* 2015.
42. National institute of Health. Quality assessment tools. 2014. Accessed from <https://www.nhlbi.nih.gov/health-pro/guidelines/in-develop/cardiovascular-risk-reduction/tools>
43. Fattore G, Ferre F, Mereaglia M, Fattore E, Agostoni C. Critical review of economic evaluation studies of interventions promoting low-fat diets. *Nutr Rev*. 2014; 72(11): 691–706. <https://doi.org/10.1111/nure.12142> PMID: 25323698
44. Nuffield Council on Bioethics. Policy process and practice. In: *Public Health: Ethical issues*. London, UK: Nuffield Council on Bioethics. 2007: 31–47.
45. Hooper L, Bartlett C, Davey Smith G, Ebrahim S. Systematic review of long term effects of advice to reduce dietary salt in adults. *BMJ*. 2002; 325(7365): 628. PMID: 12242173
46. Appel LJ. Effects of Comprehensive Lifestyle Modification on Blood Pressure Control: Main Results of the PREMIER Clinical Trial. *JAMA*. 2003; 289(16): 2083–93. <https://doi.org/10.1001/jama.289.16.2083> PMID: 12709466
47. Brunner E, White I, Thorogood M, Bristow A, Curle D, Marmot M. Can dietary interventions change diet and cardiovascular risk factors: a meta-analysis of randomized controlled trials. *Am J Public Health*. 1997; 87(9): 1415–1422. PMID: 9314790
48. Francis S, Taylor M. A social marketing theory-based diet-education program for women ages 54 to 83 years improved dietary status. *J Am Diet Assoc*. 2009; 109(12): 2052–6. <https://doi.org/10.1016/j.jada.2009.09.002> PMID: 19942023

49. Parekh S, Vandelanotte C, King D, Boyle FM. Improving diet, physical activity and other lifestyle behaviours using computer-tailored advice in general practice: a randomised controlled trial. *Int J Behav Nutr Phys Act*. 2012; 9: 108. <https://doi.org/10.1186/1479-5868-9-108> PMID: 22963354
50. Petersen K, Torpy D, Chapman I, Guha S, Clifton P, Turner K, et al. Food label education does not reduce sodium intake in people with type 2 diabetes mellitus. A randomised controlled trial. *Appetite*. 2013; 68: 147–51. <https://doi.org/10.1016/j.appet.2013.04.028> PMID: 23665299
51. Kokanović A, Mandić ML, Banjari I. Does individual dietary intervention have any impact on adolescents with cardiovascular health risks? *Medicinski Glasnik*. 2014; 11(1): 234–7. PMID: 24496370
52. Heino T, Kallio K, Jokinen E, Lagström H, Seppänen R, Välimäki I, et al. Sodium intake of 1 to 5-year-old children: the STRIP project. The Special Turku Coronary Risk Factor Intervention Project. *Acta paediatrica*. 2000; 89(4): 406–10. PMID: 10830450
53. Wang J, Olendzki BC, Wedick NM, Persuitte GM, Culver AL, Li W, et al. Challenges in sodium intake reduction and meal consumption patterns among participants with metabolic syndrome in a dietary trial. *Nutrition Journal*. 2013; 12:163 <https://doi.org/10.1186/1475-2891-12-163> PMID: 24345027
54. Cobiac LJ, Vos T, Veerman JL. Cost-effectiveness of interventions to reduce dietary salt intake. *Heart*. 2010; 96(23): 1920–5. <https://doi.org/10.1136/hrt.2010.199240> PMID: 21041840
55. Cobiac LJ, Magnus A, Lim S, Barendregt JJ, Carter R, Vos T. Which interventions offer best value for money in primary prevention of cardiovascular disease? *PLoS One*. 2012; 7: e41842. <https://doi.org/10.1371/journal.pone.0041842> PMID: 22844529
56. Nghiem N, Blakely T, Cobiac LJ, Pearson AL, Wilson N. Health and economic impacts of eight different dietary salt reduction interventions. *PLoS ONE*. 2015; 10(4): e0123915. <https://doi.org/10.1371/journal.pone.0123915> PMID: 25910259
57. Cotter J, Cotter MJ, Oliveira P, Cunha P, Polónia J. Salt intake in children 10–12 years old and its modification by active working practices in a school garden. *J Hypertens*. 2013; 31(10): 1966–71. <https://doi.org/10.1097/HJH.0b013e328363572f> PMID: 24107730
58. Katz DL, Katz CS, Treu JA, Reynolds J, Njike V, Walker J, et al. Teaching healthful food choices to elementary school students and their parents: the Nutrition Detectives™ program. *J Sch Health*. 2011; 81(1): 21–8. <https://doi.org/10.1111/j.1746-1561.2010.00553.x> PMID: 21158862
59. Aldana SG, Greenlaw RL, Diehl, Salberg A, Merrill RM, Ohmine S. The effects of a worksite Chronic disease prevention program. *J Occup Environ Med*. 2005; 47: 558–564. PMID: 15951715
60. Chen J, Wu X, Gu D. Hypertension and cardiovascular diseases intervention in the capital steel and iron company and Beijing Fangshan community. *Obes Rev*. 2008; 9 Suppl 1:142–5.
61. Levin S, Ferdowsian H, Hoover V, Green AA, Barnard ND. A worksite programme significantly alters nutrient intakes. *Public Health Nutr*. 2010; 13(10): 1629–35. <https://doi.org/10.1017/S136898000999303X> PMID: 20074388
62. Yanek LR, Becker DM, Moy TF, Gittelsohn J, Koffman DM. Project Joy: faith based cardiovascular health promotion for African American women. *Public Health Rep*. 2001; 116 Suppl 1:68–81.
63. Cappuccio FP, Kerry SM, Micah FB, Plange-Rhule J, Eastwood JB. A community programme to reduce salt intake and blood pressure in Ghana [ISRCTN88789643]. *BMC Public Health*. 2006; 6: 13. <https://doi.org/10.1186/1471-2458-6-13> PMID: 16433927
64. Takahashi Y, Sasaki S, Okubo S, Hayashi M, Tsugane S. Blood pressure change in a free-living population-based dietary modification study in Japan. *J Hypertension*. 2006; 24(3): 451–8.
65. Robare JF, Milas NC, Bayles CM, Williams K, Newman AB, Lovalekar MT, et al. The key to life nutrition program: results from a community-based dietary sodium reduction trial. *Public Health Nutr*. 2010; 13(5): 606–14. <https://doi.org/10.1017/S1368980009991583> PMID: 19781124
66. Van de Vijver S, Oti S, Addo J, de Graft-Aikins A, Agyemang C. Review of community-based interventions for prevention of cardiovascular diseases in low- and middle-income countries. *Ethn Health*. 2012; 17(6): 651–76. <https://doi.org/10.1080/13557858.2012.754409> PMID: 23297746
67. Shankar B, Brambila-Macias J, Traill B, Mazzocchi M, Capacci S. An evaluation of the UK Food Standards Agency's salt campaign. *Health Econ*. 2013; 22(2): 243–50. <https://doi.org/10.1002/hec.2772> PMID: 22223605
68. Collins M, Mason H, O'Flaherty M, Guzman-Castillo M, Critchley J, Capewell S. An economic evaluation of salt reduction policies to reduce coronary heart disease in England: A policy modeling study. *Value in Health*. 2014; 17(5): 517–24. <https://doi.org/10.1016/j.jval.2014.03.1722> PMID: 25128044
69. Gillespie D, Allen K, Guzman-Castillo M, Bandosz P, Moreira P, McGill R et al. The health equity and effectiveness of policy options to reduce dietary salt intake in England: policy forecast. *PLoS ONE*. 2015; 10(7): e0127927. <https://doi.org/10.1371/journal.pone.0127927> PMID: 26131981

70. Wilcox ML, Mason H, Fouad FM, Rastam S, al Ali R, Page TF, et al. Cost-effectiveness analysis of salt reduction policies to reduce coronary heart disease in Syria, 2010–2020. *Int J Public Health*. 2015; 60: S23–30. <https://doi.org/10.1007/s00038-014-0577-3> PMID: 24972676
71. Mason H, Shoaibi A, Ghandour R, O'Flaherty M, Capewell S, Khatib R, et al. A cost effectiveness analysis of salt reduction policies to reduce coronary heart disease in four Eastern Mediterranean countries. *PLoS ONE*. 2014; 9(1): e84445. <https://doi.org/10.1371/journal.pone.0084445> PMID: 24409297
72. Babio N, Vicent P, López L, Benito A, Basulto J, Salas-Salvadó J. Adolescents' ability to select healthy food using two different front-of-pack food labels: a cross-over study. *Public Health Nutr*. 2014; 17(6): 1403–9. <https://doi.org/10.1017/S1368980013001274> PMID: 23680067
73. Elfassy T, Yi S, Eisenhower D, Lederer A, Curtis CJ. Use of sodium information on the nutrition facts label in New York city adults with hypertension. *J Acad Nutr Diet*. 2015; 115(2): 278–83. <https://doi.org/10.1016/j.jand.2014.08.027> PMID: 25441962
74. Pietinen P, Valsta LM, Hirvonen T, Sinkko H. Labelling the salt content in foods: a useful tool in reducing sodium intake in Finland. *Public Health Nutr*. 2008; 11(4): 335–40. <https://doi.org/10.1017/S1368980007000249> PMID: 17605838
75. Temme EHM, van der Voet H, Roodenburg AJC, Bulder A, van Donkersgoed G, van Klaveren J. Impact of foods with health logo on saturated fat, sodium and sugar intake of young Dutch adults. *Public Health Nutr*. 2011; 14(4): 635–44. <https://doi.org/10.1017/S1368980010002089> PMID: 20843399
76. De Menezes EW, Lopes TDVC, Mazzini ER, Dan MCT, Godoy C, Giuntini EB. Application of Choices criteria in Brazil: Impact on nutrient intake and adequacy of food products in relation to compounds associated to the risk of non-transmissible chronic diseases. *Food Chem*. 2013; 140(3): 547–52. <https://doi.org/10.1016/j.foodchem.2013.02.031> PMID: 23601405
77. Roodenburg AJC, van Ballegooijen AJ, Dötsch-Klerk M, van der Voet H, Seidell JC. Modelling of Usual Nutrient Intakes: Potential Impact of the Choices Programme on Nutrient Intakes in Young Dutch Adults. *PLoS ONE*. 2013; 8(8): e72378. <https://doi.org/10.1371/journal.pone.0072378> PMID: 24015237
78. Chang HY, Hu YW, Yue CS, Wen YW, Yeh WT, Hsu LS, et al. Effect of potassium enriched salt on cardiovascular mortality and medical expenses of elderly men. *Am J Clin Nutr*. 2006; 83: 1289–96. PMID: 16762939
79. Murray CJ, Lauer JA, Hutubessy RC, Niessen L, Tomijima N, Rodgers A, et al. Effectiveness and costs of interventions to lower systolic blood pressure and cholesterol: a global and regional analysis on reduction of cardiovascular-disease risk. *The Lancet*. 2003; 361(9359): 717–25.
80. Rubinstein A, Colantonio L, Bardach A, Caporale J, Martí SG, Kopitowski K, et al. Estimation of the burden of cardiovascular disease attributable to modifiable risk factors and cost-effectiveness analysis of preventative interventions to reduce this burden in Argentina. *BMC Public Health*. 2010; 10: 627. <https://doi.org/10.1186/1471-2458-10-627> PMID: 20961456
81. Smith-Spangler CM, Juusola JL, Enns EA, Owens DK, Garber AM. Population strategies to decrease sodium intake and the burden of cardiovascular disease: A cost-effectiveness analysis. *Ann Intern Med*. 2010; 152(8): 481–7. <https://doi.org/10.7326/0003-4819-152-8-201004200-00212> PMID: 20194225
82. Konfino J, Mekonnen TA, Coxson PG, Ferrante D, Bibbins-Domingo K. Projected impact of a sodium consumption reduction initiative in Argentina: an analysis from the CVD policy model—Argentina. *PLoS ONE*. 2013; 8(9): e73824. <https://doi.org/10.1371/journal.pone.0073824> PMID: 24040085
83. Rubinstein A, García Martí S, Souto A, Ferrante D, Augustovski F. Generalized cost-effectiveness analysis of a package of interventions to reduce cardiovascular disease in Buenos Aires, Argentina. *Cost Eff Resour Alloc*. 2009; 7(1): 10.
84. Hendriksen MAH, Hoogenveen RT, Hoekstra J, Geleijnse JM, Boshuizen HC, van Raaij JMA. Potential effect of salt reduction in processed foods on health. *Am J Clin Nutr*. 2014; 99(3): 446–53. <https://doi.org/10.3945/ajcn.113.062018> PMID: 24335058
85. Thow AM, Downs S, Jan S. A systematic review of the effectiveness of food taxes and subsidies to improve diets: understanding the recent evidence. *Nutr Rev*. 2014; 72(9): 551–65. <https://doi.org/10.1111/nure.12123> PMID: 25091552
86. Niebylski ML, Redburn KA, Duhaney T, Campbell NR. Healthy food subsidies and unhealthy food taxation: A systematic review of the evidence. *Nutrition*. 2015; 31(6): 787–95. <https://doi.org/10.1016/j.nut.2014.12.010> PMID: 25933484
87. Mhurchu CN, Eyles H, Genc M, Scarborough P, Rayner M, Mizdrak A, et al. Effects of health-related food taxes and subsidies on mortality from diet-related disease in New Zealand: An econometric-epidemiologic modelling study. *PLoS ONE*. 2015; 10(7): e0128477. <https://doi.org/10.1371/journal.pone.0128477> PMID: 26154289

88. He FJ, Brinsden HC, Macgregor GA. Salt reduction in the United Kingdom: A successful experiment in public health. *J Hum Hypertens*. 2014a; 28(6): 345–52.
89. Mozaffarian D, Ashkan A, Benowitz NL, Bittner V, Daniels SR, Franch HA, et al. Population Approaches to Improve Diet, Physical Activity, and Smoking Habits. A Scientific Statement From the American Heart Association. *Circulation*. 2012; 126(12): 1514–1563 <https://doi.org/10.1161/CIR.0b013e318260a20b> PMID: 22907934
90. He FJ, MacGregor GA. A comprehensive review on salt and health and current experience of world-wide salt reduction programmes. *J Hum Hypertens*. 2009; 23(6): 363–84. <https://doi.org/10.1038/jhh.2008.144> PMID: 19110538
91. Pietinen P, Mannisto S, Valsta LM, Sarlio-Lähteenkorva S. Nutrition policy in Finland. *Public Health Nutr*. 2010; 13(6A): 901–6. <https://doi.org/10.1017/S1368980010001072> PMID: 20513258
92. Wang G, Labarthe D. The cost-effectiveness of interventions designed to reduce sodium intake. *J Hypertens*. 2011; 29(9): 1693–9. <https://doi.org/10.1097/HJH.0b013e328349ba18> PMID: 21785366
93. Webster JL, Dunford EK, Hawkes C, Neal BC. Salt reduction initiatives around the world. *J Hypertens*. 2011; 29(6): 1043–50. <https://doi.org/10.1097/HJH.0b013e328345ed83> PMID: 21546876
94. Wang G, Bowman BA. Recent economic evaluations of interventions to prevent cardiovascular disease by reducing sodium intake. *Curr Atheroscler Rep*. 2013; 15(9): 349. <https://doi.org/10.1007/s11883-013-0349-1> PMID: 23881545
95. He FJ, Pombo-Rodrigues S, Macgregor GA. Salt reduction in England from 2003 to 2011: its relationship to blood pressure, stroke and ischaemic heart disease mortality. *BMJ Open*. 2014b; 4(4): e004549.
96. Enkhtungalag B, Batjargal J, Chimedsuren O, Tsogzolmaa B, Anderson CS, Webster J. Developing a national salt reduction strategy for Mongolia. *Cardiovasc Diagn Ther*. 2015; 5(3): 229–37. <https://doi.org/10.3978/j.issn.2223-3652.2015.04.11> PMID: 26090334
97. Luft FC, Morris CD, Weinberger MH. Compliance to a low-salt diet. *Am J Clin Nutr*. 1997; 65(2): 698S–703S.
98. Mohan S, Campbell NRC, Willis K. Effective population-wide public health interventions to promote sodium reduction. *CMAJ*. 2009; 181(9): 605–9. <https://doi.org/10.1503/cmaj.090361> PMID: 19752102
99. He FJ & MacGregor GA. Reducing population salt intake worldwide: from evidence to implementation. *Prog Cardiovasc Dis*. 2010; 52(5): 363–82. <https://doi.org/10.1016/j.pcad.2009.12.006> PMID: 20226955
100. Wyness LA, Buttriss JL, Stanner SA. Reducing the population's sodium intake: The UK Food Standards Agency's salt reduction programme. *Public Health Nutr*. 2012; 15(2): 254–61. <https://doi.org/10.1017/S1368980011000966> PMID: 21729460
101. Otsuka R, Kato Y, Imai T, Ando F, Shimokata H. Decreased salt intake in Japanese men aged 40 to 70 years and women aged 70 to 79 years: an 8-year longitudinal study. *J Am Diet Assoc*. 2011; 111(6): 844–50. <https://doi.org/10.1016/j.jada.2011.03.020> PMID: 21616196
102. Du S, Batis C, Wang H, Zhang B, Zhang J, Popkin BM. Understanding the patterns and trends of sodium intake, potassium intake, and sodium to potassium ratio and their effect on hypertension in China. *Am J Clin Nutr*. 2014; 99(2): 334–43. <https://doi.org/10.3945/ajcn.113.059121> PMID: 24257724
103. Miura K, Ando K, Tsuchihashi T, Yoshita K, Watanabe Y, Kawarazaki H, et al. [Scientific Statement] Report of the Salt Reduction Committee of the Japanese Society of Hypertension(2) Goal and strategies of dietary salt reduction in the management of hypertension. *Hypertens Res*. 2013; 36(12): 1020–5. <https://doi.org/10.1038/hr.2013.105> PMID: 24152612
104. Asaria P, Chisholm D, Mathers C, Ezzati M, Beaglehole R. Chronic disease prevention: health effects and financial costs of strategies to reduce salt intake and control tobacco use. *Lancet*. 2007; 370(9604): 2044–53. [https://doi.org/10.1016/S0140-6736\(07\)61698-5](https://doi.org/10.1016/S0140-6736(07)61698-5) PMID: 18063027
105. Dodhia H, Phillips K, Zannou M- I, Airolidi M, Bevan G. Modelling the impact on avoidable cardiovascular disease burden and costs of interventions to lower SBP in the England population. *J Hypertens*. 2012; 30(1): 217–26. <https://doi.org/10.1097/HJH.0b013e32834d86ee> PMID: 22080224
106. Gase LN, Kuo T, Dunet D, Schmidt SM, Simon PA, Fielding JE. Estimating the potential health impact and costs of implementing a local policy for food procurement to reduce the consumption of sodium in the county of Los Angeles. *Am J Public Health*. 2011; 101(8): 1501–7. <https://doi.org/10.2105/AJPH.2011.300138> PMID: 21680933
107. Ha DA, Chisholm D. Cost-effectiveness analysis of interventions to prevent cardiovascular disease in Vietnam. *Health Policy Plan*. 2011; 26(3): 210–22. <https://doi.org/10.1093/heapol/czq045> PMID: 20843878

108. Laatikainen T, Critchley J, Vartiainen E, Salomaa V, Ketonen M, Capewell S. Explaining the Decline in Coronary Heart Disease Mortality in Finland between 1982 and 1997. *Am. J. Epidemiol.*, 2005; 162: 764–773 <https://doi.org/10.1093/aje/kwi274> PMID: 16150890
109. Van Vliet BN, Campbell NRC, Canadian Hypertension Education Program. Efforts to reduce sodium intake in Canada: why, what, and when? *Can J Cardiol.* 2011; 27(4): 437–45. <https://doi.org/10.1016/j.cjca.2011.04.012> PMID: 21801976
110. Bech-Larsen T & Aschemann-Witzel J. A Macromarketing Perspective on Food Safety Regulation The Danish Journal of Macromarketing. 2012; 32(2) 208–219
111. Lloyd-Williams F, Bromley H, Orton L, Hawkes C, Taylor-Robinson D, O'Flaherty M, et al. Smorgasbord or symphony? Assessing public health nutrition policies across 30 European countries using a novel framework. *BMC Public Health.* 2014; 14: 1195. <https://doi.org/10.1186/1471-2458-14-1195> PMID: 25413832
112. MacGregor GA, He FJ, Pombo-Rodrigues S. Food and the responsibility deal: how the salt reduction strategy was derailed by Andrew Lansley and the coalition government. *Br Med J.* 2015; 350: h1936.
113. Campos S, Doxey J, Hammond D. Nutrition labels on pre-packaged foods: a systematic review. *Public Health Nutr.* 2011; 14(8): 1496–506. <https://doi.org/10.1017/S1368980010003290> PMID: 21241532
114. Cowburn G & Stockley L. Consumer understanding and use of nutrition labelling: a systematic review. *Public Health Nutr.* 2005; 8(01): 21–8.
115. Wakefield MA, Loken B, Hornik RC. Use of mass media campaigns to change health behaviour. *The Lancet.* 2010; 376:1261–1271
116. Mols F, Haslam SA, Jetten J, Steffens NK. Why a Nudge is Not Enough: A Social Identity Critique of Governance by Stealth. *Euro J Polit Res.* 2014; 54(1): 87–98.
117. Fransen ML, Smit EG, Verlegh PWJ. Strategies and motives for resistance to persuasion: an integrative framework. *Front Psychol.* 2015; 6: 1201. <https://doi.org/10.3389/fpsyg.2015.01201> PMID: 26322006
118. Forshee RA. Innovative regulatory approaches to reduce sodium consumption: could a cap-and-trade system work? *Nutr Rev.* 2008; 66(5): 280–5. <https://doi.org/10.1111/j.1753-4887.2008.00033.x> PMID: 18454814
119. Bibbins-Domingo K, Chertow GM, Coxson PG, Moran A, Lightwood JM, Pletcher MJ, et al. Projected effect of dietary salt reductions on future cardiovascular disease. *N Eng J Med.* 2010; 362(7): 590–9.
120. Barton P, Andronis L, Briggs A, McPherson K, Capewell S. Effectiveness and cost effectiveness of cardiovascular disease prevention in whole populations: Modelling study. *BMJ.* 2011; 343: d4044. <https://doi.org/10.1136/bmj.d4044> PMID: 21798967
121. Cobiaci LJ, Veerman L, Vos T. The role of cost-effectiveness analysis in developing nutrition policy. *Annu Rev Nutr.* 2013; 33: 373–93. <https://doi.org/10.1146/annurev-nutr-071812-161133> PMID: 23642205
122. Owen L, Morgan A, Fischer A, Ellis S, Hoy A, Kelly MP. The cost-effectiveness of public health interventions. *J Public Health.* 2011; 34(1): 37–45.
123. Beckmann S, Os I, Kjeldsen S, Eide I, Westheim A, Hjermann I. Effect of dietary counselling on blood pressure and arterial plasma catecholamines in primary hypertension. *Am J hypertens.* 1995; 8(7): 704–11. PMID: 7546496
124. Tian HG, Guo ZY, Hu G, Yu SJ, Sun W, Pietinen P, et al. Changes in sodium intake and blood pressure in a community-based intervention project in China. *J Hum Hypertens.* 1995; 9(12): 959–68. PMID: 8746640
125. Cappuccio FP. Salt and cardiovascular disease. *Br Med J.* 2007; 334: 859–60.
126. Rodriguez-Fernandez R, Siopa M, Simpson SJ, Amiya RM, Breda J, Cappuccio FP. Current salt reduction policies across gradients of inequality-adjusted human development in the WHO European region: minding the gaps. *Public Health Nutr.* 2014; 17: 1894–904. <https://doi.org/10.1017/S136898001300195X> PMID: 23924617
127. Fair Society, Healthy Lives. The Marmot Review. Strategic Review of Health Inequalities in England post-2010. Published by The Marmot Review. 2010;1–238.
128. Cappuccio FP, Ji C, Donfrancesco C, Palmieri L, Ippolito R, Vanuzzo D et al. Geographic and socioeconomic variation of sodium and potassium intake in Italy: results from the MINISAL-GIRCSI programme. *BMJ Open.* 2015; 5: e007467. <https://doi.org/10.1136/bmjopen-2014-007467> PMID: 26359282
129. McGill R, Anwar E, Orton L, Bromley H, Lloyd-Williams F, O'Flaherty M, et al. Are interventions to promote healthy eating equally effective for all? Systematic review of socioeconomic inequalities in impact. *BMC Public Health.* 2015; 15: 457. <https://doi.org/10.1186/s12889-015-1781-7> PMID: 25934496

130. Lorenc T, Petticrew M, Welch V, Tugwell P. What types of interventions generate inequalities? Evidence from systematic reviews. *J Epidemiol Community Health*. 2012; 67(2): 190–3. <https://doi.org/10.1136/jech-2012-201257> PMID: 22875078
131. Orton L, Lloyd-Williams F, Taylor-Robinson D, O'Flaherty M, Capewell S. The Use of Research Evidence in Public Health Decision Making Processes: Systematic Review. *PLoS ONE*. 2011; 6(7): e21704. <https://doi.org/10.1371/journal.pone.0021704> PMID: 21818262
132. Oliver K, Innvar S, Lorenc T, Woodman J, Thomas J. A systematic review of barriers to and facilitators of the use of evidence by Policymakers. *BMC Health Services Research*. 2014; 14: 2. <https://doi.org/10.1186/1472-6963-14-2> PMID: 24383766
133. McKinlay JB & Marceau LD. Upstream healthy public policy: Lessons from the battle of tobacco. *Int J Health Serv*. 2000; 30(1): 49–69. <https://doi.org/10.2190/2V5H-RHBR-FTM1-KGCF> PMID: 10707299
134. Krieger N. Proximal, Distal, and the Politics of Causation: What's Level Got to Do With It? *Am J Public Health*. 2008; 98: 221–230. <https://doi.org/10.2105/AJPH.2007.111278> PMID: 18172144