

**Original citation:**

Goodwin, Robin, Lemola, Sakari and Ben-Ezra, Menachem. (2017) Media use and insomnia after terror attacks in France. Journal of Psychiatric Research.

**Permanent WRAP URL:**

<http://wrap.warwick.ac.uk/96341>

**Copyright and reuse:**

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**Publisher's statement:**

© 2017, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

**A note on versions:**

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: [wrap@warwick.ac.uk](mailto:wrap@warwick.ac.uk)

# Accepted Manuscript

Media use and insomnia after terror attacks in France

Robin Goodwin, Sakari Lemola, Menachem Ben-Ezra

PII: S0022-3956(17)30968-8

DOI: [10.1016/j.jpsychires.2017.12.006](https://doi.org/10.1016/j.jpsychires.2017.12.006)

Reference: PIAT 3264

To appear in: *Journal of Psychiatric Research*

Received Date: 1 September 2017

Revised Date: 1 December 2017

Accepted Date: 12 December 2017

Please cite this article as: Goodwin R, Lemola S, Ben-Ezra M, Media use and insomnia after terror attacks in France, *Journal of Psychiatric Research* (2018), doi: 10.1016/j.jpsychires.2017.12.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Title Page

## Media use and insomnia after terror attacks in France

Short title: Media use and insomnia

Professor Robin Goodwin (Corresponding author)  
Department of Psychology  
Warwick University  
Coventry, UK. CV4 7AL  
[Robin.goodwin@warwick.ac.uk](mailto:Robin.goodwin@warwick.ac.uk)  
Tel: +44 2476 522484

Dr Sakari Lemola  
Department of Psychology  
Warwick University  
Coventry, UK. CV4 7AL  
[S.Lemola@warwick.ac.uk](mailto:S.Lemola@warwick.ac.uk)

Professor Menachem Ben-Ezra  
School of Social Work,  
Ariel University, Ariel, 40700, Israel  
[menbe@ariel.ac.il](mailto:menbe@ariel.ac.il)

## Media use and insomnia after terror attacks in France

## Abstract

Direct exposure to traumatic events often precipitates sleep disorders. Sleep disturbance has also been observed amongst those indirectly exposed to trauma, via mass media. However, previous work has however focused on traditional media use, rather than contemporary social media. We tested associations between both traditional and social media consumption and insomnia symptoms following 2015 terror attacks in Paris France, controlling for location and post-traumatic symptomology. 1878 respondents, selected to represent the national French population, completed an internet survey a month after the Bataclan attacks (response rate 72%). Respondents indicated different media use, post-traumatic stress and insomnia. Controlling for demographics, location and PTSD, insomnia was associated with both traditional ( $\beta .10$   $P=.001$ ) and social media use ( $\beta .12$   $P=.001$ ). Associations between social media and insomnia were independent of traditional media use. Interventions targeted at social media may be particularly important following mass trauma.

Key words: Anxiety, Stress Disorders, Post-Traumatic, Terrorism, sleep

## Media use and insomnia after terror attacks in France

### Introduction

A large body of evidence suggest that stress, and in particular, stress from direct exposure to trauma, is a precipitating factor for sleep disturbance (Bui et al, 2012; Harvey et al, 2003; Sinha, 2016). Studies with war veterans (Lewis et al, 2009) and individuals who experienced missile (Lavie, 2001), terror attacks (Galea et al, 2002, North et al, 199) and natural disasters (Bui et al, 2012, McMillen et al, 2000) found increased difficulty in initiating sleep, returning to sleep after awakening, nightmares, and a general hyperarousal following traumatic events (Germain, 2013; Lavie, 2002; Sinha, 2016). A smaller literature has considered the impact of trauma on sleep amongst those indirectly exposed, primarily through the media. This includes associations between watching TV and dreams (following the Challenger disaster (Terr et al, 1999) , and after 9/11 (Propper et al, 2007)), terror-related TV consumption and general sleep difficulties following missile fire in Israel (Soffer-Dudek & Shahar, 2010), and reading internet news and disruptive nocturnal behaviour after the 2011 Great Japan earthquake (Bui et al, 2012). Most of this work has focused on engagement with traditional media; however, research following natural disaster suggests the immediacy and personalization offered by social media may have a greater psychological toll (Goodwin et al, 2013; 2017). As yet, the impact of these social media on sleep disturbance following trauma has been underexplored.

In November 2015 gunmen claiming allegiance to the Islamic State group attacked the Bataclan concert hall and restaurants in St. Denis, Paris, killing 130. Using data from a national survey (Goodwin et al, 2017), we consider the associations between traditional and social

media use following the attack and subsequent insomnia. Because location (physical proximity to attack) has been positively associated with distress (Goodwin et al, 2017) we control for location within vs. outside of Paris. While the relationship between mass trauma and PTSD is complex, insomnia has been associated with acute psychological distress (Sinha, 2016; Germain, 2013, Krakow et al, 2015; Schoenfeld et al, 2012) primarily via the hyperarousal component of PTSD. We therefore report both associations between media use and insomnia, and media use and insomnia, controlling for the impact of post-traumatic symptomology on insomnia. This allowed us to assess the unique association of media use on insomnia, beyond the impact of distress.

#### *Method*

A major survey organization (<https://www.surveyygo.com/>) operates with a network of international panel surveys. We drew participants from their large French panel (approx. 500 000 individuals). Potential participants were sent a web link four weeks after the attacks (week of December 13, 2015) using validated addresses. Participants were selected using random stratified sampling, using weights for key demographic elements (age, gender), to create a nationally approximate representative sample (Supplement 1). 2612 participants were sent the web link; a validation question removed 1.3% respondents, leaving 1878 who fully participated (response rate 72%; Mean age 41.1 [SD 10.8], Median age 43, age range 18-60, 54% female). This compares to a national median age of 41.2 for France in 2015, with 52% female (Statista, n.d.). Surveys took approximately 30 minutes to complete, with participants given account credit for participation which could be redeemed as cash payment. The first survey item required

informed consent to continue. Approval was from the Institutional Review Boards at Ariel University and Warwick University.

### *Measures*

*Insomnia* was assessed using the Bergen insomnia scale (Pallesen et al. 2008), a six-item, seven-point scale assessing how many days per week, over the previous month, participants had problems with sleep onset, maintenance, early morning waking, non-restorative sleep, daytime impairment and dissatisfaction with current sleep ( $\alpha = .89$ ). *PTSD* was measured using the recently proposed ICD-11 scale (Cloitre et al, 2013). This scale comprises six items on a five-point scale ( $\alpha = .91$ ). Questions asked participants how often they had felt nervous, hopeless, restless, depressed, worthless or that everything was an effort since the attacks (*none of the time to all the time*). Scales were translated to French by a French specialist in public health. To assess *media use*, respondents indicated number of hours they used each of six media sources in the weekend after the Paris attacks (Friday evening, Saturday and Sunday, 50 hours in total). Nine participants (0.5%), reporting > 50 hours, were excluded. Media use was subsequently coded into 'traditional' media (TV, Radio, Newspaper) and 'social' media use (Twitter, Facebook, Youtube).

### *Analyses*

Preliminary statistical analyses report frequencies of media types used and associations of demographic background variables with different types of media use, PTSD symptoms, and insomnia symptoms including Pearson's Correlations and t-test. A priori multi-collinearity tests

checked multi-collinearity amongst independent variables (Supplementary Table 2). Linear regressions tested associations between media use and insomnia, controlling for age, gender, location (Paris or elsewhere) and PTSD symptoms. All analyses were conducted with SPSS Statistics v. 22.

### *Results*

In the weekend after the attacks TV was the most widely used media, followed by radio and Facebook (Table 1 and Table S2). There were small positive correlations between distress and all forms of media use (Supplementary Table 2). Female respondents reported higher levels of insomnia ( $M_s$  19.08 vs. 14.46,  $t(1872) = 8.38$ ,  $P = .001$ ), but there was no significant correlation between age and insomnia ( $r(1877) = -.04$ ,  $P = .054$ ).

### *Media use and insomnia*

Figure 1 maps association of each individual media with insomnia, controlling for demographics and PTSD. Associations between media use and insomnia were strongest for the use of TV, Facebook, Youtube and Twitter.

We formed media into two larger groupings: Traditional (TV, Radio, Newspapers) and Social (Facebook, Twitter, YouTube). Table 2 presents associations between demographics, media group use and insomnia, first without then with control for PTSD symptoms. Use of both traditional media and social media were significantly associated with insomnia, even when controlling for age and sex, location and PTSD symptomatology.



*Additive contribution of social media in associations with insomnia.*

Use of social and traditional media were moderately correlated ( $r(1578) = .16$ ,  $p = .001$ ). Table 3 shows the additive effect for social media usage on insomnia, beyond the inclusion of traditional media. Social media had an additional association with insomnia, beyond traditional media use.

## *Discussion*

Research on insomnia following traumatic events has focused on direct exposure to such events (Sinha, 2016). Media consumption, however, may prove significant for the national dispersal of stress beyond those directly impacted. In our data, media use in the immediate aftermath of attacks was significantly associated with insomnia a month after the event. Further, particular types of social media were more strongly associated with insomnia than traditional media formats. This association occurred independently of concomitant PTSD symptoms.

Traumatic events stimulate a fear and stress response that stimulates the hyperarousal that underlies chronic insomnia. Mass media can prime individuals to be more fearful (Ridout et al, 2008), and think more about their mortality (Das et al, 2009). This is only poorly recognised within the current DSM-5 PTSD criteria, which does not include indirect exposure via such mass media. PTSD was higher amongst those Americans who watched more anniversary television a year after the 9/11 attacks (Bernstein et al, 2007). 'Media amplification' can explain stress responses away from the 'bulls-eye' of any event, and may be more strongly associated with stress than direct exposure. In our data exposure (as approximated through location within vs.

outside Paris) failed to have a significant additive effect on insomnia. Further, media use had an impact on psychological distress, controlling for insomnia (Supplementary Table 3), underlining the influence of this distal factor.

‘Emotional contagion’ between individuals can occur rapidly in social networks (Coviello et al, 2014; Monfort & Afzali, 2015). To the extent to which our anxiety is influenced by the anxiety of others, our results may help explain some of the adverse psychological impact of ‘second hand’ exposure to events. However, it is only recently that social media has been recognised as a particular risk following trauma (Goodwin et al, 2013). While there was some evidence of peritraumatic responses mediating the association between consumption of internet use and disrupted nocturnal behaviour following the Great Japan Earthquake (Bui et al, 2012), social media platforms such as Twitter and Facebook present a more personal and interactive media, with additional impact on an individual’s risk assessment (Lemyre et al, 2010). Social media may offer different interpretations of events to that offered by traditional media, questioning ‘official’ interpretations and suggesting alternative views on event severity and its consequences (Friedman, 2013). Social media may also be harder to avoid than traditional media, given its tendency to regularly update on personal smartphones. Previous work has focused on the impact of social media on psychological distress; we go one step further to show the impact of media use on insomnia, independent of post-traumatic response. While PTSD and insomnia may often be comorbid, this media use had an additional toll on mental well-being independent of post-traumatic stress. This suggests the need for a ‘safe space’ application, allowing individuals to filter out potentially negative information across

platforms. Such interventions could also be extended to digital televisions, with TV use also significantly associated with insomnia symptoms.

Sleep disorders after trauma present a particular challenge for clinicians, with problems often persisting after PTSD has been successfully treated (Sinha, 2016; Schoenfeld et al, 2012). While sleep disturbance is considered to be a core symptom of PTSD, insomnia symptoms often also develop in the absence of full-blown PTSD (Sinha, 2016). Sleep disturbance in the aftermath of a traumatic event often precedes the development of PTSD and predict later PTSD-symptom severity (McLay et al, 2010). Therefore, it is possible that disturbed sleep acts as a vulnerability factor for the development of clinical PTSD. While CBT has had some success in improving sleep, there is need for further research into additional interventions to address trauma-associated sleep disturbance (Schoenfeld et al, 2012).

We recognise a number of limitations. Our study had a cross-sectional design, meaning that could not assess sleep pre- and post- trauma to measure predisposing factors for later stress response (Bui et al, 2012; Krakow et al, 2015), or assess whether individuals with pre-existing insomnia symptoms were more prone for media consumption. Moreover, we cannot rule out a general impact of media use on sleep also involving effects of watching screens at night on sleep. Electronic media use in bed at night is associated with sleep disturbance (Lemola et al, 2015), with exposure to the particular spectral composition of light from flat-screens having an alerting effect (Cajochen et al, 2011) that might underlie sleep disturbances. Respondents completed self-reports, which were potentially subject to recall bias, and we lack the clinical interviews necessary to verify diagnoses for either of the two symptom groups. Because of the nature of the survey panel used we cannot be certain that the 27% who did not

respond did not differ from our respondents. Effect sizes were small, although in line with others in the literature. Past and present levels of trauma exposure and emotional regulation strategies could also be added as predictors of psychological distress and its consequences (Monfort & Afzali, 2017). The relationship between exposure to mass trauma and PTSD is complex, and other measures of psychological disorder could have been also included. As with other work on the mental health impact of social media use (Cajochen et al, 2011) we need to better understand the impact of quality as well as quantity of social media use on trauma responses and the networks within it is spread (Monfort & Afzali, 2017), particularly given the dynamic and interactive nature of so many of these media. This is likely to be particularly significant when addressing the rumours and 'fake news' that often accompanies threatening events.

Understanding the role of mass media in insomnia is likely to have significant clinical implications. Insomnia is an important precursor for posttraumatic conditions, including PTSD, depression or substance dependence (Sinha, 2016). Our work contributes to growing evidence alerting us to the potential problems associated with social media use. Such use may also detract from the formation of other relationships that might enable individuals to better deal with stressful life events (Shahya & Christakis, 2017), including the mental health challenges posed by mass trauma. Findings suggest the need for therapeutic interventions that target social media use, particularly amongst those most vulnerable to distress.

196

197

198 Authors' contributions

199 RG and MBE designed the study, obtained funding and organised data collection in France. RG  
200 and SL analysed the data; all authors drafted the initial paper and commented on and approved  
201 the final version. We would like to thank Dr. Anne-Cécile Hoyez for her help in translating the  
202 inventories, and for helpful comments from two anonymous reviewers.

203

## References

- Bastien CH, Vallieres A, Morin CM. 2004. Precipitating factors of insomnia. *Behav Sleep Med.* 2(1), 50-62.
- Bernstein KT, Ahern J, Tracy M, Boscarino JA, Vlahov D, Galea S. 2007. Television watching and risk of incident probable posttraumatic stress disorder: A prospective evaluation. *J Nerv Ment.* 195, 41-47
- Bonanno GA, Brewin CR, Kaniasty K, La Greca AM. 2010. Weighing the costs of disaster: consequences, risks, and resilience in individuals, families, and communities. *Psychol Sci Public Interest.* 11, 1-49
- Bui E, Rodgers RF, Herbert C, Franko DL, Simon N M, Birmes P, Brunet A. 2012. The impact of internet coverage of the March 2011 Japan earthquake on sleep and posttraumatic stress symptoms: an international perspective. *Am J Psychiatry.* 169, 221-222.
- Cajochen C, Frey S, Anders D, Spät, J, Bues M, Pross A, Mager, R, Wirz-Justice A, Stefani Q. 2011. Evening exposure to a light-emitting diodes (LED)-backlit computer screen affects circadian physiology and cognitive performance. *J Appl Physiol.* 110(5), 1432-1438
- Cloitre M, Garvert DW, Brewin CR, Bryant RA, Maercker, A. 2013. Evidence for proposed ICD-11 PTSD and complex PTSD: a latent profile analysis. *Eur J Psychotraumatol.* 15. doi: 10.3402/ejpt.v4i0.207006
- Coviello L, Sohn Y, Kramer ADI, Marlow C, Franceschetti M, Christakis NA, Fowler, JH. 2014. Detecting Emotional Contagion in Massive Social Networks. *PLoS One.* 9(3): e90315. doi:10.1371/journal.pone.0090315

- 226 Das E, Bushman BJ, Bezemer MD, Kerkhof P, Vermeulen IE. 2009. How terrorism news reports  
227 increase prejudice against outgroups: A terror management account. *J Exp Soc Psychol.*  
228 45(3), 453–459
- 229 Friedman S. Three Mile Island, 2011. Chernobyl and Fukushima: an analysis of traditional  
230 and new media coverage of nuclear accidents and radiation. *Bull At Sci.* 67, 55–65.
- 231 Galea S, Resnick H, Ahern J, Gold J, Bucuvalas M, Kilpatrick D, Stuber J, Vlahov D. 2002.  
232 Posttraumatic stress disorder in Manhattan, New York City, after the September 11th  
233 terrorist attacks. *J Urban Health.* 79(3),340-53.
- 234 Germain A. 2013. Sleep disturbances as the hallmark of PTSD: where are we now? *Am*  
235 *J Psychiatry.* 170(4), 372-382.
- 236 Goodwin R, Palgi Y, Hamama-Raz Y, Ben-Ezra M. 2013. In the eye of the storm or the bullseye of  
237 the media: Social media use during Hurricane Sandy as a predictor of post-traumatic  
238 stress. *J Psychiatr Res.* 47(8), 1099-1100.
- 239 Goodwin R, Kaniasty, K, Sun S, Ben-Ezra, M. 2017. Psychological distress and prejudice following  
240 terror attacks in France. *J Psychiatr Res.* 91, 111-15.
- 241 Harvey AG, Jones C, Schmidt DA. 2003. Sleep and posttraumatic stress disorder: a review. *Clin*  
242 *Psychol Rev.* 23, 377-407.
- 243 Healy ES, Kales A, Monroe L J, Bixler EO, Chamberlin K, Soldatos, CR. 1981. Onset of  
244 insomnia: Role of life-stress events. *Psychosom Med.* 43, 439–451.
- 245 Krakow BJ, Ulibarri VA, Moore BA, McIver ND. 2015. Posttraumatic stress disorder and sleep-  
246 disordered breathing: a review of comorbidity research. *Sleep Med Rev.* 24, 37-45.

- Lavie P. 2001. Sleep disturbances in the wake of traumatic events. *N Engl J Med.* 345(25), 1825-32.
- Lemola S, Perkinson-Gloor N, Brand S, Dewald-Kaufmann JF, Grob A. 2015. Adolescents' electronic media use at night, sleep disturbance, and depressive symptoms in the smartphone age. *J Youth Adolesc.* 44(2), 405-418.
- Lemyre L, Johnson C, Corneil W. 2010. Psychosocial considerations for mass decontamination. *Radiat Prot Dosimetry.* 142, 17–23.
- Lewis V, Creamer M, Failla S. 2009. Is poor sleep in veterans a function of post- traumatic stress disorder? *Mil Med.* 174(9), 948-51.
- McLay RN, Klam WP, Volert SL. 2010. Insomnia is the most commonly reported symptom and predicts other symptoms of post-traumatic stress disorder in U.S. service members returning from military deployments. *Mil Med.* 175(10), 759-62.
- McMillen JC, North CS, Smith EM. 2000. What parts of PTSD are normal: intrusion, avoidance, or arousal? Data from the Northridge, California, earthquake. *J Trauma Stress.* 13(1), 57-75.
- Monfort E, Afzali M. 2017. Traumatic stress symptoms after the November 13th 2015 Terrorist Attacks among Young Adults: The relation to media and emotion regulation. *Comp Psychiat.* 75, 68-74.
- North CS, Nixon SJ, Shariat S, Mallonee S, McMillen JC, Spitznagel EL, Smith EM. 1999. Psychiatric disorders among survivors of the Oklahoma City bombing. *JAMA.* 282(8), 755-62.
- Pallesen S, Bjorvatn B, Nordhus IH, Sivertsen B, Hjørnevik M, Morin CM. 2008. A new scale for



- measuring insomnia: the Bergen Insomnia Scale. *Percept Mot Skills*. 107(3), 691-706.
- Propper RE, Stickgold R, Keeley R, Christman SD. 2007. Is television traumatic? Dreams, stress, and media exposure in the aftermath of September 11, 2001. *Psychol Sci*. 18(4), 334-340.
- Ridout TN, Grosse AC, Appleton AM. 2008. News Media Use and Americans' Perceptions of Global Threat. *Br J Polit Sci*. 38(4), 575-593
- Schoenfeld FB, DeViva JC, Manber R. 2012. Treatment of sleep disturbances in posttraumatic stress disorder. *J Rehabil Res Dev*. 49, 729-52.
- Shakya HB, Christakis NA. Association of Facebook Use With Compromised Well-Being: A Longitudinal Study. 2017. *Am J Epidemiol*. 185(3), 203-211. doi: 10.1093/aje/kww189.
- Sinha SS. Trauma-induced insomnia: A novel model for trauma and sleep research. 2016. *Sleep Med Rev*; 25, 74-83.
- Soffer-Dudek N, Shahar G. 2010. Effect of exposure to terrorism on sleep-related experiences in Israeli young adults. *Psychiatry*. 73(3), 264-276
- Statista, 2016. Age and sex distribution of France, 2015.  
<https://www.statista.com/statistics/275391/median-age-of-the-population-in-france/> (accessed 15th November 2017)
- Terr LC, Bloch DA, Michel BA, Shi H, Reinhardt JA, Metayer S. 1999. Children's symptoms in the wake of Challenger: a field study of distant-traumatic effects and an outline of related conditions. *Am J Psychiatry*. 156(10), 1536-44.

290 Legend for figure:

291 Figure 1: Associations between individual media use and insomnia, controlling for age,  
292 sex, location and PTSD (Standardised Betas).

293

	<b>Mean</b>	<b>SD</b>
TV	18.54	14.94
Radio	4.98	7.79
Newspaper	1.98	4.53
Facebook	4.59	8.33
Twitter	0.99	4.50
Youtube	1.40	4.20

Table 1: Reported media use in the fifty hours following Bataclan attacks.

	Traditional media				Social media			
	Without PTSD		Controlling PTSD		Without PTSD		Controlling PTSD	
R <sup>2</sup>	.06		.18		.05		.17	
	Beta	T (sig)	Beta	T (sig)	Beta	T (sig)	Beta	T (sig)
Age	-.08	-3.07**	-.03	-1.48	-.00	-.03	.02	.94
Sex	.17	7.26***	.15	6.81***	.18	7.21***	.15	6.65***
Location	-.01	-.52	-.04	-1.95	-.01	-.41	-.04	-1.77
PTSD	-	-	.36	15.59***	-	-	.35	15.25***
Media	.15	6.26***	.10	4.31***	.15	5.73***	.12	4.99***

Table 2: Media and insomnia, controlling for age, sex, location and PTSD symptoms: Linear regressions. Data for sex and location was dummy coded, so that higher score indicates female, Paris location.

	Beta	T	P
<b>Model 1: F change (4, 1573) = 76.83 p=.001, R<sup>2</sup> .16</b>			
Age	-.01	-.36	.72
Sex	.15	6.64	.001
Location	-.04	-1.57	.12
PTSD	.36	15.60	.001
<b>Model 2: F change (1, 1572) = 20.33 p=.001, R<sup>2</sup> .17</b>			
Age	-.03	-1.23	.22
Sex	.15	6.50	.001
Location	-.04	-1.78	.08
PTSD	.35	14.81	.001
Traditional media	.11	4.51	.001
<b>Model 3: F change (1, 1571) = 16.16 p=.001, R<sup>2</sup> .18</b>			
Age	.00	-.00	1.00
Sex	.15	6.37	.001
Location	-.04	-1.84	.07
PTSD	.34	14.67	.001
Traditional media	.09	3.62	.001
Social media	.10	4.02	.001

Table 3. Association between social media and insomnia, controlling for traditional media use, PTSD, and demographics




Figure 1: Associations between individual media use and insomnia, controlling for age, sex, location and PTSD (Standardised Betas).

Conflict of interest: None

ACCEPTED MANUSCRIPT