

Original citation:

Hardy, Adam, Schreiber, Matthias R., Parsons, Steven G., Caceres, Claudio, Brinkworth, Carolyn, Veras, Dimitri, Gaensicke, B. T., Marsh, T. R. and Cieza, Lucas. (2016) The detection of dust around NN Ser. *Monthly Notices of the Royal Astronomical Society*, 459 (4). pp. 4518-4526

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/79470>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

This article has been accepted for publication in *Monthly Notices of the Royal Astronomical Society* © 2016 The Authors Published by Oxford University Press on behalf of the Royal Astronomical Society. All rights reserved.

Link to final published version: <http://dx.doi.org/10.1093/mnras/stw976>

A note on versions:

The version presented in WRAP is the published version or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

The detection of dust around NN Ser

Adam Hardy,^{1,2*} Matthias R. Schreiber,^{1,2} Steven G. Parsons,¹ Claudio Caceres,^{1,2}
Carolyn Brinkworth,^{3,4} Dimitri Veras,⁵ Boris T. Gänsicke,⁵ Thomas R. Marsh⁵
and Lucas Cieza^{2,6}

¹*Instituto de Física y Astronomía, Universidad de Valparaíso, Av. Gran Bretaña 1111, Valparaíso, Chile*

²*Millennium Nucleus ‘Protoplanetary Discs in ALMA Early Science’, Universidad de Valparaíso, Casilla 36-D, Santiago, Chile*

³*Spitzer Science Center, IPAC, Caltech, Pasadena, CA 91125, USA*

⁴*National Center for Atmospheric Research, Boulder, CO 80301, USA*

⁵*Department of Physics, University of Warwick, Coventry CV4 7AL, UK*

⁶*Núcleo de Astronomía de la Facultad de Ingeniería, Universidad Diego Portales, Av. Ejército 441, Santiago, Chile*

Accepted 2016 April 20. Received 2016 April 19; in original form 2016 March 16

ABSTRACT

Eclipse timing variations observed from the post-common-envelope binary (PCEB) NN Ser offer strong evidence in favour of circumbinary planets existing around PCEBs. If real, these planets may be accompanied by a disc of dust. We here present the ALMA detection of flux at 1.3 mm from NN Ser, which is likely due to thermal emission from a dust disc of mass $\sim 0.8 \pm 0.2 M_{\oplus}$. We performed simulations of the history of NN Ser to determine possible origins of this dust, and conclude that the most likely origin is, in fact, common-envelope material which was not expelled from the system and instead formed a circumbinary disc. These discs have been predicted by theory but previously remained undetected. While the presence of this dust does not prove the existence of planets around NN Ser, it adds credibility to the possibility of planets forming from common-envelope material in a ‘second-generation’ scenario.

Key words: protoplanetary discs – binaries: close – binaries: eclipsing.

1 INTRODUCTION

Post-common-envelope binaries (PCEBs) are among the most peculiar binary stars in existence. They contain at least one compact object (a white dwarf, neutron star or black hole) and a companion that orbits at very low separations. These separations are so low that the stars could not always have been in this configuration, as the compact object progenitor would have completely engulfed its companion when on its giant branch. Instead, the star’s history was likely one of a main-sequence binary with separations of the order of 1 au. The evolution of the more massive star on to the giant branch would have then caused dynamically unstable mass transfer on to its companion, resulting in a common-envelope (CE) of material. The CE surrounds the future compact object and the companion but due to drag forces within the CE, orbital energy is extracted from the binary causing the two stars to spiral inwards. When enough energy is transferred to the envelope, it will then be expelled, leaving the compact object and companion star that we can observe today (e.g. Paczynski 1976; Webbink 1984; Zorotovic et al. 2010).

Although there is general agreement that the scenario outlined above describes the basics of PCEB formation, many questions regarding PCEBs still remain. One such question which has gained

considerable attention in recent years is what effect the violent evolution of these stars might have on any circumbinary material, including planets or brown dwarfs. This is particularly relevant, as many PCEBs display a potential signature of substellar objects in orbit: almost all eclipsing PCEBs display variations in their measured eclipse timings (Zorotovic & Schreiber 2013), and these eclipse timing variations (ETVs) have been attributed to circumbinary objects periodically moving the centre of mass of the binary system (e.g. Guinan & Ribas 2001). If these ETVs are indeed due to circumbinary objects, the question emerges of whether these are first-generation objects which survived the CE evolution, or instead formed afterwards in a ‘second-generation’ scenario (Schleicher & Dreizler 2014; Völschow, Banerjee & Hsman 2014). Both scenarios face difficulties – the first-generation scenario has trouble in keeping progenitor planets in stable orbits throughout the CE evolution (Mustill et al. 2013), implying that not many first-generation planets are expected to survive. The fact that ETVs have been detected around almost all eclipsing PCEBs instead suggests that these planets must commonly form from material remaining after envelope expulsion (Zorotovic & Schreiber 2013). This second-generation scenario, however, faces its own challenges. For example, a study by Bear & Soker (2014) found that forming second-generation planets around many PCEBs would require a very efficient planet-making process, in which more than 20 per cent of the disc material goes into planets.

* E-mail: adam.hardy@postgrado.uv.cl

It is clear that there is some uncertainty surrounding how the potential planets around PCEBs formed, but there is also an ongoing discussion on whether these planets even exist. Suggestions that the planetary interpretation is incorrect include the observation that the majority of planetary models used to explain the ETVs are unstable (e.g. Horner et al. 2012, 2013) or fail drastically when confronted with more recent eclipse timing measurements (Parsons et al. 2010b; Bours et al. 2014). Further evidence comes from recent observations of the PCEB V471 Tau, in which direct imaging was carried out to search for the predicted companion (Hardy et al. 2015b). V471 Tau was the first eclipsing PCEB discovered, and therefore has a long baseline of timing measurements clearly showing ETVs (Nelson & Young 1970; Lohsen 1974). These data allowed accurate prediction of the brown dwarf's brightness and separation from the PCEB, but despite these parameters being within the capability of the direct imaging observations, no brown dwarf was detected. It is therefore highly likely that another mechanism is causing the ETVs seen in V471 Tau. The exact nature of this mechanism is uncertain, but one possibility in the case of V471 Tau is the Applegate mechanism, in which the ETVs are prescribed to periodic changes in the magnetic field of the main-sequence companion (Applegate 1992).

Whilst non-planetary explanations for ETVs in PCEBs have been gaining some ground, one PCEB which so far remains robust against the above criticisms is NN Ser. NN Ser is a relatively young PCEB with a white dwarf age of ~ 1.3 Myr (Schreiber & Gänsicke 2003). The companion to the white dwarf is an M4 type star orbiting with a period of 0.13 d (Brinkworth et al. 2006; Parsons et al. 2010a), and NN Ser displays ETVs that are well fitted by two planetary mass bodies (Beuermann et al. 2010). Unlike many other PCEBs, this planetary model has correctly predicted more recent timing measurements, and no other mechanism has yet been proposed to explain its behaviour (Beuermann, Dreizler & Hessman 2013; Marsh et al. 2014; Parsons et al. 2014). As such, NN Ser is perhaps the best PCEB with which to further test the planetary hypothesis. Its youth means that if the second-generation planetary scenario is correct, it is possible that NN Ser still possess protoplanetary disc material. If this protoplanetary disc material is already dissipated, or if the first-generation formation scenario is correct, it is further possible that the planets would be present alongside a debris disc as observed in many systems.

To test this hypothesis, we searched for dust remaining after the CE around NN Ser. We used *SOFI* and *Spitzer* to search for hot dust close to the star, and Atacama Large Millimetre/submillimetre Array (ALMA) to probe for cool dust farther out. While we detect no conclusive excess emission at IR wavelengths, we clearly detect excess emission from NN Ser with ALMA which is likely due to thermal emission from a belt of cold dust. We conclude that this dust is likely a circumbinary disc, formed of material left over from the CE.

2 OBSERVATIONS

In close binaries such as NN Ser, the main-sequence star is prone to reflection effects from the white dwarf, resulting in an excess above the stellar photosphere. This reflection effect, coupled with the systems eclipsing nature, means that the emission of NN Ser is phase-dependent. As a result, when searching for excess emission above the stellar photosphere, it is crucial that all data be taken at the same phase. In our observations, we chose this phase to be just after the end of the white dwarf eclipse by the main-sequence star. This phase was chosen as the heated face of the main-sequence star will be pointed away from us, minimizing the reflection effect. In

addition to the new data presented here, we also took the optical data from Parsons et al. (2010a), and calculated the emission at our chosen phase to give us consistent data at shorter wavelengths.

2.1 SOFI observations

J, *H* and *K* band observations of NN Ser were obtained with the Son of Isaac instrument (*SOFI*) (Moorwood, Cuby & Lidman 1998) mounted on the New Technology Telescope (NTT) at La Silla Observatory, Chile in 2010–04. The observations were made in fast photometry mode and covered almost an entire binary orbit in the *J* band, half an orbit in the *H* band and the eclipse of the white dwarf in the *Ks* band. We windowed the detector to achieve a cycle time of 10 s and offset the telescope every 10 min in order to improve sky subtraction. The *J*-band observations were slightly affected by thin clouds.

The dark current removal and flat-fielding were performed in the standard way. Sky subtraction was achieved by using observations of the sky when the target had been offset. The average sky level was then added back so that we could determine the source flux and its uncertainty with standard aperture photometry, using a variable aperture, within the ULTRACAM pipeline (Dhillon et al. 2007). A comparison star was used to account for variations in observing conditions and to flux calibrate the data by using its 2MASS magnitudes (Skrutskie et al. 2006).

2.2 Spitzer observations

The *Spitzer* data were obtained during Cycle 3, as part of programme 30070. Data were taken with the Infrared Array Camera (IRAC; see Fazio et al. 2004) with 25×100 s dithers in all four channels (3.6–8.0 μm). The data reduction for all four channels was carried out on the Corrected Basic Calibrated Data frames (CBCDs) downloaded from the *Spitzer* archive. The CBCDs were overlap-corrected and combined with the standard *Spitzer* Science Center (SSC) software MOsaic and Point Source EXtraction (*MOPEX*) (Makovoz et al. 2006), using dual outlier rejection, to create a single mosaicked image for each channel. Rejected frames were noted, and the mosaic was used to identify the position of the target. No further analysis was carried out on the mosaics.

The original downloaded CBCDs were corrected for array location dependence using the correction frames provided by the SSC. Aperture photometry was carried out on the corrected CBCD frames using an aperture radius of 3 pixels and a sky subtraction annulus from 12 to 20 pixels.

The photometry was converted from MJy sr^{-1} to mJy and aperture corrected using the standard aperture corrections provided by the SSC. No pixel phase correction was applied due to the averaging effect of the dither pattern. Additionally, no colour correction was applied since we quote the isophotal wavelengths, thereby reducing the colour dependency of the flux calibration to negligible within our uncertainties.

We rejected photometry from CBCDs that were flagged by the *MOPEX* Dual Outlier Rejection algorithm during the initial mosaic process. The quoted flux densities for each channel are the unweighted mean of the photometry from the remaining frames. The uncertainties were estimated from the rms scatter on the photometry from the individual frames, divided by the square root of the number of frames. The quoted uncertainties are either our calculated uncertainty or the IRAC instrument calibration uncertainty (Reach et al. 2005), whichever was larger.




Figure 1. ALMA image of NN Ser at 1.3 mm generated with the CLEAN algorithm using natural weighting. The emission is unresolved, but the beam size (the white ellipse) confines the emission to within 1000 au of NN Ser.

The *Spitzer* observations did not cover the orbital phase used in the optical and NIR observations. We therefore used the light curve model from Parsons et al. (2010a) and adjusted the output wavelength to match the *Spitzer* bands. We then fitted the *Spitzer* data, keeping all the parameters fixed at the same values as in Parsons et al. (2010a) except the temperature of the M star, which we allowed to vary in order to fit the amplitude of the reflection effect. We used this model to determine the brightness of the system in the *Spitzer* bands at the correct phase.

2.3 ALMA observations

The ALMA observations were conducted on 2014-04-30 and repeated on the 2015-01-22, as the rms sensitivity of the data did not reach the requested value of 25 μ J in the first instance. As no flux was detected in the first observation, we were unable to accurately account for the ALMA pointing error and line-up the different data sets. We therefore focus on the second measurement set only. The single continuum mode in Band 6 was used, implying a total bandwidth of 7.5 GHz with individual channel widths of 15.625 MHz. 39 antennas were used, with minimum and maximum baselines of 15.1 and 348.5 m, respectively.

The calibration sources associated with these observations were J1337–1257 for band-pass calibration and J1550+0527 for phase calibrations. The observations consisted of five scans of 6.87 min each, translating to a total time on the science target of 34.35 min.

Standard calibration steps were applied to the data, and the resulting visibilities were deconvolved using the CLEAN algorithm with natural weighting to create the final image (Fig. 1). To obtain the total flux at 1.3 mm, a point-source fit to the visibilities was performed using the UVMODELFIT task in CASAPY version 4.2.2, resulting in a flux value of 0.11 ± 0.03 mJy. As the emission at this wavelength is likely due to thermal emission from dust at large orbital radii (see Section 3.2), the orbital phase of the PCEB should have no effect and was not taken into account.

2.4 SED

The SED of NN Ser (Table 1) is shown in Fig. 2, along with models of a 60 000 K white dwarf (Koester 2010) and an M4-type

Table 1. SED data for NN Ser. (1) Parsons et al. (2010a). (2) This work.

Wavelength (μ m)	Flux (mJy)	σ_{Flux}	Ref.
0.154	4.16	0.01	(1)
0.227	2.59	0.01	(1)
0.354	1.20	0.01	(1)
0.474	1.011	0.003	(1)
0.621	0.748	0.004	(1)
0.758	0.604	0.004	(1)
1.23	0.43	0.09	(2)
1.64	0.3	0.1	(2)
2.15	0.2	0.1	(2)
3.56	0.117	0.005	(2)
4.50	0.088	0.005	(2)
5.74	0.056	0.011	(2)
7.87	0.039	0.022	(2)
1300	0.11	0.03	(2)




Figure 2. Spectral energy distribution of NN Ser at the end of the white dwarf eclipse. The red lines denote fluxes from a model 60 000 K white dwarf and M4 main-sequence star, with the grey lines as their sum. The blue points are the data, and the red arrows are upper limits from WISE. There is a marginal excess above the stellar photosphere in the range 0.47–0.75 μ m, but this may be the result of heating of the main-sequence star by the white dwarf. An excess at 1300 μ m is clearly detected however.

companion (Allard, Homeier & Freytag 2012). Although there are some hints of an excess in the range 0.621–1.23 μ m, this may well be the effect of a slightly non-uniform temperature across the main-sequence star caused by heating from the white dwarf. As such, there is no conclusive excess emission at wavelengths less than 8 μ m. In the ALMA band on the other hand, there is a clear excess.

3 POSSIBLE ORIGINS OF THE ALMA FLUX

3.1 Gyrosynchrotron emission

Two plausible explanations exist for the excess emission detected with ALMA, with the first being gyrosynchrotron emission from material in the magnetic field between the white dwarf and the main-sequence star. Indeed, radio emission from the PCEB V471 Tau has been detected and attributed to gyrosynchrotron emission, making a similar process plausible in NN Ser. However, the emission from V471 Tau has been measured as ~ 3 mJy at 5 GHz (Patterson, Cailault & Skillman 1993) and, if placed at the distance of NN Ser,

this emission would be reduced to ~ 0.03 mJy. Even if it is assumed that the gyrosynchrotron emission follows a flat distribution all the way to the ALMA frequency of 230 GHz, this level of emission would not explain our ALMA detection. If gyrosynchrotron were truly causing this detection, therefore, the emission from NN Ser would have to be several orders of magnitude stronger than for V471 Tau. This is not likely to be the case, although this gyrosynchrotron emission cannot be conclusively ruled out without further observations. The spectral slope of gyrosynchrotron emission is expected to be very distinct, meaning that observation at just one nearby wavelength will easily resolve this issue.

3.2 Thermal emission from dust

Alternatively, the emission may originate from a ring of dust around NN Ser. With just one SED point, the grain size distribution and surface density distribution of this disc cannot be constrained. However, the lack of clear IR excess suggests that the disc does not extend too close to the central binary and the ALMA beam size confines this emission to within 1000 au of NN Ser. Furthermore, the dust emission at this wavelength is optically thin, meaning the dust mass can be estimated by a simple equation of the form $M_{\text{dust}} = C_v \times F_v$ where C_v is a constant for a given frequency F_v (Andrews & Williams 2005).

We adopt the constant derived for 1.3 mm by Cieza et al. (2008), and use the equation

$$M_{\text{dust}} = 0.566 \times \left[\frac{F_v(1300)}{\text{mJy}} \left(\frac{d}{140 \text{ pc}} \right)^2 \right] M_{\oplus} \quad (1)$$

to a derive value of $0.8 \pm 0.2 M_{\oplus}$ for the dust mass around NN Ser, where a distance of 512 ± 43 pc was assumed (Parsons et al. 2010a). This approach contains several assumptions about the grain properties so this value should be treated with some caution. However, it allows comparison with discs around other objects, and we find that the dust mass obtained this way is similar to that of young debris discs around low-mass stars (e.g. Hardy et al. 2015a, and references therein).

4 POSSIBLE ORIGINS FOR THE DUST

We see three possible origins for the dust. First, the dust may be debris disc material which existed before the CE and survived the evolution of the central binary. Secondly, the material we observe may have existed in larger planetesimals before the CE, which then collided as a result of the binary evolution to form dust. Thirdly, the dust may have formed in the CE itself but was not successfully ejected from the system, and instead created a circumbinary disc. The violent evolution of the host binary may preclude some of these origin scenarios and to evaluate how realistic these scenarios are, we performed different simulations for dust production and survival around NN Ser. Three simulations were performed using a modified version of the *N*-body simulator MERCURY (Chambers 1999). The modifications account for the additional forces felt by particles surrounding the binary.

During the CE, particles will be affected by radiation forces from the central AGB star, combined with the stellar wind drag and a changing central mass associated with CE expulsion. We also account for radiation forces in the white dwarf–main sequence state and drag caused by the stellar wind from the M star. The luminosity of the system, and therefore radiation forces, will change as the star evolves, however, and we therefore further modify the code to

accept a time-varying luminosity input. This input was calculated using the Binary Star Evolution (BSE) code (Hurley, Tout & Pols 2002) with a white dwarf progenitor mass of $2.08 M_{\odot}$, companion mass of $0.11 M_{\odot}$, eccentricity of 0, CE efficiency of 0.25, metallicity of 0.02 and a radii at the start of Roche Lobe overflow of $194 R_{\odot}$ (Beuermann et al. 2010). All other BSE parameters were kept at their default. The simulations involved a total mass loss of $1.545 M_{\odot}$ to bring the white dwarf mass down to the measured value of $0.535 M_{\odot}$. This mass ejection was modelled as being ejected preferentially in the binary plane causing a drag for any disc material. For further details, see Appendix A.

4.1 Surviving first-generation debris disc

Debris discs have been found around main-sequence binary stars with similar separations and ages to that of the NN Ser progenitor (Trilling et al. 2007). It could therefore be that the dust we detect is a similar circumbinary debris disc that survived the CE. However, the strong radiation forces during the AGB and CE phases may remove all the small dust which dominates the luminosity in such discs.

To determine if debris disc material can survive the CE, we run simulations of a debris disc located at $r = 10, 50$ and 100 au, with the width of the disc $dr = r/2$. The minimum grain size in this disc was set at the blowout radius before the AGB commenced (i.e. $12 \mu\text{m}$), and the maximum was set at 60 km as this value had previously been found as the largest that contributes to a collisional cascade (Wyatt et al. 2007). The grain size distribution was set to follow the standard relation of particles in a collisional cascade, $n(D) = KD^{2-3q}$, with $q = 11/6$. (Tanaka, Inaba & Nakazawa 1996). We evolve this disc through 4.5 Myr of the AGB, followed by a CE with an associated mass-loss time-scale. Finally, the simulations are continued for 1.3 Myr with a white dwarf–main sequence binary.

The time-scale over which the mass loss takes place in the CE is not a well-determined parameter, as it depends on the unknown efficiency with which the CE material can extract angular momentum from the binary. We therefore leave it as a free parameter in our simulations. However, this time-scale must be shorter than the thermal time-scale of the envelope, otherwise the extracted orbital energy would be radiated away and no envelope expulsion would take place. This places a limit on the CE phase of $\sim 10^3$ yr (Webbink 1984), and we therefore perform simulations with mass-loss time-scales of between 1 and 10^3 yr.

In our simulations, we assume initially that the particles do not interact (and there are therefore no collisions) to identify what proportion of material that may have existed before the AGB is lost, and how this lost material reflects on the change in luminosity of the disc. The fractional initial disc mass which is lost as a result of CE evolution is displayed in Fig. 3 (left panel). It is apparent that for rapid mass loss (less than ~ 40 yr), almost all material is lost in all simulations, whereas a longer mass-loss time-scale can allow almost all mass to survive in the discs with initial radii of 10 and 50 au. The left panel of Fig. 3 is somewhat deceptive however, as the modelled grain size distribution places the majority of the mass in larger particles, whereas the surface area (and therefore flux) is dominated by smaller particles that are more easily lost by radiation pressure and drag. This dominance over surface area that the small grains possess means that discs which retain the bulk of their mass in large bodies may still emit considerably less at 1.3 mm. We therefore plot the fractional change in 1.3 mm flux that one would expect as a result of the CE evolution in Fig. 3 (right panel). From this panel, it becomes apparent that any dust which existed before the CE will likely be lost, reducing the luminosity of the disc




Figure 3. Left: fraction of the initial disc mass that remains after CE evolution as a function of mass loss time-scale. Disc material is lost through a combination of radiation forces, stellar wind drag and the reduction in gravitational force due to changing central mass. The three black lines reflect different initial radii for the debris disc material. Right: fraction of the initial disc luminosity which remains after the CE evolution following the same processes as the figure to the left. The much lower fractions reflect the fact that the CE evolution more efficiently removes small particles, which dominate the surface area of the emitting disc.

considerably. The most favourable configuration for luminosity still suffers an ~ 90 per cent decrease in flux, suggesting that if the dust detected indeed comprised material that survived the CE, then the progenitor disc would need a dust mass that is unrealistically high ($\sim 8 M_{\oplus}$) when compared to discs of a similar age (e.g. Panić et al. 2013; Hardy et al. 2015a). A high dust-mass disc existing around the NN Ser progenitor is made more implausible by the observations that fractional luminosity of debris discs appears to decrease with age (Rieke et al. 2005; Su et al. 2006), and that relatively few discs are observed around M-type stars older than 10 Myr, likely due to their increased stellar wind drag (Plavchan, Jura & Lipsy 2005). It is therefore very unlikely that the dust observed existed before the CE evolution.

4.2 Second-generation debris disc

As debris discs are replenished by collisions, the above simulations, in which the particles do not interact, perhaps underestimate the amount of small grains that exist after CE evolution. In fact, the rapid mass-loss which is associated with the CE expulsion can induce large eccentricities in the orbiting bodies (e.g. Veras et al. 2011), which may cause considerable collisions between planetesimals. We therefore repeat the simulations described, but this time only simulate the largest planetesimals and track their collisions. The number of collisions in these simulations will naturally depend on the number of planetesimals present and therefore the total mass of the disc. We find that no collisions occur, but our simulations are limited to low-mass discs, as modelling $\geq 20\,000$ planetesimals with MERCURY is computationally challenging. To estimate the number of collisions for higher mass discs, we instead calculate the collisional time-scale of the planetesimals which remain after the CE has been expelled. We use the prescription of Wyatt et al. (2010) to calculate the collisional rate of the largest planetesimals in our simulation, R_c :

$$R_c = K M_{\text{tot}} v_k^{8/3} a^{-3} (4\pi I_{\text{max}})^{-1} [0.54 e^{5/3} (1 - e^2)^{-4/3}] \quad (2)$$

with

$$K = 9.5 \times 10^{-6} \rho^{-1} D_{\text{max}}^{-1} Q_D^{-5/6} \quad (3)$$

where ρ is the density of the planetesimals, D_{max} the radii of the largest planetesimal in the collisional cascade in km, M_{tot} is the total mass of the disc in M_{\oplus} , v_k the Keplerian velocity of the planetesimals, a their semi-major axis, I_{max} their maximum inclination, e their eccentricity and Q_D their planetesimal strength in J kg^{-1} . The value of Q_D varies as a function of planetesimal radius and composition (e.g. Krivov, Sremčević & Spahn 2005), and we adopt a conservative value of $1 \times 10^4 \text{ J kg}^{-1}$ for rocky planetesimals of radius 60 km. The collisional rate from this equation was then inverted to calculate the collisional lifetime.

For simulations of the 50 and 100 au discs, the large semi-major axis obtained by the planetesimals causes collision time-scales $\geq 10^4$ Myr even for disc masses as high as $100 M_{\oplus}$. For the disc initially at 10 au, the collision time-scale is somewhat lower, but a large disc mass of $\sim 100 M_{\oplus}$ will still only experience frequent collisions on time-scales $\gg 20$ Myr (see Fig. 4). The dust in these discs will therefore not be replenished within the 1.3 Myr age of NN Ser. This result is similar to that of simulations of planetesimals around single AGBs, which found that the collisional time-scale can increase up to the Hubble time in some cases (Bonsor & Wyatt 2010), resulting in constant debris disc masses in the white dwarf phase.

Although the collisional time-scale in both our simulations and those around a single white dwarf are extremely long, it has been suggested that the presence of a planet on an unstable orbit can rejuvenate the collisions between planetesimals and create a debris disc (Debes & Sigurdsson 2002; Dong et al. 2010). Indeed, a second-generation debris disc which has been attributed to the influence of a planet has potentially been observed around the central star of the helix nebula (Su et al. 2007; Bílková et al. 2012). If the ETVs in NN Ser were due to first-generation planets, a debris disc caused by this mechanism remains a possibility. However, a major problem with this scenario is that progenitor systems for the first-generation planets around NN Ser have been studied in detail and found to be unstable over the main-sequence lifetime (Mustill et al. 2013). Therefore, given the increased collisional lifetime as a result of the CE and the difficulties in maintaining a first-generation planetary system to cause collisions, our results point towards the observed dust being of a different origin.




Figure 4. Collisional time-scale of a planetesimal belt initially located at 10 au, following CE evolution of differing mass-loss time-scales. The collisional time-scale depends on total disc mass, and even $\sim 100 M_{\oplus}$ of material will not cause the collisional time-scale to drop below the age of NN Ser (solid horizontal line). It is therefore unlikely that collisions can create the dust seen around NN Ser.

4.3 Remaining common-envelope material

A further possibility for the origin of the dust detected with ALMA is that it is ‘second-generation’ material, left over from the CE itself. It has been observed that the AGB phase can create significant amounts of dust (Hoogzaad et al. 2002; Lebzelter et al. 2006), and models specifically of CE evolution likewise suggest that dust formation can be very efficient in this environment (Lü, Zhu & Podsiadlowski 2013). The evolutionary history of PCEBs then implies that this dust carries a certain amount of angular momentum, which facilitates the formation of a disc of CE material. Smoothed particle hydrodynamic simulations have suggested that a large amount of the material lost from the primary can, in fact, remain bound to the system (Sandquist, Taam & Burkert 2000; Passy et al. 2012; Ricker & Taam 2012). Furthermore, dusty discs resulting from the AGB phase have been detected around both post-AGB binaries (e.g. van Winckel et al. 2009) and single-neutron stars (Wang, Chakrabarty & Kaplan 2006). However, models suggest the dust produced during the AGB will have radii of $\sim 0.3 \mu\text{m}$ (Yasuda & Kozasa 2012), and grains of this size will also be affected by radiation forces from the newly formed hot white dwarf. These grains therefore may not survive long enough to be observed around the 1.3 Myr old PCEB.

To investigate the size of grains which may survive the PCEB phase, we again run simulations using our modified version of MERCURY for a population of grains with sizes ranging from $0.01 \mu\text{m}$ to 1 cm. This simulation is commenced with the binary in its PCEB configuration, so dust is only prone to radiation forces from the white dwarf–main sequence binary and a small drag force due to a stellar wind from the main-sequence star. In this case, the grains size distribution is unknown, so we instead simulate a uniform number distribution of grains distributed logarithmically between 0.5 and 500 au. We then record the average orbital radii for each particle that survives the simulation, and plot the minimum value of the particle radii in Fig. 5.

Grains smaller than $\sim 20 \mu\text{m}$ are removed in our simulations, and this is considerably larger than the grain size believed to be generated within the AGB of $\sim 0.3 \mu\text{m}$ (Yasuda & Kozasa 2012). One might therefore expect grains made within the CE to be instantly




Figure 5. Radius of the smallest grain that can survive the 1.3 Myr that has passed since the CE of NN Ser whilst experiencing radiation forces and stellar wind drag. This is relevant to dust which was created in the CE and remained in the system as part of a circumbinary disc. These simulations do not include grain growth/reprocessing, which could allow smaller particles to be present in the disc.

lost. However, if the amount of gas and dust is large enough, the disc may be optically thick and have an inner region where radiation pressure is negligible (Takeuchi & Lin 2003; Olofsson et al. 2009). In these regions, it is conceivable that grains could grow to $100 \mu\text{m}$ in as little as $\sim 10^4$ yr (Dullemond & Dominik 2005), in the same manner as in protoplanetary discs. Models of the CE phase differ in their estimates of how much material can remain bound to the system, with values ranging from 10 per cent (Kashi & Soker 2011) to as much as 97 per cent of the CE (Passy et al. 2012). In either case, this would suggest a large amount of gas ($\gtrsim 0.15 M_{\odot}$ in the case of NN Ser) remains bound in a disc, so the existence of optically thick regions is certainly plausible. This gas might still exist alongside the detected dust, allowing one observational avenue by which this scenario can be tested.

Dust attributed to the AGB has already been detected around several post-AGB binary stars in the near and mid-IR (de Ruyter et al. 2006; van Winckel et al. 2009; Hillen et al. 2016). The post-AGB nature of these stars naturally means they are much younger than NN Ser, but studies of these objects suggest that already the grains have been through considerable reprocessing with grain sizes larger than $\sim 2 \mu\text{m}$ (Gielen et al. 2008), adding credibility to the possibility that this dust is left over AGB material. It has also been pointed out that there is some similarity between these discs and protoplanetary discs found around young stellar objects (de Ruyter et al. 2006), opening the possibility that discs around these evolved binaries may provide a new avenue with which to study protoplanetary disc evolution and dissipation.

5 PLANETS

Given the convincing planetary fit for NN Ser, an obvious question is what the detection of this dust implies for the putative planets around PCEBs. Our simulations suggest that the detected disc is most likely left over from the CE. The theory that the planets around NN Ser formed in a second-generation scenario (Völschow et al. 2014) may gain some support from this detection, as the material necessary to form these planets seems to exist. The dust detected might then be material which simply has not efficiently grown yet, or it could

be material that has grown, but since been involved in collisions. The growth of planets ≥ 1000 km in radii would stir up collisions between smaller bodies and might rejuvenate the population of small dust, as described by models of ‘self-stirring’ (Kenyon & Bromley 2004). The time-scales required to achieve this planet formation and subsequent self-stirring are also consistent with the 1.3 Myr age of NN Ser. However, although second-generation planets gain some support from our results, the detection of dust does not directly correlate with the existence of planets around NN Ser, as the dust may simply be CE material that has not yet grown into larger bodies or been ejected. Furthermore, self-stirring models can form small planets capable of stirring the disc within 1.3 Myr, but the planets predicted from the ETVs of NN Ser have masses of $6.97 M_{\text{JUP}}$ and $1.73 M_{\text{JUP}}$ (Beuermann et al. 2013). It remains uncertain if the formation of these much more massive planets can occur within the 1.3 Myr age of NN Ser. As discussed in detail in Zorotovic & Schreiber (2013), the disc instability model can form planets within this time-scale, but only at orbital separations much further than those predicted by the ETVs. Significant orbital migration would therefore need to have occurred if this were the case. Classical core accretion, on the other hand, would be able to form planets at their predicted location, but would struggle to form planets within 1.3 Myr. For example, models which estimate the formation time-scale of Jupiter within the Solar system suggest that gas giants gain the majority of their mass at times $\gtrsim 2.5$ Myr (Lissauer et al. 2009).

Another potential implication that this detection has on the field of planets around PCEBs is that these circumbinary discs can perhaps interact with the binary itself (Artymowicz et al. 1991; Kashi & Soker 2011), causing the binary to lose angular momentum and slowly change the period of its orbit. As the planets around PCEBs are inferred from fits to the ETVs, this extra angular momentum loss would need to be accounted for. Before the planetary fits of NN Ser gained credibility, the possibility of a circumbinary disc being the cause of the perceived change in angular momentum was investigated by Chen (2009). Their model relied on a disc which obtained its mass through stellar wind from the M star, and they concluded that the disc was not massive enough to cause strong angular momentum loss unless a large fraction of the wind can end up in the disc (~ 10 per cent) and the wind loss rate is ultra-high ($\sim 10^{-10} M_{\odot} \text{ yr}^{-1}$). If the disc observed here is composed of CE material then the total mass (including gas) might be very large, making it massive enough to extract angular momentum without the need of a high wind-loss rate.

Models of angular momentum loss due to circumbinary discs around cataclysmic variables are well developed (Spruit & Taam 2001; Taam & Spruit 2001), but they rely on some knowledge of the surface density and inner radius of the disc. If the disc around NN Ser follows the surface density profile of a viscous accretion disc, then the surface density might only reach significant levels at small orbital radii. At such radii, one would expect a strong IR excess in the SED which is not seen, perhaps arguing against this effect being significant. Nonetheless, the effect of this circumbinary disc on the central binary offers interesting opportunities for future work.

6 CONCLUSIONS

We present the detection of 1.3 mm flux from the PCEB NN Ser, and find most plausible explanation for this flux is thermal emission from a circumbinary disc. We run simulations of the history of NN Ser to investigate what material can survive the extra forces associated with the AGB and CE. Given the difficulties in creating/maintaining a debris disc of primordial material in our simulations, we find that

a disc of left over CE material is the most likely explanation for the detection. Such discs are predicted by theory, but have not previously been observed around a PCEB. This detection therefore adds credibility to the theory that second-generation planets might exist around NN Ser.

ACKNOWLEDGEMENTS

We would like to thank the anonymous referee for their time and helpful comments. In addition, AH, MRS, CC and LC acknowledge support from the Millennium Nucleus RC130007 (Chilean Ministry of Economy). MRS, CC, SP and LC also acknowledge support from FONDECYT (grants 1141269, 3140592, 3140585 and 1140109). DV and BTG have received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP/2007-2013)/ERC Grant Agreement n. 320964 (WD-Tracer). TRM was supported by STFC grant #ST/L000733. This work is based on observations made with ESO Telescopes at the La Silla Paranal Observatory under programme ID 085.D-0541. This work is also based, in part, on observations made with the *Spitzer Space Telescope*, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. This paper makes use of the following ALMA data: ADS/JAO.ALMA#2013.1.01342.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan) and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ.

REFERENCES

- Allard F., Homeier D., Freytag B., 2012, *R. Soc. Lond. Philos. Trans. Ser. A*, 370, 2765
 Andrews S. M., Williams J. P., 2005, *ApJ*, 631, 1134
 Applegate J. H., 1992, *ApJ*, 385, 621
 Artymowicz P., Clarke C. J., Lubow S. H., Pringle J. E., 1991, *ApJ*, 370, L35
 Bear E., Soker N., 2014, *MNRAS*, 444, 1698
 Beuermann K. et al., 2010, *A&A*, 521, L60
 Beuermann K., Dreizler S., Hessman F. V., 2013, *A&A*, 555, A133
 Bilíková J., Chu Y.-H., Gruendl R. A., Su K. Y. L., De Marco O., 2012, *ApJS*, 200, 3
 Bonsor A., Wyatt M., 2010, *MNRAS*, 409, 1631
 Bours M. C. P. et al., 2014, *MNRAS*, 445, 1924
 Brinkworth C. S., Marsh T. R., Dhillon V. S., Knigge C., 2006, *MNRAS*, 365, 287
 Burns J. A., Lamy P. L., Soter S., 1979, *Icarus*, 40, 1
 Chambers J. E., 1999, *MNRAS*, 304, 793
 Chen W.-C., 2009, *A&A*, 499, L1
 Cieza L. A., Swift J. J., Mathews G. S., Williams J. P., 2008, *ApJ*, 686, L115
 de Ruyter S., van Winckel H., Maas T., Lloyd Evans T., Waters L. B. F. M., Dejonghe H., 2006, *A&A*, 448, 641
 Debes J. H., Sigurdsson S., 2002, *ApJ*, 572, 556
 Dhillon V. S. et al., 2007, *MNRAS*, 378, 825
 Dong R., Wang Y., Lin D. N. C., Liu X.-W., 2010, *ApJ*, 715, 1036
 Dullemond C. P., Dominik C., 2005, *A&A*, 434, 971
 Fazio G. G. et al., 2004, *ApJS*, 154, 10
 Garaud P., Barrière-Fouchet L., Lin D. N. C., 2004, *ApJ*, 603, 292
 Gielen C., van Winckel H., Min M., Waters L. B. F. M., Lloyd Evans T., 2008, *A&A*, 490, 725
 Guinan E. F., Ribas I., 2001, *ApJ*, 546, L43
 Hardy A. et al., 2015a, *A&A*, 583, A66
 Hardy A. et al., 2015b, *ApJ*, 800, L24

Hillen M., Kluska J., Le Bouquin J.-B., Van Winckel H., Berger J.-P., Kamath D., Bujarrabal V., 2016, *A&A*, 588, L1

Hoogzaad S. N., Molster F. J., Dominik C., Waters L. B. F. M., Barlow M. J., de Koter A., 2002, *A&A*, 389, 547

Horner J., Hinse T. C., Wittenmyer R. A., Marshall J. P., Tinney C. G., 2012, *MNRAS*, 427, 2812

Horner J., Wittenmyer R. A., Hinse T. C., Marshall J. P., Mustill A. J., Tinney C. G., 2013, *MNRAS*, 435, 2033

Hurley J. R., Tout C. A., Pols O. R., 2002, *MNRAS*, 329, 897

Kashi A., Soker N., 2011, *MNRAS*, 417, 1466

Kenyon S. J., Bromley B. C., 2004, *AJ*, 127, 513

Koester D., 2010, *Mem. Soc. Astron. Italiana*, 81, 921

Krivov A. V., Sremčević M., Spahn F., 2005, *Icarus*, 174, 105

Lebzelter T., Posch T., Hinkle K., Wood P. R., Bouwman J., 2006, *ApJ*, 653, L145

Lissauer J. J., Hubickyj O., D'Angelo G., Bodenheimer P., 2009, *Icarus*, 199, 338

Lohsen E., 1974, *A&A*, 36, 459

Lü G., Zhu C., Podsiadlowski P., 2013, *ApJ*, 768, 193

Makovoz D., Roby T., Khan I., Booth H., 2006, in *SPIE Conf. Ser. Society of Photo-Optical Instrumentation Engineers*, p. 62740C

Marsh T. R. et al., 2014, *MNRAS*, 437, 475

Moorwood A., Cuby J.-G., Lidman C., 1998, *The Messenger*, 91, 9

Mustill A. J., Marshall J. P., Villaver E., Veras D., Davis P. J., Horner J., Wittenmyer R. A., 2013, *MNRAS*, 436, 2515

Nelson B., Young A., 1970, *PASP*, 82, 699

Olofsson J. et al., 2009, *A&A*, 507, 327

Paczynski B., 1976, in Eggleton P., Mitton S., Whelan J., eds, *IAU Symp. Vol. 73, Structure and Evolution of Close Binary Systems*. D. Reidel Publishing Co., Dordrecht, Holland, p. 75

Panić O. et al., 2013, *MNRAS*, 435, 1037

Parsons S. G., Marsh T. R., Copperwheat C. M., Dhillon V. S., Littlefair S. P., Gänsicke B. T., Hickman R., 2010a, *MNRAS*, 402, 2591

Parsons S. G. et al., 2010b, *MNRAS*, 407, 2362

Parsons S. G. et al., 2014, *MNRAS*, 438, L91

Passy J.-C. et al., 2012, *ApJ*, 744, 52

Patterson J., Caillault J.-P., Skillman D. R., 1993, *PASP*, 105, 848

Plavchan P., Jura M., Lipsky S. J., 2005, *ApJ*, 631, 1161

Reach W. T. et al., 2005, *PASP*, 117, 978

Ricker P. M., Taam R. E., 2012, *ApJ*, 746, 74

Rieke G. H. et al., 2005, *ApJ*, 620, 1010

Sandquist E. L., Taam R. E., Burkert A., 2000, *ApJ*, 533, 984

Schleicher D. R. G., Dreizler S., 2014, *A&A*, 563, A61

Schreiber M. R., Gänsicke B. T., 2003, *A&A*, 406, 305

Skrutskie M. F. et al., 2006, *AJ*, 131, 1163

Spruit H. C., Taam R. E., 2001, *ApJ*, 548, 900

Su K. Y. L. et al., 2006, *ApJ*, 653, 675

Su K. Y. L. et al., 2007, *ApJ*, 657, L41

Taam R. E., Spruit H. C., 2001, *ApJ*, 561, 329

Takeuchi T., Lin D. N. C., 2003, *ApJ*, 593, 524

Tanaka H., Inaba S., Nakazawa K., 1996, *Icarus*, 123, 450

Trilling D. E. et al., 2007, *ApJ*, 658, 1264

van Winckel H. et al., 2009, *A&A*, 505, 1221

Veras D., Wyatt M. C., Mustill A. J., Bonsor A., Eldridge J. J., 2011, *MNRAS*, 417, 2104

Veras D., Mustill A. J., Bonsor A., Wyatt M. C., 2013, *MNRAS*, 431, 1686

Veras D., Eggl S., Gänsicke B. T., 2015, *MNRAS*, 451, 2814

Völschow M., Banerjee R., Hesselman F. V., 2014, *A&A*, 562, A19

Wang Z., Chakrabarty D., Kaplan D. L., 2006, *Nature*, 440, 772

Webbink R. F., 1984, *ApJ*, 277, 355

Wyatt M. C., Smith R., Su K. Y. L., Rieke G. H., Greaves J. S., Beichman C. A., Bryden G., 2007, *ApJ*, 663, 365

Wyatt M. C., Booth M., Payne M. J., Churcher L. J., 2010, *MNRAS*, 402, 657

Yasuda Y., Kozasa T., 2012, *ApJ*, 745, 159

Zorotovic M., Schreiber M. R., 2013, *A&A*, 549, A95

Zorotovic M., Schreiber M. R., Gänsicke B. T., Nebot Gómez-Morán A., 2010, *A&A*, 520, A86

APPENDIX A: SIMULATION DETAILS

Simulations were performed using a modified version of the N -body code *MERCURY* (Chambers 1999). These modifications allowed the inclusion of Radiation forces, stellar mass loss and stellar wind drag, the details of which are described below.

A1 Radiation forces

At each time-step in the *MERCURY* simulator, each particle in the simulation felt an additional acceleration of the form (Burns, Lamy & Soter 1979):

$$\mathbf{Accl} = \frac{GM_*\beta}{r^2} \left(\hat{\mathbf{r}} - \frac{v_r \hat{\mathbf{r}} + \mathbf{v}}{c} \right), \quad (\text{A1})$$

where G is the gravitational constant, M_* the mass of the central object, β the ratio of the force of radiation pressure to gravity, r the orbital separation of the particle, $\hat{\mathbf{r}}$ the unit vector corresponding to r , c the speed of light and $v_r = \mathbf{v} \cdot \hat{\mathbf{r}}$ accounts for the Doppler shift in the radiation seen by the particle. The value of β was calculated using the equation

$$\beta = \frac{3L_* Q_{\text{pr}}}{16\pi c G M_* \rho R}, \quad (\text{A2})$$

where L_* is the luminosity of the star, ρ the density of particles, R their radius and Q_{pr} the radiation pressure efficiency which was set at 1.

A2 Stellar mass loss

The exact behaviour of how mass is lost during CE evolution is not known. We therefore assume that mass loss occurs only in the time described by the ‘mass-loss time-scale’ parameter (i.e. no mass loss occurred in the preceding RGB and AGB phases). We further assume that during this period the mass is lost linearly. We use the Bulirsch–Stoer integrator in *MERCURY* with a variable time-step, and to improve the accuracy of our simulations we apply this mass-loss as each sub-step of the Bulirsch–Stoer algorithm (e.g. Veras et al. 2013).

A3 Stellar wind drag

The accelerations on the particles due to stellar wind drag were calculated using the equations from Garaud et al. (2004), expressed in the notation of Veras, Eggl & Gänsicke (2015) as:

$$\mathbf{a}_{\text{swd}} = \begin{cases} \left(\frac{\rho_g v_s}{\rho R} \right) (\mathbf{v}_g - \mathbf{v}) & R \ll \zeta \\ \left(\frac{\rho_g B}{\rho R} \right) (\mathbf{v}_g - \mathbf{v}) |\mathbf{v}_g - \mathbf{v}| & R \gg \zeta, \end{cases} \quad (\text{A3})$$

where ρ_g is the density of the gas, ζ its mean-free path length, \mathbf{v}_g its velocity, v_s the local sound speed and B is given by the equation

$$B = \begin{cases} 9 \left[\frac{6R}{\zeta v_R} |\mathbf{v}_g - \mathbf{v}| \right]^{-1} & \text{Re} \leq 1 \\ 9 \left[\frac{6R}{\zeta v_R} |\mathbf{v}_g - \mathbf{v}| \right]^{-0.6} & 1 \leq \text{Re} \leq 800 \\ 0.165 & \text{Re} \geq 800 \end{cases} \quad (\text{A4})$$

for differing values of the Reynolds number, calculated from the equation

$$\text{Re} = \frac{6R}{\zeta v_s} |\mathbf{v}_g - \mathbf{v}|. \quad (\text{A5})$$

For the mean-free path, we use the approximate relation from Veras et al. (2015)

$$\rho_g \zeta \sim 10^{-8} \text{ kg m}^{-2}. \quad (\text{A6})$$

The stellar wind in our simulations was assumed to travel radially and with the escape velocity of the system. The mass loss during

the CE will occur preferentially in the disc plane, and we therefore confine all stellar wind to a disc of scale height $H = 0.15r$. However, this preferential direction also suggests the particles will not be expelled perfectly radially. We therefore assume only 15 per cent of the expelled mass travels radially outwards and interacts with the particles, which should ensure our drag force is a conservative estimate.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.